USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required software to e-Annotate PDFs: **Adobe Acrobat Professional** or **Adobe Reader** (version 11 or above). (Note that this document uses screenshots from **Adobe Reader DC**.)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the **Comment** tab (right-hand panel or under the Tools menu).

This will open up a ribbon panel at the top of the document. Using a tool will place a comment in the right-hand panel. The tools you will use for annotating your proof are shown below:

1. Replace (Ins) Tool – for replacing text.

Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it:

- Highlight a word or sentence.
- Click on .
- Type the replacement text into the blue box that appears.

2. Strikethrough (Del) Tool – for deleting text.

Strikes a red line through text that is to be deleted.

How to use it:

- Highlight a word or sentence.
- Click on .
- The text will be struck out in red.

experimental data if available. For ORFs to be had to meet all of the following criteria:

1. Small size (35–250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminus terminus with another coding feature; over both ends; or ORF containing a tRNA.

3. Commenting Tool – for highlighting a section to be changed to bold or italic or for general comments.

Use these 2 tools to highlight the text where a comment is then made.

How to use it:

- Click on .
- Click and drag over the text you need to highlight for the comment you will add.
- Click on .
- Click close to the text you just highlighted.
- Type any instructions regarding the text to be altered into the box that appears.

4. Insert Tool – for inserting missing text at specific points in the text.

Marks an insertion point in the text and opens up a text box where comments can be entered.

How to use it:

- Click on .
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the box that appears.

5. Attach File Tool – for inserting large amounts of text or replacement figures.

Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it:

- Click on .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

The attachment appears in the right-hand panel.

chondrial preparator
ative damage injury
re extent of membra
, malondialdehyde (
(TBARS) formation. I
used by high perform

6. Add stamp Tool – for approving a proof if no corrections are required.

Inserts a selected stamp onto an appropriate place in the proof.

How to use it:

- Click on .
- Select the stamp you want to use. (The **Approved** stamp is usually available directly in the menu that appears. Others are shown under *Dynamic*, *Sign Here*, *Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
on perfect competition, constant ret
production. In this environment goods
extra...
he...
etermined by the model. The New-Key
otaki (1987), has introduced produc
general equilibrium models with nomin
ed and...
Most of this litera

APPROVED

Drawing tools available on comment ribbon

How to use it:

- Click on one of the shapes in the **Drawing Markups** section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select *Open Pop-up Note*.
- Type any text in the red box that appears.

7. Drawing Markups Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.

For further information on how to annotate proofs, click on the **Help** menu to reveal a list of further options:

Three-dimensional geophysical characterization of the La Rambla and Zafra de Zánacara anticlines (Loranca Basin, Central Spain)

C. Ayala^{1*,†}, F.M. Rubio¹, C. Rey-Moral¹, M.I. Reguera² and C. Biete³

¹IGME (Spanish Geological Survey), La Calera n. 1, 28760 Tres Cantos, Madrid, Spain, ²IGME (Spanish Geological Survey), Ríos Rosas n. 23, 28003 Madrid, Spain, and ³Institute of Earth Sciences Jaume Almera, ICTJA, CSIC, Lluís Solé i Sabarís s/n, 08028 Barcelona, Spain

Received January 2018, revision accepted January 2019

ABSTRACT

The Zafra de Zánacara anticline (also known as the El Hito anticline), located in the Loranca Cenozoic Basin (part of the Tagus Basin, Central Spain), had been studied by several oil companies during the late 1960s and early 1970s. In 2009, within the 'Plan for selection and characterization of suitable structures of CO₂ geological storage', this anticline was selected as a potential CO₂ storage site. A preliminary three-dimensional geological model, based on five geological cross sections that were constrained with the interpretation of the available seismic profiles (that are rather old and do not have very good quality), was created. With the aim of improving the geological knowledge of the Zafra de Zánacara anticline and helping to investigate the suitability of a nearby anticline, namely La Rambla, as another structural closure that might make it a possible CO₂ storage site, a local gravity survey (~~one station every square kilometres~~) was carried out in the area, seven new geological cross sections, based on these existing seismic profiles and field geology, were built, and a new three-dimensional geological model that included both anticlines, improved through three-dimensional stochastic gravity inversion, was constructed. The densities needed for the geological formations of the model come from the analysis of rock samples, logging data from El Hito-1 drillhole and petrophysical information from Instituto Geológico y Minero de España database. The inversion has improved the knowledge about the geometry of the anticlines' traps and seals as well as the geometry of the basement relief and the structural relationship between basement and cover.

Key words: ~~CO₂ geological storage, Gravity survey, Residual Bouguer anomaly, 3D gravity inversion, Loranca Basin, Zafra de Zánacara, La Rambla.~~

INTRODUCTION

The mitigation of the greenhouse gas emissions, in particular CO₂, is being one of the main concerns in order to fight against the climate change. Since 2009, the Spanish Geological Survey has been developing a long term project called 'Plan for selection and characterization of suitable structures

of CO₂ geological storage' (IGME, 2009). The first stage of this plan has been evaluating and pre-selecting a number of possible favourable areas at a regional level whose local structures might have structural closures that make them suitable as CO₂ storage sites because its adequate geological features, good storage capacity based on preliminary calculations and the reasonable cost of developing the corresponding infrastructures.

The methodology followed to select those favourable areas can be summarized as follows:

*E-mail: c.ayala@igme.es

†Now visiting at the Institute of Earth Sciences Jaume Almera, ICTJA, CSIC, Lluís Solé i Sabarís s/n, 08028 Barcelona, Spain

1. compilation and analysis of the available existing geological and geophysical information (from IGME databases and available information from the literature);
2. constructing regional geological cross sections based on the compiled data, which include reinterpreted old seismic lines (that had to be converted into digital format) and deep drillholes;
3. building up isolines and isopachs of the main reference surfaces (top of the basement, Buntsandstein–Muschelkalk, Keuper, Lower Jurassic, Upper Jurassic, Cretaceous and Cenozoic), based on geological cross sections and geophysical information. In different areas, these reference surfaces have different degrees of resolution, depending on the geological and geophysical data available. In some cases, like in the Zafra de Zancara anticline, only some of the isolines (top of the Basement, Buntsandstein–Muschelkalk, Keuper) were considered and therefore, drawn;
4. identification of the possible formation pairs storage/seal according to lithology, thickness, depth and geometry;
5. petrophysical characterization of the storage and seal formations;
6. preliminary estimation of the CO₂ storage capacity of the selected structures.

The second stage of this plan consists of completing more detailed studies of the selected structures in order to obtain a more precise image of the structural closures that might be used as reservoir. This image could be used to calculate more accurately its storage capacity. The present work focuses on refining one of these selected structures, Zafra de Zancara anticline, and characterizes a new possible structure, namely the La Rambla anticline that was not taken into account in the first stage but we believe it is worth considering as possible storage site by analogy with its neighbouring anticline.

In these structures, two possible pairs of storage/seal formations were identified: Buntsandstein sandstones and Muschelkalk limestones as a structural closure with Keuper evaporites and shales as a seal; fractured limestones of Lower Jurassic as a structural closure with Jurassic marls as a seal. This two anticlines have been targeted because they show a rather simple tectonic structure, they are located in a suitable socioeconomic and a stable geodynamic contexts and the possible pairs storage/seal formations seem to have an adequate geometry and thickness.

In this study area, the seismic lines, some of which were already used in the first stage, are more than 40 years old with a lot of noise and poor reflectivity, which makes difficult their accurate interpretation. In addition, there is only one logged borehole in the area that could help to interpret the

seismic data, El Hito-1. Therefore, with the objective of improving the knowledge of the Zafra de Zancara anticline and characterize the La Rambla anticline in order to assess their suitability as storage site we have chosen to apply the gravity method, a cost-efficient way to study the density distribution and the geometry of the target structures. Thus, three gravimetric surveys were carried out between the end of 2009 and the beginning of 2012 acquiring a total of 577 stations. The surveys included the collection of some rock samples to obtain density values in order to constrain the density of the model. The available geological and geophysical data were used to build a preliminary three-dimensional (3D) model. Furthermore, the gravity data were processed and analysed to obtain the residual Bouguer anomaly in order to improve the preliminary 3D model through gravity inversion to obtain a final model, consistent with all the available geological and geophysical data.

In this way, it has been possible to refine the 3D architecture of the different lithological horizons and the depth-density distribution of the structures providing more insights into its structural pattern and its significance in terms of possible structural closures that might be used as CO₂ reservoirs.

GEOLOGICAL SETTING

The Tagus Basin is an intraplate basin limited to the N-NW by the Spanish Central System, to the NE by the Iberian Chain and to the SW by the Toledo Mountains. This basin is, in turn, divided into two sub-basins separated by the Altomira Range, Madrid Basin and Loranca Basin (also known as Intermediate Depression) (Fig. 1). The Madrid Basin is an intracratonic basin located in the central and western part of the Tagus Basin and filled by Cenozoic sediments overlying the Mesozoic and Paleozoic basement. The Loranca Basin, located in the easternmost part, is a piggyback basin containing approximately N-S striking, mostly west-vergent thrust-related folds formed during Late Oligocene–Early Miocene time (Díaz Molina *et al.* 1995). Although most Loranca structures do not crop out, they are identifiable in seismic profiles (Querol 1989; ITGE 1990) and affect a Mesozoic–Cenozoic cover detached over a regional décollement level formed by Middle–Upper Triassic evaporites and clays (e.g. Gómez, Díaz Molina and Lendínez 1996; Muñoz-Martín & De Vicente 1998).

In this tectonic environment, the study area (Fig. 1) is located in the southern part of the Loranca Basin, between the Altomira Range and the SW part of the Iberian Range. Its extension is about 1200 km² covering four

Figure 1 Location map of the study area, modified from Piña-Varas *et al.* (2013). CP, Catalan pre-litoral Coastal Range; RA, Aragonese branch of the Iberian Range; RC, Castilian branch of the Iberian Range; SL, Levante sector of the Iberian Range.

MAGNA geological maps 1:50,000 scale (http://mapas.igme.es/Servicios/default.aspx#IGME_MAGNA_50). In the area (Fig. 2), there are a series of anticlines with a general direction N-NW to S-SE. These anticlines have lengths between 5 and 20 km and width of approximately 3–4 km being the most relevant Zafra de Zánacara anticline, also known as the El Hito anticline (which is the longest one) and La Rambla anticline. According to Querol (1989), these anticlines mainly affect the Mesozoic rocks, from the Upper Triassic to the Upper Cretaceous and also part of the Cenozoic infill, mainly Paleogene. The level of detachment is located in the evaporitic materials of the Keuper facies (evaporites and shales) (Álvaro, Capote, Vegas 1979; Guimerá and Álvaro, 1990). Below this sedimentary cover, the interpretation of the available seismic data (from the seventies) suggest that the Paleozoic and Lower Triassic basement is deformed by a series of horsts and grabens that are related to basement anticlines (the N-S structures) and extensive faults (e.g. Valcárcel-Rodríguez, 2015).

Previous geological interpretation

In the model published in the ALGECO2 Atlas (IGME 2009), within the area of La Rambla–Zafra de Zánacara Complex, the map of basement isolines (Fig. 3a) reveals a tectonics of blocks limited by faults that, according to the seismic data, do not affect the post-Triassic cover. According to previous studies (IGME 2009), these are normal faults generated during the first stage of Triassic rifting that later were inverted. Two basement structural highs associated with the Zafra de Zánacara anticline can be observed. These structures could have closure perimeters around –1600 m and –2200 m, respectively (datum: sea level) (Fig. 3b). In the area of the La

Rambla anticline the basement shows a much lower relief, and it has not been possible to delimit structural closures in it, neither to the south, nor to the west due to lack of information (the available seismic data was scarce and not always of good quality). Thickness of the Buntsandstein–Muschelkalk (see the geometry of the top of this layer in Fig. 3b) varies between 200 and 400 m; that of the Keuper changes from tens of meters to more than 1000 m beneath some areas (Fig. 3c). Thicknesses of the remaining layers come from the geological cross sections (Fig. 4): Jurassic has a thickness of c. 1000 m, and Cretaceous is between 250 and 500 m thick. As for the Cenozoic, its thickness varies from a few tens of meters to more than 500 m in some places.

GRAVITY SURVEY

New 577 gravity stations were measured with an homogeneous distribution of approximately 1 point/km² (Fig. 2) using a Scintrex CG5 gravity meter with a standard resolution of 1 microGal and a nominal repeatability of less than 5 microGal. The Scintrex CG5 applies the luni-solar correction, and the measurements were performed in cycles of 4–6 h, and then corrected for instrumental drift. The near topographic correction (Hammer sectors B to D up to 170 m) was performed on the field. The survey was linked with the IGSN-71 network (International Gravity Standardization Network) in order to obtain the values of the observed gravity (gobs). Post-processing differential GPS has been used to determine the spatial coordinates in UTM30N and the elevation in orthometric heights. To ensure the quality of the data, and following the directions from the UNE Norm 22-611-85 (Instituto Español de Normalización, 1985), 7% of the stations have

been repeated. The root mean square (RMS) of the repetitions is calculated as

$$\text{RMS} = \pm \sqrt{\frac{\Sigma d^2}{2N}}$$

where d is the difference between the first measure and the repeated measure and N the number of repeated points. The RMS of the repetitions is displayed on Table 1.

Bouguer anomaly: Regional–residual separation

The observed gravity was corrected for the latitude, free air and Bouguer plate, and then the topographic correction was applied to obtain the complete Bouguer anomaly. The gravity data has been processed with the GRS67 (Geodetic Reference

System) formulas so the new gravity data will be compatible with the one stored in the IGME database.

We used the Bouguer correction density, i.e. reduction density, of 2.3 g/cm³ determined by the Nettleton method (Nettleton, 1971), which is based on calculation of the Bouguer anomaly with different reduction densities and selection of the density value providing the lowest correlation between the Bouguer anomaly and topography. The topographic correction was calculated using the Hammer method (Hammer, 1939) up to 21,943 m and a Digital Terrain Model grid with 100 m spacing.

The complete Bouguer anomaly (Bouguer anomaly from now on, Fig. 5a) was gridded at 500 m. The homogeneous distribution of the point data allows preserving the short wavelength component without introducing spurious anomalies.

Figure 3 Isolines of the top of formations published in the ALGECO2 Atlas (IGME, 2009). Black lines are interpreted basement faults. In each figure, a rectangle shows the study area of the present work, and discontinuous lines delimit the outline of the geological outcrops of the La Rambla (LR) and Zafra de Zancara (ZdZ) anticlines. The El Hito-1 borehole is also shown in Fig. 3(a, c) (the displayed S. Lorenzo Parrilla-1 borehole is part of the Atlas map but is outside our study area and, therefore, not mentioned on the text). All depths are in m b. s. l. (a) Top of the basement. (b) Top of Buntsandstein-Muschelkalk. (c) Top of Keuper.

Figure 4 Cross sections from ALGECO2 Atlas (IGME, 2009). Part of these cross-sections were integrated in the initial geological model. The portion of each cross section used in the modelling is outlined with a red rectangle. Note that Corte 4 (interpreted from the eastern part of seismic line DD') in the present work was projected to build up Corte 6 (see Fig. 2 for location).

The amplitude of the Bouguer anomaly ranges from c. -123 to c. -104 mGal. The pattern of the anomaly shows four lineations of relative maxima trending NW-SE with amplitudes ranging between -109 to -105 , -114 to -109 , -117 to

-115 and -115 to -112 , respectively (labelled Max1 to Max4, black lines in Fig. 5a). These relative maxima mainly coincide with outcrops of Cretaceous and Jurassic material, two of them located over the Zafra de Zancara anticline, and one on

Table 1 Root mean square (RMS) of the measures (X and Y are the horizontal coordinates, Z is the elevation)

	X (m)	Y (m)	Z (m)	Gravity (mGal)
RMS	0.36	0.38	0.09	0.08

the La Rambla anticline. A couple of prominent and round-shaped minima are also observed, one towards the NW, with amplitude of -121 to -119 mGal and another one at the NE end of the study area, with amplitude of -121 to -123 mGal (labelled Min1 and Min2 in Fig. 5a).

Because the target structures are in the uppermost part of the crust, modelling requires removing from the Bouguer anomaly a regional field of long wavelength component generated in the deeper crust in order to obtain a residual field that reflects the shallow structures. After several tests to calculate an adequate regional-residual separation, the residual Bouguer anomaly used in this work was obtained subtracting from the Bouguer anomaly a regional field (Fig. 5b) calculated using the Zeng method (Zeng, 1989). Zeng's method uses a polynomial surface adjust, but the degree of the polynomial is based on upward continuation. When we upward continue the Bouguer anomalies, we suppress the contribution of the shallow source anomalies while the anomalies of deeper bodies remain unchanged. When the height of the upward continuation is large enough, the short wavelength anomalies from shallow bodies are practically zero. It is assumed (Zeng, 1989) that the regional Bouguer anomaly is the polynomial surface which fits that particular upward continued anomaly. The advantage of this method is that allow getting the regional Bouguer anomaly in an objective and repeatable way. After upward continuing the Bouguer anomaly every 500 m up to 5000 m, we have chosen an upward continuation at 4500 m and the degree of the polynomial that adjust this upward continuation is 3. This residual (Fig. 5c) is the gravity data used as observed gravity anomaly in the inversion.

The residual Bouguer (Fig. 5c) has a range between -2.5 and 4.5 mGal and depicts more clearly the four main structural axis of NW-SE direction that are also observed in the Bouguer anomaly. These relative maxima have amplitudes ranging from 0 to 4.5 mGal, with the maximum values occurring over the axis of the anticlines. The two maxima over the Zafra de Zánacara anticline are joined at their northern end. Between the Zafra de Zánacara and La Rambla anticlines, a relative lineament of gravity minima crosses the area from NW to SE (labelled MinL2 in Fig. 5c). Another relative lineament of gravity minima (labelled MinL1 on Fig. 5c) is located

between the relative maxima on La Rambla and the relative maxima on top of the other anticline at the SW. The amplitude of the relative lows range from -2.5 to 0.5 mGal and can be related to the thick Cenozoic sediments deposited between the anticlines (Fig. 5c).

Petrophysical data

Petrophysical properties are a fundamental constraint to interpret the potential field data and minimize the number of possible solutions during the modelling process. Density was determined analysing the data coming from three different sources:

1. Twenty-one samples collected from the main geological formations that are considered of interest to be modelled (Fig. 5a). The samples have been analysed at the IGME laboratory to obtain its density. The measurements have been made on hammer-cut samples of 0.3 – 0.6 kg, by weighting them in air and water (Archimedes principle). Sensitivity of the method is about 0.01 g/cm³ and precision is on the order of 0.02 g/cm³;
2. lithological information from El Hito-1 drillhole (Lanaja y Navarro, 1987) and
3. petrophysical data for the same formations found in the literature (e.g. Pueyo *et al.* 2015).

After evaluating the petrophysical information, the densities used in the modelling are, in g/cm³ (Fig. 6): 2.28 for the Neogene-Quaternary; 2.38 for the Paleogene; 2.6 for the Upper Cretaceous, 2.55 for the Lower–Middle Cretaceous, 2.65 for the Middle Jurassic, Lower Jurassic 2 and Lower Jurassic 1; 2.5 for the Keuper; 2.56 for the Buntsandstein–Muschelkalk and 2.75 for the Paleozoic basement.

3D Gravity modelling: Forward modelling and stochastic inversion

In this work, we have carried out the modelling using 3D GeoModeller software from Intrepid Geophysics (e.g. Guillen *et al.* 2008; Gibson *et al.* 2013) and following four steps:

1. Defining the stratigraphic pile.

In our model, the stratigraphic pile (Fig. 6) defines the chronology of the stratigraphic sequence making possible to manage the relationship between the geological formations in the computation of the 3D model and through the process of inversion.

As depicted in Fig. 6, the model is made of 10 layers: Paleozoic basement; Buntsandstein–Muschelkalk; Keuper; Lower and Middle Jurassic; Lower–Middle and Up-

Figure 5 (a) Bouguer anomaly with the lineaments mentioned on the text and the location of the petrophysical samples on top; see the text for explanation of labels Max1, Max2, Max3, Max4 and Min1, Min2. (b) Regional Bouguer anomaly. (c) Residual Bouguer anomaly with the location of the cross sections used in the 3D model and the lineations of the two minimum not seen on the Bouguer anomaly, MinL1 and MinL2 (see the text for explanation). (d) 100 m Digital Terrain Model used for the topographic correction and the inversion. The four maps show the outline of the geological outcrops of the anticlines (discontinuous dark blue line). LR: La Rambla anticline; ZdZ: Zafra de Zancara anticline.

Figure 6 Stratigraphic pile with the densities used in the modelling.

per Cretaceous; Paleogene and Neogene–Quaternary. Some of the formations have been grouped into series: The Triassic series (Buntsandstein–Muschelkalk and Keuper) that supposedly were not affected by the alpine orogeny and the Cretaceous–Jurassic series that were affected and deformed as we see today.

Note that although the three Jurassic layers have the same density and cannot be distinguished in terms of gravity modelling, we have left them separately to be consistent with the geological cross sections that come from the ALGECO2 Atlas (IGME 2009) and we have used these cross sections to build up the 3D model.

2. Generating the initial geological model.

The structural model is build up in a volume that has the same horizontal dimensions as the gravity data, 20 km × 25 km, the top is located at the topographic surface (the 100 m digital elevation model; Fig. 5d) and the bottom at a depth of –3500 m, which includes the upper part of the Paleozoic basement.

The model has been created from a 100-m grid Digital Elevation Model, geological information from the El Hito-1 borehole (Lanaja and Navarro, 1987), the surface geology from the 1:50,000 National Geological Map (MAGNA) series and 12 geological cross sections that image the main geological structures (see Fig. 2 and 5c for location). The cross sections are based on IGME (2009), Biete (2010) and Biete, Roca and Hernaiz-Huerta (2012), and the reinterpretation of seismic lines DD' and C8ext specifically reworked for this work. All cross sections but C4, which has a roughly N-S direction, run in SW-NE direction, perpendicular to the geological strike. The lithologies and faults from the cross sections were digitized in GeoModeller and the network of faults affecting the different formations together with the relation-

ships between them have also been integrated in the model. The digitized cross sections within GeoModeller together with the surface geology, with the constraints of the topography and the information from the El Hito-1 borehole, are interpolated and extrapolated using co-kriging, assuming that the contact data for each interface lie on a iso-potential surface, and the orientation vectors are orthogonal to a local tangential plane to the potential field (McInerney *et al.* 2005). In this way, the relevant data that have to be taken into account in the generation of the 3D surface can be located above or below the geologic contact surface we want to generate.

3. Refining the model by means of forward gravimetric modelling.

Once generated the first 3D model, the cross sections were revised to account for possible inconsistencies and, in some cases, reinterpreted and improved by means of GeoModeller's two-dimensional forward gravity modelling tool. An additional cross section, namely CorteCay1, was built up afterwards based on the surface geological data in order to constraint the section of the model between section C8EXT and Corte 5.

To ensure the convergence in the inversion process, it is best to start off with a model whose gravimetric response has a similar pattern than the observed anomaly. For that reason, we have refined the model through forward calculation: we have computed the gravimetric response of the 3D model and compared with the observed residual gravity anomalies.

This is an iterative process that starts with computing the forward gravity anomaly then reviewing the results and modifying the model by making changes in the cross sections and then repeating the forward calculation. This process will continue until the calculated anomaly pattern resembles that of the observed anomaly. In this way, we also ensure that the results of the inversion will honour not only the gravimetric observations but also the geological inputs. A thorough description of the modelling process can be found in Gillen *et al.* (2004, 2008). In Fig. 7(a), we show the calculated residual gravity anomalies prior inversion (i.e. the calculated residual anomalies from forward modelling). The misfits prior inversion (observed minus calculated residual gravity anomaly from forward modelling) are shown in Fig. 7(c).

To make the calculations, each lithology of the 3D volume is discretized in geological 3D cells (voxels). Taking into account the spatial distribution of the gravity data and the resolution of the petrophysical data, the geological model has been discretized in cells of 250 m × 250 m × 25 m. Each cell has a density value according to its lithology.

Figure 7 (a) Calculated residual gravity anomaly prior inversion. (b) Calculated residual gravity anomaly after inversion with the liniations of the two minimum seen on the residual Bouguer anomaly (Fig. 5C) labelled MinL1 and MinL2 (see the text for explanation). (c) Misfits (observed minus calculated residual gravity anomaly) prior inversion. (d) Misfits (observed minus calculated residual gravity anomaly) after inversion. The four figures show the location of the cross sections used in the 3D model and the outline of the geological outcrops of the anticlines (discontinuous dark blue line).

Figure 8 Results of the inversion along cross section Corte 6 and DDprima (a and b respectively). The cross sections before the inversion are also included in the figure for comparison. LR and ZdZ show the location of the anticlines. The location map show the cross sections used in the 3D modelling. The sections presented in this figure are depicted in red.

4. Performing the gravimetric inversion.

GeoModeller inversion uses the same model discretization as in the forward modelling. The calculations of the gravimetric inversion are based on the Monte Carlo Markov Chain method (Tarantola, 2005) that consists of minimizing the gravimetric misfits (difference between observed and calculates gravity anomalies) by changing the lithological boundaries, the density or both parameters. For this inversion, we have set the geological constraints to allow changes in the lithological boundaries with an uncertainty of 25% and leaving the densities to change within a standard deviation of 0.02 g/cm^3 , which corresponds to the uncertainty assigned to the measurements. The probability of property change is set to 50%. Consequently, the inversion process modifies both geometry and density of the differ-

ent lithologies within the uncertainty parameters. The edge effects are avoided by expanding the geological model and by reflecting and rotating the modelled geology itself across the edges and corners of the model. The inversion has run for 50×10^6 iterations, and at the end of the inversion the root mean square of the misfits (difference between observed and calculated residual Bouguer anomaly) has been 0.32 mGal.

RESULTS

In general terms, the pattern of the calculated anomaly after the inversion (Fig. 7b) is quite similar to the one of the observed anomaly (Fig. 5c). In the areas that are well constrained by the cross sections, the misfits show values between -0.5 and $+0.5$ mGal except for small and localized misfits of higher

Figure 9 (a) Top of the basement obtained from the inversion. The figure shows the location of the cross sections used in the 3D model and the outline of the geological outcrops of the anticlines (discontinuous dark blue line). In purple dashed line: basement fault that crosscut the Buntsandstein–Muschelkalk layer. (b) Top of the Buntsandstein–Muschelkalk obtained from the inversion. The figure shows the location of the cross sections used in the 3D model and the outline of the geological outcrops of the anticlines (discontinuous dark blue line) as well as the outline of the La Rambla and Zafra de Zancara new possible structural closure in maroon. In purple dashed line: basement fault that crosscut the Buntsandstein–Muschelkalk layer. (c) Top of the Keuper obtained from the inversion. The figure shows the location of the cross sections used in the 3D model and the outline of the geological outcrops of the anticlines (discontinuous dark blue line). In purple dashed line: basement fault that crosscut the Buntsandstein–Muschelkalk layer. (d) Top of the Lower Jurassic. The figure shows the location of the cross sections used in the 3D model and the outline of the geological outcrops of the anticlines (discontinuous dark blue line) as well as the outline of the La Rambla and Zafra de Zancara new possible structural closures in maroon and dark blue, respectively.

Post-inversion Model

Figure 10 3D model depicting the target layers: Top of the basement (number 3), top of Buntsandstein–Muschelkalk (number 2) and top of Lower Jurassic (number 1). Red line: outline of the outcrops of the anticlines.

amplitude that are attributed to superficial density inhomogeneities but do not affect the geometry of the targeted layers (Fig. 7d). The most relevant misfits are the two elongated ones with amplitudes ranging between 0 and 1 mGal (labelled as 1 on Fig. 7d) and between 0 and -1 mGal (labelled as 2 on Fig. 7d) placed slightly SW of the Zafra de Zancara anticline. These misfits are located in a place with no constraints but already out of our areas of interest, and, therefore, the uncertainty associated does not affect the evaluation of the possible structural closures.

As a succinct summary of the results, we have chosen Corte 6 and DDprima (Fig. 8a and b) as the most representative of the differences between the initial and final model. Comparing both cross sections, the results show that in some places the Buntsandstein–Muschelkalk it is slightly thinner than the one from initial model (like in the La Rambla anticline), whereas in other places, its thickness has not changed during the inversion (like in some places in the Zafra

de Zancara anticline). It is worth noticing the accumulation of Buntsandstein–Muschelkalk along the westernmost basement fault of Zafra de Zancara, reaching thicknesses that can be of the order of 400 m. According to the results of the inversion, this fault cuts the Buntsandstein–Muschelkalk facies and it also seems to affect the basement. The inversion shows that the basement has some relief of antiformal shape not depicted in the initial cross sections (e.g. Fig. 8a). One of these antiforms seems to be limited by a fault (see also Fig. 8a) that can be interpreted as a reverse fault of as a normal fault that has been inverted because the basement was affected by compression (as already suggested in Biete (2010)). Note that to build up the cross section DDprima presented in this work, we reinterpreted the seismic line of the same name in the area that crosses the La Rambla anticline that overlaps with Corte 4 cross section displayed in Fig. 4. The initial geometry of our Corte 6 that runs parallel to cross section DDprima is based on that profile.

Comparing the model from ALGECO2 Atlas (IGME 2009) (depicted in Fig. 3a–c) and the model resulting from the inversion (Fig. 9a–c), we can observe several differences:

1. The basement isolines of Fig. 9(a) show a more complex basement, both with respect to the structure of La Rambla (with a basement elevation not interpreted previously), as well as to the troughs between the structural highs of the area. The results of the inversion show that below the La Rambla anticline, the basement is not as flat as previously interpreted (see Fig. 3a) but folded into smaller anticlinal structures and 50–450 m deeper than in the initial model, particularly towards the N of the anticline. In the southernmost part, with lesser constraints, the anticline is more than 1000 m deeper.

In the Zafra de Zánacara anticline, the basement folds appear better defined and in the zone between Corte 1 and Corte 6, its depth is up to 225 m shallower than in the initial model. From the results of the inversion, we suggest that the basement fault affecting not only the basement but also the Buntsandstein–Muschelkalk F_m can be extended further south for 11 km (purple dotted line in Fig. 9a and b; black dotted line on Corte 6 in Fig. 8a). However this fault does not seem to affect the Lower Jurassic.

2. In the isolines map of the Buntsandstein–Muschelkalk top (Fig. 9b), the most important changes are in its geometry more than in its thickness: the top of the Buntsandstein–Muschelkalk depicts a more defined anticlinal geometry and it is located a few tens of meters deeper in some areas. The troughs that separate them also appear better defined. The Buntsandstein–Muschelkalk in the La Rambla anticline is slightly thinner than in the initial model (from 25 to 50 m) and a few tens of meters shallower whereas in the Zafra de Zánacara anticline the thickness has remained almost unchanged except along the westernmost basement fault, where we find accumulations of Buntsandstein–Muschelkalk of about 400 m (e.g. Corte 6, Fig. 8a).

3. In the La Rambla anticline, changes in the thickness of the Keuper are of the order of tens of meters whereas in Zafra de Zánacara are closer to the ones of the initial model (see top of the Keuper in Fig. 9c)

Top of the Lower Jurassic 1 isolines (Fig. 9d) are located approximately between 0 and –1200 m (asl). Lower Jurassic 1 is deeper towards the west and shallower towards the east. The most relevant difference in depth lies at the eastern end of Corte 6, where the Lower Jurassic is c. 700 m shallower. In most of the study area, the Lower Jurassic 1 has approximately the same thickness as in the initial model. One exception is the thickness along section DDprima (Fig. 8b) in the La Rambla anticline, where the results of the inversion show a thinner

Lower Jurassic 1, which is more consistent with its thickness everywhere else in the anticline.

The most relevant changes in the thickness of the Cretaceous, Paleogene and Neogene–Quaternary layers occur in the zone between the anticlines, where these layers are several tens of meters thicker. In Zafra de Zánacara and La Rambla anticlines, their thicknesses are closer to the ones in the initial model.

DISCUSSION AND CONCLUSIONS

Comparing the initial and final model, we observe the following:

In areas with sufficient seismic coverage such as Zafra de Zánacara, the structures obtained in both models are quite similar in geometry, although the stochastic inversion process results in a slightly shallower anticlinal geometry (between 50 and 100 m). That means, according to the density profile, that the seismic model presents a slight mass deficit, which is corrected in the gravimetric inversion process.

The results of the inversion differ from the interpretation of Muñoz-Martín and De Vicente (1998) that have the view the anticlines detected from the seismic data to the east of the Altomira Range, including Zafra de Zánacara and La Rambla, are “pull up” type artifacts from the seismic interpretation due to the increase of the seismic velocities where the sedimentary cover duplicates. Our 3D modelling results show that the anticlinal structures of Zafra de Zánacara and La Rambla involve the basement thus indicating that the basement highs interpreted from the seismic reflection data are real structures and not artefacts.

The most relevant differences occur where seismic coverage is scarce or non-existent. In Zafra de Zánacara, these differences are located at the SE of the anticline: here the process of stochastic gravimetric inversion suggests the presence of a small closure in the Buntsandstein–Muschelkalk, in an area where there is no seismic coverage. The structure of the Lower Jurassic 1 has only seismic coverage along line C10, so its geometry comes exclusively from the gravimetric inversion. In addition, some of the Zafra de Zánacara folds seem to be less affected by the faults than previously thought in the sense that the throw of the faults is, in general, smaller.

In the La Rambla anticline, the lack of the seismic information to constrain the initial model has resulted in greater freedom to modify the geometry of the lithologies so the calculated gravity anomaly would fit the observed anomaly. The most relevant differences are a more complex structure especially of the basement that appears more folded and deeper

compared with the almost flat basement in the initial model. Also, the Buntsandstein–Muschelkalk is slightly thinner.

- The possibility of the southern continuation of a fault affecting the basement and the Buntsandstein–Muschelkalk, in previous models though to be affecting only the northern part of Zafra de Zánacara westernmost border, have been deduced from the inversion (purple dotted line in Fig. 9a and b).

Concerning to the study of possible structural closures that might be used as CO₂ reservoirs, the gravimetric inversion has helped to define the geometry of possible structural closures particularly in places where the information from the seismic data were inconclusive.

- In the top of the Buntsandstein–Muschelkalk (Fig. 9b), the possible candidates to be a CO₂ structural closures are delimited by the –1500 m isoline that corresponds to an approximate depth of c. 2100 m, measuring from the topography. The maximum thickness of the structural closures is c. 200 m, and they are sealed by the evaporitic material of Keuper, with thicknesses of more than 500 m. The structures in the Zafra de Zánacara anticline were already defined in the ALGECO2 project. Here our results that show better defined geometries are consistent with that of the ALGECO2. In the La Rambla anticline, it is possible to delineate up to four closed structural highs that have been deduced from the gravity inversion only.
- In the top of Lower Jurassic map (Fig. 9d), two new possible structural closures are depicted, one in the Zafra anticline, with an average thickness of 250–300 m, reaching c. 400 m in some places and another one beneath La Rambla with thickness ranging between 200 and 400 m. These potential structural closures are delimited by the –500 m isoline that corresponds to depths beneath the topography of c. 800 m in La Rambla and 1000 m in the Zafra de Zánacara anticlines. The sedimentary cover acts as a seal, having more than 1200 m in thickness.

Nonetheless, more studies (porosity, etc.) are needed to evaluate the storage capacity in order to decide whether the use of these plausible structural closures is feasible.

Our 3D model (Fig. 10) improves the geological knowledge of the ALGECO2 model in three ways:

1. It depicts the geometry of all the layers defined in the stratigraphic pile, unlike the ALGECO2 model where only the top of the basement, top of Buntsandstein and top of Keuper was obtained.
2. The surfaces of the 3D model obtained from the inversion honour all the observables (gravity, seismic, surface geology and petrophysical data) with a horizontal uncertainty

of 125 m and an uncertainty in depth of 12.5 m, whereas the surfaces from ALGECO2 (IGME 2009) were obtained by manually interpolating the cross sections based on seismic interpretation and surface geology, implementing geological constraints. Therefore, we believe that the geological surfaces obtained from the gravity inversion have more definition and better resolution than the ones previously published.

3. Our characterization of La Rambla anticline allows us to take it into account as possible CO₂ storage site since four closed structures in the Buntsandstein–Muschelkalk and one in the Lower Jurassic 1 have been identified. Moreover, one more closed structure in the Lower Jurassic 1 in the Zafra de Zánacara anticline has been recognized. These structures were not singled out during the ALGECO2 project.

ACKNOWLEDGEMENTS

This study has been supported by the Instituto para la Reestructuración de la Minería del Carbón y el Desarrollo Alternativo de las Comarcas Mineras, Ministry of Energy and Industry, Spain. We thank José Luís García-Lobón and Manuel Bernat for their help in the first stages of the modelling. We also thank Serguei Goussev and Pavol Zahorec for their constructive comments and suggestions that have helped to improve this manuscript.

ORCID

C. Ayala <https://orcid.org/0000-0001-8457-8253>

REFERENCES

- Álvarez M., Capote R. and Vegas R. 1979. Un modelo de evolución geotectónica para la Cadena Celtibérica. *Acta Geológica Hispánica* 14, 172–177.
- Biete C. 2010. *La estructura alpina del basamento hercínico en la zona del Hito (sector meridional de la Cuenca de Loranca) y su influencia en la deformación contractiva de los materiales suprayacentes: Caracterización tridimensional de la estructura*. Tesis de Máster, Universidad de Barcelona, Barcelona, Spain, 17 pp.
- Biete C., Roca E. and Hernaiz-Huerta P.P. 2012. The Alpine structure of the basement beneath the southern Loranca Basin and its influence in the thinskin contractional deformation of the overlying Mesozoic and Cenozoic cover. *Geo-Temas*, 13, 173.
- Díaz-Molina M., Arribas J., Gómez J.J. and Tortosa A. 1995. Geological modelling of a reservoir analogue: Cenozoic meander belts, Loranca Basin, Spain. *Petroleum Geoscience* 1, 43–48.
- Gibson H., Sumpton J., Fitzgerald D. and Seikel R. 2013. 3D modelling of geology and gravity data: summary workflows for mineral exploration. *East Asia: Geology, Exploration Technologies and Mines*, Bali, 2013.

- Gómez J.J., Díaz Molina M. and Lendínez A. 1996. Tectonosedimentary analysis of the Loranca Basin (Upper Oligocene-Miocene, central Spain): a 'non sequenced' foreland basin. In: *Tertiary basins of Spain* (eds. P.F. Friend and C.J. Dabrio), pp. 285–294. Cambridge University Press, Cambridge.
- Guimerà J. and Álvaro M. 1990. Structure et evolution de la compression alpine dans la Chaîne Iberique et Chaîne Cotiere Catalane. *Bulletin de la Société géologique de France* 8 (VI), 339–348.
- Guillen A., Courrioux G., Calcagno P., Lane R., Lees T. and McInerney P. 2004. Constrained gravity 3D litho-inversion applied to Broken Hill. *SEG 17th Geophysical Conference and Exhibition*, Sydney.
- Guillen A., Calcagno Ph., Courrioux G., Joly A., Ledru P. 2008. Geological modelling from field data and geological knowledge, Part II - Modelling validation using gravity and magnetic data inversion. *Physics of the Earth and Planetary Interiors* 171, 158–169. <http://doi.org/10.1016/j.pepi.2008.06.014>.
- Hammer S. 1939. Terrain corrections for gravimeter stations. *Geophysics* 4, 184–194.
- IGME. 2009. Plan de selección y caracterización de áreas y estructuras favorables para el almacenamiento geológico de CO₂ en España. SID IGME database docs n° 64.044 to 64.055.
- Instituto Español de Normalización (IRANOR). 1985. UNE Norm 22-611-85. 6 pp.
- ITGE, 1990. Documentos sobre la Geología del Subsuelo de España. Tomo III (Madrid - Depresión Intermedia). ITGE, Madrid.
- Lanaja J.M. y Navarro A. 1987. *Contribución de la exploración petrolífera al conocimiento de la Geología de España*, 465 pp. Instituto Geológico y Minero de España, Madrid.
- McInerney P., Guillen A., Courrioux G., Calcagno P. and Lees T. Building 3D geological models directly from the data? A new approach applied to Broken Hill, Australia. 2005. U. S. Geological Survey Open-File Report 2005-1428.
- Muñoz-Martín A. and De Vicente G. 1998. Cuantificación del acortamiento alpino y estructura en profundidad del extremo sur-occidental de la Cordillera Ibérica (Sierras de Altomira y Bascuñana). *Revista de la Sociedad Geológica de España* 11, 39–58.
- Nettleton L. L. 1971. *Elementary Gravity and Magnetism for Geologists and Seismologists*. Society of Exploration Geophysicists, Tulsa, OK.
- Piña-Varas P., Ledo J., Queralt P., Roca E., García-Lobón J. L., Ibarra P. and Biete C. 2013. Two-dimensional magnetotelluric characterization of the El Hito Anticline (Loranca Basin, Spain), *Journal of Applied Geophysics*, 95, 121–134. <http://doi.org/10.1016/j.jappgeo.2013.06.002>.
- Pueyo E.L., Izquierdo-Llavall E., Rodríguez-Pintó A., Rey-Moral C., Oliva-Urcia B., Casas A.M., et al. 2015. Petrophysical properties in the Iberian Range and surrounding areas (NE Spain): 1-density. *Journal of Maps*, 12, 836–844. <https://doi.org/10.1080/17445647.2015.1084545>
- Querol R. 1989. *Geología del subsuelo de la Cuenca del Tajo*, 48 pp. Escuela Técnica Superior de Ingenieros de Minas de Madrid, Madrid.
- Tarantola A. 2005. *Inverse Problem Theory and Model Parameter Estimation*, 342 pp. SIAM. Philadelphia, PA.
- Valcárcel-Rodríguez M. 2015. *Reconstrucción y restitución 3D del anticlinal de Puerta-Pareja y estudio de su evolución deformacional y cinemática en su contexto regional (cuenca de Loranca y Sierra de Altomira)*. PhD Thesis, Universitat de Barcelona, Barcelona, Spain, 285 pp.
- Zeng H. 1989. Estimation of the degree of polynomial fitted to gravity anomalies and its application. *Geophysical Prospecting*, 37: 959–973. <https://doi.org/10.1111/j.1365-2478.1989.tb02242.x>