Postprint: Plant Physiology and Biochemistry 2019 134: 53-63.

Suppression of UV-B stress induced flavonoids by biotic stress: Is there “real crosstalk”?

Dirk Schenke^{1*}, Hashlin Pascananda Utami¹, Zheng Zhou¹, María-Trinidad Gallegos² and Daguang Cai¹

1) Department of Molecular Phytopathology and Biotechnology
Institute of Phytopathology
Christian-Albrechts Universität zu Kiel
24118 Kiel
Germany

2) Department of Soil Microbiology and Symbiotic Systems
Estación Experimental del Zaidín (CSIC)
Profesor Albareda, 1
18008-Granada
Spain

* corresponding author

Abstract

Plants respond to abiotic UV-B stress with enhanced expression of genes for flavonoid production, especially the key-enzyme chalcone synthase (CHS). Some flavonoids are antioxidative, antimicrobial and/or UV-B protective secondary metabolites. However, when plants are challenged with concomitant biotic stress (simulated e.g. by the bacterial peptide flg22, which induces MAMP triggered immunity, MTI), the production of flavonoids is strongly suppressed in both *Arabidopsis thaliana* cell cultures and plants. On the other hand, flg22 induces the production of defense related compounds, such as the phytoalexin scopoletin, as well as lignin, a structural barrier thought to restrict pathogen spread within the host tissue. Since all these metabolites require the precursor phenylalanine for their production, suppression of the flavonoid production appears to allow the plant to focus its secondary metabolism on the production of pathogen defense related compounds during MTI. Interestingly, several flavonoids have been reported to display anti-microbial activities. For example, the plant flavonoid phloretin targets the *Pseudomonas syringae* virulence factors flagella and type 3 secretion system. That is, suppression of flavonoid synthesis during MTI might have also negative side-effects on the pathogen defense. To clarify this issue, we deployed an *Arabidopsis* flavonoid mutant and obtained genetic evidence that flavonoids indeed contribute to ward off the virulent bacterial pathogen *Pseudomonas syringae* pv. *tomato* (Pst) DC3000. Finally, we show that UV-B attenuates expression of the flg22 receptor FLS2, indicating that there is negative and reciprocal interaction between this abiotic stress and the plant-pathogen defense responses.

Key words:

UV-B stress, biotic stress, flg22, crosstalk, flavonoids

Introduction

Flavonoids (lat. *flavus* = yellow) are plant secondary metabolites with a diverse function in modulation of plant-environment interactions, e.g. protection against abiotic or biotic stresses (Falcone Ferreyra et al., 2012). Furthermore, they provide much to the flavor and color of fruits or vegetables and their free-radical scavenging activities suggest they have also a positive impact on human diet (Ross et al., 2002). They are produced from phenylalanine via 4-coumaroyl-CoA. The chalcone synthase (CHS) is the key-enzyme for metabolite fluxes into the flavonoid pathway (Fig. 1). This enzyme converts 4-coumaroyl-CoA or 4-dihydrocoumaroyl-CoA to naringenin chalcone or phloretin, respectively (Gosch et al., 2010). Most flavonoid pathway genes (FPGs) are highly UV-B inducible (Schenke et al., 2011) and plants lacking flavonoids are hypersensitive to UV-B stress (Li et al., 1993; Landry et al., 1995). In *Arabidopsis thaliana* cell cultures flavonoids were efficiently induced by a fluence rate of 0.53 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Schenke et al., 2011), but even a lower fluence rate of 0.1 $\mu\text{mol m}^{-2} \text{s}^{-1}$ is sufficient to induce the CHS (Brown & Jenkins, 2008; Jenkins 2014). The *Arabidopsis chs/f3h* loss of function mutant (N8585) is unable to produce flavonoids, readily visible because of the light-coloured seeds (Fig. 1) (Appelhaagen et al., 2014). This mutant exhibits the highest susceptibility towards *Heterodera schachtii*, pointing to a role of flavonoids in defense against this cyst nematode (Jones et al., 2007) and silencing of CHS genes by RNAi in *Medicago truncatula* demonstrated the need of flavonoids in nodule formation by rhizobia (Wasson et al., 2006). Structurally, chalcone synthases are closely related to stilbene synthases (STS), which are for example involved in phytoalexin production, e.g. resveratrol from grape (Schröder and Schröder, 1989; Jeandet et al., 2010). Therefore, flavonoids play a role in plant defenses towards both abiotic and biotic stress. In this paper we focus on the role of flavonoids in the *Arabidopsis thaliana* response to biotic stress exerted by the phytopathogenic bacterium *Pseudomonas syringae* pv. *tomato* (*Pst*), because there are contradicting observations concerning flavonoid production after infection, which are reviewed in this introduction.

Fig. 1 Central role of the chalcone synthase (CHS) as entry-point for the flavonoid pathway in plants. A knock-out of the single copy gene in *Arabidopsis* results in plants void of flavonoids. Phloretin is reported to inhibit the T3SS and bacterial flagella of plant-pathogenic *Pst* DC3000 (Vargas et al., 2013).

Flavonoid production is suppressed by biotic stress

Plants suffering biotic stress, either directly *via* pathogen infection or simulated by application of conserved microbe-associated molecular patterns (MAMPs) elicit a defense response, which is termed MAMP Triggered Immunity (MTI). Pathogens in turn have evolved effectors, which they translocate into the host cells in order to interfere with plant defense responses, called effector triggered susceptibility, ETS. Bacterial pathogens, such as *Pseudomonas syringae*, deliver their effectors with the Type 3 Secretion System (T3SS) into the host cells. Recognition of these effectors by the plant immune system results then in Effector Triggered Immunity, ETI (Jones and Dangl, 2006). Thus, it is advisable to treat plants with MAMPs in order to observe their pure, original and non-manipulated defense reaction. It is known for more than three decades that plants suppress the accumulation of flavonoids when facing biotic stress (Schenke et al., 2011). Especially down-regulation of the CHS expression contributes to this phenomenon and this mechanism involves prevention of histone acetylation, a hallmark for gene activation (Schenke et al., 2014). Fig 2 shows the typical suppression of UV-B induced CHS expression by the MAMP flg22, which is a 22 amino acid conserved peptide from bacterial flagella. This kind of suppression is observed in several plants (Tab. 1).

Fig. 2 Crosstalk between UV-B and biotic stress mimicked by elicitation with flg22. The CHS expression is suppressed by flg22. These data from Arabidopsis have been published in Zhou et al. (2017). Shown is the response to four different treatments: Water/VIS (C), flg22/VIS (F), UV-B/VIS (U) and both stresses simultaneously (FU). CHS gene expression level was investigated at 4h and normalized to PP2A and Actin. Error bars represent standard error of triplicate experiments, and statistical significance was checked by a two-way ANOVA.

Fig. S1 Crosstalk is functional in *Brassica napus* leaves. The typical CHS expression pattern resembles that of Arabidopsis (Fig. 2) in response to four different treatments: Water/VIS (C), flg22/VIS (F), UV-B/VIS (U) and both stresses simultaneously (FU). CHS gene expression level was investigated at 4h and normalized to PP2A and Actin. Error bars represent standard error of triplicate experiments, and statistical significance was checked by a two-way ANOVA.

Tab. 1 Flavonoid production or CHS expression suppressed in plants under biotic stress

Plant system	Suppressing biotic stress	Reference
parsley cells	<i>Phytophthora sojae</i> elicitor	Hahlbrock et al. (1981)
carrot cells	<i>Pythium aphanidermatum</i> elicitor	Gleitz et al. (1991)
sorghum plants	<i>Cochliobolus heterostrophus</i> infection	Lo & Nicholson (1998)
onion plants	<i>Botrytis allii</i> infection	McLusky et al. (1999)
<i>Glehnia littoralis</i> cells	Yeast extract elicitor	Ishikawa et al. (2005)
Arabidopsis seedlings	flg22 and elf18 elicitor	Saijo et al. (2009)
Arabidopsis cells	flg22 elicitor	Schenke et al. (2011)
<i>Brassica napus</i> seedlings	flg22 elicitor	this work (Fig. S1)

Thus, there is a growing list of examples, which suggests this suppression of flavonoid production is a typical MTI defense reaction well conserved throughout the plant kingdom. Importantly, Lozoya et al. (1991) reported that elicitor treatment of parsley cell cultures completely suppressed flavonoid production, while the effect of UV-B on elicitor induced coumarins was only slightly slowing down their accumulation. Furthermore, a time course experiment to measure CHS transcript level showed that elicitor addition was able to reduce CHS expression at any time during UV-B treatment, indicating that as soon both stresses were applied simultaneously the biotic stress is getting the upper hand (Lozoya et al., 1991). This regulation is thought to enable the plant to redirect metabolic resources from the UV-B defense response to pathogen defense, for example to increase the production of phytoalexins (e.g. coumarins such as scopoletin) or lignin (Lo & Nicholson, 1998; Schenke et al., 2011). It is supported by the observations that pathogens deploy effectors to boost anthocyanin production in order to reduce the lignin content (Tanaka et al., 2014) and that UV-B led to increased resistance towards *B. cinerea* independent of flavonoid production, but depending on enhanced sinapate accumulation required for the synthesis of syringyl-type ('defense') lignin (Demkura and Ballare, 2012). However, soluble sinapoyl-glucose might also directly inhibit the pathogen growth without the need of lignin formation as has been shown for the Arabidopsis - *Verticillium longisporum* interaction (König et al., 2014). Taken together it appears that during biotic stress (MTI) plants shut down the UV-B induced flavonoid pathway in order to produce other defense metabolites (This shall be "hypothesis one").

Flavonoid production is contributing to pathogen defense

On the other hand, it has been shown that several flavonoids can also contribute to pathogen defense responses, besides their role as UV-B protective sunscreens, having antiviral, antibacterial or antifungal activity (Cushnie & Lamb, 2005). For instance, phloretin ("Fig. 1") targets both motility and the T3SS of *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000 (Vargas et al., 2013). Accordingly, it has been observed that several flavonoids are induced in response to pathogen attack. Rutin, for example, accumulates in *Pst* infected tomato plants (Lopez-Gresa et al., 2011) or in response to biocontrol treatment of potato against infection with *Verticillium dahliae* (El Hadrami et al., 2011). Though rutin is produced as UV-B protectant, it has also antibacterial properties (van der Watt and Pretorius, 2001). It serves as a signal released by host plant roots to attract e.g. the beneficial *Bacillus subtilis* (Singh et al., 2016) or mycorrhizal fungi (Lagrange et al. 2001) facilitating root colonization. Furthermore, rutin can enhance resistance to insects in tobacco (Misra et al., 2010) as well as to *Xanthomonas oryzae* pv. *oryzae*, *Ralstonia solanacearum* and *Pst* DC3000 in rice, tobacco and Arabidopsis, respectively (Yang et al., 2016). Therefore, rutin appears to mediate priming of plant resistance by inducing the early salicylic acid (SA) signalling pathway, quite similar to its aglycon quercetin (Yang et al., 2016; Jia et al., 2010). Quercetin exhibits anti-quorum sensing effects in *Pseudomonas aeruginosa* and reduces biofilm formation (Pejin et al., 2015). Other antimicrobial flavonoids are the rice phytoalexin sakuranetin, which is produced in response to the fungus *Pyricularia oryzae* (Kodama et al., 1992) or catechin, that inhibits infection hyphae formation by *Alternaria alternata* appressoria on strawberry leaves (Yamamoto et al., 2000) as well as rust infection of poplar (Ullah et al., 2017). It has been shown that catechin also interferes with *Pseudomonas aeruginosa* quorum sensing (Vandeputte et al., 2010). Taxifolin from kempas (*Koompassia malaccensis*) has antimicrobial activity against *Streptococcus sobrinus* (Kuspradini et al., 2009), while luteolinidin accumulated in sugar cane in

response to *Colletotrichum falcatum* (Malathi et al., 2008). Luteolin accumulated in infected sorghum and affected spore germination of *Colletotrichum sublineolum* (Du et al., 2010). However, the strongest evidence for flavonoids as phytoalexins comes from isoflavone-derived pterocarpanes, which are described for the *Fabaceae* family (legumes), such as medicarpin, pisatin, glyceollin or phaseolin. Legumes contain generally a greater number of chalcone and stilbene synthase family proteins, which points to functional divergence. Samac and Graham (2007) concluded from other studies that suppression of the anthocyanin branch in soybean after infection with avirulent *P.syringae* pv *glycinea* and concomitant up-regulation of the flavone and isoflavone branch enhanced production of antimicrobial isoflavones, which is supported by Yu et al. (2003) describing that isoflavonoid synthesis was increased by a combination of ectopic expression of maize transcription activators and inhibition of the competing anthocyanin pathway. This suggests that flavonoid suppression is also present in legumes. It is likely that some chalcone/stilbene synthase copies are required to cope with biotic stress (phytoalexin and/or phytoanticipin production), some for UV-B protection and some to establish symbiotic interactions, e.g. with rhizobia or mycorrhiza (Steinkellner et al., 2007; Hassan and Mathesius, 2012). Such manifold involvement of flavonoids in plant-environment interactions is probably the evolutionary driving force to increase CHS/STS copy numbers in order to respond to both biotic and abiotic stress. These are just a few examples summarized in figure 3. Interestingly, some pathogens are able to degrade flavonoids, such as *Verticillium dahliae* metabolizing rutin (El Hadrami et al., 2015) and the rice blast fungus *Pyricularia oryzae* metabolizing the phytoalexin sakuranetin (Katsumata et al., 2017) or deploy flavonoid efflux pumps as shown for *Pst* (Vargas et al., 2011). In conclusion, plants produce flavonoids in response to infection because of their antimicrobial properties ("hypothesis two").

Fig. 3 Overview of selected flavonoid family members reported to have antimicrobial properties. The black skeleton structure is typical for each family, while the red substitutions are specific for the given example. For details see text.

Aim of this study

Clearly both hypotheses are logically straight-forward, but also somehow mutually exclusive. Regarding pathogen defense, is the production of flavonoids now disadvantageous (“hypothesis one”) or advantageous (“hypothesis two”) for the plant? We try here to resolve this question by deploying the *Arabidopsis chs/f3h* double mutant completely lacking flavonoids and compared growth of the bacterial pathogen *Pst* between mutant and wild-type plants. Finally, we searched for indications whether the suppression of UV-B induced flavonoids by biotic stress is part of a unidirectional or reciprocal crosstalk, involving also suppression of biotic stress responses by UV-B.

Results

In a first experiment, we challenged the *Arabidopsis chs/f3h* double mutant void of flavonoids with different bacterial strains (Fig. 4). The results indicate that flavonoids contribute to restrict the bacterial growth in *Arabidopsis*. The *Pst* T3SS mutants grow less well than the virulent strain DC3000, which confirms the role of effector proteins to establish a compatible interaction and that the applied system is generally working.

Fig. 4 Flavonoids dampen bacterial growth in *Arabidopsis*. Comparison of *Pst* DC3000 (WT), *hrpL* (transcription factor for virulence genes; Castillo-Lizardo et al., 2015) and *hrcQ* (T3SS-component; Badel et al., 2006) growth rate in *Arabidopsis* Ler-0 (WT) and *chs/f3h* mutant. Growth rates of wild-type (DC3000) and mutant (*hrpL* and *hrcQ*) *Pst* strains were analysed by qPCR of biomass marker genes. Statistical significance was checked by t-test (*: $p \leq 0.05$ **: $p \leq 0.01$) based upon 3 biological replicates.

To analyse the effect of flavonoids on *Pst*, we analysed bacterial gene expression in response to phloretin as well as naringenin by RNAseq (Tab. S1). Notably, the bacteria respond to these treatments with induction of multidrug efflux pumps, while the T3SS was not expressed

In order to further support the role of flavonoids in plant defense against pathogens we investigated the effect of UV-B on *Pst* growth in *Arabidopsis* and infected UV-B pre-treated *Arabidopsis* plants with *Pst* (Fig. 5). However, we observed that the UV-B pre-treatment of *Arabidopsis* had no significant effect on the subsequent infection with two *Pst* strains, including the wild-type DC3000. Only for the *hrcQ* mutant the UV-B pre-treatment suppressed bacterial growth.

Fig. 5 Effects of UV-B pre-treatment on bacterial growth in Arabidopsis Ler-0 (WT) and *chs/f3h* mutant plants. Growth rates of wild-type (DC3000) and mutant (*hrpL* and *hrcQ*) *Pst* strains were analysed by qPCR of biomass marker genes. Statistical significance was checked by t-test (*: $p \leq 0.05$) based upon 3 biological replicates.

To investigate whether flavonoid pathway genes (FPGs) are generally suppressed in response to biotic stress and to search for antagonistically regulated defense genes we deployed a meta-analysis of publicly available gene expression data (Fig. S2 and S3) using GENEVESTIGATOR (Hruz et al., 2008). In Arabidopsis the FPGs were generally suppressed by biotic stress (Fig. S2A), but two of these stress treatments showed a strong up-regulation of the key marker gene CHS. The up-regulation in response to virulent *Pst* DC3000 was investigated in more detail, because of its relevance to this work (Fig. 6).

Fig. 6 Meta-analysis by GENEVESTIGATOR of the Arabidopsis – *Pst* interaction with respect to suppression of FPGs. The *Pst hrcC* mutant with defective T3SS is non-pathogenic and could resemble MTI, while *Pst* DC3000 is a virulent strain causing disease via ETS. The avirulent *Pst avrRpt2* strain provokes the Effector Triggered Immunity (ETI). Similar treatments were selected and indicate that CHS up-regulation is also time dependent, taking effect after 24h from inoculation.

Fig. S2 Meta-analysis of 140 selected public gene expression experiments involving Arabidopsis treatments by GENEVESTIGATOR (perturbations). A) flavonoid pathway genes (FPGs), B) Pattern Recognition Receptor genes triggering MTI, C) genes involved in defense signalling and D) genes for lignin or phytoalexin production (secondary metabolism).

Fig. S3 Meta-analysis of 60 selected public gene expression experiments involving Arabidopsis mutants by GENEVESTIGATOR (perturbations). A) flavonoid pathway genes (FPGs), B) Pattern Recognition Receptor genes triggering MTI, C) genes involved in defense signalling and D) genes for lignin or phytoalexin production (secondary metabolism).

Additionally, two genes were identified in Arabidopsis as antagonistically regulated to the FPGs by treatments (Fig. S2) and in mutant backgrounds (Fig. S3). From the tested pathogen defense associated genes, the flg22-binding receptor like kinase FLS2 (Gómez-Gómez & Boller, 2000) and the glucosyltransferase UGT72E1 involved in monolignol 4-O-glucoside production (Lanot et al., 2006) were down-regulated when FPGs were up-regulated and *vice versa*. This tempted us to investigate the expression of both genes in our Arabidopsis system, observing a negative effect of UV-B on *FLS2* but not on *UGT72E1* expression (Fig. 7), being indicative of a real reciprocal crosstalk between UV-B and flg22 signalling.

Fig. 7 Expression analysis of *FLS2* (A) and *UGT72E1* (B) in response to flg22 and UV-B alone or in combination. Gene expression levels were normalized to Actin. Gene expression was investigated after 4h in response to four different treatments: Water/VIS (C), flg22/VIS (F), UV-B/VIS (U) and both stresses simultaneously (FU). Error bars represent standard error of triplicate experiments, and statistical significance was checked by a two-way ANOVA. However, the two-tailed p-value of an unpaired t-test equals 0.0462 for *FLS2* expression when comparing only the C and U treatment.

Furthermore, we investigated differential CHS expression in response to biotic stress within other plant systems and selected several chalcone and stilbene synthase family proteins for a GENEVESTIGATOR meta-analysis (Fig. S4). In Arabidopsis only the CHS showed striking differential expression in response to biotic stress and elicitor application, while the other three chalcone and stilbene synthase genes were not strongly regulated in response to pathogens (Fig. S4A). Apparently, in almost all plant species investigated, some CHS/STS copies are suppressed in response to biotic stress.

Fig. S4 Meta-analysis of chalcone and stilbene synthase family protein expression in Arabidopsis and several other plant species by GENEVESTIGATOR with focus on biotic stress (and light conditions when available). The dots mark CHS orthologous genes as identified by GENEVESTIGATOR. A) Arabidopsis, B) tomato, C) *Physcomitrella patens*, D) barley, E) rice, F) maize, G) soybean, H) *Medicago truncatula* and I) wheat.

Discussion

Flavonoids contribute in Arabidopsis to fend off pathogenic bacteria

Some aspects about suppression of flavonoid pathway genes (FPGs) during flg22-triggered MTI had not been dealt with so far. We have investigated if flavonoids really play also a role in the pathogen defense system *Arabidopsis thaliana* – *Pseudomonas syringae* pv. tomato DC3000 as suggested by Vargas et al. (2013) by deploying the flavonoid deficient Arabidopsis mutant *chs/f3h*. Reduced bacterial growth on the mutant would support hypothesis 1 (that less flavonoids means more resources for defense metabolites such as phytoalexins and lignin), while a better growth rate would support hypothesis 2 (that also flavonoids are produced against pathogens, thus less flavonoids means reduced anti-microbial activity). It appears that flavonoids indeed contribute to plant protection against the bacterial pathogen *Pst* DC3000, because bacterial growth was enhanced in the mutant (Fig. 4). We got also indications that UV-B pre-treatment could suppress bacterial growth, though only for the *Pst hrcQ* mutant (Fig. 5). Because this effect was lost in the *ch3/f3h* mutant, it is an indication that UV-B induced flavonoids were indeed involved in suppression of bacterial growth, supporting hypothesis 2. However, the majority of reports describing effects of UV-B on plant-pathogen interactions showed that disease symptoms rather increased (Tab. 2). Though we now show that flavonoids indeed contribute to pathogen defense in Arabidopsis, there is still ample evidence for suppression of flavonoid production during plant – pathogen interactions (Tab. 1), including Arabidopsis (Zhou et al., 2017).

Tab. 2 Effects of UV-B light on plant-pathogen interactions

Plant-pathogen interaction	Effect of UV-B light	Reference
susceptible cucumber vs. <i>Colletotrichum lagenarium</i> or <i>Cladosporium cucumerinum</i>	UV-B pre-treatment increased disease severity	Orth et al. (1990)
sugar beet vs. <i>Cercospora beticola</i>	disease severity increased in combined exposure to UV-B	Panagopoulos et al. (1992)
susceptible wheat vs. <i>Puccinia recondita</i>	UV-B exposure increased severity of rust disease	Biggs et al. (1984) in Manning & Tiedemann (1995)
Arabidopsis vs. <i>Botrytis cinerea</i>	Increased plant resistance, independent of flavonoids, but involving sinapate production	Demkura and Ballare (2012)

Maybe it would be helpful to look on this crosstalk from a pathogen point of view. Several pathogens deploy effectors to manipulate their host and some effectors of the biotrophic pathogen *Ustilago maydis* can induce flavonoid/anthocyanin production thereby withdrawing resources from lignin formation (Tanaka et al., 2014). Thus, also *Pst* might theoretically deploy some of its T3SS effectors to influence flavonoid production. For example, it has been shown that the effector HopZ1 promotes *P. syringae* multiplication in soybean by targeting the enzyme 2-hydroxyisoflavanone dehydratase (GmHID1), which is required to produce the isoflavanoid and phytoalexin precursor daidzein (Zhou et al., 2011). Mechanistically, up-regulation of FPGs by *Pst* DC3000 infection could be achieved by effectors, reported to induce for example the Arabidopsis NCED3 gene resulting in elevated ABA levels (de Torres-Zabala et al., 2007+2009). ABA in turn appears to have a positive impact on FPG expression (Fig. S2 and discussion below). Our meta-analysis focusing on the Arabidopsis – *Pst* interaction indicates suppression of FPGs in response to infection with bacterial T3SS-deficient (*hrcC*) mutants, resembling MTI (Fig. 6). Infection with *Pst* DC3000, which deploys effectors, up-regulates FPG expression but this effect is limited to the key-enzyme CHS. Thus, such a putative T3SS effector has to be able to discriminate between the otherwise highly co-regulated FPGs. In this case CHS up-regulation might be a strategy of *Pst* DC3000 to interfere with lignin and phytoalexin production, similarly to that of *Ustilago maydis*. In another meta-analysis (Fig. S4) we found CHS up-regulation also in soybean (infected with *Phakopsora pachyrhizi*) or wheat (infected with *Puccinia graminis*) pointing to crosstalk manipulation. Gene duplication events (as observed e.g. for CHS family members in legumes) could allow differential expression of the multiple alleles under diverse conditions, especially

noticeable in the two legumes soybean and *Medicago truncatula*. That is, copies, which are predominantly induced by biotic stress, might enable the production of flavonoid phytoalexins in the presence of microbes (= activated MTI). Only for tomato and *Physcomitrella patens* there was no obvious CHS suppression by biotic stress treatments, but this might be due to the very limited number of such experiments in the database. Fig. 6 suggests furthermore, that the host plant in turn reinstalls suppression of flavonoid production by down-regulation of FPGs in response to ETI, which is resembled by infection with *Pst* expressing the *avrRpt2* gene (Tsuda et al., 2009). Thus, it is also possible that single up-regulation of the CHS in response to virulent *Pst* DC3000 is a plant reaction to favour production of compounds such as phloretin (Fig. 1), which targets the bacterial T3SS and flagella (Vargas et al., 2013), while no other flavonoids are produced. Here we show by RNAseq that bacteria respond to phloretin as well as naringenin with induction of multidrug efflux pumps in conditions where the T3SS is not expressed (Tab. S1). This indicates that *Pst* defense against plant flavonoids is mainly attributed to T3SS-independent expression of efflux pumps and not to effector deployment. Among these efflux pumps we detected MexAB-OprM, which is required for counteracting the detrimental effects of these flavonoids on *Pst* DC3000 and full virulence (Vargas et al., 2011). Considering that life began in aquatic environments with no atmosphere screening space radiation, phenylpropanoid metabolites such as flavonoids first served solely as UV-B protectants. Motile pathogens, such as bacteria posed later a new threat, thus - before developing something completely new - modified flavonoids were used to ward them off. Later, when plants began to conquer the land, they evolved another branch of the phenylpropanoid pathway that produced lignin. Upon pathogen recognition plants had now to decide whether to restrict the invasion with flavonoids or lignin. Probably lignin was more effective, hence the suppression of flavonoid production. Thus, it appears that CHS activation to produce antibacterial flavonoids is an ancient defense response, substituted with an evolutionary newer plant strategy to suppress the FPGs during MTI and ETI in order to induce more effective defenses, such as lignin and phytoalexins.

Crosstalk between UV-B and biotic stress appears to be reciprocal

As stated in Schenke et al. (2011), there was no apparent and statistically significant influence of UV-B on the expression of the flg22-inducible marker gene *FRK1*, hence it appeared that suppression is rather unidirectional and not reciprocal, which would be an important prerequisite to consider this regulation “real crosstalk” (Jansen et al. (THIS ISSUE)). A meta-analysis of Arabidopsis expression data (Fig. S2) generally confirmed that up-regulation of pathogen defense genes correlated with suppression of FPGs, whereas the opposite was not observable when focusing on the *FRK1* gene. In order to detect conversely regulated pathogen defense genes, 200 Arabidopsis perturbation experiments were selected upon co-regulation of six FPGs with GENEVESTIGATOR. Figure S2 shows that the FPGs are also inducible by sucrose and it is known that this effect can be attenuated by concomitant flg22 or elf18 treatment as well (Saijo et al., 2009). Thus, experiments with Arabidopsis grown on sucrose-containing medium have been excluded from our meta-analysis. In 60 experiments most of the 6 FPGs showed up-regulation, while in 140 cases these genes were collectively down-regulated (Fig. S2A, S3A). The FPGs were generally suppressed by biotic stress, but five treatments showed somehow up-regulation of the key marker gene CHS. However, 4 of these treatments came from very early time-points with no clear induction of pathogen defense responses and CHS up-regulation by *G. orontii* at 6h (study 2) stands even in contrast to a comparable study showing no significant regulation pointing to treatment effects in study #2. Such effects could be for example light-induction when control plants were directly harvested after the night period and treated plants are exposed additionally to light conditions during infection. When comparing FPG expression from these 60 experiments with 3 sets containing each six genes involved in MTI (Fig. S2B, S3B; Pattern Recognition- and Co-Receptors, PRRs), involved in defense signalling, including *FRK1*, (Fig. S2C, S3C) or being involved in secondary metabolism for phytoalexin or lignin production (Fig. S2D, S3D), the *FRK1* was suppressed only 7 times when FPGs were up-regulated. However, two other genes were suppressed 40 times (*FLS2*, encoding the flagellin receptor) or 33 times (*UGT72E1*, thought to be involved in lignin metabolism). Thus, we conducted an expression analysis of these two genes (Fig. 7) observing that these pathogen defense related genes were up-regulated in response to flg22 as expected. However, only the *FLS2* gene was negatively affected by UV-B and the co-treatment showed that biotic stress signalling initiated by flg22 application could override the negative effect of UV-B. In parsley cell cultures it was shown at the metabolite level that UV-B

had also an attenuating effect on elicitor-induced coumarin production (Lozoya et al., 1991). Therefore, the negative effect of UV-B on basal FLS2 expression as suggested by the meta-analysis in Fig. S2B and confirmed in Fig. 7, respectively, adds a novel aspect to this now reciprocal crosstalk. It is further supported by the meta-analysis in Fig. S3B, showing that in the *uvr8* KO background FLS2 expression was increased. This points to a potential involvement of the UV-B receptor UVR8 in this crosstalk when it comes to UV-B effects on the biotic stress response. Because all FPGs are strongly down-regulated in the *uvr8* mutant (Fig. S3A), a further suppression by biotic stress cannot be investigated. An antagonistically regulation between FLS2 and the FPGs might still be biologically relevant, since attenuation of the flg22-induced MTI *via* dampening FLS2 expression under high UV-B levels could contribute to re-install an efficient UV-protection by enabling the production of flavonoids again.

Furthermore, plants can detect additional bacterial MAMPs, such as elf18 or lipopolysaccharides. However, other MAMP receptors, like EFR for elf18 recognition, are not so obviously down-regulated when FPGs are up-regulated (Fig. S2), whereas elf18-dependent MTI is also able to suppress the FPGs (Serrano et al., 2012). Therefore, testing long-time UV-B exposed plants for decreased FLS2 protein levels will be required in order to evaluate the effect of UV-B on overall flg22-induced defenses. Nevertheless, FLS2 activation can in turn inhibit light-induced stomatal opening (Zhang et al., 2008), supporting there is real crosstalk. Maybe it is critical to use the term crosstalk when arguing on different levels (e.g. stress perception, signal transduction, phytohormones, transcription factors/gene regulation or secondary metabolites). In this system we observe effects on the very end of UV-B signalling (expression of structural FPGs for metabolite production) and on the very beginning of flg22 signalling (on the stress perception level). Also a comparison on the level of secondary metabolism is complicated, since UV-B induced flavonoid secondary metabolites can contribute to both, UV-B protection (Li et al., 1993) as well as protection towards biotic stress exerted e.g. by infection with *Pst* (Fig. 3 and 4). Thus, this dual function of flavonoids makes this system additionally a nice example for cross-resistance (Jansen et al. (THIS ISSUE)).

Additional to induction of flavonoids, UV-B has been reported to trigger expression of PR1 and PDF1.2 in *Arabidopsis* *via* ROS production, albeit clear activation of PDF1.2 required unnaturally extensive (30h) UV-B irradiation (Mackerness et al., 1999). This observation is further challenged by a report from Serrano et al. (2012), who demonstrated that naringenin pre-treatment attenuated the flg22-induced activation of PR-1 and PR-2 genes and that PR expression was increased at least in the CHS mutant line SALK_020583. Thus, the plant would also benefit in this respect from CHS suppression during MTI. Intriguingly, Falcone Ferreyra et al. (2015) demonstrated that the *Arabidopsis* *dmr6* mutant is not a salicylic acid (SA) 3-hydroxylase, but it is involved in the production of flavones, such as apigenin. They suggest that pathogens could induce flavones to decrease SA levels, because the *dmr6* mutant has been shown to accumulate more SA. Whether flavone synthesis inhibits SA production or competes for a shared precursor (coumaric acid) remains to be elucidated. Furthermore, oxidative damage by UV-C has the potential to increase plant resistance, for example in the *Arabidopsis* - *Hyaloperonospora parasitica* interaction (Kunz et al., 2008). Flavonoids can scavenge such induced reactive oxygen species (Hideg et al., 2013), thus they might be also able to attenuate the oxidative burst, an important pathogen defense reaction that triggers the hypersensitive response (HR) against biotrophic pathogens. Consequently, those side effects may contribute to an altered plant resistance towards pathogens uncoupled from flavonoid production. If flavonoids in *Arabidopsis* are suppressed by biotic stress, but also contribute to plant defense it might be an option for the plant to store sufficient amounts as preformed phytoanticipins in the vacuole (Dixon & Pasinetti, 2010), which could allow a synthesis of both hypotheses. In order to visualize what is known and what we could expect from the crosstalk between the abiotic UV-B stress response and biotic bacterial infection Fig. 8 was drawn to give a quick overview of the above discussed findings.

and in grape ABA contributes to UV-B protection (Berli et al., 2010). Furthermore, it appears there is a mutually antagonistic relationship between ABA and SA (Yasuda et al., 2008), which could also contribute to the regulation of FPGs as described above. Thus, the involvement of ABA could be beneficial for flavonoid production, while compromising plant immunity (Cao et al., 2011). Intriguingly, UV-B irradiation also induces ABA release from human cells (Bruzzzone et al., 2011), suggesting that ABA is involved in a very ancient defense response to UV-B stress.

We conclude there is reciprocal crosstalk between abiotic and biotic stress signalling on the production of flavonoids, because of the several arguments presented here. Furthermore, flavonoids play a role in plant protection against both UV-B and pathogen infection, indicating there is potential for cross-resistance. Although the plant's pathogen defense response appears to prevail by suppressing the UV-B induced *FPG* up-regulation, a negative effect under rising UV-B levels depends not only on the inhibited *de novo* synthesis of flavonoids, but most likely also on storage capacities and flavonoid turn-over. Thus, the temporary suppression of flavonoid production by biotic stress might be bridged for a certain time. However, in light of the observed stratospheric ozone decline (Ball et al., 2018), this crosstalk could impact the plant's UV-B resistance / fitness under a continuous exposure to high UV-B levels.

Material & Methods

Plant treatment with UV-B and flg22

Four *Brassica napus* seedlings were grown in pots with a soil/sand mixture under short day conditions (8 h light) and after two weeks transferred for two days to darkness in order to completely suppress *CHS* mRNA levels (Zhou et al., 2017). After two days in darkness two pots were sprayed with water, while the remaining two pots were sprayed with 1 mM flg22. To let flg22 taking effect the sprayed plants were incubated for two hours in darkness and were then exposed to UV-B or VIS-light as control for four h. A three mm thick glass plate was used to constrict the UV-B radiation emitted by two broadband UV-B lamps with an emission spectrum from 290 to 315 nm (Philips TL 20W/12 RS) and two PROTEC.CLASS lamps (PLSL 18W/21) for concomitant white light supply (12.69 m²/s) to natural level which is sufficient to induce photomorphogenic UV-B response (0.53 μmol m²/s), while the control plants exposed only to VIS light were shielded with two additional layers of polyester plastic foil (Folanorm SF/AS 0.13 mm, Folex GmbH, Cologne, Germany). Thus, each biological replicate consists of the 4 treatments Water / VIS-light control (C), flg22 treatment (F), UV-B treatment (U) and the co-treatment flg22/UV-B (F/U). The treatment of *Arabidopsis thaliana* seedlings was similar as described in Zhou et al. (2017).

RNA isolation

Leaf material from *Brassica napus* was harvested into 2 ml tubes containing ca. 20 zirconium beads (Zircosil 1.2-1.7 mm, Mühlmeier mahltechnik, Germany), immediately flash-frozen in liquid nitrogen and stored at -80°C for later RNA isolation. Total RNA was extracted with 1ml TRIzol® reagent (Thermo Fisher Scientific) according to the manufacturer's instructions using a Precellys® Evolution ball-mill (bertin technologies) grinding for 3 x 20 sec at 6,800 rpm. Samples were incubated for 5 min at room temperature. Thereafter, 200 μl chloroform was added, vortexed vigorously and centrifuged at 10,000 rpm for 15 min at 4°C. The clear supernatant was transferred into a sterile 1.5 ml tube and incubated at 4°C for 30 min before being centrifuged at 12,000 rpm for 15 min at 4°C. Then, the supernatant was discarded and RNA precipitate was washed twice with 1 ml 80% DEPC-ethanol and 1 ml 100% ethanol followed by centrifugation at 12,000 rpm for 5 min at 4°C and discard the supernatant. The RNA pellet was dried in the air stream and stored at 4°C after being dissolved in 50 μl DEPC-water.

cDNA synthesis and qPCR

1 µg total RNA was treated with RNase-free DNaseI (Fermentas) and then transcribed in a volume of 20 µl with the RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific) into first strand cDNA according to suppliers' instructions. 2 µl of a 1:10 diluted cDNA preparation was mixed with 18 µl master mix as described in the manual of the Maxima SYBR Green/ROX qPCR Master Mix (Thermo Scientific). PCR was performed on a CFX96 Touch Real-Time PCR Detection System (Bio-Rad) using the following conditions: 3 min 95°C; 40 x 15 sec 95°C, 30 sec 59°C, 30 sec 72°C, melting curve from 65°C to 95°C. Gene expression was determined using the delta C_T Method to calculate the fold-induction (compared to dark control samples) or the relative transcription levels (RTL) according to Pfaffl (2001). The data were normalized to PP2A and Actin as reference genes. Each data point is based on three independent biological replicates measured as two technical replicates each. The sequences of the gene-specific primer used in this study are given in supplementary table 2. Statistical analysis was carried out using a 2-Way ANOVA according to Minitab software (MINITAB, 2000).

Comparative transcriptome analysis

For RNA isolation, cells of *Pst* DC3000 were grown in LB with rifampicin (10 µg/ml) and incubated at 25°C for 4-5 hours to an OD₆₆₀=0.5 in the absence or in the presence of sublethal concentrations of phloretin and naringenin [0.2 mM in dimethyl sulphoxide (DMSO)]. Cells were harvested, washed with sarkosyl 0.1% and cell pellets were immediately frozen in liquid nitrogen and stored at -80°C until RNA isolation. RNA was isolated for RNAseq, using TRI ReagentLS (Molecular Research Center, Cincinnati, OH, USA) as described before (Vargas et al., 2011). Residual DNA was removed with the RNase-free DNase I set (Roche). For comparative transcriptome analysis, library generation and sequencing was performed by Macrogen Inc. (Seoul, Republic of Korea) using Illumina HiSeq2000 to produce reads of 100 bp. Reads were clipped and trimmed to remove low quality nucleotides, Illumina adapters and rRNAs by using SeqTrimNext (Falgueras et al., 2010). Filtered reads were aligned to the *Pst* DC3000 reference genome with Bowtie (Langmead et al., 2009). SAMtools (Li et al., 2009) utilities were used to convert sam files to bam format, and to sort, count reads in order to analyse differential expression with RobiNA (Lohse et al., 2009). RNAseq data were deposited in the public database ArrayExpress at EBI under the assigned accession number E-MTAB-6426.

Bacterial preparation

Three strains of *Pseudomonas syringae* pv. *tomato* (*Pst*) were used in this study. Besides the wild-type DC3000, two mutant strains hrpL (transcriptional factor for virulence genes; Castillo-Lizardo et al., 2015) and hrcQ (T3SS-component; Badel et al., 2006) were cultured at 28°C on solid NYGA medium (5g/L bactopectone, 3 g/L yeast extract, 20ml/L glycerol, 15 g/L agar) supplemented with different antibiotics; for DC3000 50mg/l Kanamycin and 50mg/l Rifampicin, for HrpL 100 mg/l Kanamycin and for HrcQ 50mg/l Spectinomycin. For infection assays a single colony was used to inoculate a fresh NYGA plate with the appropriate antibiotic. After 2 days 5 mL of 10mM MgCl₂ were used to float off the bacteria and the suspension was then transferred into a 15 mL tube and centrifuged at 5000 rpm for 5 minutes. The supernatant was discarded and the bacterial pellet re-suspended with 2 mL MgCl₂. The OD600 was determined and set to 0.2 via dilution with 10mM MgCl₂ yielding approximately 1 x 10⁸ colony forming units/mL (cfu/mL). Before spray inoculation Silwet was added to a final concentration of 0,04%.

Arabidopsis thaliana treatment

Some plant-pathogenic fungi require UV light for sporulation or apothecia formation, leading to stronger disease symptoms on various vegetables under sunlight in greenhouse experiments without UV-absorbing (cut-off at 390 nm) vinyl films (Sasaki et al., 1985). On the other hand, UV-B reduced blister blight disease by *Exobasidium vexans* in tea plants, because the fungus is UV-B sensitive (Gunasekera et al., 1997). Thus, it is necessary to consider potential side-effects of UV-B on the pathogen itself. Orth et al. (1990) came to the conclusion that UV-B affected more the host plant than the fungal pathogen, because disease severity was similar in plants that received only UV-B pre-treatment and those that received additional UV-B treatment after the inoculation. This suggests that a pre-treatment with UV-B might be sufficient to observe effects. Since it is known that UV-B also damages *Pseudomonas syringae* (Miller et al., 2001), we decided

to infect UV-B pre-treated *Arabidopsis* plants with *Pst*, assuming that they would have a higher flavonoid content than the untreated control. *Arabidopsis thaliana* Ler-0 (wild-type) and the flavonoid deficient mutant *chs/f3h* (N8585) were grown in pots with a soil/sand mixture under short day conditions (8 h light) for eight weeks. For the UV-B pre-treatment the plants were exposed for three days to six hours attenuated UV-B as described above, while the control plants were shielded additionally by two layers' polyester plastic foil. Before spray inoculation the plants were watered well and kept under a hood to increase the humidity for increased stomata opening. The plants were then sprayed with the bacterial suspensions and 10mM MgCl₂ Solution including 0.04% Silwet as control until all leaves were dripping wet. The inoculated plants were placed back under a hood in the growth chamber (22 °C, short-day conditions) for 24h. Then the hood was removed and the plants were left in the growth chamber until harvest at three days past inoculation (dpi). Samples were collected from two plants (three leaf discs each) for each of the three independent biological replicates, thus the final values represent six plants or 18 leaf discs in total.

Bacterial growth quantification by qPCR

At 3 dpi leaf disc samples were collected to quantify bacterial growth by qPCR. Three leaf discs from two plants of each independent biological replicate were pooled in a 2 ml tube containing ca. 20 zirconium beads, flash-frozen with liquid nitrogen and stored at -80°C for later DNA isolation. DNA extraction was carried out with the GeneJet plant genomic DNA Purification Kit (Thermo scientific). 350 µL of lysis buffer A were added to the frozen samples, which were then ground using the ball-mill as described above. The remaining steps followed according to the kits instructions yielding 100 µL final DNA solution. Subsequently, the DNA concentration was adjusted to 4ng/µL and about 20 ng of DNA were subjected to qPCR analysis. The DNA-based quantification of *Pst* growth in *Arabidopsis* tissue was carried out as described by Ross et al. (2016). Here the PerfeCta SYBR Green Fast Mix (Quantabio) was used and the primer for biomass analysis are given in supplementary table 2. The CFX96 Touch Real-Time PCR Detection System (Bio-Rad) was programmed as followed: 3 min 95°C; 50 x 10 sec 95°C, 10 sec 62°C, 10 sec 72°C, melting curve from 65°C to 95°C for the bacterial biomass determination and 3 min 95°C; 50 x 10 sec 95°C, 20 sec 56°C, 20 sec 72°C, melting curve from 65°C to 95°C for measuring the plant biomass.

Meta-analysis of public expression data

For the meta-analysis with GENEVESTIGATOR (Hruz et al., 2008) in Fig. S2 and S3 only experiments were chosen, which were based on *Arabidopsis* WT plants. Based on their conserved co-regulation six flavonoid pathway genes (set A) were chosen to select 150 perturbation experiments in which these genes were 52 times up- and 98 times down-regulated. Some pathogenesis related experiments were included to address if there is also up-regulation of CHS in response to pathogens in *Arabidopsis* and one elicitor experiment with only weak effects on suppression of this pathway. Three more sets of pathogen defense-related genes were compared to the FPGs. The set B consisted of six known genes for pattern recognition, set C of six genes for defense signalling and set C of six gene involved in production of secondary metabolites for pathogen defense.

Acknowledgements

We would like to thank C. Ramos (Universidad de Málaga, Spain) and E. López-Solanilla (CBPG, Spain) for supplying us the two avirulent *hrpL* and *hrcQ_b-U Pst* mutants, respectively. Hashlin Pascananda Utami was supported by a scholarship from the Indonesian Endowment Fund for Education (LPDP) and Zheng Zhou from the Chinese Scholarship Council (CSC).

Tab. S2 Primer used in this study

Gene	Identifier	forward	reverse	[bp]
Actin2	BnaA10g22340D	TCTACAACGAGCTCCGTGTTG	TGTGAGACACACCATCACCA	216

PP2A	BnaA06g33370D	CAATGACGATGACGATGAGGTG	ATGCTCAACCAAGTCACTCTCC	208
CHS	BnaA03g04590D	TTGTCTGCTCCGAGATCA	CCCTCAAGTGTCATCTATG	219
Actin2	AT3G18780	ACCTTGCTGGACGTGACCTTACTGAT	GTTGTCTCGTGGATTCCAGCAGCTT	298
PP2A	AT1G13320	CAATGACGATGACGATGAGGTG	ATGCTCAACCAAGTCACTCTCC	208
FLS2	AT5G46330	CTGCTCGGATTCTAGGTTTC	GCAAACCTCTGGAGCTAAGTATCC	94
UGT72E1	AT3G50740	TGCGGAGCAGATGATGAACG	CGCCCTCGTAATCACTCCCT	97
Biomass marker genes to measure bacterial growth according to Ross et al., (2016)				
RHIP1	AT4G26410	GAGCTGAAGTGGCTTCCATGAC	GGTCCGACATACCCATGATCC	81
oprF	PSPTO_2299	AACTGAAAAACACCTTGGGC	CCTGGGTTGTTGAAGTGGTA	304

Table legends

Table 1

Flavonoid production or CHS expression suppressed in plants under biotic stress

Table 2

Effects of UV-B light on plant-pathogen interactions

Table S1

Differential gene expression analysis by RNAseq of Pst DC3000 grown in the presence of DMSO and phloretin or naringenin. IDs of upregulated genes encoding efflux pump components are highlighted.

Table S2

Primer used in this study

Figure legends

Figure 1

Central role of the chalcone synthase (CHS) as entry-point for the flavonoid pathway in plants. A knock-out of the single copy gene in Arabidopsis results in plants void of flavonoids. Phloretin is reported to inhibit the T3SS and bacterial flagella of plant-pathogenic Pst DC3000 (Vargas et al., 2013).

Figure 2

Fig. 2 Crosstalk between UV-B and biotic stress mimicked by elicitation with flg22. The CHS expression is suppressed by flg22. These data from Arabidopsis have been published in Zhou et al. (2017). Shown is the response to four different treatments: Water/VIS (C), flg22/VIS (F), UV-B/VIS (U) and both stresses simultaneously (FU). CHS gene expression level was investigated at 4h and normalized to PP2A and Actin. Error bars represent standard error of triplicate experiments, and statistical significance was checked by a two-way ANOVA.

Figure 3

Overview of selected flavonoid family members reported to have antimicrobial properties. The black skeleton structure is typical for each family, while the red substitutions are specific for the given example. For details see text.

Figure 4

Flavonoids dampen bacterial growth in Arabidopsis. Comparison of *Pst* DC3000 (WT), *hrpL* (transcription factor for virulence genes; Castillo-Lizardo et al., 2015) and *hrcQ* (T3SS-component; Badel et al., 2006) growth rate in Arabidopsis Ler-0 (WT) and *chs/f3h* mutant. Growth rates of wild-type (DC3000) and mutant (*hrpL* and *hrcQ*) *Pst* strains were analysed by qPCR of biomass marker genes. Statistical significance was checked by t-test (*: $p \leq 0.05$ **: $p \leq 0.01$) based upon 3 biological replicates.

Figure 5

Effects of UV-B pre-treatment on bacterial growth in Arabidopsis Ler-0 (WT) and *chs/f3h* mutant plants. Growth rates of wild-type (DC3000) and mutant (*hrpL* and *hrcQ*) *Pst* strains were analysed by qPCR of biomass marker genes. Statistical significance was checked by t-test (*: $p \leq 0.05$) based upon 3 biological replicates.

Figure 6

Meta-analysis by GENEVESTIGATOR of the Arabidopsis – *Pst* interaction with respect to suppression of FPGs. The *Pst hrcC* mutant with defective T3SS is non-pathogenic and could resemble MTI, while *Pst* DC3000 is a virulent strain causing disease via ETS. The avirulent *Pst avrRpt2* strain provokes the Effector Triggered Immunity (ETI). Similar treatments were selected and indicate that CHS up-regulation is also time dependent, taking effect after 24h from inoculation.

Figure 7

Expression analysis of FLS2 (A) and UGT72E1 (B) in response to flg22 and UV-B alone or in combination. Gene expression levels were normalized to Actin. Gene expression was investigated after 4h in response to four different treatments: Water/VIS (C), flg22/VIS (F), UV-B/VIS (U) and both stresses simultaneously (FU). Error bars represent standard error of triplicate experiments, and statistical significance was checked by a two-way ANOVA. However, the two-tailed p-value of an unpaired t-test equals 0.0462 for FLS2 when comparing only the C and U treatment.

Figure 8

Summary of different aspects in the interaction between abiotic UV-B and biotic pathogen stress responses. UV-B induces the production of flavonoids via activation of CHS and other FPGs, while pathogens trigger the production of phytoalexins (PAX), lignin or SA via HCT, CRR2 or BA2H, respectively. Reactive Oxygen Species (ROS) can be either produced in response to low or high doses of UV-B or during the oxidative burst in response to infection. Both hypotheses are incorporated in the centre and potential links for suppression of biotic stress responses are indicated by dashed lines, suggesting reciprocal interaction between both abiotic and biotic stress signal transduction pathways.

Figure S1

Crosstalk is functional in *Brassica napus* leaves. The typical CHS expression pattern resembles that of Arabidopsis (Fig. 2) in response to four different treatments: Water/VIS (C), flg22/VIS (F), UV-B/VIS (U) and both stresses simultaneously (FU). CHS gene expression level was investigated at 4h and normalized to PP2A and Actin. Error bars represent standard error of triplicate experiments, and statistical significance was checked by a two-way ANOVA.

Figure S2

Meta-analysis of 140 selected public gene expression experiments involving Arabidopsis treatments by GENEVESTIGATOR (perturbations). A) flavonoid pathway genes (FPGs), B) Pattern Recognition Receptor genes triggering MTI, C) genes involved in defense signalling and D) genes for lignin or phytoalexin production (secondary metabolism).

Figure S3

Meta-analysis of 60 selected public gene expression experiments involving Arabidopsis mutants by GENEVESTIGATOR (perturbations). A) flavonoid pathway genes (FPGs), B) Pattern Recognition Receptor genes triggering MTI, C) genes involved in defense signalling and D) genes for lignin or phytoalexin production (secondary metabolism).

Figure S4

Meta-analysis of chalcone and stilbene synthase family protein expression in Arabidopsis and several other plant species by GENEVESTIGATOR with focus on biotic stress (and light conditions when available). The dots mark CHS orthologous genes as identified by GENEVESTIGATOR. A) Arabidopsis, B) tomato, C) *Physcomitrella patens*, D) barley, E) rice, F) maize, G) soybean, H) *Medicago truncatula* and I) wheat.

720 **References**

- 721 Appelhagen I, Thiedig K, Nordholt N, Schmidt N, Huep G, Sagasser M and Weisshaar B (2014) Update on
722 transparent testa mutants from *Arabidopsis thaliana*: characterisation of new alleles from an isogenic
723 collection. *Planta* 240(5), 955-70. doi: 10.1007/s00425-014-2088-0.
- 724 Badel JL, Shimizu R, Oh HS and Collmer A (2006) A *Pseudomonas syringae* pv. tomato avrE1/hopM1
725 mutant is severely reduced in growth and lesion formation in tomato. *Mol Plant Microbe Interact.* 19(2), 99-
726 111. doi: 10.1094/MPMI -19-0099.
- 727 Ball WT, Alsing J, Mortlock DJ, Staehelin J, Haigh JD, Peter T, Tummon F, Stübi R, Stenke A, Anderson J,
728 Bourassa A, Davis SM, Degenstein D, Frith S, Froidevaux L, Roth C, Sofieva V, Wang R, Wild J, Yu P,
729 Ziemke JR and Rozanov EV (2018) Evidence for a continuous decline in lower stratospheric ozone
730 offsetting ozone layer recovery, *Atmos. Chem. Phys.* 18, 1379-1394.
- 731 Berli FJ, Moreno D, Piccoli P, Hespanhol-Viana L, Silva MF, Bressan-Smith R, Cavagnaro JB and Bottini
732 R (2010) Absciscic acid is involved in the response of grape (*Vitis vinifera* L.) cv. Malbec leaf tissues to
733 ultraviolet-B radiation by enhancing ultraviolet-absorbing compounds, antioxidant enzymes and membrane
734 sterols. *Plant Cell Environ.* 33(1), 1-10. doi: 10.1111/j.1365-3040.2009.02044.x.
- 735 Brown BA and Jenkins GI (2008) UV-B Signaling Pathways with Different Fluence-Rate Response Profiles
736 Are Distinguished in Mature *Arabidopsis* Leaf Tissue by Requirement for UVR8, HY5, and HYH. *Plant*
737 *Physiol.* 146(2), 576-588. doi: 10.1104/pp.107.108456.
- 738 Bruzzone S, Basile G, Mannino E, Sturla L, Magnone M, Grozio A, Salis A, Fresia C, Vigliarolo T, Guida L,
739 De Flora A, Tossi V, Cassia R, Lamattina L and Zocchi E (2012) Autocrine abscisic acid mediates the UV-
740 B-induced inflammatory response in human granulocytes and keratinocytes. *J Cell Physiol.* 227(6), 2502-
741 10. doi: 10.1002/jcp.22987.
- 742 Cao FY, Yoshioka K and Desveaux D (2011) The roles of ABA in plant-pathogen interactions. *J Plant Res.*
743 124(4), 489-99. doi: 10.1007/s10265-011-0409-y.
- 744 Castillo-Lizardo MG, Aragón IM, Carvajal V, Matas IM, Pérez-Bueno ML, Gallegos MT, Barón M and Ramos
745 C (2015) Contribution of the non-effector members of the HrpL regulon, *iaaL* and *matE*, to the virulence of
746 *Pseudomonas syringae* pv. tomato DC3000 in tomato plants. *BMC Microbiol.* 15:165. doi: 10.1186/s12866-
747 015-0503-8.
- 748 Cushnie TP and Lamb AJ (2005) Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 26(5), 343-56.
- 749 de Torres-Zabala M, Truman W, Bennett MH, Lafforgue G, Mansfield JW, Rodriguez Egea P, Bögre L and
750 Grant M (2007) *Pseudomonas syringae* pv. tomato hijacks the *Arabidopsis* abscisic acid signalling pathway
751 to cause disease. *EMBO J.* 26(5), 1434-43.
- 752 de Torres Zabala M, Bennett MH, Truman WH and Grant MR (2009) Antagonism between salicylic and
753 abscisic acid reflects early host-pathogen conflict and moulds plant defence responses. *Plant J.* 59(3), 375-
754 86. doi: 10.1111/j.1365-313X.2009.03875.x.
- 755 Demkura PV and Ballaré CL (2012) UVR8 mediates UV-B-induced *Arabidopsis* defense responses against
756 *Botrytis cinerea* by controlling sinapate accumulation. *Mol Plant.* 5(3), 642-52. doi: 10.1093/mp/sss025.
- 757 Dixon RA and Pasinetti GM (2010) Flavonoids and isoflavonoids: from plant biology to agriculture and
758 neuroscience. *Plant Physiol.* 154(2), 453-57. doi: 10.1104/pp.110.161430.
- 759 Du Y, Chu H, Wang M, Chu IK and Lo C (2010) Identification of flavone phytoalexins and a pathogen-
760 inducible flavone synthase II gene (*SbFNSII*) in sorghum. *J Exp Bot.* 61(4), 983-94. doi: 10.1093/jxb/erp364.
- 761 El Hadrami A, Adam LR and Daayf F (2011) Biocontrol treatments confer protection against *Verticillium*
762 *dahliae* infection of potato by inducing antimicrobial metabolites. *Mol Plant Microbe Interact.* 24(3), 328-35.
763 doi: 10.1094/MPMI-04-10-0098.

764 El Hadrami A, Islam MR, Adam LR and Daayf F (2015) A cupin domain-containing protein with a
765 quercetinase activity (VdQase) regulates *Verticillium dahliae*'s pathogenicity and contributes to
766 counteracting host defenses. *Front Plant Sci.* 6:440. doi: 10.3389/fpls.2015.00440.

767 Falcone Ferreyra ML, Rius SP and Casati P (2012) Flavonoids: biosynthesis, biological functions, and
768 biotechnological applications. *Front Plant Sci* 3:222. doi: 10.3389/fpls.2012.00222.

769
770 Falcone Ferreyra ML, Emiliani J, Rodriguez EJ, Campos-Bermudez VA, Grotewold E and Casati P (2015)
771 The Identification of Maize and Arabidopsis Type I FLAVONE SYNTHASEs Links Flavones with Hormones
772 and Biotic Interactions. *Plant Physiol.* 169(2), 1090-107. doi: 10.1104/pp.15.00515.

773 Falgueras, J., Lara, A.J., Fernández-Pozo, N., Cantón, F.R., Pérez-Trabado, G., Claros, M.G. 2010.
774 SeqTrim: a high-throughput pipeline for pre-processing any type of sequence read. *BMC Bioinformatics.*
775 11:38. doi: 10.1186/1471-2105-11-38.

776 Gleitz J, Schnitzler JP, Steimle D and Seitz HU (1991) Metabolic changes in carrot cells in response to
777 simultaneous treatment with ultraviolet light and a fungal elicitor. *Planta.* 184(3), 362-7. doi:
778 10.1007/BF00195338.

779 Gosch C, Halbwirth H and Stich K (2010) Phloridzin: biosynthesis, distribution and physiological relevance
780 in plants. *Phytochemistry* 71(8-9), 838-43. doi: 10.1016/j.phytochem.2010.03.003.

781 Gunasekera TS, Paul ND and Ayres PG (1997) The effects of ultraviolet-B (UV-B: 290–320 nm) radiation
782 on blister blight disease of tea (*Camellia sinensis*). *Plant Pathology* 46(2), 179-185. doi: 10.1046/j.1365-
783 3059.1997.d01-216.x.

784 Hahlbrock K, Lamb CJ, Purwin C, Ebel J, Fautz E and Schäfer E (1981) Rapid Response of Suspension-
785 cultured Parsley Cells to the Elicitor from *Phytophthora megasperma* var. *sojae*: INDUCTION OF THE
786 ENZYMES OF GENERAL PHENYLPROPANOID METABOLISM. *Plant Physiol.* 67(4), 768-73.

787 Hassan S and Mathesius U (2012) The role of flavonoids in root-rhizosphere signalling: opportunities and
788 challenges for improving plant-microbe interactions. *J Exp Bot.* 63(9), 3429-44. doi: 10.1093/jxb/err430.

789 Hideg E, Jansen MA and Strid A (2013) UV-B exposure, ROS, and stress: inseparable companions or
790 loosely linked associates? *Trends Plant Sci.* 18(2), 107-15. doi: 10.1016/j.tplants.2012.09.003.

791 Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W and Zimmermann
792 P (2008) Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. *Adv*
793 *Bioinformatics* 2008:420747. doi: 10.1155/2008/420747.

794 Ishikawa A, Kitamura Y, Ozeki Y, Itoh Y, Yamada A and Watanabe M (2005) Post-stress metabolism
795 involves umbelliferone production in anthocyanin-producing and nonproducing cells of *Glehnia littoralis*
796 suspension cultures. *J Plant Physiol.* 162(6), 703-10.

797 Jansen et al. (THIS ISSUE) Interactive effects of UV-B radiation in a complex environment

798 Jeandet P, Delaunois B, Conreux A, Donnez D, Nuzzo V, Cordelier S, Clément C and Courot E (2010)
799 Biosynthesis, metabolism, molecular engineering, and biological functions of stilbene phytoalexins in plants.
800 *Biofactors.* 36(5), 331-41. doi: 10.1002/biof.108.

801 Jenkins GI (2014) The UV-B Photoreceptor UVR8: From Structure to Physiology. *Plant Cell.* 26(1), 21-37.
802 doi: 10.1105/tpc.113.119446.

803 Jia Z, Zou B, Wang X, Qiu J, Ma H, Gou Z, Song S and Dong H (2010) Quercetin-induced H₂O₂
804 mediates the pathogen resistance against *Pseudomonas syringae* pv. *Tomato* DC3000 in *Arabidopsis*
805 *thaliana*. *Biochem Biophys Res Commun.* 396(2), 522-7. doi: 10.1016/j.bbrc.2010.04.131.

806 Jones JD and Dangl JL (2006) The plant immune system. *Nature.* 444(7117), 323-329.

807 Jones JT, Furlanetto C and Phillips MS (2007) The role of flavonoids produced in response to cyst
808 nematode infection of *Arabidopsis thaliana*. *Nematology* 9(5), 671-677. doi: 10.1163/156854107782024875

809 Katsumata S, Hamana K, Horie K, Toshima H and Hasegawa M (2017) Identification of Sternbin and
 810 Naringenin as Detoxified Metabolites from the Rice Flavanone Phytoalexin Sakuranetin by *Pyricularia*
 811 *oryzae*. *Chem Biodivers.* 14(2). doi: 10.1002/cbdv.201600240.

812 Kodama S, Miyakawa J, Akatsuka T and Kiyosawa S (1992) Sakuranetin, a flavanone phytoalexin from
 813 ultraviolet-irradiated rice leaves. *Phytochemistry* 31(11), 3807–3809.

814 König S, Feussner K, Kaever A, Landesfeind M, Thurow C, Karlovsky P, Gatz C, Polle A and Feussner I
 815 (2014) Soluble phenylpropanoids are involved in the defense response of *Arabidopsis* against *Verticillium*
 816 *longisporum*. *New Phytol.* 202(3), 823-837. doi: 10.1111/nph.12709.

817 Kunz BA, Dando PK, Grice DM, Mohr PG, Schenk PM and Cahill DM (2008) UV-induced DNA damage
 818 promotes resistance to the biotrophic pathogen *Hyaloperonospora parasitica* in *Arabidopsis*. *Plant Physiol.*
 819 148(2), 1021-1031. doi: 10.1104/pp.108.125435.

820 Kuspradini H, Mitsunaga T, Ohashi H (2009) Antimicrobial activity against *Streptococcus sobrinus* and
 821 glucosyltransferase inhibitory activity of taxifolin and some flavanonol rhamnosides from kempas
 822 (*Koompassia malaccensis*) extracts. *J Wood Sci.* 55(4), 308-313.

823 Lagrange H, Jay-Allmand C and Lapeyrie F (2001) Rutin, the phenolglycoside from eucalyptus root
 824 exudates, stimulates *Pisolithus hyphal* growth at picomolar concentrations. *New Phytol.* 149(2), 349-355.
 825 doi: 10.1046/j.1469-8137.2001.00027.x.

826 Landry LG, Chapple CC and Last RL (1995) *Arabidopsis* mutants lacking phenolic sunscreens exhibit
 827 enhanced ultraviolet-B injury and oxidative damage. *Plant Physiol.* 109(4), 1159-66.

828 Langmead B, Trapnell C, Pop M and Salzberg SL (2009) Ultrafast and memory-efficient alignment of short
 829 DNA sequences to the human genome. *Genome Biol* 10(3), R25. doi: 10.1186/gb-2009-10-3-r25.

830 Lanot A, Hodge D, Jackson RG, George GL, Elias L, Lim EK, Vaistij FE and Bowles DJ (2006) The
 831 glucosyltransferase UGT72E2 is responsible for monolignol 4-O-glucoside production in *Arabidopsis*
 832 *thaliana*. *Plant J.* 48(2), 286-95.

833 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G and Durbin R (2009)
 834 The sequence alignment/map format and SAMtools. *Bioinformatics* 25(16), 2078-2079. doi:
 835 10.1093/bioinformatics/btp352.

836 Li J, Ou-Lee TM, Raba R, Amundson RG and Last RL (1993) *Arabidopsis* Flavonoid Mutants Are
 837 Hypersensitive to UV-B Irradiation. *Plant Cell.* 5(2), 171-179.

838 Liu CJ, Deavours BE, Richard SB, Ferrer JL, Blount JW, Huhman D, Dixon RA and Noel JP (2006)
 839 Structural basis for dual functionality of isoflavonoid O-methyltransferases in the evolution of plant defense
 840 responses. *Plant Cell.* 18(12), 3656-69.

841 Lo SC and Nicholson RL (1998) Reduction of light-induced anthocyanin accumulation in inoculated
 842 sorghum mesocotyls. Implications for a compensatory role in the defense response. *Plant Physiol.* 116(3),
 843 979-89.

844 Lohse M, Bolger AM, Nagel A, Fernie AR, Lunn JE, Stitt M and Usadel B (2012) RobiNA: a user-friendly,
 845 integrated software solution for RNA-Seq-based transcriptomics. *Nucleic Acids Res.* W622-7. doi:
 846 10.1093/nar/gks540.

847 López-Gresa MP, Torres C, Campos L, Lisón P, Rodrigo I, Bellés JM and Conejero V (2011) Identification
 848 of defence metabolites in tomato plants infected by the bacterial pathogen *Pseudomonas syringae*. *Environ.*
 849 *Exp. Bot.* 74(1), 216-228. doi: 10.1016/j.envexpbot.2011.06.003.

850 Lozoya E, Block A, Lois R, Hahlbrock K and Scheel D (1991) Transcriptional repression of light-induced
 851 flavanoid synthesis by elicitor treatment of cultured parsley cells. *Plant J.* 1(2), 227-234.

852 Mackerness SAH, Surplus SL, Blake P, John CF, Buchanan-Wollaston V, Jordan BR and Thomas B (1999)
853 Ultraviolet-B-induced stress and changes in gene expression in *Arabidopsis thaliana*: role of signalling
854 pathways controlled by jasmonic acid, ethylene and reactive oxygen species. *Plant Cell Environ.* 22(11),
855 1413-1423.

856 Malathi P, Viswanathan R, Padmanaban P, Mohanraj D, Ganesh Kumar V and Salin KP (2008) Differential
857 accumulation of 3-deoxy anthocyanidin phytoalexins in sugarcane varieties varying in red rot resistance in
858 response to *Colletotrichum falcatum* infection. *Sugar Tech* 10(2), 154-157.

859 Manning WJ and v. Tiedemann A (1995) Climate Change: Effects of increased atmospheric carbon dioxide
860 (CO₂), ozone (O₃), and ultraviolet-B (UV-B) Radiation on plant diseases. *Environmental Pollution*, 88(2),
861 219-245.

862 McLusky SR, Bennett MH, Beale MH, Lewis MJ, Gaskin P and Mansfield JW (1999) Cell wall alterations
863 and localized accumulation of feruloyl-3'-methoxytyramine in onion epidermis at sites of attempted
864 penetration by *Botrytis allii* are associated with actin polarisation, peroxidase activity and suppression of
865 flavonoid biosynthesis. *Plant J.* 17(5), 523-534.

866 Miller CD, Mortensen WS, Braga GU and Anderson AJ (2001) The *rpoS* gene in *Pseudomonas syringae* is
867 important in surviving exposure to the near-UV in sunlight. *Curr Microbiol.* 43(5), 374-377.

868 Misra P, Pandey A, Tiwari M, Chandrashekar K, Sidhu OP, Asif MH, Chakrabarty D, Singh PK, Trivedi PK,
869 Nath P and Tuli R (2010) Modulation of transcriptome and metabolome of tobacco by *Arabidopsis*
870 transcription factor, *AtMYB12*, leads to insect resistance. *Plant Physiol.* 152(4), 2258-68. doi:
871 10.1104/pp.109.150979.

872 Orth AB, Teramura AH and Sisler HD (1990) Effects of Ultraviolet-B Radiation on Fungal Disease
873 Development in *Cucumis sativus*. *American Journal of Botany* 77(9), 1188-1192.

874 Panagopoulos I, Bornman JF and Björn LO (1992) Response of sugar beet plants to ultraviolet-B (280-320
875 nm) radiation and *Cercospora* leaf spot disease. *Physiol. Plantarum.* 84(1), 140-145. doi: 10.1111/j.1399-
876 3054.1992.tb08776.x

877 Pejin B, Ciric A, Markovic JD, Glamoclija J, Nikolic M, Stanimirovic B and Sokovic M (2015) Quercetin
878 Potently Reduces Biofilm Formation of the Strain *Pseudomonas aeruginosa* PAO1 in vitro. *Curr Pharm*
879 *Biotechnol.* 16(8), 733-737.

880 Rakitin VY, Karyagin VV, Rakitina TY, Prudnikova ON, and Vlasov PV (2008) UV-B Stress-Induced ABA
881 Production in *Arabidopsis thaliana* Mutants Defective in Ethylene Signal Transduction Pathway. *Russ J*
882 *Plant Physiol* 55(6), 854-856. doi: 10.1134/S1021443708060174

883 Ross A and Somssich IE (2016) A DNA-based real-time PCR assay for robust growth quantification of the
884 bacterial pathogen *Pseudomonas syringae* on *Arabidopsis thaliana*. *Plant Methods.* 12, 48.

885 Saijo Y, Tintor N, Lu X, Rauf P, Pajerowska-Mukhtar K, Häweker H, Dong X, Robatzek S and Schulze-
886 Lefert P (2009) Receptor quality control in the endoplasmic reticulum for plant innate immunity. *EMBO J.*
887 28(21), 3439-3449.

888 Samac DA and Graham MA (2007) Recent advances in legume-microbe interactions: recognition, defense
889 response, and symbiosis from a genomic perspective. *Plant Physiol.* 144(2), 582-587.

890 Sasaki T, Honda Y, Umekawa M and Nemoto M (1985) Control of certain diseases of greenhouse
891 vegetables with ultraviolet-absorbing vinyl film. *Plant Dis.* 69, 530-533.

892 Schenke D, Böttcher C and Scheel D (2011) Crosstalk between abiotic UV-B stress and biotic (*flg22*) stress
893 signaling in *Arabidopsis* prevents flavonol accumulation in favor of pathogen defense compound production.
894 *Plant, Cell Environ.* 34(11), 1849-1864. doi: 10.1111/j.1365-3040.2011.02381.x.

895 Schenke D, Cai D and Scheel D (2014) Suppression of UV-B stress responses by *flg22* is regulated at the
896 chromatin level via histone modification. *Plant, Cell Environ.* 37(7), 1716-1721. doi: 10.1111/pce.12283.

897 Schröder J and Schröder G (1990) Stilbene and chalcone synthases: related enzymes with key functions
898 in plant-specific pathways. *Z Naturforsch C*. 45(1-2), 1-8.

899 Serrano M, Kanehara K, Torres M, Yamada K, Tintor N, Kombrink E, Schulze-Lefert P and Saijo Y (2012)
900 Repression of sucrose/ultraviolet B light-induced flavonoid accumulation in microbe-associated molecular
901 pattern-triggered immunity in *Arabidopsis*. *Plant Physiol*. 158(1), 408-422. doi: 10.1104/pp.111.183459.

902 Singh A, Gupta R and Pandey R (2016) Rice Seed Priming with Picomolar Rutin Enhances Rhizospheric
903 *Bacillus subtilis* CIM Colonization and Plant Growth. *PLoS One*. 11(1):e0146013. doi:
904 10.1371/journal.pone.0146013.

905 Steinkellner S, Lenzemo V, Langer I, Schweiger P, Khaosaad T, Toussaint JP and Vierheilig H (2007)
906 Flavonoids and strigolactones in root exudates as signals in symbiotic and pathogenic plant-fungus
907 interactions. *Molecules*. 12(7), 1290-306.

908 Tanaka S, Brefort T, Neidig N, Djamei A, Kahnt J, Vermerris W, Koenig S, Feussner K, Feussner I and
909 Kahmann R (2014) A secreted *Ustilago maydis* effector promotes virulence by targeting anthocyanin
910 biosynthesis in maize. *Elife*. 3, e01355. doi: 10.7554/eLife.01355.

911 Tsuda K, Sato M, Stoddard T, Glazebrook J and Katagiri F (2009) Network properties of robust immunity
912 in plants. *PLoS Genet*. 5(12), e1000772. doi: 10.1371/journal.pgen.1000772.

913 Ullah C, Unsicker SB, Fellenberg C, Constabel CP, Schmidt A, Gershenzon J and Hammerbacher A (2017)
914 Flavan-3-ols Are an Effective Chemical Defense against Rust Infection. *Plant Physiol*. 175(4), 1560-1578.
915 doi: 10.1104/pp.17.00842.

916 van der Watt E and Pretorius JC (2001) Purification and identification of active antibacterial components in
917 *Carpobrotus edulis* L. *J Ethnopharmacol*. 76(1), 87-91.

918 Vandeputte OM, Kiendrebeogo M, Rajaonson S, Diallo B, Mol A, El Jaziri M and Baucher M (2010)
919 Identification of catechin as one of the flavonoids from *Combretum albiflorum* bark extract that reduces the
920 production of quorum-sensing-controlled virulence factors in *Pseudomonas aeruginosa* PAO1. *Appl*
921 *Environ Microbiol*. 76(1), 243-53. doi: 10.1128/AEM.01059-09.

922 Vargas P, Felipe A, Michán C and Gallegos MT (2011) Induction of *Pseudomonas syringae* pv. tomato
923 DC3000 MexAB-OprM multidrug efflux pump by flavonoids is mediated by the repressor PmeR. *Mol Plant*
924 *Microbe Interact*. 24(10), 1207-19. doi: 10.1094/MPMI-03-11-0077.

925 Vargas P, Farias GA, Nogales J, Prada H, Carvajal V, Barón M, Rivilla R, Martín M, Olmedilla A and
926 Gallegos MT (2013) Plant flavonoids target *Pseudomonas syringae* pv. tomato DC3000 flagella and type
927 III secretion system. *Environ Microbiol Rep*. 5(6), 841-50. doi: 10.1111/1758-2229.12086.

928 Wasson AP, Pellerone FI and Mathesius U (2006) Silencing the flavonoid pathway in *Medicago truncatula*
929 inhibits root nodule formation and prevents auxin transport regulation by rhizobia. *Plant Cell* 18(7), 1617-
930 29.

931 Yamamoto M, Nakatsuka S, Otani H, Kohmoto K and Nishimura S (2000) (+)-catechin acts as an infection-
932 inhibiting factor in strawberry leaf. *Phytopathology* 90(6), 595-600. doi: 10.1094/PHYTO.2000.90.6.595.

933 Yang W, Xu X, Li Y, Wang Y, Li M, Wang Y, Ding X and Chu Z (2016) Rutin-Mediated Priming of Plant
934 Resistance to Three Bacterial Pathogens Initiating the Early SA Signal Pathway. *PLoS One*. 11(1),
935 e0146910. doi: 10.1371/journal.pone.

936 Yasuda M, Ishikawa A, Jikumaru Y, Seki M, Umezawa T, Asami T, Maruyama-Nakashita A, Kudo T,
937 Shinozaki K, Yoshida S and Nakashita H (2008) Antagonistic interaction between systemic acquired
938 resistance and the abscisic acid-mediated abiotic stress response in *Arabidopsis*. *Plant Cell* 20(6), 1678-92.
939 doi: 10.1105/tpc.107.054296.

940
941 Yu O, Shi J, Hession AO, Maxwell CA, McGonigle B and Odell JT (2003) Metabolic engineering to increase
942 isoflavone biosynthesis in soybean seed. *Phytochemistry*. 63(7), 753-763.

- 943 Zhou H, Lin J, Johnson A, Morgan RL, Zhong W and Ma W (2011) *Pseudomonas syringae* type III effector
944 HopZ1 targets a host enzyme to suppress isoflavone biosynthesis and promote infection in soybean. *Cell*
945 *Host & Microbe* 9(3), 177-186. doi: 10.1016/j.chom.2011.02.007.
- 946 Zhou Z, Schenke D, Miao Y and Cai D (2017) Investigation of the crosstalk between the flg22 and the UV-
947 B-induced flavonol pathway in *Arabidopsis thaliana* seedlings. *Plant Cell Environ.* 40(3), 453-458. doi:
948 10.1111/pce.12869.