**Ontogenetic changes in the feeding functional response of the marine
copepod *Paracartia grani***

Manuel Olivares*, Enric Saiz, Albert Calbet

Instituto de Ciencias del Mar (ICM, CSIC), Passeig Marítim de la Barceloneta
37-49, E-08003, Barcelona, Spain

*Corresponding author: molivares@icm.csic.es

Running page head: Feeding patterns through ontogeny of copepods

Key words: zooplankton, *Acartia*, nauplii, development, grazing, ingestion,
clearance rate, functional trait

Abstract

The feeding activity of copepods is crucial for the structuring and functioning of marine ecosystems. The quantification of feeding rates of different copepod life stages across a range of prey densities (functional response) is essential knowledge for the improvement of plankton dynamic models. In this study, we conducted experiments to compare the feeding functional responses of nauplii, copepodites, and adults of the marine copepod *Paracartia grani* (formerly *Acartia grani*) when grazing on the flagellate *Rhodomonas salina*. We found that all the copepod stages followed a sigmoid curve in their functional responses (Holling type III model), indicating a metabolic threshold constraining the foraging effort at low prey densities. The maximum clearance rates of nauplii and copepodites increased with body mass with a power scaling factor of 1, but the maximum clearance rates of adults did not follow the pattern observed for juvenile stages, likely because of the relatively small prey size used in the experiments. Copepod maximum ingestion rates, however, showed allometric scaling along ontogeny that was closer to the typical 0.75 power law and seemed to not be so dependent on prey size.

1. Introduction

Zooplankton constitute the main link between primary producers (phytoplankton) and higher trophic levels in marine food webs (Calbet 2001, Calbet & Saiz 2005). They are a key component of the biological pump and have a major role in nutrient recycling in the photic layer (Steinberg et al. 2000, Alcaraz et al. 2010, Turner 2015). Within the marine mesozooplankton, copepods typically account for up to 35-70% of the biomass and represent more than 90% of the abundance (Longhurst 1985). Consequently, the resource exploitation conducted by copepods is crucial for the structuring and functioning of marine ecosystems.

Functional responses of copepods are defined as the relationship between their vital rates (e.g., feeding rates, egg production rates) and prey availability (Solomon 1949, Holling 1966). Because of the major role of copepods in the zooplankton, copepod functional responses have great implications for plankton population dynamics (Runge 1988, Gentleman & Neuheimer 2008, Kiørboe et al. 2018). The quantification of functional responses in both field and laboratory studies enables the development of mathematical models to forecast the response of plankton communities to changing species densities (Morozov 2010, Morozov et al. 2012). These predictive models are intrinsically very sensitive to the parameterization of functional responses (e.g., the shape of the functional response), and, for this reason, the proper analysis of copepod functional responses becomes an imperative task to improve our capability to assess potential variations in the future composition of plankton communities and the biogeochemical fluxes that occur through them (Gentleman & Neuheimer 2008, Wollrab & Diehl 2015, Egilmez & Morozov 2016).

Traditionally, the study of feeding activity patterns in copepods has been limited mostly to adults, particularly females. In this respect, feeding functional responses have been well documented for adults of key marine copepod species using a wide variety of prey types, and the records are much more numerous for females than for males (Støttrup & Jensen 1990, Isari & Saiz 2011, Gonçalves et al. 2014, Saiz et al. 2014). Females have been preferred over males in grazing experiments mainly because of their longer life span, higher abundances in natural populations, and fundamental role in recruitment (Kouwenberg 1993, Kiørboe 2006, Kiørboe et al. 2015). However, the evidence of distinct feeding behaviours between the sexes (e.g., lower feeding rates in males) demonstrates that gender differences should be taken into account when analysing feeding activity patterns in marine copepods (Lampitt 1978, Uye & Kayano 1994, Saage et al. 2009, Chen et al. 2010, van Someren Gréve et al. 2017b).

Similar to males, juvenile copepods (i.e., nauplii and copepodites) have been greatly neglected in previous research addressing copepod feeding patterns. Juvenile copepods outnumber adults throughout the year in plankton communities (Webber & Roff 1995, Calbet et al. 2001, Turner 2004). They play a key ecological role as major grazers of phytoplankton and small heterotrophs. Therefore, the study of their feeding activity is fundamental to evaluating the grazing pressure on the smaller plankton fraction in marine ecosystems (Lučić et al. 2003, Turner 2004, Almeda et al. 2011). The first investigations that focused attention on the feeding patterns of the early stages of copepods focused on their rates of grazing on phytoplankton (Mullin & Brooks 1967, Paffenhöfer 1971, Fernández 1979). In response to the increasing interest in

this topic, further investigations were carried out on other aspects, such as their foraging behaviour, prey size spectrum, and prey selection (Berggreen et al. 1988, Paffenhöfer 1998, Henriksen et al. 2007, Helenius & Saiz 2017). In terms of feeding functional responses, Fernández (1979) was the first author to provide empirical data on naupliar grazing rates of *Calanus pacificus* across a wide range of prey types and food concentrations. Since then, feeding functional responses have been described for the juvenile copepod stages of other genera, such as *Acartia/Paracartia* (Henriksen et al. 2007, Ismar et al. 2008, Helenius & Saiz 2017), *Oithona* (Henriksen et al. 2007, Almeda et al. 2010, Saiz et al. 2014), and *Centropages* (López et al. 2007).

Despite this valuable information, we think that the current literature on marine copepods has not completely addressed the changes in feeding functional response through the entire copepod life cycle. For instance, Almeda et al. (2010) analysed feeding functional responses up to early copepodites, and Saiz et al. (2014) worked with nauplii and adults but not with copepodites. Berggreen et al. (1988) reported the changes in the prey size spectrum of *Acartia tonsa* from nauplii to adults, but in their study, the feeding functional responses were not analysed. Thus, it appears that we still lack a complete record regarding how feeding functional patterns change through ontogeny in marine copepods. To the best of our knowledge, only Santer and van den Bosch (1994) have studied this issue, with the freshwater species *Cyclops vicinus*.

Thus, the main goal of this study was to assess the ontogenetic changes in the feeding functional response of marine copepods. To accomplish our objective, we conducted grazing experiments in the laboratory with different development stages of the calanoid copepod *Paracartia grani* (formerly *Acartia grani*) and a

broad range of prey concentrations of the flagellate *Rhodomonas salina*. *P. grani* is a common neritic species that has been observed from high to middle latitudes in the eastern Atlantic and across the Mediterranean Sea (Boyer et al. 2012). In our study, we compared feeding functional patterns among different *P. grani* life stages (nauplii, copepodites, adults) and sexes (males and females). We expected that the values of the fundamental parameters defining the feeding functional response (maximum clearance and ingestion rates, half-saturation constants) would be influenced by changes along ontogeny in the allometric scaling of metabolic requirements (Ikeda et al. 2001, Saiz & Calbet 2007) and variations in morphological and behavioural traits (Paffenhöfer 1998).

2. Materials & methods

2. 1. Experimental cultures

For the experiments, the cryptophyte *Rhodomonas salina* was used as prey [equivalent spherical diameter (ESD): 6.9 μm , C content: 38 pg cell⁻¹, N content: 7.2 pg cell⁻¹; Helenius and Saiz (2017)]. A culture of *R. salina* was grown in f/2 medium (Guillard 1983) and kept in the exponential growth phase by diluting 1/3 of the culture daily. For the copepods, a cohort of *Paracartia grani* was obtained from a stock culture maintained at the Instituto de Ciencias del Mar (ICM, CSIC) for more than 10 years (Saiz et al. 2015). To produce the cohort, eggs were siphoned out from the bottom of the copepod stock culture and inoculated into 20-L polycarbonate tanks filled with 0.1 μm filtered seawater. Approximately 24 hours after the egg addition, unhatched eggs were removed from the bottom with a siphon to ensure the uniformity of the cohort. The abundance of copepods in the cohort tank started with ca. 15,000 nauplii, and decreased as

copepods were taken out for experiments (ca. 1,000 adults by the end). The cohort was fed *ad libitum* at concentrations ranging from 7 (nauplii) to 20 (adults) ppm of *R. salina*. *R. salina* is a nutritionally sufficient prey that has been commonly used in the maintenance of *Acartia* cultures (Støttrup & Jensen 1990, Broglio et al. 2003, Saiz et al. 2015). Both the phytoplankton and the copepod cultures used for the experiments were kept in the laboratory in a cold room at 19±1°C and under a 10 h:14 h L:D photoperiod.

2. 2. Functional response experiments

Six feeding functional response experiments were carried out successively using different copepod life stages from the same *P. grani* cohort, encompassing two experiments conducted with nauplii, two experiments with copepodites, and two experiments with adults (one with males and one with females). Prior to each experiment, copepods were collected from the cohort using a 60-µm (nauplii and copepodites) or a 200-µm (adults) mesh and placed in filtered seawater for 2.5-3 hours while setting up the experiment. For each functional response experiment, six prey concentrations of exponentially growing *R. salina* were prepared. The range of prey concentrations was based on previous knowledge (Calbet & Alcaraz 1997, Helenius & Saiz 2017) to cover limiting to satiating food conditions (500-12,000 cells mL⁻¹ for NII-III, 500-14,000 cells mL⁻¹ for NV-VI, 500-16,000 cells mL⁻¹ for CI-II, 1,000-20,000 cells mL⁻¹ for CIII-IV, and 1,000-24,000 cells mL⁻¹ for adults). The prey concentrations were measured with a Beckman Coulter Multisizer III particle counter. Eight bottles were filled with each prey concentration: two served as initial bottles (only prey), three as control bottles (only prey), and three as experimental bottles (prey and

copepods). Bottle volumes ranged from 72 mL for early nauplii to 620 mL for adults. All the bottles were filled following a three-step procedure (one-third of the bottle was sequentially filled each time) to ensure the homogeneity of prey densities among the bottles. Inorganic nutrients (5 mL L^{-1} of f/2 solution, i.e., a final concentration equivalent to f/400) were also added to each bottle to compensate for any differential effect of copepod nutrient excretion on algal growth. Once the bottles were filled with the prey suspensions, the copepods were added to the bottles either as aliquots (nauplii and copepodites) or individually with a pipette (adults). For the aliquots, the concentration of nauplii/copepodites in the filtered seawater was estimated on the basis of replicated counts of subsamples drawn with an automatic pipette, and then the aliquot volume of the copepod suspension required to reach the desired copepod concentration was added to each experimental bottle. To compensate for dilution in the experimental bottles, the same volume of filtered seawater was added to the initial and control bottles. Copepod adults were transferred to suspensions of the respective prey concentration to avoid any dilution effects, and then they were individually removed and transferred to the experimental bottles. Given that different copepod stages (sizes) and different prey concentrations were tested, the number of copepods added to each bottle was adjusted based on unpublished previous data to result in a reduction of ca. 30% of the prey concentration during incubation. Thus, the number of individuals incubated in the bottles ranged from low copepod abundance to high copepod abundance across the different prey concentrations (80 to 240 NII-III, 40 to 160 NV-VI, 40 to 220 CI-II, 25 to 90 CIII-IV, 10 to 35 adult males, and 10 to 30 adult females). All bottles were sealed with plastic film to avoid air bubbles and

capped. The initial bottles were sacrificed at the very beginning of the incubation periods to assess the actual initial prey concentrations using the Coulter counter. Some extra copepods were preserved in 4% formaldehyde for the measurement of initial copepod size. The control and experimental bottles were mounted on a rotating plankton wheel (0.2 rpm) and incubated for 20-24 hours at $19\pm 1^{\circ}\text{C}$ and under a 10 h:14 h L:D cycle ($\text{PAR } 15\text{-}34 \mu\text{mol m}^{-2} \text{s}^{-1}$). After the incubations, the suspensions from the control and experimental bottles were filtered through a submerged sieve with a 60- μm (nauplii and copepodites) or a 200- μm (adults) mesh to separate the copepods from the prey. The final prey concentrations in each bottle were measured, as they were for the initial bottles, with the Coulter counter. The experimental copepods were checked for activity under a stereomicroscope and then fixed with formaldehyde (4% final concentration) for quantification, stage determination, and final body size measurements. Prey sizes were calculated as the geometric mean between initial and final prey volumes measured with the particle counter. The initial and final copepod sizes were determined by photographing 30-40 nauplii or copepodites and 20-25 adults with an inverted microscope and measuring their body (nauplii) or prosome (copepodites, adults) lengths with the ImageJ software (Schneider et al. 2012). Copepod sizes were calculated for each prey concentration as the arithmetic mean between their initial and final mean lengths.

2. 3. Data analysis

Average prey concentrations and copepod feeding rates (clearance and ingestion rates) were determined according to equations in Frost (1972).

Holling type III curves were fitted to the copepod feeding rates as a function of the prey concentration using the software package SigmaPlot version 14.0. The curve fitting was carried out by non-linear regression following the equations in Helenius and Saiz (2017):

$$F = \frac{I_{\max} C}{C^2 + K_m^2} \quad (1)$$

and

$$I = \frac{I_{\max} C^2}{C^2 + K_m^2} \quad (2)$$

where F is the clearance rate, I is the ingestion rate, $I_{\max}$ is the maximum ingestion rate, C is the prey concentration, and K_m is the half-saturation constant. Estimates of the maximum ingestion rate ($I_{\max}$), half-saturation constant (K_m) and prey concentration at which 90% of the maximum ingestion rate was reached (C_{90}) were obtained directly from the Holling type III model. Maximum clearance rates ($F_{\max}$) were calculated as described in Helenius and Saiz (2017):

$$F_{\max} = \frac{I_{\max}}{2K_m} \quad (3)$$

where $F_{\max}$ is the maximum clearance rate, $I_{\max}$ is the maximum ingestion rate, and K_m is the half-saturation constant.

Carbon-specific feeding rates were calculated using the conversion factor 0.221 $\mu\text{g C } \mu\text{m}^{-3}$ for *R. salina* (Helenius & Saiz 2017) and the length-weight relationships for *P. grani*: $W = 3.24 \times 10^{-7} L^{2.34}$ for nauplii and $W = 1.12 \times 10^{-10} L^{3.58}$ for later stages (Saiz et al. unpubl. data), where W is the copepod mass in $\mu\text{g C}$, and L is the body (nauplii) or prosome (copepodites and adults) length in μm .

The relationships between copepod body mass and maximum feeding rates (clearance and ingestion rates) were adjusted to a power function. The power fits were obtained from linear regression after the log transformation of the variables using the software KaleidaGraph 4.5.4. To improve the robustness of the fitting, for this analysis, we added additional estimates of the maximum ingestion rates of *P. grani* obtained from unpublished experiments (Olivares et al. unpubl. data) conducted under similar conditions.

3. Results

The average size of the copepods used in the functional response experiments ranged from 139 μm in early nauplii to 1052 μm in adult females (Table 1). The mean prey size was very similar in all experiments (range 7.1-7.4 μm ESD). All stages of *Paracartia grani* showed a Holling type III feeding functional response when grazing on *Rhodomonas salina*. Clearance rates peaked at an intermediate prey concentration and decreased at lower and higher concentrations (Fig. 1). The maximum clearance rates increased with stage/size from 0.36 $\text{mL ind}^{-1} \text{d}^{-1}$ for early nauplii to 15.9 $\text{mL ind}^{-1} \text{d}^{-1}$ for adult females; the maximum clearance rates of adult males were, however, lower than the estimates for late copepodites and females (Table 1). Ingestion rates increased with prey concentration following a sigmoid curve and became steady after reaching food saturation (Fig. 1). The maximum ingestion rates increased from ca. 1600 $\text{cells ind}^{-1} \text{d}^{-1}$ (ca. 0.08 $\mu\text{g C ind}^{-1} \text{d}^{-1}$) in early nauplii to ca. 218,000 $\text{cells ind}^{-1} \text{d}^{-1}$ (ca. 9.12 $\mu\text{g C ind}^{-1} \text{d}^{-1}$) in adult females (Table 1). The maximum ingestion rates of females were approximately three times higher than those of males.

Figure 2 shows the power fits between copepod body mass and copepod maximum clearance rates ($r^2 = 0.991$, $p < 0.01$; Fig. 2a) and maximum ingestion rates ($r^2 = 0.967$, $p < 0.001$; Fig. 2b). The maximum clearance rates of nauplii and copepodites increased with body mass with a scaling factor of 1.09 ± 0.07 , whereas the fit between body mass and maximum ingestion rates had an exponent of 0.83 ± 0.05 considering all the copepod life stages. The maximum clearance rates of adults did not follow the general pattern observed for nauplii and copepodites.

In terms of mass-specific rates, the maximum daily ingestion ranged from ca. 210% of the body carbon in nauplii to 265% of the body carbon in early copepodites and then declined to 68% and 124% of the body carbon in adult males and females, respectively (Fig. 3, Table 1). Adults became satiated at much higher prey concentrations than copepodites and nauplii. The half-saturation concentrations (K_m) were 3773 cells mL^{-1} (ca. $170 \mu\text{g C L}^{-1}$) and 6877 cells mL^{-1} (ca. $290 \mu\text{g C L}^{-1}$) for males and females, respectively, and less than 2300 cells mL^{-1} (ca. $110 \mu\text{g C L}^{-1}$) for juvenile stages (Table 1). The K_m values for copepodites (CI-II and CIII-IV) were lower than those for nauplii. The food concentrations at which 90% of satiation (i.e., the maximum ingestion rate) was attained (C_{90}) followed a similar pattern to that of K_m : decreasing from nauplii to copepodites and substantially increasing in adults (Table 1).

4. Discussion

4. 1. Ontogenetic changes in the feeding functional response

We examined the changes in the feeding functional response of the marine copepod *Paracartia grani* through ontogeny and focused on relevant ecological

aspects, such as the shape and the key parameters of the functional response (i.e., $I_{\max}$, $F_{\max}$, K_m). Previous studies have also addressed ontogenetic changes in the feeding patterns of marine copepods, but they focused on other aspects of feeding, such as the prey size spectrum (Berggreen et al. 1988), or did not analyse functional responses through the whole developmental range from nauplii to adults (Almeda et al. 2010, Saiz et al. 2014).

The functional response of an organism shows its capability to exploit environmental resources and the potential flux of energy into secondary production (somatic growth, egg production). We found that the feeding functional response of *P. grani* grazing on *Rhodomonas salina* consistently followed a Holling type III pattern in all life stages. Holling type III responses have been traditionally linked to predator-prey systems where predators may control prey densities by switching to alternative prey types according to the relative prey abundances, exploiting the more abundant resources and thereby ensuring a basal level of prey diversity in the system (Gismervik & Andersen 1997, Van Leeuwen et al. 2007). However, Holling type III responses are not restricted to multiple-prey systems, as they have been reported in numerous single-prey experiments (Kiørboe et al. 2018). In our study, we conducted experiments with one algal species and inferred a clear type III response in all cases. The interpretation of type III responses in single-prey scenarios is that predators stop actively searching for prey when the energy gained by feeding would not compensate for the energy invested in capturing prey (in our study, this threshold would correspond to approximately 2,000 cells mL⁻¹ for nauplii and copepodites, and 4,000-6,000 cells mL⁻¹ for adults). Accordingly, this foraging strategy has been proposed to be adaptive for copepods associated

with productive environments where prey availability is high, such as coastal environments (Paffenhöfer & Stearns 1988). A recent analysis by Kiørboe et al. (2018) noted that in single-prey environments, Holling type III responses are more typical of copepods with feeding strategies that involve higher metabolic costs (for instance, creating feeding currents to enhance the encounter rate with prey), while less active swimmers, such as ambush-feeding copepods, generally show a Holling type II response. Adults of *Acartia* species (*sensu lato*) normally produce feeding currents when feeding on relatively small prey (Jonsson & Tiselius 1990, Saiz & Kiørboe 1995). The fact that we observed the same feeding functional pattern throughout the whole ontogeny of *P. grani* suggests that active foraging occurs not only in adults but also in the juvenile stages of this marine species. Nauplii of *P. grani* are known to perform continuous hopping movements during foraging and show a type III response for a wide range of prey types (Henriksen et al. 2007, Helenius & Saiz 2017). There are no detailed reports regarding the underlying mechanisms involved in *P. grani* nauplii feeding, but in the case of the related species *Acartia tonsa*, nauplii are able to detect a prey item remotely and then capture it by using temporary feeding currents generated by the second antennae and mandibles (Paffenhöfer et al. 1996, Bruno et al. 2012). Although previous studies (Almeda et al. 2010, Saiz et al. 2014) and our results indicate that the feeding functional response follows the same pattern through copepod development, Santer and van den Bosch (1994) reported ontogenetic differences in the freshwater copepod species *Cyclops vicinus*. They found that the naupliar and early copepodite stages of *C. vicinus* exhibited a Holling type II response, whereas late copepodites and adults responded in more of a

rectilinear pattern. However, their observations must be taken with caution because the rectilinear response in late copepodites and adults might be a consequence of not reaching satiating conditions at the experimental food concentrations. Moreover, it is often difficult to accurately distinguish between type II and type III responses in datasets either because very low food concentrations might not be included in the experimental range or because the large variance in counts at low concentrations masks a type III response.

4. 2. *Changes in the maximum clearance rates through ontogeny*

The maximum clearance rate that we estimated for nauplii II-III ($0.36 \text{ mL ind}^{-1} \text{ d}^{-1}$) was very similar to the value reported by Helenius and Saiz (2017) for the same copepod species, stage, and prey species ($0.33 \text{ mL ind}^{-1} \text{ d}^{-1}$). These values were also comparable to the maximum clearance rates estimated by Henriksen et al. (2007) for *P. grani* nauplii II-III grazing on larger prey organisms (0.39 and $0.41 \text{ mL ind}^{-1} \text{ d}^{-1}$ for *Thalassiosira weissflogii* and *Heterocapsa* sp., respectively).

The maximum clearance rates of *P. grani* increased with body mass from nauplii to late copepodites to the power of 1.09, which is in accordance with the scaling factor of 0.93 determined for the clearance rates of the closely related species *A. tonsa* when grazing on *Rhodomonas baltica* (Berggreen et al. 1988) and the value of 1 reported for maximum clearance rates across various zooplankton taxa (Kiørboe 2011, Kiørboe & Hirst 2014). Maximum clearance rates describe the capability of a predator to locate and capture prey at low prey densities when factors such as handling time or gut filling are not relevant. Compared to nauplii, copepodites have more developed feeding appendages

that can produce larger and stronger feeding currents to capture prey and scan larger volumes of water (Paffenhöfer 1971). Therefore, a higher feeding efficiency in copepodites may result in a steeper increase in maximum clearance rates with body mass compared to the typical allometric scaling factor of 0.75 found for other parameters, such as maximum ingestion rates (see next subsection).

The maximum clearance rates of adults did not follow the general pattern observed in juvenile stages, as the maximum clearance rates of males and females (ca. 9 and 16 mL ind⁻¹ d⁻¹, respectively) were not higher than that of late copepodites (ca. 16 mL ind⁻¹ d⁻¹). We think that this discordance in adult clearance rates is due to the relatively small size of the prey, *R. salina*. It is known that the optimal prey size of marine copepods changes with body size and through ontogeny (Berggreen et al. 1988, Landry & Fagerness 1988, Saiz et al. 2014). It could be argued that the use of the same prey type throughout development may affect the interpretation of our results. In fact, our estimates of the maximum clearance rates for adult females were 4-7 times lower than those previously reported for *P. grani* females feeding on larger prey organisms, such as *T. weissflogii* [ca. 13 µm ESD, 70 mL fem⁻¹ d⁻¹; Saiz et al. (1992)], *Heterocapsa* sp. [ca. 13 µm ESD, 66 mL fem⁻¹ d⁻¹; Isari et al. (2013)], and *Oxyrrhis marina* [ca. 17 µm ESD, 111 mL ind⁻¹ d⁻¹; Calbet et al. (2007)]. In Table 1, we can also see that adults showed the highest K_m and C_{90} values through development, indicating that the discrepancy between optimal prey size and predator size was the greatest among all life stages. Therefore, it is likely that if we had used larger prey in the experiments with adults, the maximum clearance

rates would have conformed to the power-of-1 scaling pattern throughout development.

4. 3. *Changes in the maximum ingestion rates through ontogeny*

Although the maximum ingestion rate that we estimated for nauplii II-III was comparable to the value reported by Helenius and Saiz (2017) for the same copepod species, stage, and prey species (1623 and 1234 cells ind⁻¹ d⁻¹, respectively), our estimate of the carbon-specific maximum ingestion rate (ca. 213% body C d⁻¹) was much higher than the value found in their study (74% body C d⁻¹). This large difference in mass-specific rates relies mostly on the fact that the carbon biomass that we extrapolated for our nauplii (34 ng C ind⁻¹) was substantially lower than the measured biomass in their study (62 ng C ind⁻¹), even though the average naupliar sizes were very similar (139 and 143 µm, respectively). Our estimate, nonetheless, was closer to the maximum daily ingestions reported by Henriksen et al. (2007) for *P. grani* nauplii feeding on *T. weissflogii* and *Heterocapsa* sp. (185% and 299% body C d⁻¹, respectively). In this regard, Henriksen et al. (2007) determined that *P. grani* nauplii, with an average body length of 165 µm, contained 40 ng C per animal, which indicates that the carbon content values measured by Helenius and Saiz (2017) might have been anomalously high.

The maximum daily ration that we estimated for adult females (124% body C) fell within the range of 103-188% d⁻¹ found in the literature for the closely related species *A. tonsa* at same experimental temperature and when provided with the same prey species (Kiørboe et al. 1985, Toudal & Riisgård 1987, Støttrup & Jensen 1990, Thor & Wendt 2010). This value was also close to the

specific rates reported for *P. grani* feeding on other prey, such as *Heterocapsa* sp. [ca. 13 μm ESD, 147% d^{-1} , Isari et al. (2013)] and *Oxyrrhis marina* [ca. 17 μm ESD, 149% d^{-1} , Calbet et al. (2007)]. It is worth noting that across prey sizes, the maximum mass-specific ingestion rates show a much narrower range of variability in comparison with the maximum clearance rates, which can differ by a factor of up to 7 (see previous subsection). In this regard, Frost (1972) previously reported that the maximum clearance rates of *Calanus pacificus* varied between 91 and 223 $\text{mL ind}^{-1} \text{d}^{-1}$ across a range of prey diameters of between 11 and 87 μm , whereas the maximum ingestion rates remained fairly constant between 24.2 and 27.1 $\mu\text{g C ind}^{-1} \text{d}^{-1}$. Maximum clearance rates are very dependent on the predator perception of prey, which is particularly sensitive to prey size and motility (Saiz et al. 2014, Almeda et al. 2018). In contrast, the maximum ingestion rate appears to be not greatly influenced by prey size (Frost 1972) and is more constrained to factors such as copepod gut volume, digestion time and gut turnover rate (Hassett & Landry 1988, Henriksen et al. 2007). Handling times, as a parameter in Holling equations, are typically also considered for the satiation effect, but in copepods they are generally too short to constrain food intake (Tiselius et al. 2013, Kiørboe et al. 2016). The fact that specific maximum ingestion rates are not very dependent on prey size helps to explain the exponent of 0.83 in our allometric analysis. This exponent, although slightly higher, was not significantly different from the expected value of 0.75 (t-test, $p > 0.05$) according to the three-quarter power law accepted for most organisms, including marine calanoid copepods (West & Brown 2005, Saiz & Calbet 2007, Kiørboe & Hirst 2014). Thus, it appears that the use of a

single prey size in our experiments did not result in any substantial bias in the estimation of maximum ingestion rates.

In relation to sex differences, the mass-specific maximum ingestion rates of males were ca. 1.8 times lower than those of females (68% and 124% of body $C\ d^{-1}$, respectively). Sex-related differences in feeding rates have been previously reported for the same copepod species (Saiz et al. 1992) and other copepod species (Saage et al. 2009, van Someren Gréve et al. 2017b). On the one hand, some authors have attributed the relatively lower feeding rates of males to size differences between the sexes (Bautista et al. 1988, van Someren Gréve et al. 2017b). However, our allometric analysis (Fig. 2) shows that the maximum ingestion rates of *P. grani* males were lower than expected for a copepod of their size. In fact, the feeding rates of late copepodites were also higher than those of males. On the other hand, previous studies state that in some copepod species, adult males are more active swimmers than females and spend a substantial fraction of their time searching for a mate to the detriment of feeding (Kiørboe 2007, Almeda et al. 2017, van Someren Gréve et al. 2017a). We do not know whether sex-related differences in swimming activity could explain the low feeding rates found in *P. grani* males, but it seems unlikely given that the males and females of the related species *A. tonsa* appear to display similar swimming patterns (Bagøien & Kiørboe 2005). We think that the differences in maximum feeding rates between the sexes may not rely on their swimming behaviour but rather on differences in the allocation of energy to the reproductive process. In this regard, *P. grani* females can exhibit high reproductive output under optimal food conditions that would require higher ingestion rates (Saiz et al. 2015, Saiz et al. 2017).

459

460 4. 4. Conclusions

461 Our study provides empirical data about the ontogenetic changes that marine
462 copepods show in the exploitation of food resources. The insights obtained can
463 help improving the modelling capability to estimate energy transfer through
464 copepod populations with different stage compositions in pelagic food webs. We
465 have shown that the functional response of the copepod *P. grani* when feeding
466 on *R. salina* followed a common pattern (Holling type III) from early nauplii to
467 adults. The maximum clearance rates increased with body mass with a power
468 scaling factor of 1, except for the adult stages, that deviated from that common
469 allometric trend, presumably due to the relatively small prey size. The maximum
470 ingestion rates scaled to body mass with a factor of 0.8, close to the 0.75 typical
471 value found in many animals. In this case, prey size did not seem to
472 substantially affect our estimates of maximum ingestion rates along ontogeny.
473 We suggest that future studies addressing feeding activity of copepods should
474 also consider the changes in prey size spectrum along ontogeny. Besides, we
475 think that plankton dynamic models would benefit from assessments in multiple
476 resource scenarios, in which selectivity patterns and switching responses can
477 modulate the foraging behaviour of marine copepods.

478 **Acknowledgements**

479 We thank Kaiene Griffell for assisting with culture maintenance. This study was
480 framed within the project FERMI (CGL2014-59227-R, MINECO/AEI/FEDER,
481 UE) and is a contribution of the Marine Zooplankton Ecology Group (2017 SGR
482 87). Manuel Olivares is supported financially by an FPU grant (FPU15/01747)
483 from the Spanish Ministry of Education, Culture and Sport.

484 **Literature cited**

- 485 Alcaraz M, Almeda R, Calbet A, Saiz E, Duarte CM, Lasternas S, Agustí S, Santiago R, Movilla J,
 486 Alonso A (2010) The role of arctic zooplankton in biogeochemical cycles: respiration
 487 and excretion of ammonia and phosphate during summer. *Polar Biol* 33:1719-1731
- 488 Almeda R, Augustin CB, Alcaraz M, Calbet A, Saiz E (2010) Feeding rates and gross growth
 489 efficiencies of larval developmental stages of *Oithona davisae* (Copepoda, Cyclopoida).
 490 *J Exp Mar Biol Ecol* 387:24-35
- 491 Almeda R, Calbet A, Alcaraz M, Saiz E, Trepát I, Arin L, Movilla J, Saló V (2011) Trophic role and
 492 carbon budget of metazoan microplankton in northwest Mediterranean coastal
 493 waters. *Limnol Oceanogr* 56:415-430
- 494 Almeda R, van Someren Gréve H, Kiørboe T (2017) Behavior is a major determinant of
 495 predation risk in zooplankton. *Ecosphere* 8 (2):e01668
- 496 Almeda R, van Someren Gréve H, Kiørboe T (2018) Prey perception mechanism determines
 497 maximum clearance rates of planktonic copepods. *Limnol Oceanogr* 63:2695-2707
- 498 Bagøien E, Kiørboe T (2005) Blind dating-mate finding in planktonic copepods. III.
 499 Hydromechanical communication in *Acartia tonsa*. *Mar Ecol Prog Ser* 300:129-133
- 500 Bautista B, Rodríguez V, Jiménez F (1988) Short-term feeding rates of *Acartia grani* in natural
 501 conditions: diurnal variation. *J Plankton Res* 10:907-920
- 502 Berggreen U, Hansen B, Kiørboe T (1988) Food size spectra, ingestion and growth of the
 503 copepod *Acartia tonsa* during development: implications for determination of
 504 copepod production. *Mar Biol* 99:341-352
- 505 Boyer S, Arzul I, Bonnet D (2012) Some like it hot: *Paracartia grani* (Copepoda: Calanoida)
 506 arrival in the Thau lagoon (south of France-Mediterranean Sea). *Mar Biodivers Rec* 5:1-
 507 5
- 508 Broglio E, Jonasdóttir SH, Calbet A, Jakobsen HH, Saiz E (2003) Effect of heterotrophic versus
 509 autotrophic food on feeding and reproduction of the calanoid copepod *Acartia tonsa*:
 510 relationship with prey fatty acid composition. *Aquat Microb Ecol* 31:267-278
- 511 Bruno E, Andersen Borg CM, Kiørboe T (2012) Prey detection and prey capture in copepod
 512 nauplii. *PLOS ONE* 7:e47906
- 513 Calbet A (2001) Mesozooplankton grazing effect on primary production: A global comparative
 514 analysis in marine ecosystems. *Limnol Oceanogr* 46:1824-1830
- 515 Calbet A, Alcaraz M (1997) Growth and survival rates of early developmental stages of *Acartia*
 516 *grani* (Copepoda: Calanoida) in relation to food concentration and fluctuations in food
 517 supply. *Mar Ecol Prog Ser* 147:181-186
- 518 Calbet A, Garrido S, Saiz E, Alcaraz M, Duarte CM (2001) Annual zooplankton succession in
 519 coastal NW Mediterranean waters: the importance of the smaller size fractions. *J*
 520 *Plankton Res* 23:319-331
- 521 Calbet A, Saiz E (2005) The ciliate-copepod link in marine ecosystems. *Aquat Microb Ecol*
 522 38:157-167
- 523 Calbet A, Saiz E, Barata C (2007) Lethal and sublethal effects of naphthalene and 1,2-
 524 dimethylnaphthalene on the marine copepod *Paracartia grani*. *Mar Biol* 151:195-204
- 525 Chen M-R, Kâ S, Hwang J-S (2010) Diet of the copepod *Calanus sinicus* Brodsky, 1962
 526 (Copepoda, Calanoida, Calanidae) in northern coastal waters of Taiwan during the
 527 northeast monsoon period. *Crustaceana* 83:851-864
- 528 Egilmez HI, Morozov A (2016) Tri-trophic plankton models revised: importance of space, food
 529 web structure and functional response parametrisation. *Math Model Nat Phenom*
 530 11:16-33
- 531 Fernández F (1979) Nutrition studies in the nauplius larva of *Calanus pacificus* (Copepoda:
 532 Calanoida). *Mar Biol* 53:131-147

533 Frost BW (1972) Effects of size and concentration of food particles on the feeding behavior of
534 the marine planktonic copepod *Calanus pacificus*. Limnol Oceanogr 17:805-815

535 Gentleman WC, Neuheimer AB (2008) Functional responses and ecosystem dynamics: how
536 clearance rates explain the influence of satiation, food-limitation and acclimation. J
537 Plankton Res 30:1215-1231

538 Gismervik I, Andersen T (1997) Prey switching by *Acartia clausi*: experimental evidence and
539 implications of intraguild predation assessed by a model. Mar Ecol Prog Ser 157:247-
540 259

541 Gonçalves RJ, van Someren Gréve H, Couespel D, Kiørboe T (2014) Mechanisms of prey size
542 selection in a suspension-feeding copepod, *Temora longicornis*. Mar Ecol Prog Ser
543 517:61-74

544 Guillard RRL (1983) Culture of phytoplankton for feeding marine invertebrates. In: Jr CJB (ed)
545 Culture of Marine Invertebrate Animals Selected readings. Hutchinson Ross Publ. C.,
546 Stroudsburg, Pennsylvania

547 Hassett RP, Landry MR (1988) Short-term changes in feeding and digestion by the copepod
548 *Calanus pacificus*. Mar Biol 99:63-74

549 Helenius LK, Saiz E (2017) Feeding behaviour of the nauplii of the marine calanoid copepod
550 *Paracartia grani* Sars: Functional response, prey size spectrum, and effects of the
551 presence of alternative prey. PLOS ONE 12:e0172902

552 Henriksen CI, Saiz E, Calbet A, Hansen BW (2007) Feeding activity and swimming patterns of
553 *Acartia grani* and *Oithona davisae* nauplii in the presence of motile and non-motile
554 prey. Mar Ecol Prog Ser 331:119-129

555 Holling CS (1966) The functional response of invertebrate predators to prey density. Mem
556 Entomol Soc Can 98:5-86

557 Ikeda T, Kanno Y, Ozaki K, Shinada A (2001) Metabolic rates of epipelagic marine copepods as a
558 function of body mass and temperature. Mar Biol 139:587-596

559 Isari S, Antó M, Saiz E (2013) Copepod foraging on the basis of food nutritional quality: can
560 copepods really choose? PLOS ONE 8:e84742

561 Isari S, Saiz E (2011) Feeding performance of the copepod *Clausocalanus lividus* (Frost and
562 Fleminger, 1968). J Plankton Res 33:715-728

563 Ismar SMH, Hansen T, Sommer U (2008) Effect of food concentration and type of diet on
564 *Acartia* survival and naupliar development. Mar Biol 154:335-343

565 Jonsson PP, Tiselius P (1990) Feeding behaviour, prey detection and capture efficiency of the
566 copepod *Acartia tonsa* feeding on planktonic ciliates. Mar Ecol Prog Ser 60:35-44

567 Kiørboe T (2006) Sex, sex-ratios, and the dynamics of pelagic copepod populations. Oecologia
568 148:40-50

569 Kiørboe T (2007) Mate finding, mating, and population dynamics in a planktonic copepod
570 *Oithona davisae*: There are too few males. Limnol Oceanogr 52:1511-1522

571 Kiørboe T (2011) How zooplankton feed: mechanisms, traits and trade-offs. Biol Rev Camb
572 Philos Soc 86:311-339

573 Kiørboe T, Ceballos S, Thygesen UH (2015) Interrelations between senescence, life-history
574 traits, and behavior in planktonic copepods. Ecology 96:2225-2235

575 Kiørboe T, Gonçalves RJ, Couespel D, van Someren Gréve H, Saiz E, Tiselius P (2016) Reply to
576 comment: Prey perception in feeding-current feeding copepods. Limnol Oceanogr
577 61:1169-1171

578 Kiørboe T, Hirst AG (2014) Shifts in mass scaling of respiration, feeding, and growth rates
579 across life-form transitions in marine pelagic organisms. Am Nat 183:E118-130

580 Kiørboe T, Møhlenberg F, Hamburger K (1985) Bioenergetics of the planktonic copepod *Acartia*
581 *tonsa*: relation between feeding, egg production and respiration, and composition of
582 specific dynamic action. Mar Ecol Prog Ser 26:85-97

583 Kiørboe T, Saiz E, Tiselius P, Andersen KH (2018) Adaptive feeding behavior and functional
584 responses in zooplankton. Limnol Oceanogr 63:308-321

585 Kouwenberg JHM (1993) Sex ratio of calanoid copepods in relation to population composition
 586 in the northwestern mediterranean. *Crustaceana* 64:281-299
 587 Lampitt RS (1978) Carnivorous feeding by a small marine copepod. *Limnol Oceanogr* 23:1228-
 588 1231
 589 Landry MR, Fagerness VL (1988) Behavioral and morphological influences on predatory
 590 interactions among marine copepods. *Bull Mar Sci* 43:509-529
 591 Longhurst AR (1985) The structure and evolution of plankton communities. *Prog Oceanogr*
 592 15:1-35
 593 López E, Anadón R, Harris RP (2007) Functional responses of copepod nauplii using a high
 594 efficiency gut fluorescence technique. *Mar Biol* 150:893-903
 595 Lučić D, Njire J, Morović M, Precali R, Fuks D, Bolotin J (2003) Microzooplankton in the open
 596 waters of the northern Adriatic Sea from 1990 to 1993: the importance of copepod
 597 nauplii densities. *Helgol Mar Res* 57:73-81
 598 Morozov A, Poggiale J-C, Cordoleani F (2012) Implementation of the zooplankton functional
 599 response in plankton models: State of the art, recent challenges and future directions.
 600 *Prog Oceanogr* 103:80-91
 601 Morozov AY (2010) Emergence of Holling type III zooplankton functional response: bringing
 602 together field evidence and mathematical modelling. *J Theor Biol* 265:45-54
 603 Mullin MM, Brooks ER (1967) Laboratory culture, growth rate, and feeding behavior of a
 604 planktonic marine copepod. *Limnol Oceanogr* 12:657-666
 605 Olivares M, Saiz E, Calbet A (unpubl. data)
 606 Paffenhöfer G-A (1971) Grazing and ingestion rates of nauplii, copepodids and adults of the
 607 marine planktonic copepod *Calanus helgolandicus*. *Mar Biol* 11:286-298
 608 Paffenhöfer G-A (1998) On the relation of structure, perception and activity in marine
 609 planktonic copepods. *J Mar Syst* 15:457-473
 610 Paffenhöfer G-A, Stearns DE (1988) Why is *Acartia tonsa* (Copepoda: Calanoida) restricted to
 611 nearshore environments? *Mar Ecol Prog Ser* 42:33-38
 612 Paffenhöfer G-A, Strickler JR, Lewis KD, Richman S (1996) Motion behavior of nauplii and early
 613 copepodid stages of marine planktonic copepods. *J Plankton Res* 18:1699-1715
 614 Runge JA (1988) Should we expect a relationship between primary production and fisheries?
 615 The role of copepod dynamics as a filter of trophic variability. *Hydrobiologia* 167-
 616 168:61-71
 617 Saage A, Vadstein O, Sommer U (2009) Feeding behaviour of adult *Centropages hamatus*
 618 (Copepoda, Calanoida): Functional response and selective feeding experiments. *J Sea*
 619 *Res* 62:16-21
 620 Saiz E, Alcaraz M, Paffenhöfer G-A (1992) Effects of small-scale turbulence on feeding rate and
 621 gross-growth efficiency of three *Acartia* species (Copepoda, Calanoida). *J Plankton Res*
 622 14:1085-1097
 623 Saiz E, Calbet A (2007) Scaling of feeding in marine calanoid copepods. *Limnol Oceanogr*
 624 52:668-675
 625 Saiz E, Calbet A, Griffell K (2017) Sex-dependent effects of caloric restriction on the ageing of
 626 an ambush feeding copepod. *Sci Rep* 7:12662
 627 Saiz E, Calbet A, Griffell K, Bersano JGF, Isari S, Solé M, Peters J, Alcaraz M (2015) Ageing and
 628 caloric restriction in a marine planktonic copepod. *Sci Rep* 5
 629 Saiz E, Griffell K, Calbet A (unpubl. data)
 630 Saiz E, Griffell K, Calbet A, Isari S (2014) Feeding rates and prey : predator size ratios of the
 631 nauplii and adult females of the marine cyclopoid copepod *Oithona davisae*. *Limnol*
 632 *Oceanogr* 59:2077-2088
 633 Saiz E, Kiørboe T (1995) Predatory and suspension feeding of the copepod *Acartia tonsa* in
 634 turbulent environments. *Mar Ecol Prog Ser* 122:147-158

- Santer B, van den Bosch F (1994) Herbivorous nutrition of *Cyclops vicinus*: the effect of a pure algal diet on feeding, development, reproduction and life cycle. J Plankton Res 16:171-195
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nat Methods 9:671-675
- Solomon ME (1949) The natural control of animal populations. J Anim Ecol 18:1-35
- Steinberg DK, Carlson CA, Bates NR, Goldthwait SA, Madin LP, Michaels AF (2000) Zooplankton vertical migration and the active transport of dissolved organic and inorganic carbon in the Sargasso Sea. Deep Sea Res I 47:137-158
- Støttrup JG, Jensen J (1990) Influence of algal diet on feeding and egg-production of the calanoid copepod *Acartia tonsa* Dana. J Exp Mar Biol Ecol 141:87-105
- Thor P, Wendt I (2010) Functional response of carbon absorption efficiency in the pelagic calanoid copepod *Acartia tonsa*. Limnol Oceanogr 55:1779-1789
- Tiselius P, Saiz E, Kiørboe T (2013) Sensory capabilities and food capture of two small copepods, *Paracalanus parvus* and *Pseudocalanus* sp. Limnol Oceanogr 58:1657-1666
- Toudal K, Riisgård HU (1987) Acute and sublethal effects of cadmium on ingestion, egg production and life-cycle development in the copepod *Acartia tonsa*. Mar Ecol Prog Ser 37:141-146
- Turner JT (2004) The importance of small planktonic copepods and their roles in pelagic marine food webs. Zool Stud 43:255-266
- Turner JT (2015) Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's biological pump. Prog Oceanogr 130:205-248
- Uye S-i, Kayano Y (1994) Predatory feeding behavior of *Tortanus* (Copepoda: Calanoida): life-stage differences and the predation impact on small planktonic crustaceans. J Crustac Biol 14:473-483
- Van Leeuwen E, Jansen VAA, Bright PW (2007) How population dynamics shape the functional response in a one-predator–two-prey system. Ecology 88:1571-1581
- van Someren Gréve H, Almeda R, Kiørboe T (2017a) Motile behavior and predation risk in planktonic copepods. Limnol Oceanogr 62:1810-1824
- van Someren Gréve H, Almeda R, Lindegren M, Kiørboe T (2017b) Gender-specific feeding rates in planktonic copepods with different feeding behavior. J Plankton Res 39:631-644
- Webber MK, Roff JC (1995) Annual structure of the copepod community and its associated pelagic environment off Discovery Bay, Jamaica. Mar Biol 123:467-479
- West GB, Brown JH (2005) The origin of allometric scaling laws in biology from genomes to ecosystems: towards a quantitative unifying theory of biological structure and organization. J Exp Biol 208:1575-1592
- Wollrab S, Diehl S (2015) Bottom-up responses of the lower oceanic food web are sensitive to copepod mortality and feeding behavior. Limnol Oceanogr 60:641-656

Table legends

Table 1. Parameters of the feeding functional responses of *Paracartia grani* on *Rhodomonas salina*. Maximum clearance rates ($F_{\max}$), maximum ingestion rates ($I_{\max}$), half-saturation constants (K_m), and prey concentrations at which 90% of the maximum ingestion rates were reached (C_{90}) are shown for the different copepod stages and sizes (N = nauplii, C = copepodites, Ad = adults). Means $\pm$ SE are provided.

Figure legends

Fig. 1. Functional responses of (a) nauplii II-III, (b) nauplii V-VI, (c) copepodites I-II, (d) copepodites III-IV, (e) adult males, and (f) adult females of *Paracartia grani* feeding on *Rhodomonas salina*. Holling type III curves were fitted to the clearance and ingestion rates as a function of the average prey concentration. Data in brackets are excluded from the model.

Fig. 2. (a) Maximum clearance rates and (b) maximum ingestion rates of *Paracartia grani* feeding on *Rhodomonas salina* with increasing copepod body mass (W). Power function fits and corresponding equations are shown. Note that for the clearance rates, only data on nauplii and copepodites are used for the fit. For the ingestion rates, data from the functional response experiments (■) and additional unpublished trials (▲) are pooled together.

Fig. 3. Carbon-specific functional responses of (a) nauplii and (b) copepodites and adults of *Paracartia grani* feeding on *Rhodomonas salina*. Holling type III curves were fitted to the carbon-specific ingestion rates as a function of the average prey concentration. Means and SE bars are shown. Data in brackets are excluded from the model.

Table 1. Parameters of the feeding functional responses of *Paracartia grani* on *Rhodomonas salina*. Maximum clearance rates ($F_{\max}$), maximum ingestion rates ($I_{\max}$), half-saturation constants (K_m), and prey concentrations at which 90% of the maximum ingestion rates were reached (C_{90}) are shown for the different copepod stages and sizes (N = nauplii, C = copepodites, Ad = adults). Means $\pm$ SE are provided.

Copepod stage	Copepod size (μm)	$F_{\max}$ ($\text{mL ind}^{-1} \text{d}^{-1}$)	$I_{\max}$ ($\text{cells ind}^{-1} \text{d}^{-1}$)	$I_{\max}$ (% body C d^{-1})	K_m (cells mL^{-1})	C_{90} (cells mL^{-1})
NII-III	139 $\pm$ 2	0.36 $\pm$ 0.02	1623 $\pm$ 46	213 $\pm$ 6.5	2272 $\pm$ 138	6815
NV-VI	238 $\pm$ 2	1.29 $\pm$ 0.06	5623 $\pm$ 185	207 $\pm$ 6.5	2188 $\pm$ 136	6563
CI-II	457 $\pm$ 3	6.25 $\pm$ 0.34	21649 $\pm$ 659	265 $\pm$ 7.6	1733 $\pm$ 120	5198
CIII-IV	632 $\pm$ 2	15.7 $\pm$ 0.98	52761 $\pm$ 1290	195 $\pm$ 5.4	1684 $\pm$ 121	5051
Ad Males	922 $\pm$ 1	8.98 $\pm$ 0.46	67772 $\pm$ 1858	68 $\pm$ 1.9	3773 $\pm$ 243	11318
Ad Females	1052 $\pm$ 3	15.9 $\pm$ 0.57	218457 $\pm$ 6401	124 $\pm$ 3.3	6877 $\pm$ 368	20632

Fig. 1

Fig. 2

Fig. 3