**The role of fire history, land-use, and vegetation structure on the response of
Mediterranean lizards to fire**

Tiago Pinto^{1,2}, Bruno Moreira^{1,3}, Helena Freitas¹, Xavier Santos^{4,*}

¹ Centre for Functional Ecology, Department of Life Sciences, University of Coimbra.
Calçada Martim de Freitas, 3000-456 Coimbra, Portugal.

² CIBIO-UE, Research Centre in Biodiversity and Genetic Resources, University of
Évora. Departament of Biology, Mitra. 7002-554 Évora, Portugal.

³ CIDE-CSIC, Campus IVIA, Ctra. Náquera Km. 4.5, 46113 Montcada, Valencia,
Spain.

⁴ CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO
Laboratório Associado, Universidade do Porto, Campus Agrário de Vairão, R. Padre
Armando Quintas 7, 4485-661 Vairão, Portugal.

* Corresponding author: Xavier Santos, e-mail: xsantossantiro@gmail.com

Abstract: Fire is a critical component of terrestrial ecosystems and essential to
understand the composition and diversity of communities in fire-prone regions.
Nevertheless, fire does not act alone, and other factors such as land-use type and
vegetation structure can also operate at different scales, prompting the response to fire
of ectotherm communities such as reptiles. We have evaluated the influence of
environmental factors on the abundance, diversity, and richness of reptile communities
in Serra da Estrela Natural Park (northern Portugal), at both the landscape and micro-
habitat scales. In spring 2014 and 2015, 20 transects were surveyed eight times, the
vegetation structure and composition were measured, and the extent of land-use types

around each transect calculated in this Mediterranean community. At the landscape scale, reptile abundance was higher in natural woodlands and more complex habitats (higher vegetation heterogeneity and plant richness), and reptile evenness was affected by vegetation heterogeneity and time since fire. By contrast, species richness was not related to any environmental factor measured in this study. Only two lizard species were common in the park, *Psammodromus algirus* and *Podarcis guadarramae*. Their relative abundances increased with contrasting habitat variables as the former is a ground-dwelling lizard specialist that prefers heterogeneous (unburnt) habitats and the second is a rock lizard that selects open (burnt) stands. At the micro-habitat scale, the probability of reptile presence, reptile abundance, and reptile richness increased with plant richness and decreased with tree cover. This study highlights the importance of several environmental factors, that operate from micro-habitat to landscape scales to understand the response of reptiles to fire.

Keywords: fire, reptile responses, habitat structure, land-use.

1. Introduction

Fire is a fundamental driver of many terrestrial biomes, including Mediterranean type ecosystems (Bond et al., 2005; Bowman et al., 2009, Pausas and Keeley, 2009), and many species have adaptive traits that enable them to persist in these fire-prone habitats (Keeley et al., 2012, Santos and Cheylan, 2013). In the Mediterranean basin, vegetation structure (e.g. heterogeneity, plant diversity, vegetation cover), land-use and fire history (e.g. fire recurrence and time since fire), synergistically interact and shape species habitats (Blondel et al., 2010), with important consequences for the diversity and abundance of different animal species (Moreira and Russo, 2007), but particularly for

51 taxa with low-dispersal ability such as reptiles (Santos et al., 2014; Badiane et al.,
52 2017). Reptiles are sensitive to alterations in habitat attributes such as vegetation
53 structure (e.g. heterogeneity and cover of the different vegetation strata) because they
54 are ectotherms, narrow-ranging species, and characterized by low vagility and dispersal
55 rates (Huey, 1982; Valentine and Schwarzkopf, 2008; Azor et al., 2015).

56 There is evidence that fire history characteristics (e.g. fire recurrence and time since
57 fire) are not the only drivers affecting reptile communities in fire-prone ecosystems, and
58 other processes, like habitat structure, biogeographic affinity, and food availability,
59 operating at different spatial scales, might be equally important (Driscoll and
60 Henderson, 2008; Pastro et al., 2013; Ferreira et al., 2016b). Besides the direct impact
61 of fire on animals (i. e. mortality; Smith et al., 2001; Couturier et al., 2011), shifts in
62 vegetation along postfire successional stages are important in order to characterize
63 reptile communities in fire-prone ecosystems (Driscoll et al., 2012; Santos and Cheylan,
64 2013; Smith et al., 2013; Santos et al., 2016). The complex relationship between fire
65 and vegetation structure (Nimmo et al., 2014) makes it advisable to incorporate
66 vegetation factors into fire-history models to better understand the response of reptiles
67 to fire (Santos et al., 2016).

68 At the landscape scale, vegetation heterogeneity (Nimmo et al., 2012) and land-use
69 type can be key factors explaining total abundance and species richness in reptile
70 communities (Ribeiro et al., 2009; Azor et al. 2015; Santos et al. 2016). At the micro-
71 habitat scale, characteristics such as vegetation cover or plant richness can be important
72 because they influence the habitat by affecting the availability of shelters, food, or sun
73 exposure (Webb and Shine, 1998, Kim and Holt, 2012, Elzer et al., 2013). These factors
74 can operate in concert with fire history to shape reptile communities in fire-prone
75 ecosystems according to the specific life-history traits of each reptile species (Ferreira et

al., 2016b; Santos et al., 2016). For this reason, landscape- and micro-habitat-scale factors have to be integrated to attain a comprehensive view of the composition of reptile communities in fire-prone ecosystems.

This study was conducted in the Serra da Estrela Natural Park (northern Portugal), and performed at two different scales: at the landscape scale, we analyzed the effect of fire history (fire recurrence and time since fire), vegetation heterogeneity, and land-use (main land-use type and land-use diversity) on the reptile abundance, richness and evenness. At the micro-habitat scale, we analyzed the role of plant richness and vegetation cover (tree, shrub, and herbaceous cover) on the probability of reptile presence, reptile abundance and evenness.

We hypothesize that species richness, reptile abundance and evenness are higher in natural areas (woodlands and scrublands) and in areas with higher land-use diversity (landscape scale) as well as in locations with higher plant diversity and lower tree cover (micro-habitat scale). Furthermore, we also expect that fire history (number of fires and time since fire) will affect the reptile community by altering the habitat structure (Ferreira et al., 2016b; Santos et al., 2016).

2. Materials and methods

2.1. Fire regime in Portugal and in the study area

Portugal has the highest fire incidence in Europe (Nunes et al., 2005; Catry et al., 2006; Oliveira et al., 2012), and fire is considered one of the most important agents of landscape change (Silva et al., 2011). Fire frequency and intensity have increased remarkably since the 1960s due to a combination of socioeconomic and environmental conditions such as rural abandonment, and conversion of marginally productive agriculture to eucalypt and pine plantations (Moreira et al., 2001, 2011; Fernandes et al.,

2013). The climate is warm temperate characterized by hot, dry summers and cool, wet winters; likewise, natural vegetation is typically evergreen, resistant to drought, and pyrophytic (Nunes et al., 2005). Thus, both socioeconomic and environmental conditions lead to a high accumulation of fuel leading to a high risk of fire (Moreira et al., 2001, 2009). Catry et al. (2009) concluded that human activities were the primary cause of wildfires since about 60 % of ignitions have occurred in areas of high population density and close to roads.

Serra da Estrela is a mountainous area located in the east-central region of Portugal (Supplementary materials Figure S1), characterized by a high elevational gradient (250 to 1993 m). The mean annual rainfall ranges from 1000 mm in the lowest elevations to approximately 2500 mm at highest elevations (SNIRH, 2015), and the mean annual temperature ranges from 22°C to 4°C at the lowest and highest elevations, respectively. The Serra da Estrela Natural Park (hereafter SENP) is the largest protected area in Portugal (1.0106 km²) and encloses most of this mountain area. The park is a landscape mosaic shaped by a long fire history (Connor et al., 2012) and multiple land uses (including woodlands, scrublands, coniferous plantations, and agriculture areas) arranged across an elevational gradient.

The reptile community at the SENP is composed of 20 species, i.e. 12 lizards and 8 snakes (Lesparre and Crespo, 2008; Supplementary materials Table S2), from both Mediterranean and Atlantic origins, (following Sillero et al., 2009).

2.2. Site selection and environmental characterization

Twenty sites were selected across a fire-history gradient (Supplementary materials Figure S1). The fire history was provided by the Instituto da Conservação da Natureza e Florestas (ICNF). Each site was characterized according to two fire-history variables,

time since last fire (ranging from 0 to >40 years) and fire recurrence (number of fires, ranging from 0 to 6 fires over a period of 40 years; Supplementary materials Table S3). Sites were a minimum of 600 m apart (mean distance and standard deviation between pairs: 17 ± 10 km) and sites with a similar fire history were placed as far as possible from each other to avoid spatial autocorrelation.

Land-use was examined in 200-m buffers around each transect. Raw data were obtained from the Portuguese land-cover map (IGP, 2008) and reclassified to five land-use types: monocultures (pine or eucalyptus plantations), agriculture fields, natural woodlands, scrublands, and urban areas (Supplementary materials Figure S1). Our interest was in the effects of these major land-use types on the abundance and diversity of Mediterranean reptiles (Ribeiro et al., 2009; Azor et al., 2015). We calculated the land-use diversity as the Simpson diversity index of all land uses within the 200-m buffer, and according to the commonest land-use types (i.e. monoculture, scrubland and natural woodland; Supplementary materials Figure S4). The spatial distribution of the transects by land-use classification was clustered, with the majority of monocultures located on the western part and scrubland areas on the eastern part of the study area.

Elevation was considered during site selection due to its influence on reptile presence and distribution (McCain, 2010; Kutt et al., 2011). Transects were selected within the elevational range of 250-1250 m, to avoid sampling on mountaintops that are characterized by the presence of alpine reptile species (i.e. *Iberolacerta monticola* and *Coronella austriaca*) and the absence of fires.

To characterize vegetation structure and heterogeneity, each transect was divided into 20-m sections. In each section, the vegetation within 2 m on either side of the transect (“left” and “right”) was identified to the species level (only for perennial species, because their constancy over the sampling period of the study). For each

transect, the vegetation heterogeneity was accessed through a similarity index (Baselga et al., 2007) ranging between 0 and 1, with values approaching 1 indicating greater dissimilarity (higher vegetation heterogeneity). This index provided an estimate of overall compositional heterogeneity among sections, regardless of differences in species richness. For this, we divided the total number of plant species in each 20 m section by the sum of the minimum values of species not shared between each pair (left and right) of sections. For each transect and at each 20-m section, we described the vegetation structure according to the extent of tree, shrub, and herbaceous layers, ranging each layer from 0% to 100% cover.

2.3. Reptile sampling

Transects were placed on trails to facilitate reptile detectability (see a similar procedure in Santos and Cheylan, 2013), with an average length of 1 km (SD: 159 m; range 770 – 1250 m) and were as linear as possible to avoid recording the same individuals along the transect. Between May and September of 2014, and May and June of 2015, each transect was visited eight times (four visits in spring 2014, 1 visit in autumn 2014, and 3 visits in spring 2015) for 45 minutes each, walking slowly at a constant speed. During each visit, all transects were surveyed within 4-5 days, and visits were at least a week and a half apart. The surveys were made on sunny days and during the hours of reptile activity (from 09h00 to 18h00). Searches included open sites and we also turned rocks and other refuges to maximize observations (Santos and Cheylan, 2013; Santos et al., 2016). All the reptiles recorded were identified to the species level, except the two skink species of the genus *Chalcides* (*C. striatus* and *C. bedriagai*), that were pooled into one group because they were difficult to distinguish during the surveys. All reptile sightings were georeferenced and the perpendicular distance to the transect was noted to access

differences in detectability (e.g. due to differences in vegetation cover; see a similar treatment in Santos and Cheylan, 2013). We found no difference in reptile detectability related to time since fire ($P = 0.27$) or fire recurrence ($P = 0.35$).

2.4. Statistical analysis

For statistical analyses, we used “Visit” as the reptile sampling unit (20 transects * 8 visits per transect = 160 sampling units) to model the total abundance of reptiles, as well as the abundance of the two most common reptile species, the lizards *Psammmodromus algirus* and *Podarcis guadarramae*. Due to the low detectability of many reptile species, we pooled the data of visits into two “Year” units (20 transects * 2 years = 40 samples), since four visits per year more accurately describe the reptile richness and diversity of the transects.

The community metrics examined include the total number of reptiles (abundance), the number of different reptile species observed (richness), and the Pielou’s evenness as a measure of reptile diversity. Evenness quantifies the degree of numerical equality among reptile abundances in each sampling unit; therefore, higher evenness scores indicate more complex (heterogeneous) communities in terms of species composition and relative abundances.

At the landscape level, we tested the effects of the fire recurrence and time since fire (fire history), vegetation heterogeneity, plant richness, land-use type (monocultures, scrublands, and natural woodland) and land-use diversity on the three reptile community metrics. Although recent studies have suggested that General Additive Models (GAM) are more suitable than General Linear Models (GLM) for modelling the response of reptile species to environmental factors and fire (Hu et al. 2013, 2016; Santos et al., 2016), a preliminary test with our data indicated a linear model would be

more suitable (smoothing terms in GAMs had $\text{edf} = 1$; Thomas et al. 2015). Thus, we used Generalized Linear Mixed Models (GLMM) with Poisson distribution (for reptile abundance and species richness) and Gaussian distribution (for diversity) and used “Visit” (for reptile abundance) and “Year” (for species richness and evenness) as random effects. Altitude was also included as a random effect using three levels of elevation (< 600 , $600-900$ and > 900 m). The levels of elevation were chosen according to the climate in the PNSE. Below 900 m, the climate is typically meso-Mediterranean, while between 900 and 1250 m the climate is supra-Mediterranean. We divided the meso-Mediterranean climate level to have a similar number of transects in each level. The vegetation composition also changes along this altitude gradient.

Fire tends to modify the composition of reptile communities (Santos and Cheylan, 2013). For this reason, we computed differences in reptile communities (using species-specific abundance data) between pairs of sampling transects using the Bray-Curtis similarity index. This index allowed a comparison of the number and abundance of reptile species shared between transects. From the pairwise similarity matrix, we performed an Analysis of Similarity, ANOSIM (Primer, 2001), which gives a general R-value and allows comparisons in reptile communities between transect classes. For the ANOSIM, transects were classified according to the dominant land-use type (monoculture, scrubland and natural woodland), and time-since-fire interval, namely recently burned (< 5 years since the last fire), medium (5-10 years since the last fire), long (10-20 years since the last fire) and long-unburned (> 20 years since the last fire) (Supplementary materials Table S3). GLMM and ANOSIM are complementary approaches: GLMM examined the effect of environmental variables on reptile metrics and species-specific abundances of two lizard species, whereas ANOSIM compared matrices of species abundances between transect classes.

Only two lizard species were found in high numbers to allow specific statistical analyses. Thus, the environmental and fire variables were used to test their effects on the abundance of the two commonest lizard species of the study area, *Psammodromus algirus* and *Podarcis guadarramae*. We first ran GLMM with all the factors, selected the best model according to AIC values using backward selection via “drop1” command, and re-ran the best model with the selected environmental and fire variables to examine the significance effect on lizard abundances.

At the micro-habitat scale, we used geo-referenced reptile locations to test the effects of plant richness and vegetation cover (tree, shrub, and herbaceous cover) at each 20-m section on the probability of reptile presence, reptile abundance, and reptile richness. For this, we used GLMM with “Transect” as the random effect, and binomial distribution (for probability of reptile presence) or Poisson distribution (for reptile abundance and species richness).

All the analyses were performed using the R software packages ggplot2, car, MuMIn, gamm4, and lme4 (R Development Core Team, 2015).

3. Results

In total, we observed 433 reptiles belonging to 14 species (70% of the total species occurring in the Natural Park; see Supplementary Materials Table S2). The ground-dwelling lizard *Psammodromus algirus* was the most abundant species representing 83% of all sightings and was detected in all transects. The second most common species was the wall lizard *Podarcis guadarramae*, representing ca. 6% of the sightings and detected in 55% of surveyed transects.

At the landscape scale, vegetation heterogeneity and plant richness positively influenced reptile abundance and evenness (Table 1; Figure 1). Likewise, reptile

abundance differed among land-use types, as natural woodlands had the highest abundance whereas monocultures showed the lowest abundance (GLMM with "Visit" as random effect; $F_{2,14} = 3.76$, $P = 0.05$; Figure 2). Reptile evenness declined with time since fire, while fire recurrence had no significant effect on any of the reptile metrics (Figure 1). Also, none of the variables considered exerted any effect on species richness (Table 1). Regarding differences in the reptile community composition (species presence and abundance), the ANOSIM showed differences between the three land-use types ($R = 0.155$, $P = 0.044$), especially between natural and scrubland transects ($R = 0.22$, $P = 0.036$). Differences were also observed according to time-since-fire intervals ($R = 0.227$, $P = 0.013$), especially between recently and medium-burnt transects ($R = 0.445$, $P = 0.007$).

The abundance of *P. algirus* was positively affected by both vegetation heterogeneity and the plant richness in the transects, whereas time since fire and fire recurrence did not significantly affect its abundance (Table 2; Figure 3). The abundance of *P. gadarramae* was affected by fire recurrence (positive effect) and the time since fire (negative effect; Table 2); that is, this lizard was more abundant in open habitats as those produced by recurrent and recent fires. Moreover, *P. gadarramae* was negatively affected by vegetation heterogeneity and positively affected by plant richness (Table 2; Figure 3).

At the micro-habitat scale, both the probability of reptile occurrence, reptile abundance and species richness were positively related to plant richness and negatively related to tree cover (Table 3; Figure 4), suggesting a major effect of the micro-habitat on reptiles.

4. Discussion

This study combined the use of variables at the landscape and micro-habitat scales to understand the response of a Mediterranean reptile community in a fire-prone region of the Iberian Peninsula. We acknowledge that the sampling methodology and effort were not enough for keeping adequate data for a complete view of the overall reptile community response to fire. In our study area, many species (especially snakes) are rare and difficult to detect. However, our results are reasonably adequate for common lizard species (see similar conclusions in Australian Banksia woodlands, Hu et al., 2016).

Despite the fire history of the study area, fire frequency and time since fire only partially explained the reptile species community metrics. Specifically, reptile evenness decreased with the time since fire. However, species-specific analyses demonstrated opposing responses between lizard species. This result suggests that Mediterranean reptiles can present diverse and complex responses to fire (see also similar conclusions in Pastro et al., 2013) that can be masked by the relevance of habitat factors, and corroborates the sensitivity of reptiles to changes in habitat complexity (Santos et al., 2016). Additional variables such as fire return interval, percentage of burned area and fire intensity/severity should be included in future modeling efforts.

4.1. Community-metric results

Most of the reptiles observed in our study were Mediterranean lizard species (Sillero et al., 2009). These lizards typically prefer open habitats (Ferreira et al., 2017) in part due to their physiological capacity to reduce water loss (Ferreira et al., 2016a). Accordingly, we could expect positive responses at recently burnt habitats as previously observed in other regions (Driscoll and Henderson, 2008; Ashton and Knipps, 2011; Driscoll et al., 2012). In our study, reptile (primarily lizards) abundance was strongly related to habitat quality both in terms of the heterogeneity and plant richness (Fischer et al., 2004; Pastro

et al., 2013), as well as land-use type. This result confirms that non-natural land uses, with low plant diversity, such as monocultures can be inadequate for Mediterranean reptiles (Azor et al., 2015) most likely due to their low thermal quality (Schreuder and Clusella-Trullas, 2016). Thus, the strong effect of land use (e. g., pine plantation) can be masking fire effects. Accordingly, in regions dominated by monocultures, land use can be a stronger driver of reptile abundance/community composition than fire history.

Surprisingly, none of the variables studied turned out to affect reptile richness. This may be due to species replacement between transects of different fire histories (Santos et al., 2016), or because the analysis was confounded by the overall low species richness on transects. We acknowledge that other sampling methods such as drift fences might be less biased towards a few abundant lizard species. However, the active searching was effective in other Mediterranean areas giving good results in terms of species richness and diversity (e. g. Santos and Cheylan, 2013).

We found a significant association between reptile evenness, fire history (time since fire) and habitat (vegetation heterogeneity) variables, as long-unburnt sites held less complex reptile communities (lower evenness) than did recently burnt ones. This result contrast with other studies showing that long-unburnt natural woodlands had more diverse and rich reptile communities than burnt sites (Santos and Cheylan, 2013; Ferreira et al., 2016b). In our study area (SENP), long-unburnt sites were often monocultures with a dense canopy and low understory that held a low reptile diversity. This exemplifies the importance of the interaction effect of fire history and land use to understand the reptile responses to fire in highly human-modified landscapes such as the Mediterranean basin.

4.2. Micro-habitat scale effects on community metrics

The micro-habitat scale (i.e. habitat characteristics at the location where reptiles were observed within the transect, at each 20 m section) more clearly explained the patterns of reptile occurrence, than the analysis at broader scale, a pattern demonstrated in previous studies (Driscoll and Henderson, 2008; Santos and Poquet, 2010). Our results showed that the probability of reptile presence increased with the richness of plant species. A similar result was observed for reptile abundance at the landscape scale and confirmed that reptile community patterns in response to environmental disturbance can operate at different scales (Ferreira et al., 2017). In parallel to the importance of solar radiation for reptile thermoregulation (Webb and Shine, 2008; Schreuder and Clusella-Trullas, 2016), the presence of different plant species can offer different niches for reptiles and can increase food provision (herbaceous plants and scrubs attract many pollinators; Potts et al. 2003), shelter, and protection against predators (Santos et al., 2016).

Contrary to plant richness, tree cover had a negative effect on all the reptile community metrics (Ferreira et al., 2017). Other studies have also demonstrated a strong connection between thermal properties of the habitat and reptile habitat selection in the Mediterranean (Ferreira et al., 2017) and elsewhere (Martín and Salvador, 1995; Meik et al., 2002), and finally reptile performance (Sears and Angilletta, 2015). Our study indicates that these environmental properties at the micro-habitat scale are relevant to understand the occurrence patterns of reptiles in fire-prone regions.

4.3. Species-specific responses: *P. algirus* and *P. guadarramae* abundance

Psammodromus algirus responded positively to the habitat heterogeneity, increasing its abundance while the habitat increased its complexity. This lizard is a ground-dwelling specialist and habitat generalist (Carrascal and Díaz, 1989; Salvador, 2015). It is able to

successfully adapt to areas affected by fire (Santos and Poquet, 2010) and it is also common in long-unburnt transects, both in natural woodlands and in monocultures (this study).

Contrary to *P. algirus*, the abundance of *Podarcis guadarramae* was driven by a number of variables, and confirmed its habitat specialization in rocky spots (Diego-Rasilla and Perez-Mellado, 2003). The extreme structural changes caused by recurrent (or recent) fires provided conditions (open rocky spots) that favored this wall lizard (Ferreira, 2016b; Ferreira et al., 2018). Thus, habitat specialization merits consideration in evaluating the effects of fire on reptiles (Hu et al., 2016; Santos et al., 2016).

4.4. Conservation remarks

The study site has been historically affected by fires, and fire frequency, intensity and extent have increased remarkably in recent years. However, we found that fire history factor (particularly fire recurrence) had little impact on the reptile communities. Indeed, the rock lizard *P. guadarramae* was more abundant in open habitats produced by recurrent and recent fires. On the contrary, habitat characteristics such as vegetation complexity (positive effect) and land transformation to monocultures (negative effect) were likely important drivers of reptile occurrence in fire-prone areas. Monocultures have long been a source of economic income in SENP as well as in many other areas of northern Portugal. However, monocultures are inadequate habitat for Mediterranean reptiles (Azor et al., 2015), and if they replace late successional (mature) habitats, local extinctions of some reptile species can occur (Driscoll et al., 2012; Valentine et al., 2012). The SENP region supports the highest herpetofaunal biodiversity in Portugal (only paralleled by Peneda-Gerês National Park) and thus, should be a national priority for the conservation of the reptile biodiversity. The development of management programs to

promote pyrodiversity could be an adequate practice to enhance reptile diversity, as previously observed in other vertebrate taxa (Tingley et al., 2016; Kelly and Brotons, 2017).

Acknowledgments

We thank to J. Conde for the support provided in the beginning of this study. ICNF provided the permit for sampling reptiles in the Serra da Estrela Natural Park (number 22327). We thank the comments of two anonymous reviewers that improved an early version of the manuscript. This study was partially funded by a British Herpetological Society Student Grant Scheme awarded to TP. BM was supported by the Fundação para a Ciência e a Tecnologia [grant number SFRH/BPD/90277/2012].

References

- Ashton, K.G., Knipps, A.C.S., 2011. Effects of fire history on amphibian and reptile assemblages in Rosemary Scrub. *J. Herpetol.* 45, 497–503.
- Azor, J.S., Santos, X., Pleguezuelos, J.M., 2015. Conifer-plantation thinning restores reptile biodiversity in Mediterranean landscapes. *For. Ecol. Manage.* 354, 185-189.
- Badiane, A., Matos, C., Santos, X., 2017. Uncovering environmental, land-use and fire effects on the distribution of a lowdispersal species, the Hermann's tortoise *Testudo hermanni*. *Amph.-Rept.* 38, 67-77.
- Baselga, A., Jiménez-Valverde, A., Niccolini, G., 2007. A multiple-site similarity measure independent of richness. *Biol. Letters* 3, 642-645.
- Blondel, J., Aronson, J., Bodiou, J.-Y., Boeuf, W., 2010. The Mediterranean region. Biological biodiversity in space and time. Oxford University Press, Oxford.

401 Bond, W.J., Woodward, F.I., Midgley, G.F., 2005. The global distribution of
 402 ecosystems in a world without fire. *New Phytol.* 165, 525-538.

403 Bowman, D.M.J.S., Balch, J.K., Artaxo, P., Bond, W.J., Carlson, J.M., Cochrane, M.A.,
 404 D'Antonio, C.M., DeFries, R.S., Doyle, J.C., Harrison, S.P., Johnston, F.H.,
 405 Keeley, J.E., Krawchuk, M.A., Kull, C.A., Marston, J.B., Moritz, M.A., Prentice,
 406 I.C., Roos, C.I., Scott, A.C., Swetnam, T.W., van der Werf, G.R., Pyne, S.J.,
 407 2009. Fire in the Earth System. *Science* 324(5926), 481-484.

408 Carrascal, L.M., Díaz, J.A., 1989. Thermal ecology and spatio-temporal distribution of
 409 the Mediterranean lizard *Psammmodromus algirus*. *Ecography* 12, 137-143.

410 Catry, F., Rego, F.C., Bugalho, M.N., Lopes, T., Silva, J.S., Moreira, F., 2006. Effects of
 411 fire on tree survival and regeneration in a Mediterranean ecosystem. *Forest Ecol.*
 412 *Managem.* 234, S197.

413 Catry, F.X., Rego, F.C., Bação, F.F., Moreira, F., 2009. Modelling and mapping wildfire
 414 ignition risk in Portugal. *Int. J. Wildl. Fire* 18, 921-931.

415 Connor, S.E., Araújo, J., van der Knaap, W.O., van Leeuwen, J.F., 2012. A long-term
 416 perspective on biomass burning in the Serra da Estrela, Portugal. *Quarter. Science*
 417 *Rev.* 55, 114-124.

418 Couturier, T., Cheylan, M., Guérette, E., Besnard, A., 2011. Impacts of a wildfire on the
 419 mortality rate and small-scale movements of a Hermann's tortoise *Testudo*
 420 *hermanni hermanni* population in southeastern France. *Amph.-Rept.* 32, 541-545.

421 Diego-Rasilla, F.J., Perez-Mellado, V., 2003. Home range and habitat selection by
 422 *Podarcis hispanica* (Squamata, Lacertidae) in Western Spain. *Folia Zool.*
 423 *Praha*, 52, 87-98.

424 Driscoll, D.A., Henderson, M.K., 2008. How many common reptile species are fire
 425 specialists? A replicated natural experiment highlights the predictive weakness of
 426 a fire succession model. *Biol. Conserv.* 141, 460-471.

427 Driscoll, D.A., Smith, A.L., Blight, S., Maindonald, J., 2012. Reptile responses to fire
 428 and the risk of post-disturbance sampling bias. *Biodivers. Conserv.* 21, 1607-1625.

429 Elzer, A.L., Pike, D.A., Webb, J.K., Hammill, K., Bradstock, R.A., Shine, R., 2013.
 430 Forest-fire regimes affect thermoregulatory opportunities for terrestrial ectotherms.
 431 *Austral Ecol.* 38, 190-198.

432 Farji-Brener, A.G., Corley, J.C., Bettinelli, J., 2002. The effects of fire on ant
 433 communities in north-western Patagonia: the importance of habitat structure and
 434 regional context. *Divers. Dist.* 8, 235-243.

435 Fernandes, P.M., Davies, G.M., Ascoli, D., Fernández, C., Moreira, F., Rigolot, E.,
 436 Stoof, C.R., Veja, J.A., Molina, D., 2013. Prescribed burning in southern
 437 Europe: developing fire management in a dynamic landscape. *Frontiers Ecol.*
 438 *Envir.* 11(s1).

439 Ferreira, C., Santos, X., Carretero, M.A., 2016a. Does ecophysiology mediate reptile
 440 responses to fire regimes? Evidence from Iberian lizards. *PeerJ* 4, e2107.

441 Ferreira, D., Brito, J.C., Santos, X., 2018. Long-interval monitoring reveals changes in
 442 the structure of a reptile community in a biogeographic transition zone. *Basic and*
 443 *Applied Herpetology*. In press.

444 Ferreira, D., Mateus, C., Santos, X., 2016b. Responses of reptiles to fire in transition
 445 zones are mediated by bioregion affinity of species. *Biodivers. Conserv.* 25, 1543-
 446 1557.

447 Ferreira, D., Žagar, A., Santos, X., 2017. Uncovering the rules of (reptile) species
 448 coexistence in transition zones between bioregions. *Salamandra* 53, 193-200.

449 Fischer, J., Lindenmayer, D.B., Cowling, A., 2004. The challenge of managing multiple
 450 species at multiple scales: reptiles in an Australian grazing landscape. *J. Appl.*
 451 *Ecol.* 41, 32-44.

452 Hu, Y., Kelly, L.T., Gillespie, G.R., Jessop, T.S. 2016. Lizard responses to forest fire
 453 and timber harvesting: Complementary insights from species and community
 454 approaches. *For. Ecol. Manage.* 379, 206–215.

455 Hu, Y., Urlus, J., Gillespie, G., Letnic, M., Jessop, T.S., 2013. Evaluating the role of
 456 fire disturbance in structuring small reptile communities in temperate
 457 forests. *Biodivers. Conserv.* 22, 1949-1963.

458 Huey, R.B., 1982. Temperature, physiology, and the ecology of reptiles, in: Gans, C.,
 459 Pough, F.H., Eds., *Biology of the Reptilia Vol. 12, Physiology (C)*. Academic
 460 Press, London, pp. 25-91.

461 IGP, 2008. Carta de Ocupação do Solo – COS 2007_N5 (1:25,000) (Land cover map –
 462 COS 90). Portuguese Geographic Institute (IGP),
 463 < <http://www.igeo.pt/produtos/CEGIG/COS.htm> >.

464 Keeley, J.E., Bond, W.J., Bradstock, R.A., Pausas, J.G., Rundel, P.W., 2012. Fire in
 465 Mediterranean Ecosystems: Ecology, Evolution and Management. Cambridge
 466 University Press, Cambridge.

467 Kelly, L.T., Brotons, L., 2017. Using fire to promote biodiversity. *Science* 355 (6331),
 468 1264-1265.

469 Kim, T.N., Holt, R.D., 2012. The direct and indirect effects of fire on the assembly of
 470 insect herbivore communities: examples from the Florida scrub habitat. *Oecologia*
 471 168, 997-1012.

472 Kutt, A.S., Bateman, B.L., Vanderduys, E.P., 2011. Lizard diversity on a rainforest–
 473 savanna altitude gradient in north-eastern Australia. *Austral. J. Zool* 59, 86-94.

474 Lesparre, D., Crespo, E.G., 2008. A herpetofauna do Parque Natural da Serra da Estrela.
 475 CISE, Seia, Portugal.

476 Lindenmayer, D.B., Wood, J.T., MacGregor, C., Michael, D.R., Cunningham, R.B.,
 477 Crane, M., Montague-Drake, R., Brown, D., Muntz, R., Driscoll, D.A., 2008. How
 478 predictable are reptile responses to wildfire? *Oikos* 117, 1086-1097.

479 Martín, J., Salvador A., 1995. Microhabitat selection by the Iberian rock lizard *Lacerta*
 480 *monticola*: effects on density and spatial distribution of individuals. *Biol.*
 481 *Conserv.* 79, 303–307.

482 McCain, C.M., 2010. Global analysis of reptile elevational diversity. *Global Ecol*
 483 *Biogeog* 19, 541-553.

484 Meik, J.M., Jeo, R.M. Mendelson J.R., Jenks K.E., 2002. Effects of bush encroachment
 485 on an assemblage of diurnal lizard species in central Namibia. *Biol. Conserv.* 106,
 486 29–36.

487 Moreira, F., Rego, F.C., Ferreira, P.G., 2001. Temporal (1958–1995) pattern of change
 488 in a cultural landscape of northwestern Portugal: implications for fire
 489 occurrence. *Land. Ecol.* 16, 557-567.

490 Moreira, F., Russo, D., 2007. Modelling the impact of agricultural abandonment and
 491 wildfires on vertebrate diversity in Mediterranean Europe. *Landscape Ecol.* 22,
 492 1461-1476.

493 Moreira, F., Vaz, P., Catry, F., Silva, J.S., 2009. Regional variations in wildfire
 494 susceptibility of land-cover types in Portugal: implications for landscape
 495 management to minimize fire hazard. *Int. J. Wildl. Fire* 18, 563-574.

496 Moreira, F., Viedma, O., Arianoutsou, M., Curt, T., Koutsias, N., Rigolot, E., Barbati,
 497 A., Corona, P., Vaz, P., Mouillot, F., 2011. Landscape–wildfire interactions in

498 southern Europe: implications for landscape management. *J. Envir. Manage.* 92,
499 2389-2402.

500 Nimmo, D.G., Kelly, L.T., Farnsworth, L.M., Watson, S.J., Bennett, A.F., 2014. Why
501 do some species have geographically varying responses to fire history? *Ecography*
502 37, 805-813.

503 Nimmo, D.G., Kelly, L.T., Spence-Bailey, L.M., Watson, S.J., Haslem, A., White, J.G.,
504 Clarke, M.F., Bennett, A.F., 2012. Predicting the century-long post-fire responses of
505 reptiles. *Global Ecol. Biogeog.* 21, 1062-1073.

506 Nimmo, D.G., Mac Nally, R., Cunningham, S.C., Haslem, A., Bennett, A.F., 2015. Vive
507 la résistance: reviving resistance for 21st century conservation. *TREE* 30, 516-
508 523.

509 Nunes, M.C., Vasconcelos, M.J., Pereira, J.M., Dasgupta, N., Alldredge, R.J., Rego,
510 F.C., 2005. Land cover type and fire in Portugal: do fires burn land cover
511 selectively? *Land. Ecol.* 20, 661-673.

512 Oliveira, S.J., Pereira, J.M.C., Carreiras, J.M. 2012. Fire frequency analysis in Portugal
513 (1975-2005) using Landsat-based burnt area maps. *Int. J. Wildl. Fire* 21, 48-60.

514 Pastro, L.A., Dickman, C.R., Letnic, M., 2013. Effects of wildfire, rainfall and region
515 on desert lizard assemblages: the importance of multi-scale processes. *Oecologia*
516 173, 603-614.

517 Pausas, J.G., 2004. Changes in fire and climate in the eastern Iberian Peninsula
518 Mediterranean Basin. *Climatic Change* 63, 337–350.

519 Pausas, J.G., Fernández-Muñoz, S., 2012. Fire regime changes in the Western
520 Mediterranean Basin: from fuel-limited to drought-driven fire regime. *Climatic*
521 *Change* 110, 215-226.

522 Pausas, J.G., Keeley, J.E., 2009. A burning story: the role of fire in the history of
523 life. *BioScience* 59, 593-601.

524 Potts, S.G., Vulliamy, B., Dafni, A., Ne'eman, G., O'Toole, C., Roberts, S., Willmer, P.,
525 2003. Response of plant-pollinator communities to fire: changes in diversity,
526 abundance and floral reward structure. *Oikos*, 101, 103-112.

527 PRIMER-E, 2001. Primer 5 for Windows version 5.2.2. PRIMER-E Ltd.

528 R Core Team, 2015. R: A language and environment for statistical computing. R
529 Foundation for Statistical Computing, Vienna, Austria. < [http://www.R-](http://www.R-project.org/)
530 [project.org/](http://www.R-project.org/)>.

531 Ribeiro, R., Santos, X., Sillero, N., Carretero, M.A., Llorente, G.A., 2009. Biodiversity
532 and land uses: Is agriculture the biggest threat in reptiles' assemblages? *Acta*
533 *Oecol.* 35, 327-334.

534 Salvador, A., 2015. Lagartija colilarga – *Psammodromus algirus*, in: Salvador, A.,
535 Marco, A. (Eds.), *Enciclopedia Virtual de los Vertebrados Españoles*. Museo
536 Nacional de Ciencias Naturales, Madrid.

537 Santos, X., Badiane, A., Matos, C., 2016. Contrasts in short-term and long-term
538 responses of Mediterranean reptile species to fire and habitat structure. *Oecologia*
539 180, 205-2016.

540 Santos, X., Cheylan, M., 2013. Taxonomic and functional response of a Mediterranean
541 reptile assemblage to a repeated fire regime. *Biol. Conserv.* 168, 90-98.

542 Santos, X., Mateos, E., Bros, V., Brotons, L., De Mas, E., Herraiz, J.A., Herrando, S.,
543 Miño, A., Olmo-Vidal, J.M., Quesada, J., Ribes, J., Sabaté, S., Sauras-Yera, T.,
544 Serra, A., Vallejo, V.R., Viñolas, A., 2014. Is response to fire influenced by
545 dietary specialization and mobility? A comparative study with multiple animal
546 assemblages. *PloS ONE* 9(2), e88224.

547 Santos, X., Poquet, J.M., 2010. Ecological succession and habitat attributes affect the
 548 postfire response of a Mediterranean reptile community. *Eu. J. Wild. Res.* 56,
 549 895-905.

550 Schreuder, E., Clusella-Trullas, S., 2016. Exotic trees modify the thermal landscape and
 551 food resources for lizard communities. *Oecologia* 182, 1213–1225.

552 Sears, M.W., Angilletta Jr, M.J., 2015. Costs and benefits of thermoregulation revisited:
 553 both the heterogeneity and spatial structure of temperature drive energetic costs.
 554 *Am. Nat.* 185, E94–E102.

555 Segura, C., Feriche, M., Pleguezuelos, J.M., Santos, X., 2007. Specialist and generalist
 556 species in habitat use: implications for conservation assessment in snakes. *J. Nat.*
 557 *Hist.* 41, 2765-2774.

558 Sillero, N., Brito, J. C., Skidmore, A.K., Toxopeus, A.G., 2009. Biogeographical
 559 patterns derived from remote sensing variables: the amphibians and reptiles of the
 560 Iberian Peninsula. *Amph.-Rept.* 30, 185-206.

561 Silva, J.S., Vaz, P., Moreira, F., Catry, F., Rego, F.C., 2011. Wildfires as a major driver
 562 of landscape dynamics in three fire-prone areas of Portugal. *Landscape Urban*
 563 *Planning* 101, 349-358.

564 Smith, A.L., Bull, C.M., Driscoll, D.A., 2013. Successional specialization in a reptile
 565 community cautions against widespread planned burning and complete fire
 566 suppression. *J. Appl. Ecol.* 50, 1178-1186.

567 Smith, L.J., Holycross, A.T., Painter, C.W., Douglas, M.E., 2001. Montane rattlesnakes
 568 and prescribed fire. *Southwest. Nat.* 46, 54-61.

569 SNIRH, 2015. Sistema Nacional de Informação de Recursos Hídricos. <www.snirh.pt>

570 Thomas, R., Vaughan, I., Lello, J., 2015. Data analysis with R statistical software. A
 571 guidebook for scientists. Wales, Eco-explore.

572 Tingley, M.W., Ruiz-Gutiérrez, V., Wilkerson, R.L., Howell, C.A., Siegel, R.B., 2016.
 573 Pyrodiversity promotes avian diversity over the decade following forest fire. *Proc.*
 574 *Royal Soc. B* 283, 20161703.
 575 Valentine, L.E., Reaveley, A., Johnson, B., Fisher, R., Wilson, B.A., 2012. Burning in
 576 banksia woodlands: how does the fire-free period influence reptile
 577 communities? *PLoS ONE* 7, e34448.
 578 Valentine, L.E., Schwarzkopf, L., 2008. Effects of Weed-Management Burning on
 579 Reptile Assemblages in Australian Tropical Savannas. *Conserv. Biol.* 23, 103-
 580 113.
 581 Webb, J.K., Shine, R., 2008. Differential effects of an intense wildfire on survival of
 582 sympatric snakes. *J. Wild. Manage.* 72, 1394-1398.
 583

Legends of Tables and Figures

Table 1. Results of the GLMM analysis for three reptile-community metrics: Reptile abundance, Species richness, and Pielou's evenness. Only the results for the best model selected are presented.

Table 2. Results of the GLMM analysis for the abundance of *Psammodromus algirus* and *Podarcis gadarramae*. Only results for the best model selected are presented.

Table 3. Results of the GLMM analysis for the Probability of reptile presence, reptile abundance and species richness in relation to vegetation cover (tree, scrub and herb) and plant richness. Only results for the best model selected are presented.

Figure 1. Significant Generalized Linear Mixed Models' association between reptile metrics (abundance and evenness) and predictive variables plant richness, time-since-fire and vegetation heterogeneity) at Serra de Estrela Natural Park.

Figure 2. Mean and standard error scores of the reptile abundance per transect in relation to land-use type.

Figure 3. Significant Generalized Linear Mixed Models' association between *P. algirus* and *P. gadarramae* abundances and plant richness, vegetation heterogeneity, time since fire, fire recurrence, and plant-richness predictive variables at Serra de Estrela Natural Park.

Figure 4. Relationship between plant richness and reptile abundance (a), and reptile richness (b) at the micro-habitat scale.

604 **Table 1.**

	Reptile abundance			Species richness			Pielou's evenness		
	Estimate	Z value	P	Estimate	Z value	P	Estimate	t value	P
Fire recurrence		n.s.			n.s.			n.s.	
Time since fire		n.s.			n.s.		-0.01	-2.25	0.020
Vegetation heterogeneity	1.13	1.89	0.060		n.s.		-1.16	-2.03	0.040
Plant richness	0.06	2.64	0.008		n.s.			n.s.	
Land use type		n.s.			n.s.			n.s.	
Land-use diversity		n.s.			n.s.		0.51	1.60	n.s.

605

606 **Table 2.**

	<i>P. algirus</i> abundance			<i>P. guadarramae</i> abundance		
	Estimate	Z value	<i>P</i>	Estimate	Z value	<i>P</i>
Fire recurrence		n.s.		0.49	2.10	0.04
Time since fire		n.s.		-0.06	-2.61	0.009
Vegetation heterogeneity	1.94	3.09	0.002	-6.55	-2.55	0.010
Plant richness	0.06	2.71	0.007	0.42	2.61	0.009
Land-use type		n.s.			n.s.	
Land-use diversity		n.s.			n.s.	

607

609 **Table 3.**

	Reptile Presence			Reptile Abundance			Species Richness		
	Estimate	Z value	P	Estimate	Z value	P	Estimate	Z value	P
Tree Cover	-0.48	-1.82	0.07	n.s.			-0.29	-1.72	0.09
Scrub Cover		n.s.		n.s.			n.s.		
Herb Cover	-0.54	-1.84	0.07	n.s.			n.s.		
Plant Richness	0.12	2.55	0.01	0.06	2.21	0.03	0.07	2.57	0.01

610

The role of fire history, land-use and vegetation structure on the response of Mediterranean lizard communities to fire

Tiago Pinto, Bruno Moreira, Helena Freitas, Xavier Santos

Supplementary Materials

Supplementary materials Figure S1. Location of the Serra da Estrela Natural Park in Portugal (A), and distribution of the sites selected (black triangles) according to the land uses (B), number of fires (C) and time since the last fire (D).

Supplementary materials Table S2. Number of individuals (N) and percentage of transects where each reptile species was observed.

Family	Species	N	%
Anguidae	<i>Anguis fragilis</i>	1	5
Amphisbaenidae	<i>Blanus cinereus</i>	1	5
Scincidae	<i>Chalcides</i> sp.	14	30
Lacertidae	<i>Lacerta schreiberi</i>	4	10
Lacertidae	<i>Podarcis carbonelli</i>	1	5
Lacertidae	<i>Podarcis guadarramae</i>	23	55
Lacertidae	<i>Psammodromus algirus</i>	361	100
Lacertidae	<i>Psammodromus hispanicus</i>	2	10
Lacertidae	<i>Timon lepidus</i>	17	45
Colubridae	<i>Coronella girondica</i>	3	15
Colubridae	<i>Malpolon monspessulanus</i>	1	5
Colubridae	<i>Hemorrhois hippocrepis</i>	1	5
Colubridae	<i>Rhinechis scalaris</i>	2	10
Viperidae	<i>Vipera latastei</i>	2	10

Supplementary materials Table S3. Location, fire-history, elevation and land-use type of the 20 surveyed transects.

Transect code	Longitude	Latitude	TSF intervals (years)	Fire recurrence	Elevation (m)	Land-Use
0F2	-7712606	40386645	>40	0	686	Natural woodland
0F3	-7533153	40513008	>40	0	677	Natural woodland
0F4	-7518362	40496794	>40	0	1097	Monoculture
A1	-7513324	4054926	20-38	2	464	Scrubland
A2	-780024	40303941	20-38	1	437	Monoculture
A3	-7856275	40291665	20-38	1	524	Monoculture
B1	-7594615	40475461	10-20	1	822	Natural woodland
B2	-758698	40464968	10-20	3	983	Natural woodland
B4	-7714982	40332528	10-20	1	857	Monoculture
C1_1	-7772071	40301344	5-10	1	470	Scrubland
C1_3	-7686587	40302533	5-10	1	978	Natural woodland
C1_4	-7690771	4029921	5-10	1	769	Natural woodland
C2_3	-7518873	40542699	5-10	4	510	Scrubland
C2_4	-7597737	40456767	5-10	3	417	Scrubland
D1_1	-7650357	40402264	<5	1	1027	Scrubland
D1_2	-7726969	40366441	<5	1	581	Monoculture
D1_5	-7867992	40413481	<5	1	272	Natural woodland
D2_4	-7597737	40456767	<5	6	964	Scrubland
D2_5	-7540406	40507338	<5	3	852	Scrubland
D2_6	-7668131	40407816	<5	4	1018	Scrubland

Supplementary materials Figure S4. Pictures of transects in two different land-use types, namely monoculture and scrubland.