Delineation of the olive pollen proteome and its allergenome unmasks cyclophilin as a relevant cross-reactive allergen

Pablo San Segundo-Acosta^{‡, #}, Carmen Oeo-Santos^{‡, #}, Sara Benedé[‡], Vivian de los Ríos[§], Ana Navas[¶], Berta Ruiz-Leon[¶], Carmen Moreno[¶], Carlos Pastor-Vargas[¢], Aurora Jurado[¶], Mayte Villalba^{‡, ^}, Rodrigo Barderas^{*, ^, &}

[#], Both authors contributed equally. [^], Co-senior authors

[‡]Departamento de Bioquímica y Biología Molecular, Facultad de Ciencias Químicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain

[§]Centro de Investigaciones Biológicas, CSIC, E-28040 Madrid, Spain

[¶]Hospital Universitario Reina Sofía de Córdoba, E-14004 Córdoba, Spain

[¢]Hospital Fundación Jiménez Díaz, E-28040 Madrid, Spain

[&]Chronic Disease Programme (UFIEC), Instituto de Salud Carlos III, Majadahonda E-28220, Madrid, Spain

^{*}To whom correspondence should be addressed

***Corresponding Author:** Rodrigo Barderas, Functional Proteomics Unit, Chronic Disease Programme (UFIEC), Instituto de Salud Carlos III. Majadahonda, Madrid. Spain. 28220.

E-mail: r.barderasm@isci.es

Abstract

Olive pollen is a major allergenic source worldwide for its extensive cultivation. We have combined available genomic data with a comprehensive proteomic approach to get the annotated olive tree (*Olea europaea* L.) pollen proteome and define its complex allergenome. A total of 1,907 proteins were identified by LC-MS/MS using predicted protein sequences from its genome. Most proteins (60%) were predicted to possess catalytic activity and be involved in metabolic processes. In total, 203 proteins belonging to 47 allergen families were found in olive pollen. A peptidyl-prolyl *cis-trans* isomerase –cyclophilin- produced in *Escherichia coli*, was found as a new olive pollen allergen (Ole e 15). Most Ole e 15-sensitized patients were children (63%) and showed strong IgE recognition to the allergen. Ole e 15 shared high sequence identity with other plant, animal and fungal cyclophilins and presented high IgE cross-reactivity with pollen, plant food and animal extracts.

KEYWORDS: *Olive pollen proteome, allergenome, in-depth proteomics, allergen, cyclophilin, cross-reactivity*

INTRODUCTION

The pollen grain has aroused a lot of interest not only for its role as male gametophyte in reproduction (1), but also for its major role as allergen source. Proteomic studies have shown that proteins acting in pollen formation and germination are mainly involved in energy production, defense and protein synthesis and processing (2-4). Moreover, as remarked above, some pollen proteins are capable of triggering airway inflammation and antigenic responses, which in some individuals derive in allergy (5-8). For this reason, proteomic analysis of pollen can be used for allergen identification, as it has been done for several allergenic sources (9-13). However, there is limited genomic information of tree species with allergenic pollen, reducing the potential number of these studies.

Recently, the complete genome of the wild olive (*Olea europaea* subsp. *europaea* var. *sylvestris* -also denominated as oleaster-), a closely related variant of cultivated olive (*O. europaea* subsp. *europaea* var. *europaea*) has been reported (14). Cultivated olive belongs to the Oleaceae family (order Lamiales) and has been traditionally cultivated in the Mediterranean Basin, being the main crop in some areas of Spain, Italy and Greece (15). In addition, olive trees are extensively cultivated worldwide for olive oil and olive fruit production. Several regions of these countries reach extremely high olive pollen counts in the pollination season, becoming olive pollen a major allergenic health problem (16). Until now, thirteen olive pollen allergens, Ole e 1 to Ole e 12 and Ole e 14, have been identified (17-19). However, some allergens remain unidentified due to technical restrictions and limited genomic and proteomic data. Moreover, very little is known about other proteins present in this pollen, which although not allergenic, may well influence the immune response of patients by promoting inflammation (20-22).

In this work, we have performed a comprehensive proteomic analysis of cultivated olive pollen, taking advantage of the wild olive genomic data. By an in-depth bottom-up proteomic approach, olive pollen proteome has been in-depth identified using a Q-exactive mass spectrometer, and functionally catalogued according to Gene Ontology (GO) terms. We next extended our analysis to the definition of the olive pollen allergenome, identifying its allergen

1
2
3 families by searching the proteomic data with the Allergome, World-Health Organization and
4
5 International Union of Immunological Societies (WHO/IUIS) and AllFam (the database of
6
7 allergen families) databases (23-25). Among them, we finally focused the study on the
8
9 verification of the allergenicity of four proteins non-previously studied in olive pollen because
10
11 of their potential involvement in cross-reactivity with other allergenic sources. We found
12
13 cyclophilin, a peptidyl-prolyl *cis-trans* isomerase as a new relevant olive pollen allergen.
14
15 Enzyme-linked immunosorbent assay (ELISA), western blot (WB), and Immunoglobulin E
16
17 (IgE) inhibition assays revealed its significance, high potency and its wide range cross-reactivity
18
19 with pollen and plant-derived food, dust mites and its human counterparts.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

Protein extracts

Commercial olive pollen from Allergon (Batch N°032509801, Ängelholm, Sweden) collected in California (US) from a controlled set of cultivated olive trees belonging to different cultivars was used for protein extract preparation for proteomics. Three parallel protein extractions were performed and used later as independent biological replicates. Briefly, for each biological replicate, 300 mg of olive pollen were separately resuspended in 6 mL of extraction buffer (100 mM Tris-HCl pH 7.3, 150 mM NaCl, 2 mM EDTA, 10% Glycerol, 1% Triton X-100, and 1 mM Phenylmethanesulfonyl fluoride (PMSF) with 1x Complete Protease Inhibitor Cocktail (Roche, Basel, Switzerland). After homogenization, the resultant emulsion was subjected to mild stirring for 90 min at room temperature. Then, it was transferred to 2 mL tubes and centrifuged at 12,000 x g for 20 min at 4°C. The insoluble material was resuspended again in 1.5 mL of extraction buffer and stirred for 1 h. After centrifugation, both supernatants were pooled together and a sample analyzed by SDS-PAGE.

Pollen protein extracts and plant-derived food protein extracts were prepared for the other assays according to established protocols (26), and their concentration measured using the Lowry protein assay.

SDS-PAGE, trypsin digestion and LC-MS/MS analysis

Firstly, 40 µg of each extract replicate were separated by 15% SDS-PAGE and stained with Coomassie Brilliant Blue G 250 (Merck, Darmstadt, Germany). Then, whole lanes were cut into four (replicate 1) or six slices (replicates 2 and 3) prior to in-gel trypsin digestion (27, 28). Then, excised bands were cut in small pieces and separately destained with 50 mM ammonium bicarbonate (ABC)/50% acetonitrile (ACN), dehydrated with ACN and dried. Samples were then reduced with 10 mM dithiothreitol in 25 mM ABC and alkylated with iodoacetamide to a final concentration of 50 mM. Then, gel pieces were dried, rehydrated with 12.5 ng/mL porcine trypsin (Thermo Fisher Scientific, Waltham, MA, US) in 50 mM ABC and incubated overnight

1
2
3 at 37 °C. Peptides were extracted using 100% ACN and 0.5% trifluoroacetic acid, purified using
4 a Zip Tip with 0.6 µL C18 resin (Millipore, Sigma-Aldrich Química SL, Madrid, Spain), and
5 dried. Finally, samples were reconstituted in 5 µL 0.1% formic acid/2% ACN before the
6 analysis by nLC-MS/MS in the Proteomics and Genomics Facility of the Centro de
7 Investigaciones Biológicas (CIB-CSIC, Madrid, Spain), a member of ProteoRed-ISCI
8 network.
9
10

11 All peptide separations were carried out on an Easy-nLC 1000 nano system (Thermo Fisher
12 Scientific). For each analysis, the sample was loaded into a precolumn Acclaim PepMap 100
13 (Thermo Fisher Scientific) and eluted in a RSLC PepMap C18, 15 cm long, 50 µm inner
14 diameter and 2 µm particle size (Thermo Fisher Scientific). The mobile phase flow rate was 300
15 nL/min using a gradient of 0.1% formic acid in water (solvent A) and 0.1% formic acid and
16 100% ACN (solvent B). The gradient profile was set as follows: 5%-35% solvent B for 45 min,
17 35%-100% solvent B for 5 min, 100% solvent B for 10 min. Four microliters of each sample
18 were injected.
19
20

21 MS analysis was performed using a Q-Exactive mass spectrometer (Thermo Fisher
22 Scientific). For ionization, 2000 V of liquid junction voltage and 270 °C capillary temperature
23 was used. The full scan method employed a m/z 400–1500 mass selection, an Orbitrap
24 resolution of 70,000 (at m/z 200), a target automatic gain control (AGC) value of $3e6$, and
25 maximum injection times of 100 ms. After the survey scan, the 10 most intense precursor ions
26 were selected for MS/MS fragmentation. Fragmentation was performed with normalized
27 collision energy of 27 and MS/MS scans acquired with a starting mass of m/z 100. AGC target
28 was $2e5$, resolution of 17,500 (at m/z 200), intensity threshold of $8e3$, isolation window of 2
29 m/z units and maximum IT was 100 ms. Charge state screening was enabled to reject
30 unassigned, singly charged, and greater than or equal to seven protonated ions. A dynamic
31 exclusion time of 20s was used to discriminate against previously selected ions.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data processing

MS data were analyzed with Proteome Discoverer (version 1.4.1.14, Thermo Fisher Scientific) using standardized workflows. Mass spectra raw files were searched against the NCBI *Olea europaea* var. *sylvestris* predicted protein database (taxid 158386, version 11/2017, 49,613 RefSeq protein sequence entries) using SEQUEST search engine. Precursor and fragment mass tolerance were set to 10 ppm and 0.02 Da, respectively, allowing 2 missed cleavages, carbamidomethylation of cysteines as a fixed modification, methionine oxidation and threonine, tyrosine and serine phosphorylation as a variable modification. Identified peptides were filtered using Percolator algorithm with a q-value threshold of 0.01 (High Confidence Filter settings, FDR <1%) (29). The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository, with the data set identifiers PXD012334 (Replicate 1), PXD012279 (Replicate 2), and PXD012280 (Replicate 3).

Experimental and statistical design

The experimental and statistical design for each experiment of the study are described in each subsection and depicted in Figure 1 and Figure S1. Only proteins containing at least two isoform-specific peptides in one of the replicates, or with one high-confidence isoform-specific peptide but detected in the three replicates, were considered in the study. To identify proteins belonging to allergen families, the full sequences of the identified proteins after LC-MS/MS were retrieved using NCBI Batch Entrez. Then, proteins were locally blasted (E-value threshold 1.0×10^{-15} and 1 Blast hit) against all entries of the Allergome database (4,331 entries; UniProt release 10/2018) using the Blast2GO 5 PRO software (5.2.4 version) (30). Sequence similarity was calculated by Blast2GO, and sequence identity was calculated using Clustal Omega (31). Only those proteins meeting the following criteria were considered as potential allergens: matching with an allergen accepted by the WHO/IUIS Allergen Nomenclature Sub-committee (25) with at least 50% similarity; matching directly with proteins belonging to the same allergen families as proteins meeting the previous criteria, with at least 80% similarity; matching with proteins already identified as potential allergens in the Oleaceae family.

Gene Ontology (GO) annotation and prediction of secreted proteins

To perform the functional GO annotation and KEGG analysis of each protein, we used the Blast2GO 5 PRO software (5.2.4 version). Firstly, we created a database with the curated NCBI RefSeq predicted proteomes from rice (*Oriza sativa*, taxid:4530), arabidopsis (*Arabidopsis thaliana*, taxid: 3702), latex (*Hevea brasiliensis*, taxid: 3981), barrelclover (*Medicago truncatula*, taxid: 3880), black cottonwood (*Populus trichocarpa*, taxid: 3694), peach (*Prunus persica*, taxid: 3760), tomato (*Solanum lycopersicum*, taxid:4081), potato (*Solanum tuberosum*, taxid: 4113) and common grape vine (*Vitis vinifera*, taxid: 3603). GO annotations for these curated species are deposited in Ensembl Plants (32), and accessible through Blast2GO by BioMart (33). Then, olive pollen identified proteins were locally blasted against this database (maximum number of BLAST hits: 1; minimal E-value: 1.0E-50; BLAST descriptor annotator activated, HPS length cut off: 50, rest of parameters: default). GO mapping (GOA version 09.2018, default settings) and annotation (taxonomy filter: taxa: 33090, Viridiplantae) were performed. Additionally, InterProScan (IPS, default settings) analysis was run (34), and the annotated IPS GOs were merged with previous GOs. Kyoto Encyclopedia of Genes and Genomes (KEGG) Enzyme codes were also annotated. Further information about molecular pathways was obtained by KEGG analysis (35). For Fisher's exact test enrichment analysis of molecular function and biological process GOs, the obtained olive pollen annotation was tested against the whole *O. europaea* var. *sylvestris* predicted proteome, and annotated using the same parameters as described above for pollen proteins. Only GO terms with p-value<1.0E-10 were considered as enriched.

Secreted proteins were predicted using the SignalP 4.1 (Organism group: eukaryotes; D-cutoff value 0.45 for Signal P-noTM and TM networks) (36).

Analytical procedures

1D SDS-PAGE (15% or 17% gels) were stained with Coomassie Blue R-250 (Sigma-Aldrich, St.Louis, MO, US) or transferred onto nitrocellulose membranes. Molecular mass calculations

were performed using unstained protein biomarkers SM0431 (Fermentas, Waltham, MA, US) or pre-stained protein molecular weight markers (Precision Plus Protein All Blue and Prestained Protein Standards low range from Bio-Rad, Hercules, CA, US).

For two-dimensional gel electrophoresis (2DE), isoelectric focusing of 80 µg of olive pollen extract was achieved under reducing conditions with 3 mM tributylphosphine in a PROTEAN IEF Cell (Bio-Rad) using pH 3-10, 7 cm length linear ReadyStrip IPG gels (Bio-Rad). After isoelectrofocusing, proteins were separated by 17% SDS-PAGE under reducing conditions in the presence of 50 mM DTT and 3.7% iodoacetamide and transferred onto nitrocellulose membranes (Amersham Biosciences, Little Chalfont, UK) according to Towbin method for WB.

Protein concentration of the purified proteins was calculated by measuring the absorbance at 280 nm using a DU-7 spectrometer (Beckman, Barcelona, Spain) after theoretical extinction coefficient calculation with the ProtParam tool from ExPASy. The concentration was calculated using the experimental extinction coefficient ($E^{0.1\%}$) of 0.616 obtained after amino acid analysis of 5 µg and 10 µg of purified protein in duplicates with a BioChrom 30 Amino Acid analyzer (Harvard Bioscience, Inc., Holliston, MA, US).

Patients' sera and antibodies

A total of 146 sera of patients allergic to olive pollen were used to carry out the study. All samples were handled anonymously according to the ethical and legal guidelines of the Complutense University of Madrid (CEI PI 49) and Reina Sofia University Hospital (ref. 3033). Written informed consent was obtained from all patients. Sera from olive pollen allergic patients according to their positive olive pollen skin prick test and positive profile by ImmunoCAP 250 (Thermo Fisher Scientific, Uppsala, Sweden) were obtained from two different Spanish populations (Córdoba and Madrid, Spain) according to standard operating procedures. Values >0.35 kU/L by ImmunoCAP were considered positive. Skin prick tests (SPT) were performed according to standard procedures. A wheal diameter <3 mm was considered negative. All patients included in this study reported an immediate IgE-mediated

allergic reaction and a positive SPT with olive pollen. As a summary of the pollen sensitization of each population: sera from Madrid (n=42) were all co-sensitized to *Lolium perenne*, whereas sera from Córdoba (n= 104) were sensitized almost exclusively to olive pollen (Table S1). All patients presented asthma or rhinoconjunctivitis.

Rabbit polyclonal antiserum against rOle e 15 was obtained by injections of the protein (100 µg) in complete Freud's adjuvant, accomplishing the ethics guidelines of the Fundación Jiménez Díaz Hospital (Madrid, Spain). Horseradish peroxidase-labeled goat polyclonal antibody against rabbit IgG was purchased from Bio-Rad. Horseradish peroxidase-labeled mouse anti-human IgE was purchased from Southern Biotech (Birmingham, AL, US).

Circular Dichroism spectroscopic analyses, structural 3D modelling and alignments

The CD spectrum of rOle e 15 (0.2 µg/µL) in 20 mM phosphate buffer, pH 7.0 was recorded in the far-UV (190-260 nm) on a JASCO J-715 spectropolarimeter (Japan Spectroscopic Co., Tokyo, Japan) using a 0.1 cm optical-path quartz cuvette and at 20°C, 85°C and after cooling down to 20°C. The deconvolution of the obtained spectra was performed using the CDNN software. Thermal unfolding was followed measuring the ellipticity at 226 nm heating (20 to 85°C) or cooling (85 to 20°C) at 0.5°C/min with a computer controlled water-bath. Far UV spectra were baseline subtracted and represented as mean residue molecular ellipticity.

Sequence identity and similarity of the Ole e 15 amino acid sequence against other plant, animal and fungal cyclophilins were calculated using DIALIGN (Bielefeld Bioinformatics Service, Universität Bielefeld, Germany). The results were visualized using the GeneDoc program. Known cyclophilin structures were downloaded from PDB. Structural models in PDB format were generated using the ExPASy Swiss-Model tool with human (PDB: 1OCA) and Cat r 1 (PDB: 2mc9) as templates. Final 3D-structure models were visualized with PyMOL (Schrödinger, LCC, New York, US). The electrostatic surface potential of rOle e 15 was calculated by the Adaptive Poisson Boltzmann Solver (APBS) program (37) and visualized with PyMOL.

Immunological characterization

Indirect ELISA was performed in 96-well plates (Costar, Corning, New York, US) coated with 0.1 µg recombinant protein per well, according to previously optimized protocols (18, 38), using individual serum samples diluted 1:10.

WB analysis of human IgE bound to alternatively 0.5 µg rOle e 15, or 40 µg of olive pollen extract, immobilized onto nitrocellulose membranes was performed incubating with individual human sera (1:10) and horseradish peroxidase-labeled mouse anti-human IgE Fc (1:1,000). Binding of specific polyclonal antisera against rOle e 15 was detected by goat anti-rabbit IgG horseradish peroxidase-labeled antibody (1:3,000). Chemiluminescent signal was developed using ECL-Western blotting reagent (Amersham Bioscience) or WesternBright™ QUANTUM (Advansta Inc., San Jose, California, US) reagents.

For the IgE inhibition experiments with indicated protein extracts, a pool of sera (n=5 for WB; n=6 for ELISA) was pre-incubated with 5 mg/mL for WB; or alternatively 0.25 and 2.5 mg/mL for ELISA of the inhibiting extracts for 2 h at room temperature and shaking. The following steps of the ELISA or WB experiments were performed as described above and the results shown as percentage of inhibition (18, 38). For the IgE inhibition experiments using purified proteins, serum was pre-incubated with 10 µg of rOle e 15, nOle e 1 or both. IgE reactivity was visualized as previously described (18, 35). Inhibition values were calculated as percentage in comparison to the non-inhibited control using the formula $[(1 - OD_{492\text{ nm}} \text{ with inhibitor} / OD_{492\text{ nm}} \text{ without inhibitor}) * 100]$ by ELISA and as signal lost in comparison to a non-inhibited control (No inhibition) by WB after densitometry of the bands using the Quantity One 1-D Analysis software (Bio-Rad Laboratories Inc., Madrid, Spain).

Cloning of the putative allergens from *O. europaea* pollen, expression and purification

Total RNA purification from pollen was performed using TRIzol and RNeasy Plant MiniKit (Qiagen, Hilden, Germany). Total RNA quality and concentration were analyzed using 2100 Bioanalyzer (Agilent Genomics). cDNA amplification was performed using the SMARTer®

RACE 5'/3' kit (Clontech, TaKara Bio Company, Mountain View, CA,US) or the SuperScript III First-Strand Synthesis System (Thermo Fisher Scientific). Different degenerate and non-degenerate oligonucleotides (Table S2) were synthesized based on the available proteomic and genomic data. PCR products were cloned into the pCR2.1 plasmid using the TOPO TA Cloning kit (Invitrogen, Groningen, The Netherlands) and sequenced. Then, specific overlapping oligonucleotides (Table S2) were designed to subclone the cDNAs encoding the complete proteins into the pET28a plasmid (Novagen, Billerica, MA, USA) as N-terminal His₆-tagged proteins using, alternatively, the NEBuilder[®] HiFi DNA Assembly Mix (New England Biolabs, Ipswich, Massachusetts, US) or the InFusion[®] HD Cloning Kit (Clontech) following the manufacturer instructions.

The pET28a/cDNA constructs were used to transform BL21 (DE3) *E. coli* cells. 10 mL 30 µg/mL kanamycin/LB cultures containing the transformed cells were grown overnight and then, diluted ten times and grown until OD_{600nm} reached 0.7. Finally, protein production was induced with 0.4 mM isopropyl thio-β-D-thiogalactoside (IPTG) and grown at 30°C 230 rpm for 16 h. Then, the cultures were centrifuged at 6,000 x g for 20 min at 4°C. Proteins were purified by gravity-flow chromatography from the soluble fraction of cell lysates using Ni-NTA agarose (Qiagen) following the manufacturer instructions with minor modifications. Briefly, cell pellets corresponding to 250 mL *E. coli* culture were resuspended in 10 mL lysis buffer (300 mM NaCl, 1 mM PMSF, 50 mM phosphate buffer pH 8.0), distributed in 1 mL aliquots and disrupted by five cycles of subsequent freezing in liquid N₂ for 40 s and thawing at 42°C in a water bath for 5 min. Then, soluble fractions obtained by centrifugation at 6,000 x g for 20 min at 4°C were incubated 16 h at 4°C in a 15 mL conical centrifuge tube with 500 µL Ni-NTA (CliniSciences, Nanterre, France) agarose previously pre-incubated three times with 5 mL lysis buffer. Then, Ni-NTA agarose was washed five times for 10 min at 4 °C with 5 mL wash buffer (10 mM imidazole, 300 mM NaCl, 50 mM phosphate buffer pH 8.0) recovering each time the resin by centrifugation at 4,000 x g for 10 min at 4°C. Proteins were then eluted with 5 mL elution buffer (500 mM imidazole, 300 mM NaCl, 50 mM phosphate buffer pH 8.0) for 20 min at 4°C on rotation. To avoid the presence of traces of Ni-NTA agarose in the eluted fractions, a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

final centrifugation step was performed at 12,000 x g for 5 min at 4°C. Finally, purified proteins were desalted onto PD10 desalting columns (GE Healthcare, Chicago, IL, US) with 20 mM ABC, quantified, aliquoted, lyophilized and stored at -20°C until use.

RESULTS

Definition of the olive pollen proteome by proteomics and bioinformatics

In the present study, we have used an in-depth bottom-up proteomic approach to analyze the proteomic profile of olive pollen, to classify its allergen families and define its allergenome, and to characterize new olive pollen allergen candidates (Figure 1).

Figure 1. Workflow of the study to delineate the olive pollen proteome and its allergenome, and describe Ole e 15 as a new relevant allergen implicated in cross-reactivity. 3D-models of the proteins were created using 1sev.1.A, 5.nue.1.B, 1xsx.1.A and 2mc9 PDB structures for malate dehydrogenase (mitochondrial), malate dehydrogenase (cytosolic), enolase and cyclophilin, respectively.

First, three olive pollen protein extracts replicates were fractionated by SDS-PAGE and protein bands in-gel trypsin digested. Then, the extracted peptides were analyzed by LC-MS/MS with a Q-exactive mass spectrometer. Next, the quality of the MS results was investigated calculating the mean and SD of the mass error in ppm in the three replicates. For replicate 1 the mean±SD mass error was 0.66 and 1.84, for replicate 2 was 2.39 and 1.64, and for replicate 3 was 2.41 and 1.68. As the data showed a good quality, we proceeded to subsequent analyses.

From the three biological replicates, a total of 1,907 different proteins were identified (Table S3 and Figure S1). Protein sequences identified from olive pollen were functionally mapped and

annotated with the corresponding GO terms using the Blast2GO software after local protein BLAST with NCBI RefSeqs sequences from plant species with curated GO annotations from Ensembl Plant (*see* Methods). A total of 1,738 pollen proteins were annotated with GO terms after combining GO mapping and InterProScan data (Figure 2a and Table S4).

Figure 2. Classification of the olive pollen proteome according to the annotated GO terms after Blast2GO analysis and identification of potential allergens. (a) Summary table with the general results after Blast2GO and SignalP 4.1 bioinformatic analysis. (b) Biological process, molecular function and cellular component GO term distribution on the level 2 of complexity. (c) Number of proteins of each enzyme class after GO-Enzyme Code mapping. (d) Table summarizing the main results after protein Blast against the Allergome database and application of criteria for the identification of potential allergens and allergen families.

Cellular component terms revealed that identified olive pollen proteins were widely distributed among the cell, with more than 500 proteins associated to membranes. Metabolic processes were the most abundant in biological process terms (Figure 2b and Figure S2). Regarding molecular function, cyclic compound, ion and protein binding activities, and hydrolase activity, were highly represented (Figure S2). To be noticed, 1,176 out of the 1,907 identified proteins displayed catalytic activity (61.7%), being hydrolases, oxidoreductases and transferases the most abundant enzyme classes found (Figure 2c and Table S4). KEGG pathway analysis showed that 7 out of the 10 most enriched pathways were related to metabolism (Figure

S3a). To analyze the enrichment of certain GO terms in pollen, a Fisher's exact test was performed comparing the annotated GO terms of the pollen proteome with that of the genome-derived proteome (Figure S3b and Table S4). GO terms related to salt stress, salt absorption and amino acid and sugar metabolism were over-represented in pollen. Moreover, 239 (12.5 %) of the identified pollen proteins were predicted as secreted by SignalP 4.1 (Table S5).

Olive pollen allergenome delineation reveals the presence of 47 allergen families

Next, the delineation of the olive pollen allergenome was investigated to identify putative allergens in this allergenic source with high clinical relevance (17). Firstly, the whole wild olive genome-predicted proteome and the olive pollen identified proteome were locally blasted against the Allergome database. A total of 3,469 and 363 hits were retrieved, respectively. Then, a set of criteria (*see* Methods) were manually applied to filter the results and identify potential allergens. Of the total of 2,446 potential allergens belonging to 76 different allergen families in the wild olive predicted-proteome, 203 proteins belonging to 47 allergen families were present in pollen (Figure 2d, Table 1, Table S6 and Table S7). Among them, 10 families previously reported as actual olive pollen allergens were observed (17). Moreover, sixty of these proteins (29.6%) were predicted to be secreted by SignalP 4.1 (Table S5), which is one the most common features of allergenic proteins.

Table 1. Olive pollen allergenome definition by in-depth proteomics and bioinformatics using the Allergome database.

Accession Number*	Protein name	Homologous allergen in the IUIS database	Uniprot Code	Organism	Amino acid similarity/identity (%)	E-value	Allergen family name	All Fam Code	Route of sensitization	Total potential olive pollen allergens identified in the family****
XP_022892128.1	Major pollen allergen. Ole e 1-like	Ole e 1	P19963	<i>O. europaea</i> (Cultivated olive)	99/95	7.4E-107	Ole e 1-like	AF087	Inhalation	17
XP_022882512.1	Profilin-3	Ole e 2	A4GDU2	<i>O. europaea</i> (Cultivated olive)	100/98	0	Profilin	AF051	Inhalation	4
XP_022850789.1	Polcalcin	Ole e 3	O81092	<i>O. europaea</i> (Cultivated olive)	100/98	0	EF-Hand	AF007	Inhalation	5
XP_022849259.1	Superoxide dismutase [Cu-Zn] 2	Ole e 5	Q8L5E0	<i>O. europaea</i> (Cultivated olive)	100/100	0	Cu/Zn Superoxide Dismutase	AF020	Inhalation	2
XP_022896469.1	Non-specific lipid transfer protein	Ole e 7	(18)**	<i>O. europaea</i> (Cultivated olive)	81/74	-	Prolamin	AF050	Inhalation	3
XP_022899478.1	Glucan endo-1,3-beta-D-glucosidase	Ole e 9	D8VPP5	<i>O. europaea</i> (Cultivated olive)	100/99	0	Beta-1,3-glucanase and X8 domain	AF063	Inhalation	7
XP_022888359.1	Pectinesterase 1	Ole e 11	A4GDU2	<i>O. europaea</i> (Cultivated olive)	99/99	0	Pectin methylesterase	AF026	Inhalation	9
XP_022894573.1	Isoflavone reductase-like protein	Ole e 12	E1U332	<i>O. europaea</i> (Cultivated olive)	100/99	0	Isoflavone reductase	AF123	Inhalation	5
XP_022859834.1	Protein P21-like	Act d 2	L7TRX2	<i>A. deliciosa</i> (Kiwifruit)	74/57	1.69E-97	Thaumatococcus-like Protein (Ole e 13)	AF060	Ingestion	1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

XP_022872535.1	Exopolygalacturonase-like	Pla a 2	Q6H9K0	<i>P. xacerifolia</i> (London planetree)	73/51	2.15E-142	Polygalacturonase (Ole e 14)	AF057	Inhalation	4
XP_022846733.1	Enolase 2	Hev b 9	Q9LEI9	<i>H. brasiliensis</i> (Latex)	96/89	0	Enolase	AF031	Contact	4
XP_022888607.1	Glutathione S-transferase L3-like	Bet v 8	A0A096XCA4	<i>B. verrucosa</i> (Birch)	82/69	3.76E-121	Gutathione S- transferase	AF010	Inhalation	1
XP_022880502.1	Cysteine proteinase inhibitor 1-like	Act d 4	Q6TPK4	<i>A. deliciosa</i> (Kiwifruit)	77/51	9.97E-34	Cystatin	AF005	Ingestion	1
XP_022876210.1	Probable cysteine protease RD19B	Act d 1	A5HIJ3	<i>A. deliciosa</i> (Kiwifruit)	91/80	0	Papain-like cysteine protease	AF030	Ingestion	9
XP_022857847.1	5-Methyltetrahydropteroyltriglutamate- homocysteine methyltransferase	Sal k 3	C1KEU0	<i>S. kali</i> (Russian thistle)	95/89	0	Cobalamin- independent methionine synthase	AF152	Inhalation	7
XP_022854141.1	Probable pectate lyase P59	Art v 6	A0PJ16	<i>A. vulgaris</i> (Mugwort)	65/49	1.63E-40	Pectate lyase	AF073	Inhalation	3
XP_022867183.1	Superoxide dismutase [Mn], mitochondrial-like isoform X2	Hev b 10	P35017	<i>H. brasiliensis</i> (Latex)	90/83	7.5E-145	Fe/Mn superoxide dismutase	AF019	Contact	2
XP_022880113.1	NAD(P)H dehydrogenase (quinone) FQR1 isoform X1	Alt a 7	P42058	<i>A. alternata</i> (Alternaria)	71/55	1.68E-56	NADPH- dependent FMN reductase	AF053	Inhalation	7
XP_022853324.1	60S acidic ribosomal protein P2B- like	Pru du 5	Q8H2B9	<i>P. dulcis</i> (Almond)	95/85	1.94E-31	60S acidic ribosomal protein	AF070	Ingestion	1
XP_022854082.1	Luminal-binding protein 5-like	Cor a 10	Q9FSY7	<i>C. avellana</i> (Common Hazel)	97/92	0	Heat shock protein HSP70	AF002	Ingestion	22
XP_022897418.1	Peroxiredoxin-2	Mala f 3	P56578	<i>M. furfur</i>	60/45	5.9E-44	Redoxin	AF131	Inhalation	5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

XP_022855565.1	Soluble inorganic pyrophosphatase 6, chloroplastic	Der f 32	A0A088SCP3	<i>D. farinae</i> (Human dust mite)	67/47	2.04E-74	Inorganic pyrophosphatase	AF205	Inhalation	1
XP_022885711.1	Thioredoxin H-type	Tri a 25	Q9LDX4	<i>T. aestivum</i> (Wheat)	76/57	9.12E-47	Thioredoxin	AF023	Ingestion	5
XP_022875019.1	Actin-depolymerizing factor -like	Der f 31	A0A088SAY1	<i>D. farinae</i> (Human dust mite)	59/38	1.32E-23	Actin-depolymerizing factor-homolog	AF201	Inhalation	6
XP_022889510.1	Ferritin-2, chloroplastic-like isoform X2	Der f 30	L7UZ91	<i>D. farinae</i> (Human dust mite)	56/39	1.7E-19	Ferritin	AF198	Inhalation	1
XP_022851627.1	Aldehyde dehydrogenase family 2 member B7, mitochondrial-like	Cla h 10	P40108	<i>C. herbarum</i>	68/56	2.04E-74	Aldehyde dehydrogenase	AF040	Inhalation	5
XP_022867276.1	Aspartic proteinase-like	Aed a 11	Q03168	<i>A. aegypti</i> (Yellow fever mosquito)	72/44	1.28E-84	Eukaryotic Aspartyl Protease	AF004	Contact	2
XP_022896740.1	Peptidyl-prolyl <i>cis-trans</i> isomerase 1	Hev b Cyp	F8RW92	<i>H. brasiliensis</i> (Latex)	97/91	1.58E-109	Cyclophilin	AF038	Inhalation/Contact	7
XP_022855094.1	Ribosomal protein L3-2	Asp f 23	B0XSL2	<i>A. fumigatus</i>	73/69	1.9E-105	Ribosomal Protein L3	AF058	Inhalation	1
XP_022895071.1	Glyceraldehyde-3-phosphate dehydrogenase, cytosolic	Tri a 34	C7C4X1	<i>T. aestivum</i> (Wheat)	95/87	5.9E-101	Glyceraldehyde 3-phosphate dehydrogenase	AF184	Inhalation	7
XP_022882931.1	Subtilisin-like protease SBT4.15	Cuc m 1	Q39547	<i>C. melo</i> (Melon)	64/47	0	Subtilisin-like serine protease	AF021	Ingestion	5
XP_022862800.1	Malate dehydrogenase, mitochondrial	MDH cit I	P17783	<i>C. lanatus</i> (Watermelon)	97/86	0	Lactate/Malate dehydrogenase	AF014	Ingestion/Inhalation	9
XP_022874613.1	Malate dehydrogenase, cytosolic	-	(37)***	<i>F. excelsior</i>	-	-				

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

				(Ash)						
XP_022877964.1	Probable serine/threonine-protein kinase PBL23	Sal k 2	Q8L5K9	<i>S. kali</i> (Russian thistle)	60/36	3.73E-53	Protein kinase	AF017	Inhalation	2
XP_022876859.1	Reactive Intermediate Deaminase A, chloroplastic-like	Der f 34	A0A1J1DL12	<i>D. farinae</i> (Human dust mite)	58/30	8.23E-18	RidA/YjgF/L-PSP family	AF213	Inhalation	1
XP_022847291.1	Stress-related protein-like	Hev b 3	F8SKC3	<i>H. brasiliensis</i> (Latex)	66/50	3.67E-58	Rubber elongation factor	AF125	Contact	2
XP_022891594.1	Aldehyde oxidase GLOX-like	Art an 7	A0A2H4HHY6	<i>A. annua</i> (Sweet wormwood)	57/39	2.27E-132	Galactose oxidase	-	Inhalation	1
XP_022843917.1	Cytochrome c	Cur l 3	Q96VP3	<i>C. lunata</i>	78/56	1.77E-44	Cytochrome c	AF006	Inhalation	2
XP_022889919.1	Elongation factor 1-delta	Pen c 24	Q69BZ7	<i>P. citrinum</i>	78/40	3.83E-24	Eukaryotic elongation factor 1	AF011	Inhalation	2
XP_022844073.1	Serpin-ZX-like	Tri a 33	P93692	<i>T. aestivum</i> (Wheat)	72/55	1.34E-144	Serpin serine protease inhibitor	AF018	Inhalation	1
XP_022891403.1	3-oxoacyl-[acyl-carrier-protein] reductase 4-like	Alt a 8	P0C0Y4	<i>A. alternata</i> (Alternaria)	55/32	1.82E-30	Short chain dehydrogenase	AF028	Inhalation	3
XP_022890941.1	Triosephosphate isomerase, cytosolic-like, partial	Tri a 31	Q9FS79	<i>T. aestivum</i> (Wheat)	91/80	0	Triosephosphate isomerase	AF032	Inhalation	4
XP_022863167.1	Vicilin-like seed storage protein At2g28490, partial	Coc n 1	A0A0S3B0K0	<i>C. nucifera</i> (Coconut tree)	70/52	2.03E-108	Cupin	AF045	Ingestion	1
XP_022856299.1	Major allergen Pru ar 1-like	Pru ar 1	O50001	<i>P. armeniaca</i> (Apricot tree)	79/67	1.46E-63	Bet v 1 family	AF069	Ingestion	1
XP_022861507.1	Eukaryotic translation initiation factor 3 subunit l-like	For t 2	B2ZPG7	<i>F. taiwana</i>	68/48	7.37E-112	WD-40 Repeat	AF142	Contact	1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

				(Biting midge)						
XP_022899451.1	Fructose-bisphosphate aldolase, cytoplasmic isozyme-like	Sal s 3	C0H9I1	<i>S. salar</i> (Atlantic salmon)	74/60	2.48E-145	Fructose bisphosphate aldolase class I	AF197	Ingestion	5
XP_022874390.1	Acid beta-fructofuranosidase 2, vacuolar-like	Sola l 2	Q8RVW4	<i>S. lycopersicum</i> (Tomato)	80/60	0	Glycoside hydrolase family 32	AF052	Ingestion	3
XP_022889771.1	Calnexin homolog 1-like	Pen ch 31	Q2TL59	<i>P. chrysogenum</i>	67/44	5.75998E-93	Calreticulin	AF055	Inhalation	6

MS/MS spectra for the 5 proteins with single peptide entries in the three biological replicates (Glutathione S-transferase L3-like, Isoflavone reductase-like protein, Peptidyl-prolyl cis-trans isomerase CYP21-like, Pollen allergen Che a 1-like and Superoxide dismutase [Cu-Zn] 2-like isoform X3) are shown in Figure S4.

* NCBI Accession numbers of the *Olea europaea* var. *sylvestris* proteins sharing the highest sequence identity with Allergome allergens of the indicated allergen families (see also Table S7).

** Ole e 7 sequence has been recently published and is not yet deposited in Uniprot (18).

*** Cytosolic malate dehydrogenase from ash (*F. excelsior*) pollen, a member of the Oleaceae family, was suggested as potential allergen in (39).

**** Isoallergens, allergen variants and different allergenic proteins belonging to the same family (see also Table S6 and Table S7 for a full description of the allergens identified).

One of the four selected allergen candidates, cyclophilin from olive pollen, shows significant IgE reactivity

After proteomic analysis, four potential allergens among the 47 allergen families identified from olive pollen (cyclophilin, enolase, and cytosolic and mitochondrial malate dehydrogenases) were selected for validation to assess their IgE reactivity. Three different cDNA clones from cyclophilin and one cDNA clone for the rest of the proteins were obtained (Table S8). All proteins shared >95% sequence identity with predicted proteins in the *Olea europaea* var. *sylvestris* database.

We then proceeded to express and purify the potential allergens to analyze their IgE seroreactivity. The predicted mature forms of the proteins were expressed in *E. coli* as His₆-tagged N-terminal proteins. A purity > 95% for all proteins was achieved (Figure 3a). Proteins were subsequently analyzed by ELISA using sera from olive pollen allergic patients from two populations from Spain -Madrid and Córdoba-, and five non-atopic individuals.

Figure 3. Analysis of the purified recombinant proteins and IgE-reactivity to rOle e 15. (a) Analysis of 0.5 µg purified recombinant proteins by Coomassie Blue staining after 15% SDS-PAGE under reducing conditions. Purified natural Ole e 1 was used as control. (b) Dot plots representing the strength of the IgE reactivity to rOle e 15. The dotted line indicates the response threshold (O.D. values > 0.1). (c) Bar graphs representing the prevalence of Ole e 15-sensitized patients in both populations (left) and the pediatric prevalence among rOle e 15-sensitized patients (right). (d) Prevalence data are also displayed.

Only olive pollen cyclophilin showed IgE recognition, and thus, was named Ole e 15 according to WHO/IUIS (Figure 3b). No significant differences in the recognition frequency were observed between both patients' populations, with prevalence values of 14.3% and 12.5% in Madrid and Córdoba, respectively (Figure 3c).

Most of the allergic patients showed high IgE levels against the allergen, especially in the population from Córdoba. Furthermore, we observed a higher number of pediatric patients than

adults among the Ole e 15 sensitized patients. In total, 8 out of the 13 allergic patients detected in Córdoba (61.5 %) and 4 out of the 6 allergic patients detected in Madrid (66.7 %) were pediatric (Figure 3c).

Recombinant Ole e 15 shows the typical folding, shares high sequence identity with other plant, animal and fungal cyclophilins and is involved in cross-reactivity

Next, we proceeded to further physicochemically and immunologically characterize this new olive pollen allergen. To this end, we initially assessed by mass spectrometry and CD its correct molecular mass and folding, respectively. First, an experimental molecular mass of 19,408 Da was observed for His₆-tagged rOle e 15, in agreement with the expected theoretical molecular mass without the initial Met (19,391 Da), which is processed when proteins are produced in bacteria. Far-UV CD spectra and percentages of secondary structure of rOle e 15 were in agreement with those described for other cyclophilins. In addition, thermal denaturation experiments for rOle e 15 revealed a T_m of 47°C, being unable to recover its structure cooling afterwards (Figure S5).

Ole e 15 amino acid sequence alignment with other plant, animal and fungal allergenic cyclophilins showed a high degree of conservation, with identities of 94%, 68% and 65% with Sola l 5, Der f 29 and Mala s 6, respectively (Figure S6). Molecular modelling revealed a predicted structure with the typical plant cyclophilin folding (Figure 4a-c and Figure S6), consisting of a β-barrel of seven anti-parallel strands, two surrounding α-helices and a characteristic stretch of seven amino acids called the divergent loop, which is mainly present in plants (40). Moreover, the predicted amino acids defining the active surface in the cyclophilin family were present (Figure 4a) (41). Its theoretical surface electrostatic potential was mainly basic, but showed a neutral potential in the putative active surface region (Figure 4b). In addition, other cyclophilins were modelled and compared to Ole e 15 (Figure 4c), showing an almost identical folding among them except for the divergent loop.

Figure 4. Prediction and comparison of the Ole e 15 structure with other cyclophilins, and analysis of its potential IgE- cross-reactivity. (a) 3D-model of Ole e 15 structure using that of Cat r 1 (PDB: 2mc9) as template. Putative catalytic residues (R62, F67, M68, N70, A110, F120, W128, L129, and H133) and the divergent loop (KSGKPLH) sequence are labeled. (b) predicted electrostatic potential of Ole e 15 using Adaptive Poisson Boltzmann Solver (APBS) program, considering 2 and 78 as dielectric constant values for protein and solvent, respectively. Active site region is highlighted with a circle. Electropositively- and electronegatively-charged regions are colored blue and red, respectively. Neutral regions are colored white. (c) Comparison of the structures of plant (Ole e 15, Bet v 7 and Cat r 1), fungal (Mala s 6) and animal (human PPIA, Der f 29, *C. elegans* and *A. aegypti*) cyclophilins. Plant and animal cyclophilins were predicted using structures of Cat r 1 (PDB: 2cm9) and human PPIA (PDB: 1OCA) as ExPASy Swiss-Model tool templates, respectively. All structures were generated by PyMOL. (d) Analysis of the IgE inhibition to rOle e 15 was performed by ELISA using an equivolumetric sera pool of olive pollen patients and 25 and 250 µg of the indicated extracts as inhibitors. (e) Analysis of the IgE binding to Ole e 15 was performed by WB using an equivolumetric sera pool and 500 µg of the indicated protein extracts as inhibitors. All these data suggested a potential implication of olive pollen cyclophilin in cross-reactivity, as it has been shown for other plant and fungal cyclophilins (40, 42, 43).

To address this question, we performed IgE inhibition assays with a pool of sera from olive pollen allergic patients, using pollen, plant-derived food, dust mite and human lung protein extracts as inhibitors (Figure 4d-e).

Remarkably, IgE inhibition was observed with all pollen and plant-derived food protein extracts from related and unrelated species, together with dust mite and human lung or cell protein extracts, either by ELISA or WB. The highest inhibition was achieved with extracts from different types of tree pollens from the Oleaceae family and birch. Regarding plant-derived food extracts, an IgE inhibition higher than 60% was observed for all extracts analyzed by ELISA, whereas by WB the IgE inhibition was lower for nuts extracts (up to 65% for pistachio (*P. vera*) than for fruit and seeds extracts (up to 98% for sesame (*S. indicum*) or pumpkin (*C. maxima*). Interestingly, a 40% and 80% IgE inhibition by ELISA and 55% and 96% by WB were also achieved by using human dust mite (*D. pteronyssinus*) and human (*H. sapiens*) lung protein extracts, respectively.

The identification of natural Ole e 15 is hampered by Ole e 1, the main allergen in the olive pollen extract

Finally, we assessed the behavior of natural Ole e 15 (nOle e 15) in the olive pollen protein extract to determine why this relevant allergen implicated in pollen-pollen, pollen-plant food, or pollen-dust mite cross-reactivity has remained elusive in previous investigations. The most abundant protein in olive pollen extract is the major allergen Ole e 1, which after SDS-PAGE, appears as two main bands of 17 to 18.5 kDa for the non-glycosylated and N-glycosylated isoforms, respectively, with a pI interval (5.0 to 9) for the different isoforms of this allergen. On the other hand, a theoretical mass of 18.1 kDa and 8.7 pI were predicted for natural Ole e 15, which led us to hypothesize that the presence of Ole e 1 could have masked the identification of Ole e 15 in the extract.

To address this question, WB analysis after 1D and 2D SDS-PAGE of the olive pollen extract with anti-Ole e 1 and anti-Ole e 15 pAb and IgE inhibition assays confirmed that nOle e

15 is present as a monomeric protein with an experimental pI ~8.4 and apparent molecular mass very similar to glycosylated Ole e 1 (Figure 5a-b). As a consequence, reactivity to nOle e 15 in a cosensitized patient with reactivity to Ole e 1 would remain hidden because of the high protein content of Ole e 1 and its prevalence, and might only be detected if the serum is previously inhibited with Ole e 1 (Figure 5c).

Figure 5. Analysis of Ole e 15 in the olive pollen extract. (a, b) WB analysis after SDS-PAGE of olive pollen extract (40 μ g) using anti-Ole e 1 and anti-Ole e 15 pAbs. (a) 1D SDS-PAGE. (B) 2D SDS-PAGE. (c) IgE-reactivity analysis after SDS-PAGE and transference to membranes of olive pollen extract. Strips were incubated with the serum #14, sensitized against Ole e 1 and Ole e 15, after either no inhibition or its IgE inhibition with alternatively 10 μ g of purified nOle e 1, rOle e 15 or both. No signal was detected without serum incubation (No sera). IgE-reactivity of a serum sensitized to Ole e 15 but not to Ole e 1 was also analyzed (Serum #4).

Discussion

Olive pollen is an important allergenic source in those regions with extensive cultivars for olive oil production. The increasing cultivation of this tree in China, India, Australia, Russia, California (US) and multiple countries of South America has led to a growing number of olive pollen allergic patients. Therefore, it will become a top allergenic health problem worldwide in the next years; and thus, the definition of its allergenome will significantly improve the clinical management of olive pollen allergic patients.

Here, to delineate the cultivated olive pollen proteome and its allergenome, we took advantage of the recently reported wild olive genome in combination with a high performance bottom-up LC-MS/MS approach to optimize protein identification. The number of sequences obtained represents the 3.8% of the complete genome-predicted proteome and is in agreement with published data regarding transcriptomic analysis in olive pollen, in which 1,976 transcripts encoding complete proteins, including isoforms, were found (44). Then, we performed functional annotation of the identified proteins, using a newer version of Blast2GO software and a more up-to-date Gene ontology (GOA) database than the previously used for the genome (14). Our annotation results are the expected for pollen protein functions in germination and tube growth, extremely regulated multistep processes in which secreted or membrane proteins, and enzymes, play a major role (45). In fact, it is also in concordance with other pollen proteomic studies in which most of the proteins were annotated to metabolic processes (46-49). Collectively, we have performed a comprehensive in-depth proteomic study of the olive tree allergenic pollen, and elucidated the distribution of its proteins according to their associated biological function and location, which can be helpful for further studies related to olive pollen biology. Nevertheless, it must be considered that the obtained data is highly dependent on technical and biological issues. For instance, it is known that protein extraction methods in plant proteomics can be determinant to detect hydrophobic membrane proteins and low abundant proteins (50, 51). In addition, it has been reported that the use of different solid-phase extraction methods (i.e. ZipTip vs SOLAμ) after in-gel trypsin digestion and prior to nLC-MS/MS analysis can increase the coverage in the proteome (52). Moreover, previous studies have demonstrated

that pollen grains from different olive cultivars differ in the quantity and number of proteoforms, including those of important allergens like Ole e 1, Ole e 2 and Ole e 5 (53-55). This, together with environmental factors like geographic variability in the reached pollen counts, the coexistence of other allergen sources and pollution, as well as human factors like diet, alters the sensitization patterns to olive pollen, especially regarding minor allergens (56). Therefore, further proteomic studies analyzing pollen grains from common cultivars using alternative protein extraction methods would add relevant biological and clinical information to the data here presented.

We focused our study on the identification of allergenic proteins in olive pollen to complete its allergenome. From the 76 allergen families present in the genome-predicted proteome, 47 allergen families were identified in pollen, with 20 non-previously described in this source. Importantly, all of the previously reported allergen families were found (17-19, 57). Only the allergen Ole e 6 was not observed. Ole e 6 is present among the potential allergens identified from the genome-predicted proteome (Table S6), and its absence in our proteomic data could be due to either its extremely low abundance in pollen (46), or because of the stringent filters used.

An important feature for some allergenic sources is the high degree of heterogeneity in the same allergen family. According to the WHO/IUIS guidelines for allergen nomenclature, allergenic proteins from the same species are considered isoallergens when they have similar molecular size and a sequence identity of at least 67%; or allergen variants (isoforms) when the proteins display a sequence identity >90%. It has been shown that isoallergens and allergen variants found in pollen can display different IgE reactivity (19, 38, 53-55, 59). Moreover, in some sources, proteins of the same allergen family, but not considered isoallergens, can be allergenic (60). In our study, we found more than one protein in 34 of the 47 identified allergen families. In this sense, the Ole e 1-like allergen family was the most heterogeneous (Table 1 and Table S7). Of the 17 proteins found in this family, eight were Ole e 1 variants and, five were Ole e 1 isoallergens. This high degree of Ole e 1 polymorphism had already been described and can vary among different olive cultivars (55, 61). The further study of these allergen polymorphisms might help to identify hypoallergenic molecules, which can be used for the

development of safer immunotherapy strategies. On the other hand, we observed interesting potential allergens which remain to be studied. For example, the presence of a thaumatin-like protein (TLP) similar to Act d 2 was found (Table 1 and Table S7). A member of the TLP family had been described as an olive fruit allergen (Ole e 13) (62), but the presence of other TLPs in olive pollen was unknown. Thus, this finding suggests this protein family could also be allergenic in olive pollen allergic patients, and may be implicated in cross-reactivity with fruits. Additionally, we observed the presence of several allergen families commonly found in the pollen of weeds, grasses and trees (63), such as pectate lyase, papain-like cysteine protease, or cyclophilin. These data validated our workflow to identify potential allergenic proteins, defining the most complete allergenic proteomic profile of olive pollen up to date.

The next goal of this work consisted of the validation of the allergenic potential of some of the newly identified candidate allergens. We selected those protein families reported to be allergenic in at least two allergenic sources, including one non-pollinic, to identify those olive pollen allergens implicated in cross-reactivity. MDH (cytoplasmic and mitochondrial), enolase, and cyclophilin were selected for validation. After their cloning, expression and purification, only cyclophilin was able to bind IgEs from the allergic patients tested. Cyclophilin was named Ole e 15 according to the WHO/IUIS. Remarkably, cytoplasmic and mitochondrial MDH had been previously described as potential allergenic proteins in watermelon, soybean seeds, ash pollen, or yeast; enolase in latex, ragweed pollen, or molds; and cyclophilin in carrot, tomato, birch pollen, molds, and dust mites (9, 39, 43, 64-72). Then, it cannot be discarded that these olive pollen proteins could be reported as allergenic if tested against other allergic populations or if they were expressed in heterologous systems different from *E. coli*, which would provide post-translational modifications and complex folding.

The last objective of our study was the characterization of olive pollen cyclophilin. Recombinant protein structure exhibited a folding with similar secondary structure as Cat r 1 by far-UV CD (40), indicating that the protein was correctly folded. Regarding its immunological behavior, we observed that the prevalence of Ole e 15-sensitized patients was about 13% among olive pollen allergic patients from Córdoba and Madrid, with a majority of them showing high

IgE levels against the allergen. Although Ole e 15 sensitization was more frequent in pediatric patients (17.4%) than in adults (7.8%) in both populations, the correlation between Ole e 15 sensitization and pediatric patients was not statistically significant. However, this trend will be assessed in further studies with higher cohorts. After inhibition experiments, we found that Ole e 15 is highly implicated in cross-reactivity with pollen, plant derived food and dust mites. Human cyclophilin, which shows 70% sequence identity with Ole e 15, was also able to inhibit the IgE binding to the recombinant protein. Our results are in agreement with a previous study in which plant cyclophilins were predicted to cross-react with animal cyclophilins, due to a high degree of surface conservation (40). This confirmed the broad cross-reactivity of cyclophilins, and extended the physicochemical and immunological knowledge of this pan-allergen to olive pollen. Moreover, as Ole e 15 shares similar physicochemical properties with Ole e 1, the identification of cyclophilin as an allergen has remained elusive by standard techniques. Therefore, the in-depth olive pollen proteomic analysis allowed the identification of this relevant allergen, and suggests that its presence should be assessed in natural preparations to prevent the overestimation of the number of patients allergic to Ole e 1.

In summary, we have performed the most comprehensive proteomic profiling of olive pollen and defined its allergen families. We have also unmasked Ole e 15, a relevant olive pollen allergen belonging to the cyclophilin family with similar physicochemical features to the main olive pollen allergen. After cloning, expression and characterization of Ole e 15, we found that it is a highly potent cross-reactive allergen with plant and animal cyclophilins.

Supporting information:

The following supporting information is available free of charge at ACS website

<http://pubs.acs.org>

Table S1. Information of the olive pollen allergic patients used in the study.

Table S2. List of the oligonucleotides used in the study.

Table S3. Proteins identified in olive pollen by LC-MS/MS.

Table S4. Olive pollen proteins annotated with GO terms.

Table S5. Olive pollen proteins predicted as secreted.

Table S6. Wild olive genome predicted proteins belonging to allergen families.

Table S7. Olive pollen proteins identified by LC-MS/MS belonging to allergen families.

Table S8. Information of the potential allergens cloned.

Figure S1. Summary of the LC-MS/MS olive pollen identified proteins in the three replicates.

Figure S2. Number of GO terms annotated of biological process, molecular function, and cellular component.

Figure S3. KEGG pathways and GO terms enrichment.

Figure S4. Mass spectra corresponding to proteins identified with a single peptide.

Figure S5. Circular dichroism analyses of rOle e 15.

Figure S6. Amino acid sequence alignment of Ole e 15 with other allergenic cyclophilins.

Acknowledgments

We thank the excellent technical support of Sara Abián. This work was supported by grants co-founded by Fondo Europeo de Desarrollo Regional -FEDER-: SAF2014-53209-R to M.V. and R.B., and SAF2017-86483-R to M.V. from the Ministerio de Economía y Competitividad, and the Thematic Networks and Co-operative Research Centres: RIRAAF Network RD12/0013/0015 and ARADyAL (RD16/0006/0014) from the Instituto de Salud Carlos III (ISCIII). B.R., A.J., A.N, and C.M. acknowledge PI-01119-2016 from the Consejería de Salud (Junta de Andalucía) and the Alergosur Foundation. R.B. also acknowledges the financial support of the PI17CIII/00045 grant from the AES-ISCIII program. P.SS.A. FPU predoctoral contract is supported by the Spanish Ministerio de Educación, Cultura y Deporte. S.B. was a fellow of the Juan de la Cierva program of the Ministerio de Economía y Competitividad (Spain). C.O-S. was supported by a contract of the Programa Operativo de Empleo Juvenil y la Iniciativa de Empleo Juvenil (YEI) with the participation of the Consejería de Educación, Juventud y Deporte de la Comunidad de Madrid y del Fondo Social Europeo.

Disclosure of potential conflict of interest

All authors declare no conflicts of interest.

References

1. McCormick, S., Pollen. *Curr Biol* **2013**, 23, (22), R988-90.
2. Chaturvedi, P.; Ghatak, A.; Weckwerth, W., Pollen proteomics: from stress physiology to developmental priming. *Plant Reprod* **2016**, 29, (1-2), 119-32.
3. Ghatak, A.; Chaturvedi, P.; Paul, P.; Agrawal, G. K.; Rakwal, R.; Kim, S. T.; Weckwerth, W.; Gupta, R., Proteomics survey of Solanaceae family: Current status and challenges ahead. *J Proteomics* **2017**, 169, 41-57.
4. Fila, J.; Radau, S.; Matros, A.; Hartmann, A.; Scholz, U.; Feciková, J.; Mock, H.-P.; Čapková, V.; Zahedi, R. P.; Honys, D., Phosphoproteomics profiling of tobacco mature pollen and pollen activated *in vitro*. *Mol Cell Proteomics* **2016**, 15, (4), 1338-1350.
5. Esch, R. E., Grass pollen allergens. *Clin Allergy Immunol* **2008**, 21, 107-26.
6. Asam, C.; Hofer, H.; Wolf, M.; Aglas, L.; Wallner, M., Tree pollen allergens-an update from a molecular perspective. *Allergy* **2015**, 70, (10), 1201-11.
7. Gadermaier, G.; Hauser, M.; Ferreira, F., Allergens of weed pollen: an overview on recombinant and natural molecules. *Methods* **2014**, 66, (1), 55-66.
8. Mas, S.; Torres, M.; Garrido-Arandia, M.; Salamanca, G.; Castro, L.; Barral, P.; Purohit, A.; Pauli, G.; Rodriguez, R.; Batanero, E.; Barderas, R.; Villalba, M., Ash pollen immunoproteomics: identification, immunologic characterization, and sequencing of 6 new allergens. *J Allergy Clin Immunol* **2014**, 133, (3), 923-6 e3.
9. An, S.; Chen, L.; Long, C.; Liu, X.; Xu, X.; Lu, X.; Rong, M.; Liu, Z.; Lai, R., *Dermatophagoides farinae* allergens diversity identification by proteomics. *Mol Cell Proteomics* **2013**, 12, (7), 1818-28.
10. Choopong, J.; Reamtong, O.; Sookrung, N.; Seesua, W.; Indrawattana, N.; Sakolvaree, Y.; Chaicumpa, W.; Tungtrongchitr, A., Proteome, allergenome, and novel allergens of house dust mite, *Dermatophagoides farinae*. *J Proteome Res* **2016**, 15, (2), 422-30.
11. Liu, X.Y.; Yang, K. Y.; Wang, M.Q.; Kwok, J. S.L.; Zeng, X.; Yang, Z.; Xiao, X.J.; Lau, C. P.Y.; Li, Y.; Huang, Z.m.; Ba, J.g.; Yim, A. K.Y.; Ouyang, C.Y.; Ngai, S.M.; Chan, T.F.; Leung, E. L.H.; Liu, L.; Liu, Z.G.; Tsui, S. K.W., Highquality assembly of *Dermatophagoides pteronyssinus* genome and transcriptome reveals a wide range of novel allergens. *J Allergy Clin Immunol* **2018**, 141, (6), 22682271.e8.
12. Chan, T.F.; Ji, K.M.; Yim, A. K.Y.; Liu, X.Y.; Zhou, J.W.; Li, R.Q.; Yang, K. Y.; Li, J.; Li, M.; Law, P. T.W.; Wu, Y.L.; Cai, Z.L.; Qin, H.; Bao, Y.; Leung, R. K.K.; Ng, P. K.S.; Zou, J.; Zhong, X.J.; Ran, P.X.; Zhong, N.S.; Liu, Z.G.; Tsui, S. K.W., The draft genome, transcriptome, and microbiome of *Dermatophagoides farinae* reveal a broad spectrum of dust mite allergens. *J Allergy Clin Immunol* **2015**, 135, (2), 539-548.
13. Nugraha, R.; Kamath, S. D.; Johnston, E.; Zenger, K. R.; Rolland, J. M.; O'Hehir, R. E.; Lopata, A. L., Rapid and comprehensive discovery of unreported shellfish allergens using large-scale transcriptomic and proteomic resources. *J Allergy Clin Immunol* **2018**, 141, (4), 1501-1504.e8.
14. Unver, T.; Wu, Z.; Sterck, L.; Turktas, M.; Lohaus, R.; Li, Z.; Yang, M.; He, L.; Deng, T.; Escalante, F. J.; Llorens, C.; Roig, F. J.; Parmaksiz, I.; Dundar, E.; Xie, F.; Zhang, B.; Ipek, A.; Uranbey, S.; Erayman, M.; Ilhan, E.; Badad, O.; Ghazal, H.; Lightfoot, D. A.; Kasarla, P.; Colantonio, V.; Tombuloglu, H.; Hernandez, P.; Mete, N.; Cetin, O.; Van Montagu, M.; Yang, H.; Gao, Q.; Dorado, G.; Van de Peer, Y., Genome of wild olive and the evolution of oil biosynthesis. *Proc Natl Acad Sci U S A* **2017**, 114, (44), E9413-E9422.
15. Diez, C. M.; Trujillo, I.; Martinez-Urdiroz, N.; Barranco, D.; Rallo, L.; Marfil, P.; Gaut, B. S., Olive domestication and diversification in the Mediterranean Basin. *New Phytol* **2015**, 206, (1), 436-47.
16. D'Amato, G.; Lobefalo, G., Allergenic pollens in the southern Mediterranean area. *J Allergy Clin Immunol* **1989**, 83, (1), 116-22.
17. Villalba, M.; Rodriguez, R.; Batanero, E., The spectrum of olive pollen allergens. From structures to diagnosis and treatment. *Methods* **2014**, 66, (1), 44-54.
18. Oeo-Santos, C.; Mas, S.; Benede, S.; Lopez-Lucendo, M.; Quiralte, J.; Blanca, M.; Mayorga, C.; Villalba, M.; Barderas, R., A recombinant isoform of the Ole e 7 olive pollen

- allergen assembled by de novo mass spectrometry retains the allergenic ability of the natural allergen. *J Proteomics* **2018**, 187, 39-46.
19. Oeo-Santos, C.; Mas, S.; Quiralte, J.; Colas, C.; Blanca, M.; Fernandez, J.; Feo-Brito, F.; Villalba, M.; Barderas, R., A hypoallergenic polygalacturonase isoform from olive pollen is implicated in pollen-pollen cross-reactivity. *Int Arch Allergy Immunol* **2018**, 177, (4), 290-301.
20. Boldogh, I.; Bacs, A.; Choudhury, B. K.; Dharajiy, N.; Alam, R.; Hazra, T. K.; Mitra, S.; Goldblum, R. M.; Sur, S., ROS generated by pollen NADPH oxidase provide a signal that augments antigen-induced allergic airway inflammation. *J Clin Invest* **2005**, 115, (8), 2169-79.
21. Kauffman, H. F.; Tamm, M.; Timmerman, J. A.; Borger, P., House dust mite major allergens Der p 1 and Der p 5 activate human airway-derived epithelial cells by protease-dependent and protease-independent mechanisms. *Clin Mol Allergy* **2006**, 4, 5.
22. Matsumura, Y., Role of allergen source-derived proteases in sensitization via airway epithelial cells. *J Allergy (Cairo)* **2012**, 2012, 903659.
23. Mari, A.; Rasi, C.; Palazzo, P.; Scala, E., Allergen databases: Current status and perspectives. *Curr Allergy Asthma Rep* **2009**, 9, (5), 376-383.
24. Radauer, C.; Bublin, M.; Wagner, S.; Mari, A.; Breiteneder, H., Allergens are distributed into few protein families and possess a restricted number of biochemical functions. *J Allergy Clin Immunol* **2008**, 121, (4), 847-52 e7.
25. Radauer, C.; Nandy, A.; Ferreira, F.; Goodman, R. E.; Larsen, J. N.; Lidholm, J.; Pomes, A.; Raulf-Heimsoth, M.; Rozynek, P.; Thomas, W. R.; Breiteneder, H., Update of the WHO/IUIS Allergen Nomenclature Database based on analysis of allergen sequences. *Allergy* **2014**, 69, (4), 413-9.
26. Barral, P.; Suarez, C.; Batanero, E.; Alfonso, C.; Alche Jde, D.; Rodriguez-Garcia, M. I.; Villalba, M.; Rivas, G.; Rodriguez, R., An olive pollen protein with allergenic activity, Ole e 10, defines a novel family of carbohydrate-binding modules and is potentially implicated in pollen germination. *Biochem J* **2005**, 390, (Pt 1), 77-84.
27. Pelaez-Garcia, A.; Barderas, R.; Batlle, R.; Vinas-Castells, R.; Bartolome, R. A.; Torres, S.; Mendes, M.; Lopez-Lucendo, M.; Mazzolini, R.; Bonilla, F.; Garcia de Herreros, A.; Casal, J. I., A proteomic analysis reveals that Snail regulates the expression of the nuclear orphan receptor Nuclear Receptor Subfamily 2 Group F Member 6 (Nr2f6) and interleukin 17 (IL-17) to inhibit adipocyte differentiation. *Mol Cell Proteomics* **2015**, 14, (2), 303-15.
28. Pelaez-Garcia, A.; Barderas, R.; Mendes, M.; Lopez-Lucendo, M.; Sanchez, J. C.; Garcia de Herreros, A.; Casal, J. I., Data from proteomic characterization of the role of Snail1 in murine mesenchymal stem cells and 3T3-L1 fibroblasts differentiation. *Data Brief* **2015**, 4, 606-13.
29. Kall, L.; Canterbury, J. D.; Weston, J.; Noble, W. S.; MacCoss, M. J., Semi-supervised learning for peptide identification from shotgun proteomics datasets. *Nat Methods* **2007**, 4, (11), 923-5.
30. Gotz, S.; Garcia-Gomez, J. M.; Terol, J.; Williams, T. D.; Nagaraj, S. H.; Nueda, M. J.; Robles, M.; Talon, M.; Dopazo, J.; Conesa, A., High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Res* **2008**, 36, (10), 3420-35.
31. Madeira, F.; Park, Y. M.; Lee, J.; Buso, N.; Gur, T.; Madhusoodanan, N.; Basutkar, P.; Tivey, A. R. N.; Potter, S. C.; Finn, R. D.; Lopez, R., The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res* **2019**.
32. Zerbino, D. R.; Achuthan, P.; Akanni, W.; Amode, M. R.; Barrell, D.; Bhai, J.; Billis, K.; Cummins, C.; Gall, A.; Giron, C. G.; Gil, L.; Gordon, L.; Haggerty, L.; Haskell, E.; Hourlier, T.; Izuogu, O. G.; Janacek, S. H.; Juettemann, T.; To, J. K.; Laird, M. R.; Lavidas, I.; Liu, Z.; Loveland, J. E.; Maurel, T.; McLaren, W.; Moore, B.; Mudge, J.; Murphy, D. N.; Newman, V.; Nuhn, M.; Ogeh, D.; Ong, C. K.; Parker, A.; Patricio, M.; Riat, H. S.; Schuilenburg, H.; Sheppard, D.; Sparrow, H.; Taylor, K.; Thormann, A.; Vullo, A.; Walts, B.; Zadissa, A.; Frankish, A.; Hunt, S. E.; Kostadima, M.; Langridge, N.; Martin, F. J.; Muffato, M.; Perry, E.; Ruffier, M.; Staines, D. M.; Trevanion, S. J.; Aken, B. L.; Cunningham, F.; Yates, A.; Flicek, P., Ensembl 2018. *Nucleic Acids Res* **2018**, 46, (D1), D754-d761.

33. Durinck, S.; Spellman, P. T.; Birney, E.; Huber, W., Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat Protoc* **2009**, 4, (8), 1184-91.
34. Jones, P.; Binns, D.; Chang, H. Y.; Fraser, M.; Li, W.; McAnulla, C.; McWilliam, H.; Maslen, J.; Mitchell, A.; Nuka, G.; Pesseat, S.; Quinn, A. F.; Sangrador-Vegas, A.; Scheremetjew, M.; Yong, S. Y.; Lopez, R.; Hunter, S., InterProScan 5: genome-scale protein function classification. *Bioinformatics* **2014**, 30, (9), 1236-40.
35. Kanehisa, M.; Goto, S., KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* **2000**, 28, (1), 27-30.
36. Petersen, T. N.; Brunak, S.; von Heijne, G.; Nielsen, H., SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* **2011**, 8, (10), 785-6.
37. Baker, N. A.; Sept, D.; Joseph, S.; Holst, M. J.; McCammon, J. A., Electrostatics of nanosystems: application to microtubules and the ribosome. *Proc Natl Acad Sci U S A* **2001**, 98, (18), 10037-41.
38. Mas, S.; Oeo-Santos, C.; Cuesta-Herranz, J.; Diaz-Perales, A.; Colas, C.; Fernandez, J.; Barber, D.; Rodriguez, R.; de Los Rios, V.; Barderas, R.; Villalba, M., A relevant IgE-reactive 28kDa protein identified from *Salsola kali* pollen extract by proteomics is a natural degradation product of an integral 47kDa polygalaturonase. *Biochim Biophys Acta* **2017**, 1865, (8), 1067-1076.
39. Poncet, P.; Senechal, H.; Clement, G.; Purohit, A.; Sutra, J. P.; Desvaux, F. X.; Wal, J. M.; Pauli, G.; Peltre, G.; Gougeon, M. L., Evaluation of ash pollen sensitization pattern using proteomic approach with individual sera from allergic patients. *Allergy* **2010**, 65, (5), 571-80.
40. Ghosh, D.; Mueller, G. A.; Schramm, G.; Edwards, L. L.; Petersen, A.; London, R. E.; Haas, H.; Gupta Bhattacharya, S., Primary identification, biochemical characterization, and immunologic properties of the allergenic pollen cyclophilin cat R 1. *J Biol Chem* **2014**, 289, (31), 21374-85.
41. Davis, T. L.; Walker, J. R.; Campagna-Slater, V.; Finerty, P. J., Jr.; Paramanathan, R.; Bernstein, G.; MacKenzie, F.; Tempel, W.; Ouyang, H.; Lee, W. H.; Eisenmesser, E. Z.; Dhe-Paganon, S., Structural and Biochemical Characterization of the Human Cyclophilin Family of Peptidyl-Prolyl Isomerases. *PLOS Biol* **2010**, 8, (7), e1000439.
42. Glaser, A. G.; Limacher, A.; Fluckiger, S.; Scheynius, A.; Scapozza, L.; Cramer, R., Analysis of the cross-reactivity and of the 1.5 Å crystal structure of the *Malassezia sympodialis* Mala s 6 allergen, a member of the cyclophilin pan-allergen family. *Biochem J* **2006**, 396, (1), 41-9.
43. Cadot, P.; Nelles, L.; Srahna, M.; Dilissen, E.; Ceuppens, J. L., Cloning and expression of the cyclophilin Bet v 7, and analysis of immunological cross-reactivity among the cyclophilin A family. *Mol Immunol* **2006**, 43, (3), 226-35.
44. Carmona, R.; Zafra, A.; Seoane, P.; Castro, A. J.; Guerrero-Fernández, D.; Castillo-Castillo, T.; Medina-García, A.; Cánovas, F. M.; Aldana-Montes, J. F.; Navas-Delgado, I.; Alché, J. d. D.; Claros, M. G., ReprOlive: a database with linked data for the olive tree (*Olea europaea* L.) reproductive transcriptome. *Front Plant Sci* **2015**, 6, (625).
45. Dresselhaus, T.; Franklin-Tong, N., Male-Female crosstalk during pollen germination, tube growth and guidance, and double fertilization. *Mol Plant* **2013**, 6, (4), 1018-1036.
46. Kawakami, T.; Darby, B. J.; Ungerer, M. C., Transcriptome resources for the perennial sunflower *Helianthus maximiliani* obtained from ecologically divergent populations. *Mol Ecol Resour* **2014**, 14, (4), 812-9.
47. Sheoran, I. S.; Sproule, K. A.; Olson, D. J.; Ross, A. R.; Sawhney, V. K., Proteome profile and functional classification of proteins in *Arabidopsis thaliana* (Landsberg erecta) mature pollen. *Sexual Plant Reproduction* **2006**, 19, (4), 185-196.
48. Zhang, J.; Wu, L. S.; Fan, W.; Zhang, X. L.; Jia, H. X.; Li, Y.; Yin, Y. F.; Hu, J. J.; Lu, M. Z., Proteomic analysis and candidate allergenic proteins in *Populus deltoides* CL. "2KEN8" mature pollen. *Front Plant Sci* **2015**, 6, 548.
49. Pablos, I.; Eichhorn, S.; Briza, P.; Asam, C.; Gartner, U.; Wolf, M.; Ebner, C.; Bohle, B.; Arora, N.; Vieths, S.; Ferreira, F.; Gadermaier, G., Proteomic profiling of the weed feverfew, a neglected pollen allergen source. *Sci Rep* **2017**, 7, (1), 6049.

50. Niu, L.; Yuan, H.; Gong, F.; Wu, X.; Wang, W., Protein extraction methods shape much of the extracted proteomes. *Front Plant Sci* **2018**, 9, (802).
51. Karthikaichamy, A.; Deore, P.; Rai, V.; Bulach, D.; Beardall, J.; Noronha, S.; Srivastava, S., Time for multiple extraction methods in proteomics? A comparison of three protein extraction methods in the Eustigmatophyte Alga *Microchloropsis gaditana* CCMP526. *Omics* **2017**, 21, (11), 678-683.
52. Schmelter, C.; Funke, S.; Treml, J.; Beschnitt, A.; Perumal, N.; Manicam, C.; Pfeiffer, N.; Grus, F. H., Comparison of two solid-phase extraction (SPE) methods for the identification and quantification of porcine retinal protein markers by LC-MS/MS. *Int J Mol Sci* **2018**, 19, (12).
53. Jimenez-Lopez, J. C.; Morales, S.; Castro, A. J.; Volkmann, D.; Rodriguez-Garcia, M. I.; Alche, J. D., Characterization of profilin polymorphism in pollen with a focus on multifunctionality. *PLoS One* **2012**, 7, (2), e30878.
54. Alche, J. D.; Castro, A. J.; Jimenez-Lopez, J. C.; Morales, S.; Zafra, A.; Hamman-Khalifa, A. M.; Rodriguez-Garcia, M. I., Differential characteristics of olive pollen from different cultivars: biological and clinical implications. *J Investig Allergol Clin Immunol* **2007**, 17 Suppl 1, 17-23.
55. Hamman-Khalifa, A.; Castro, A. J.; Jimenez-Lopez, J. C.; Rodriguez-Garcia, M. I.; Alche Jde, D., Olive cultivar origin is a major cause of polymorphism for Ole e 1 pollen allergen. *BMC Plant Biol* **2008**, 8, 10.
56. Barber, D.; Moreno, C.; Ledesma, A.; Serrano, P.; Galan, A.; Villalba, M.; Guerra, F.; Lombardero, M.; Rodriguez, R., Degree of olive pollen exposure and sensitization patterns. Clinical implications. *J Investig Allergol Clin Immunol* **2007**, 17 Suppl 1, 11-6.
57. Rejon, J. D.; Delalande, F.; Schaeffer-Reiss, C.; Alche, J. D.; Rodriguez-Garcia, M. I.; Van Dorselaer, A.; Castro, A. J., The pollen coat proteome: at the cutting edge of plant reproduction. *Proteomes* **2016**, 4, (1).
58. Barral, P.; Tejera, M. L.; Trevino, M. A.; Batanero, E.; Villalba, M.; Bruix, M.; Rodriguez, R., Recombinant expression of Ole e 6, a Cys-enriched pollen allergen, in *Pichia pastoris* yeast: detection of partial oxidation of methionine by NMR. *Protein Expr Purif* **2004**, 37, (2), 336-43.
59. Seutter von Loetzen, C.; Reuter, A.; Spiric, J.; Schulenburg, T.; Bellinghausen, I.; Volker, E.; Vogel, L.; Rosch, P.; Schiller, D., Quality and potency profile of eight recombinant isoallergens, largely mimicking total Bet v 1-specific IgE binding of birch pollen. *Clin Exp Allergy* **2019**, 49, (5), 712-723.
60. Dubiela, P.; Kabasser, S.; Smargiasso, N.; Geiselhart, S.; Bublin, M.; Hafner, C.; Mazzucchelli, G.; Hoffmann-Sommergruber, K., Jug r 6 is the allergenic vicilin present in walnut responsible for IgE cross-reactivities to other tree nuts and seeds. *Sci Rep* **2018**, 8, (1), 11366.
61. Villalba, M.; Batanero, E.; Monsalve, R. I.; Gonzalez de la Pena, M. A.; Lahoz, C.; Rodriguez, R., Cloning and expression of Ole e I, the major allergen from olive tree pollen. Polymorphism analysis and tissue specificity. *J Biol Chem* **1994**, 269, (21), 15217-22.
62. Torres, M.; Alvarez-Garcia, E.; Bartra, J.; Alcantara, M.; Palomares, O.; Villalba, M.; Rodriguez, R., The allergenic structure of the thaumatin-like protein Ole e 13 is degraded by processing of raw olive fruits. *J Investig Allergol Clin Immunol* **2014**, 24, (3), 162-8.
63. Pablos, I.; Wildner, S.; Asam, C.; Wallner, M.; Gadermaier, G., Pollen Allergens for Molecular Diagnosis. *Curr Allergy Asthma Rep* **2016**, 16, (4), 31.
64. Pastor, C.; Cuesta-Herranz, J.; Cases, B.; Perez-Gordo, M.; Figueredo, E.; de las Heras, M.; Vivanco, F., Identification of major allergens in watermelon. *Int Arch Allergy Immunol* **2009**, 149, (4), 291-8.
65. Kim, J.-H.; Ahn, K.-M.; Kim, W.; Han, Y.; Kim, Y.-R.; Lee, S.; Kim, H.-Y., Identification of novel IgE-binding soybean allergens using serological analysis of a recombinant cDNA expression library (SEREX). *Food Sci Biotechnol* **2014**, 23, (4), 1037-1042.
66. Yoshimi, O.; Masanobu, K.; Hiroshi, Y.; Akemi, S.; Eiko, S.-K.; Susumu, T.; Takashi, H.-O.; Tomoko, Y.; Katsuhisa, U.; Hideyo, Y.; Kazuo, A.; Ikunoshin, K.; Kazutoh, T., Two-dimensional electrophoresis of *Malassezia* allergens for atopic dermatitis and isolation of Mal

- f 4 homologs with mitochondrial malate dehydrogenase. *Eur J Biochem* **1999**, 261, (1), 148-154.
67. Wagner, S.; Breiteneder, H.; Simon-Nobbe, B.; Susani, M.; Krebitz, M.; Niggemann, B.; Brehler, R.; Scheiner, O.; Hoffmann-Sommergruber, K., Hev b 9, an enolase and a new cross-reactive allergen from hevea latex and molds. Purification, characterization, cloning and expression. *Eur J Biochem* **2000**, 267, (24), 7006-14.
68. Zhao, F.; Elkelish, A.; Durner, J.; Lindermayr, C.; Winkler, J. B.; Rusmally, R. F.; Behrendt, H.; Traidl-Hoffmann, C.; Holzinger, A.; Kofler, W.; Braun, P.; von Toerne, C.; Hauck, S. M.; Ernst, D.; Frank, U., Common ragweed (*Ambrosia artemisiifolia* L.): allergenicity and molecular characterization of pollen after plant exposure to elevated NO₂. *Plant Cell Environ* **2016**, 39, (1), 147-64.
69. Kespohl, S.; Raulf, M., Mould allergens: Where do we stand with molecular allergy diagnostics?: Part 13 of the series Molecular Allergology. *Allergo J Int* **2014**, 23, (4), 120-125.
70. Fujita, C.; Moriyama, T.; Ogawa, T., Identification of cyclophilin as an IgE-binding protein from carrots. *Int Arch Allergy Immunol* **2001**, 125, (1), 44-50.
71. Fluckiger, S.; Fijten, H.; Whitley, P.; Blaser, K.; Cramer, R., Cyclophilins, a new family of cross-reactive allergens. *Eur J Immunol* **2002**, 32, (1), 10-7.
72. Welter, S.; Dolle, S.; Lehmann, K.; Schwarz, D.; Weckwerth, W.; Worm, M.; Franken, P., Pepino mosaic virus infection of tomato affects allergen expression, but not the allergenic potential of fruits. *PLoS One* **2013**, 8, (6), e65116.

For TOC Only