Can landscape characteristics help explain the different trends of Cantabrian brown bear subpopulations?

Cindy Lamamy¹, Giulia Bombieri², Alejandra Zarzo-Arias², Enrique González-Bernardo² and Vincenzo Penteriani^{2,3*}

¹*Université de Liège - Gembloux Agro-Bio Tech, Département BIOSE : Biosystem Engineering, Gestion des forêts et des espaces naturels 2, Passage des Déportés, 5030 Gembloux, Belgium*

²*Research Unit of Biodiversity (UMIB, UO-CSIC-PA), Oviedo University — Campus Mieres, Mieres, Spain*

³*Pyrenean Institute of Ecology (IPE), C.S.I.C., Avda. Montañana 1005, 50059 Zaragoza, Spain*

*Corresponding author: penteriani@ipe.csic.es; +34650432446

Abstract

A central challenge in animal conservation is to understand how a population may respond to different habitat characteristics, which may affect their growth and viability. At the beginning of the 20th century, the Cantabrian brown bear *Ursus arctos* population (north-western Spain) was separated into western and eastern subpopulations. Today, brown bears in the Cantabrian Mountains are recovering and the two subpopulations are reconnected. However, the western portion of the population represents ca. 90% of the entire population, the number of females with cubs-of-the-year has also shown a more rapid increase in the western subpopulation than in the eastern one and mean litter size is significantly larger in the west. By comparing the characteristics of the landscape used by brown bears in the western vs. eastern sectors of the population, we intended highlighting focal elements of landscape composition and structure that may help explain the differences in numbers and fecundity of these two subpopulations. We suggest that habitat use alone might not have the expected role in potentially explaining differences between subpopulations. Both the current positive trend of the Cantabrian population and our results seems to show that the dynamics affecting these subpopulations might be more complex than previously believed and cannot be understood on the basis of habitat analyses only. Suspicions may arise around direct human influences (e.g. persistence of poaching and/or bad practices during hunting) on the different trends exhibited by the two sectors of this endangered bear population.

Key words: habitat fragmentation, habitat use, human-modified landscapes, human-dominated landscapes, *Ursus arctos*

ACKNOWLEDGEMENTS

We thank the Administrations of the Gobierno del Principado de Asturias and the Junta de Castilla y León for providing the bear database. In particular, we would like to thank Teresa Sánchez Corominas, Pedro García-Rovés, Paloma Peón Torre and Víctor Vázquez of the Principado de Asturias, and María Ángeles Osorio Polo and David Cubero of the Junta de Castilla y León, for their continuous assistance during this study. The suggestions of two anonymous referees largely improved the manuscript. During this research, V.P. was financially supported by the Excellence Project CGL2017-82782-P financed by the Spanish Ministry of Science, Innovation and Universities, the Agencia Estatal de Investigación (AEI) and the Fondo Europeo de Desarrollo Regional (FEDER, EU). V.P. was also funded by a GRUPIN research grant from the Regional Government of Asturias (Ref.: IDI/2018/000151). G.B. was financially supported by a collaboration contract with the MUSE – Museo delle Scienze of Trento (Italy).

INTRODUCTION

The size and intrinsic dynamics (e.g. fecundity and mortality) of animal populations may be related to different properties of the landscape, such as habitat quality, vegetation composition and fragmentation of key habitat patches, i.e. the size and isolation of habitat patches, as well as the amount of habitat available (Fahrig 2003; Pardini et al. 2005; Honorato et al. 2015). Indeed, habitat loss and fragmentation may also decrease the viability of animal populations because in small populations the influence of stochastic processes may increase (Pardini et al. 2005; Penteriani et al. 2005a, b, 2008). Thus, a central challenge in animal conservation and management is to understand how a population (or a subpopulation) may respond to different habitat characteristics, which may affect their growth and viability. To this end, habitat analyses at different scales may help provide complementary information, from individual requirements and their responses to local environmental variability to population patterns and processes (Martin et al. 2012).

Brown bears *Ursus arctos* inhabiting the Cantabrian Mountains (north-western Spain) represent the westernmost population of the species' range in Europe. Of the four isolated brown bear populations in western Europe, i.e. Italian Apennines and eastern Alps, Pyrenees (France and eastern Spain) and Cantabrian, the latter represents the only population showing a trend to natural recovery (Gonzalez et al. 2016). At the beginning of the 20th century, the Cantabrian brown bear population was separated into western and eastern subpopulations by a 50–100 km wide strip of land, with bears in the two subpopulations declining in number until the mid-1990s (Palomero et al. 2007; Gonzalez et al. 2016). Today, brown bears in the Cantabrian Mountains are showing a positive trend and the two subpopulations are reconnected (Pérez et al. 2010; Gonzalez et al. 2016). However, the western portion of the population is considered to represent

ca. 90% of the entire population, for which a genetic census conducted in 2006 estimated approximately 203 bears (confidence interval CI 95% = 168–260) in the west and only 19 bears (CI 95% = 12–40) in the east (Pérez et al. 2014). The number of females with cubs-of-the-year (FCOY), a proxy of the population trend, has also shown a more rapid increase in the western subpopulation than in the eastern one (6 vs. 34 FCOY in 2016 and an average of 4.3 vs. 25 FCOY during the ten year period 2007–2016; unpublished data from Gobierno del Principado de Asturias, Junta de Castilla y León, Gobierno de Cantabria and Xunta de Galicia; Palomero et al. 2007). Mean litter size is 1.8 ± 0.2 cubs ($n = 596$ cubs) in the west and 1.3 ± 0.6 cubs ($n = 99$ cubs) in the east, with the former being significantly larger than the latter (Mann-Whitney test $Z = -4.66$, $P = 0.0001$, $n = 29$, 27) (Penteriani et al. 2018). Thus, despite the consistent positive trend in population size of the western subpopulation of Cantabrian brown bears, the eastern subpopulation has had a substantially smaller population increase (Martínez Cano et al. 2016). Such trends have characterised these two populations over the last forty years (Palomero et al. 2007; FAPAS/FIEP 2017).

The contrast in population growth between the western and eastern subpopulations might suggest high mortality rates in the eastern subpopulation and/or a relatively limited dispersal of bears out of the western sector. Indeed, some previous studies have suggested: (1) that the western sector is a zone of high human impact but with good habitat quality, whereas the eastern zone shows low human impact but suboptimal bear habitats (Wiegand et al. 1998); and (2) a narrow connection between the two subpopulations, apparently a consequence of the development of communication infrastructures (García et al. 2007; Pérez et al. 2010; San Miguel et al. 2012; Mateo-Sánchez et al. 2014b). However, even if female philopatry has been highlighted as one of the causes of the slow rate of recolonization of the eastern sector

(Palomero et al. 2007), this cannot explain the apparent lower reproductive rates or higher mortality in this subpopulation, which is clearly exhibiting a slower increase than the western subpopulation. Earlier studies have also indicated (3) that the principal mechanism determining changes in FCOY is the effect of environmental factors (e.g. resources availability) on mortality rates, as individual-based modelling seems to point out (Martínez Cano et al. 2016); and (4) a prevalence of low quality food-items in the eastern sector of the Cantabrian Mountains (Clevenger et al. 1992; Naves et al. 2003, 2006). For example, during the hyperphagic season, eastern bears consume a high proportion of less energetic foods (e.g. graminoids and forbs), which may engender a lower accumulation of fat and, consequently, could restrict the duration of hibernation and result in less reproductive success than the western subpopulation (Naves et al. 2006).

Here, we compared the characteristics of the landscape used by Cantabrian brown bears in the western *vs.* eastern sectors of the population, with the main aim of highlighting focal elements of landscape composition and structure that may help explain the above-cited differences in numbers and fecundity of these two bear subpopulations. A better knowledge of the habitat used by individuals of both these subpopulations can offer crucial insights for the conservation and management of this small, isolated and endangered brown bear population at the limit of the species' occidental range.

METHODS

Study area.—Our habitat analyses encompassed most of the Cantabrian range currently occupied by brown bears (Asturias, León and Palencia provinces). The Cantabrian

Mountains are one of the main mountain range systems in Spain, stretching over 300 km across the northern part of the country, running east-west parallel to the Atlantic coast. Maximum elevation is 2648 m a.s.l. and average elevation is around 1100 m (Martínez Cano et al. 2016). As a general rule, the eastern sector of the Cantabrian Mountains is characterised by higher average altitudes (1300 m a.s.l.) and lower average slopes (21%) than the western area (700 m a.s.l. and 34%) (Naves et al. 2003). The Cantabrian Mountains are characterized by an oceanic climate, with pronounced differences on opposite slopes of the mountain range (Pato and Obeso 2012; Mateo-Sánchez et al. 2014b; Rocas-Díaz et al. 2014): northern slopes exhibit mild winters and rainy summers, with fewer days of frost and less snow accumulation in winter, while on southern slopes winters are longer and colder, with ample snow and fresh and short summers without drought. The greater thermal amplitude and lower rainfall is accentuated towards the east due to the higher average altitude of the range. More specifically, average rainfall ranges from more than 900 mm per year in the lowest zones to more than 2000 mm at the highest elevations in the west; due to the Foehn effect, rainfall drops to 700 mm per year in some areas of the eastern sector.

Subalpine matorral (*Juniperus communis*, *Vaccinium uliginosum*, *V. myrtillus* and *Arctostaphylos uva-ursi*) dominates mountain areas above the treeline (~1700 m) (Martínez Cano et al. 2016). Sites below 1500 m a.s.l. mainly consist of deciduous forests of oak *Quercus petraea*, beech *Fagus sylvatica* and chestnut *Castanea sativa*, with bilberry dominating the understory (Pato and Obeso 2012). In some areas, former forests have been converted into pasture and brushwood (*Genista*, *Cytisus*, *Erica* and *Calluna*) through human activities (Naves et al. 2003, 2006).

Throughout the entire study area, bears are surrounded by a matrix of urbanised and cultivated areas, with a high density of transport routes (Mateo-Sánchez et al.

2016). There are differences between the areas occupied by humans in both bear subpopulations, with an average population density of 4.9 inhabitants km⁻¹ in the municipalities occupied by the eastern subpopulation and more than double that figure (10.9 inhabitants km⁻¹) in the western area (INE 2017). Throughout the study area the main economic activity is the extensive breeding of livestock, particularly cattle, followed by mining and tourism, as well as other activities such as hunting and timber harvest (Fernández-Gil et al. 2006). Cattle density (bovine, equine, ovine and caprine), equated to 'heads of adult cattle' (UGM), does not differ substantially between the municipalities where both bear subpopulations are located (18.5 UGM km⁻¹ in the eastern area and 15.6 UGM km⁻¹ in the western one) (Real Decreto 1131/2010).

Brown bear data.—The locations of brown bears were obtained from: (1) direct bear observations that were georeferenced by personnel of the Principado de Asturias and Junta de Castilla y León, primarily by the Patrulla Oso, i.e. the Bear Patrol, of the Principado de Asturias and of the Junta de Castilla y León, as well as by all the other guards of both regional governments, by the Asturian Foundation for the Conservation of Wildlife (FAPAS, Fondo para la Protección de los Animales Salvajes), the FOA (Fundación Oso de Asturias) and the Brown Bear Foundation (FOP, Fundación Oso Pardo); (2) remotely triggered cameras that were randomly placed by the FAPAS and Bear Team during the last twenty years, mainly in forested areas where bears are less visible (FAPAS/FIEP 2017) and (3) personal georeferenced observations of the authors. Thus, the long-term monitoring of the Cantabrian population, which started at the beginning of the nineteen-nineties, is essentially based on direct sightings (i.e. direct observation and pictures obtained from camera traps) and the location of indirect signs of presence, i.e. footprints, fur and scats, as well as records of damage caused by bears to livestock, beehives, crops, and human activities and infrastructures. Consequently, locations were both the result of systematic observations and random observations. For the western sector of the Cantabrian Mountains it was possible to collect 7194 bear locations (from 1992 to 2017), whereas for the eastern sector 1541 bear locations were available (from 1985 to 2017; Fig. 1). We collected data following the guidelines of the American Society of Mammalogists (Sikes and The Animal Care and Use Committee of the American Society of Mammalogists 2016).

Landscape characterisation.—The delimitation of the bear subpopulations can be artificially represented by the A-66 highway from Oviedo to León (Fig. 1). In order to clearly distinguish the main habitat of the two subpopulations, brown bear locations within 3000 metres of both sides of the highway were removed from the analyses.

Moreover, because we had relatively few observations between 1985 and 2000 (5%) and the landscape characteristics might have been considerably different before 2000, we removed such observations from our analyses and only included the most recent data (2000-2017). The multi-scale approach to determining bear habitat use was based on four different spatial scales, i.e. circular plots with radii of 500, 1000, 2000 and 3000 m, respectively, around each bear location (see also Penteriani et al. in press). These plots covered most of the bear's movements in the study area (Clevenger and Purroy 1988; Penteriani et al. 2017). This approach is based on Johnson's (1980) model, which assumes that animals make decisions about basic resources (e.g. breeding sites, cover and food) at retrogressively smaller scales and in a hierarchical manner, ranging from local factors within habitat patches to broad scale determinants of home range selection and population distribution at landscape levels (Gastón et al. 2017). To understand relevant habitat patterns as perceived by animals in complex landscapes, it is necessary to identify the scales at which particular environmental variables influence habitat selection and, therefore, it is recommended that habitat modelling follows a multi-scale approach (Gastón et al. 2017). In fact, what may seem appropriate at one scale may have little importance at another, making the results dependent on the scale (Ciarniello et al. 2007). Indeed, habitat relationships are highly sensitive to the scale at which explanatory variables are assessed for brown bears in the Cantabrian Mountains (Mateo-Sánchez et al. 2014a).

The landscape characteristics of both brown bear subpopulations of the Cantabrian Mountains were characterized by three categories of variables: (1) landscape composition; (2) landscape structure (i.e. fragmentation); and (3) human footprint, i.e. landscape variables associated with humans. As to landscape composition, several variables were taken into account at the above-mentioned four spatial scales. The

altitude of the plot centres (m a.s.l.) was obtained from an altitude raster layer of the Iberian Peninsula (<http://www.diva-gis.org>) and calculated by the QGIS software (QGIS Geographic Information System 2015) using the "Point Sampling" tool. Ruggedness was determined from the total length (m) of the linear development of the elevation curves (*\$length* function of the QGIS software). The same approach was applied to obtain river density. The landscape GIS layers extracted from the Centro de Descargas del CNIG (IGN; Cartografía Temática Ambiental of the Principado de Asturias; Hojas del Mapa de Vegetación, Litología, Roquedos y Hábitat del Oso. 2011. Escala 1:25000. © Principado de Asturias, Spain) were reclassified into eight habitat types: (1) human settlements and infrastructures (which belong to the human footprint category); (2) crops; (3) fruit trees; (4) forest (i.e. hardwood and mixed forests); (5) conifer forests; (6) pasture; (7) shrubland; and (8) rocky areas. The *Intersect* tool was used to obtain the percentage of land use in each plot. This function extracts the information contained in the plots, and then the *\$area* function determines the area (%) occupied by each habitat type in the different plots that we analysed.

Linear structures belonging to the human footprint category, such as road and trail densities, were analysed using the same *\$length* function. In addition, the shortest distances from human infrastructures were calculated by using the *Hub* distance function.

Statistical analyses.—Statistical analyses only included those habitat parameters with less than 20% of zeros in all the buffers considered. The variables ‘conifer’ and ‘forest’ were highly correlated ($r = 0.85$) and, therefore, we excluded the variable ‘conifer’, which represents a lower quality habitat for bears than hardwood or mixed forests (Clevenger et al. 1997).

To explore possible differences in landscape characteristics between the two subpopulations, we built GLMMs with a binomial distribution which included the location of bears either to the east (0) or west (1) subpopulation as the response variable and the landscape parameters as predictor variables. Specifically, for each of the four spatial scales, we constructed a set of competing models that included all possible combinations of predictor variables, from the simplest null model (intercept model only) to a complete model that included all the landscape parameters.

The best competing model or set of models was chosen based on Akaike's information criterion (AICc). Models with a $\Delta AICc$ below 2 were considered as equally competitive. Values of $\Delta AICc$ and weighted AICc, indicating the probability that the model selected was the best among the competing candidates (Table 1), were calculated as well. Parameter coefficients and the relative importance value (RIV) of each explanatory variable were generated by employing model averaging on the 95% confidence set (Burnham and Anderson 2002). Models were run in R v. 3.5.1 statistical software (R Foundation for Statistical Computing 2018) using the *lme4* package (Bates et al. 2015). Model generation and model averaging were performed using the *MuMIn* package (Bartoń 2013).

Furthermore, in order to characterize landscape structure in the western and eastern sectors, we created two raster files using QGIS software and the *LecoS* (Landscape ecology Statistics) tool (Jung 2016). This tool allows the overlay of a polygon and a raster in order to study the fragmentation of the landscape. The east and west sectors were defined with the external points of the bear locations. A polygon for each sector was therefore set in this way to overlay with the land cover raster. The following indexes were calculated for forest and shrubland classes in Fragstats software (McGarigal et al. 2012): Edge density (ED; equal to the sum of the lengths (m) of all

edge segments in the landscape, divided by the total landscape area (m²), multiplied by 10,000 to facilitate the display of the results), Largest Patch Index (LPI; at the class level quantifies the percentage of total landscape area comprised by the largest patch), Mean Patch Size (MPS), Mean Nearest-Neighbour distance (MNN; defined using simple Euclidean geometry as the mean shortest straight-line distance between the focal patch and its nearest neighbour of the same class) and Number of patches (NP). At the landscape scale, Shannon (SHDI) and Simpson (SIDI) diversity indices were calculated.

RESULTS

Our habitat analyses showed some relevant differences in the landscape composition and structure used by brown bears within each subpopulation (Fig. 2). At all scales, the best model always included altitude, ruggedness, percentage of forest and percentage of shrubland. At the 1000 and 2000 m scales, the best model also included river length. At the largest scale (3000 m), road length, trail length and rocky areas were also included in the best model together with all the above-mentioned variables (Table 1). The importance of these variables in explaining the difference between the two subpopulations was also supported by their values of p, CI and RIV (Table 2). Specifically, at each spatial scale, western brown bears used areas characterised by higher values of ruggedness and shrubland than eastern bears, which are generally located at higher altitudes and in more forested areas (Table 2; see also SD1 and Fig. 2). At the largest spatial scale (3000 m), western bears used more rocky landscapes and inhabited areas characterised by a greater human footprint, i.e. areas with longer road and trail lengths, compared to eastern bears.

From a structural point of view (Table 3): (1) forest cover was found to be more fragmented in the western sector, as revealed by the larger number of patches as well as the lower values of largest patch index and mean patch size compared to the east; similarly, (2) shrubland also seemed to be more fragmented in the west, as demonstrated by the larger number of patches; (3) western and eastern sectors showed similar values for edge density, mean nearest-neighbour distance and landscape diversity (calculated by Shannon's and Simpson's indexes).

DISCUSSION

The best models for brown bear habitat use in the Cantabrian Mountains were at all scales characterised by altitude, ruggedness and percentage of forest and shrubland, which have the potential to most effectively separate the western from eastern bear habitat use. As a consequence, if habitat use plays a role, at least partially, in determining the different trends observed in the two subpopulations, these four landscape descriptors might be the most important in explaining such dissimilarities. These variables may be even more relevant than the human footprint considered by our analyses, which was shown to be higher in the denser and more productive western subpopulation. And this, despite the fact that habitat suitability models seems to indicate that brown bears avoid human infrastructures and agricultural areas in the Cantabrian Mountains (Martin et al. 2012). This may suggest that human pressure in terms of infrastructures (road and trail lengths) may not be the factor limiting the growth of the eastern population. Indeed, the presence of roads and trails may have minimal influence on bear distribution as long as they do not represent impassable barriers (Mateo-Sánchez et al. 2015). However, when measuring only indirectly human footprint

through proxies, these parameters do not necessarily represent the intensity of the pressure that man actually exerts on a territory. Thus, such a possibility might still be interpreted with caution and deserves further attention.

As also recently highlighted by Martínez Cano et al. (2016), higher fecundity and population size in the western sector might be explained in part by the habitat used by bears: (1) the western subpopulation inhabits a sector characterised by a greater availability in shelters, such as rocky or rugged areas, as also suggested by Mateo-Sánchez et al. (2014, 2016). Indeed, in the Cantabrian Mountains, a human-modified landscape largely dominated by the presence of human infrastructures, rocky and rugged areas may allow bears to more easily avoid human presence and activities. Also, (2) shrubland, which is more frequently used by western brown bears, might provide an additional source of shelter and trophic resources. Because of the importance of habitat connectivity for brown bears in the Cantabrian Mountains (Mateo-Sánchez et al. 2014a), shrubland may also increase matrix permeability and habitat continuity in the western sector, where this habitat of high value for bears links the more fragmented forest patches of the western landscape matrix.

Forest cover provides a high quality habitat where brown bears can shelter and feed (Clevenger et al. 1992b, Mattson et al. 2001; Schrag et al. 2008). However, even if the use of forested landscapes in the eastern sector is higher than in the west, it is the eastern subpopulation which exhibits fewer bears and lower fecundity. This result might suggest that: (1) forest cover, which is still sufficiently large for bear requirements in the west, does not represent a limiting factor for both subpopulations; and (2) heterogeneity of landscape composition is also important, yet previous multi-scale models have shown the value of having large areas of forest intermixed with shrubland (Mateo-Sánchez et al. 2014a). Landscape fragmentation is often perceived as negative

due to habitat loss and the breaking apart of habitat (Fahrig 2003). However, habitat fragmentation is not always unfavourable for species. Fragmentation is more than just habitat loss because, by creating small and isolated patches, it also changes the properties of the remaining habitat resulting in a more complex habitat landscape (Fahrig 2003), which can have a positive effect on biodiversity (Huffaker 1958). Actually, the higher human presence and more fragmented forest habitat in the west do not seem to have the expected negative effect on this subpopulation. The same amount of a habitat subdivided into several smaller areas may ensure better habitat distribution and heterogeneity in the matrix (e.g. shorter distances between patches of the same category) (Fahrig 2003) and limits the effect of local disturbances (Roff 1974a, b; den Boer 1981). For example, the different arrangement of shrub and forest habitats in the western vs. the eastern sectors might have an impact on the ability to access diverse trophic resources. As a final conclusion, we suggest that landscape characteristics alone might not have the expected role in potentially explaining differences between subpopulations. Indeed, even if some habitat patches offer additional shelter and food to the western subpopulation, characterised by a higher bear density and fecundity, western bears also use less forested areas (a well-known brown bear high-quality habitat for both food and cover) as well as areas with a higher presence of human infrastructures.

Fifteen years ago, the Naves et al. (2003) analysis of the Cantabrian subpopulations drew a not-too-optimistic picture of the condition of brown bear habitat in northern Spain. Indeed, the authors asserted that: (a) the two subpopulations occupied a high proportion of suboptimal habitats; and (b) over the long term, the Cantabrian Mountains could be a demographic sink due to the combination of high human impact and low natural habitat quality. However, both the current positive trend of the

Cantabrian population and our results seems to show that the dynamics affecting the Cantabrian brown bear subpopulations might be more complex than previously believed and cannot be understood on the basis of habitat analyses only. Indeed, numbers of bears and, particularly, FCOY have increased in both subpopulations since the beginning of this century, even if at a slower rate in the eastern sector of the Cantabrian Mountains. Moreover, since trend data on both subpopulations have been available (i.e. the last forty years), the eastern subpopulation has always been shown to be 1/3 – 1/4 that of the western one (Palomero et al. 2007; FAPAS/FIEP 2017). If the habitat use displayed by the two brown bear subpopulations of the Cantabrian Mountains only offers limited help in explaining the differences in number and fecundity exhibited by the evolution of the two subpopulations, other factors might be acting on them. Given that (a) previous population models have partially failed in forecasting the dynamics and trends of brown bears in the Cantabrian Mountains and (b) patterns of increase and fecundity continue to be different between the west and the east (Gonzalez et al. 2016; Penteriani et al. 2018), we suggest that future analyses should be focused more on how the human environment can achieve coexistence with bears than on the natural landscape (Zarzo-Arias et al. 2018). If habitat alone cannot help explain the different properties of the two Cantabrian bear subpopulations, then suspicions may arise around direct human influences on the different trends exhibited by the two sectors of this endangered bear population. For instance, persistence of poaching and/or bad practices during wild boar *Sus scrofa* hunting might be more pronounced in the eastern sector of the Cantabrian Mountains, engendering a higher mortality of more vulnerable bear classes, such as cubs and subadults.

SUPPLEMENTARY DATA

Supplemental Data SD1. Values (mean $\pm$ SD, range) of the significantly different habitat variables (see Table 2) that were used to characterise the western (1) vs. eastern (0) sectors of the Cantabrian Mountains inhabited by brown bears.

Spatial scale	Parameters	East(0) West(1)	Mean $\pm$ SD (range)
500 m			
	Altitude (m a.s.l.)	0	1294 $\pm$ 203.9 (647 - 2184)
		1	1062 $\pm$ 341.3 (112 - 1987)
	Ruggedness (m)	0	2637 $\pm$ 1097.0 (0 - 6084)
		1	4119 $\pm$ 976.6 (0 - 8925)
	% forest	0	48.0 $\pm$ 32.7 (0 - 98.4)
		1	43.7 $\pm$ 29.7 (0 - 98.4)
	% shrubland	0	26.4 $\pm$ 28.6 (0 - 98.5)
		1	35.6 $\pm$ 28.5 (0 - 98.4)
1000 m			
	Altitude (m a.s.l.)	0	1294 $\pm$ 203.9 (647 - 2184)
		1	1062 $\pm$ 341.3 (112 - 1987)
	Ruggedness (m)	0	10756 $\pm$ 3951.0 (0 - 21355)
		1	16298 $\pm$ 3002.8 (0 - 28749)
	River length (m)	0	2165 $\pm$ 1424 (0 - 6718.0)
		1	1759 $\pm$ 1141.4 (0 - 6334)
	% forest	0	46.8 $\pm$ 27.2 (0 - 98.5)
		1	41.5 $\pm$ 22.5 (0 - 98.4)
	% shrubland	0	27.4 $\pm$ 24.2

			(0 - 98.2)
	1		37.7 ± 22.0 (0 - 98.4)
2000 m			
Altitude (m a.s.l.)	0		1294 ± 203.9 (647 - 2184)
	1		1062 ± 341.3 (112 - 1987)
Ruggedness (m)	0		43417 ± 14626 (0 - 83810)
	1		63474 ± 10310 (2971 - 87081)
River length (m)	0		8127 ± 3535 (0 - 17507)
	1		6009 ± 2629 (0 - 17155)
% shrubland	0		29.7 ± 20.7 (0 - 94.1)
	1		43.2 ± 16.9 (0 - 92.3)
3000 m			
Altitude (m a.s.l.)	0		1294 ± 203.9 (647 - 2184)
	1		1062 ± 341.3 (112 - 1987)
Ruggedness (m)	0		97059 ± 32574 (0 - 197058)
	1		140315 ± 21228 (17600 - 184031)
River length (m)	0		17976 ± 6561 (0 - 32792)
	1		12802 ± 4459 (0 - 32309)
Road length (m)	0		5999 ± 5078 (0 - 24268)
	1		7460 ± 5148 (0 - 31303)
Trail length (m)	0		1609 ± 2375 (0 - 14783)
	1		4831 ± 4550 (0 - 366195)
% forest	0		40.8 ± 17.8 (0 - 82.7)

		1	36.2± 15.2 (0 - 90.8)
		0	30.8 ± 19.0 (0 - 80.2)
	% shrubland	1	45.1 ± 14.5 (0 - 86.8)
		0	5.9 ± 9.0 (0 - 62.2)
	% rocky areas	1	4.2 ± 4.7 (0 - 38.2)

389

390

REFERENCES

- 1131/2010 RD (2010) Real Decreto 1131/2010, de 10 de septiembre, por el que se establecen los criterios para el establecimiento de las zonas remotas a efectos de eliminación de ciertos subproductos animales no destinados a consumo humano generados en explotaciones ganaderas.
- Bartoń K (2013) Model selection and model averaging based on information criteria (AICc and alike). *Compr R Arch Netw* 1:13.
- Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw* 67:1–48. doi: doi:10.18637/jss.v067.i01
- Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical information-theoretic approach. Springer, Berlin
- Ciarniello LM, Boyce MS, Seip DR, Heard DC (2007) Grizzly bear habitat election is scale dependent. *Ecol Appl* 17:1424–1440.
- Clevenger AP, Purroy FJ (1988) El oso en León. Universidad de León, Servicio de Publicaciones, León, Spain
- Clevenger AP, Purroy FJ, Campos MA (1997) Habitat assessment of a relict brown bear *Ursus arctos* population in northern Spain. *Biol Conserv* 80:17–22.
- Clevenger AP, Purroy FJ, Pelton MR (1992) Food habits of brown bears (*Ursus arctos*) in the Cantabrian Mountains, Spain. *J Mammal* 73:415–421.
- den Boer P (1981) On the survival of populations in a heterogeneous and variable environment. *Oecologia* 50:39–53.
- Fahrig L (2003) Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol*

413 Syst 34:487–515. doi: 10.1146/annurev.ecolsys.34.011802.132419

414 FAPAS/FIEP (2017) El Oso. Pirineos y Cordillera Cantábrica. Gráficas Muñiz, Gijón,
 415 Spain

416 Fernández-Gil A, Naves J, Delibes M (2006) Courtship of brown bears *Ursus arctos* in
 417 northern Spain: phenology, weather, habitat and durable mating areas. *Wildlife*
 418 *Biol* 12:367–373. doi: 10.2981/0909-6396(2006)12[367:COBBUA]2.0.CO;2

419 García P, Lastra J, Marquínez J, Nores C (2007) Detailed model of shelter areas for the
 420 Cantabrian brown bear. *Ecol Inform* 2:297–307. doi: 10.1016/j.ecoinf.2007.08.003

421 Gastón A, Ciudad C, Mateo-Sánchez MC, et al (2017) Species’ habitat use inferred
 422 from environmental variables at multiple scales: How much we gain from high-
 423 resolution vegetation data? *Int J Appl Earth Obs Geoinf* 55:1–8. doi:
 424 10.1016/j.jag.2016.10.007

425 Gonzalez EG, Blanco JC, Ballesteros F, et al (2016) Genetic and demographic recovery
 426 of an isolated population of brown bear *Ursus arctos* L., 1758. *PeerJ* 4:e1928. doi:
 427 10.7717/peerj.1928

428 Honorato R, Crouzeilles R, Ferreira MS, Grelle CE V (2015) The effects of habitat
 429 availability and quality on small mammals abundance in the Brazilian Atlantic
 430 Forest. *Nat Conserv* 13:133–138. doi: 10.1016/j.ncon.2015.11.010

431 Huffaker C (1958) Experimental studies on predation: dispersion factors and predator-
 432 prey oscillations. *Hilgardia* 27:343–383.

433 INE (2017) Cifras Oficiales de Población de los Municipios Españoles: Revisión del
 434 Padrón Municipal.

435 Johnson D (1980) The comparison of usage and availability measurements for
 436 evaluating resource preference. *Ecology* 61:65–71.

437 Jung M (2016) A python plugin for automated landscape ecology analysis. *Ecol Inform*
 438 31:18–21.

439 Martin J, Revilla E, Quenette PY, et al (2012) Brown bear habitat suitability in the
 440 Pyrenees: Transferability across sites and linking scales to make the most of scarce
 441 data. *J Appl Ecol* 49:621–631. doi: 10.1111/j.1365-2664.2012.02139.x

442 Martínez Cano I, González Taboada F, Naves J, et al (2016) Decline and recovery of a
 443 large carnivore : environmental change and long- term trends in an endangered
 444 brown bear population. *Proc R Soc B* 283:20161832. doi: 10.1098/rspb.2016.1832

445 Mateo-Sánchez MC, Balkenhol N, Cushman S, et al (2015) A comparative framework
 446 to infer landscape effects on population genetic structure: are habitat suitability
 447 models effective in explaining gene flow? *Landsc Ecol* 30:1405–1420. doi:
 448 10.1007/s10980-015-0194-4

449 Mateo-Sánchez MC, Cushman S a., Saura S (2014a) Scale dependence in habitat
 450 selection: the case of the endangered brown bear (*Ursus arctos*) in the Cantabrian
 451 Range (NW Spain). *Int J Geogr Inf Sci* 28:1531–1546. doi:
 452 10.1080/13658816.2013.776684

453 Mateo-Sánchez MC, Cushman SA, Saura S (2014b) Connecting endangered brown bear
 454 subpopulations in the Cantabrian Range (north-western Spain). *Anim Conserv*
 455 17:430–440. doi: 10.1111/acv.12109

456 Mateo-Sánchez MC, Gastón A, Ciudad C, et al (2016) Seasonal and temporal changes
 457 in species use of the landscape: how do they impact the inferences from multi-scale

458 habitat modeling? *Landsc Ecol* 31:1261–1276. doi: 10.1007/s10980-015-0324-z

459 Mattson DJ, Kendall KC, Reinhart DP (2001) Whitebark pine, grizzly bears and red
460 squirrels. In: Tomback DF, Arno SF, Keane RE (eds) Whitebark pine
461 communities: ecology and restoration. Island Press, Washington DC., pp 121–136

462 McGarigal K, Cushman SA, Ene E (2012) FRAGSTATS v4: Spatial Pattern Analysis
463 Program for Categorical and Continuous Maps.

464 Naves J, Fernández-Gil A, Rodríguez C, Delibes M (2006) Brown Bear Food Habits At
465 the Border of Its Range: a Long-Term Study. *J Mammal* 87:899–908. doi:
466 10.1644/05-MAMM-A-318R2.1

467 Naves J, Wiegand T, Revilla E, Delibes M (2003) Endangered species constrained by
468 natural and human factors: the case of brown bears in northern Spain. *Conserv Biol*
469 17:1276–1289.

470 Palomero G, Ballesteros F, Nores C, et al (2007) Trends in Number and Distribution of
471 Brown Bear Females with Cubs-of-the-year in the Cantabrian Mountains, Spain.
472 *Ursus* 18:145–157. doi: 10.2192/1537-6176(2007)18[145:TINADO]2.0.CO;2

473 Pardini R, Marques S, Souza D, et al (2005) The role of forest structure , fragment size
474 and corridors in maintaining small mammal abundance and diversity in an Atlantic
475 forest landscape. *Biol Conserv* 124:253–266. doi: 10.1016/j.biocon.2005.01.033

476 Pato J, Obeso JR (2012) Growth and reproductive performance in bilberry (*Vaccinium*
477 myrtillus) along an elevation gradient. *Ecoscience* 19:59–68. doi: 10.2980/19-1-
478 3407

479 Penteriani V, Del Mar Delgado M, López-Bao JV, et al (2017) Patterns of movement of

480 released female brown bears in the Cantabrian Mountains, northwestern Spain.
 481 *Ursus* 28:165–170. doi: 10.2192/URSU-D-16-00012.1

482 Penteriani V, Otalora F, Ferrer M (2005a) Floater survival affects population
 483 persistence. The role of prey availability and environmental stochasticity. *Oikos*
 484 108:523–534. doi: 10.1111/j.0030-1299.2005.13514.x

485 Penteriani V, Otalora F, Ferrer M (2008) Floater mortality within settlement areas can
 486 explain the Allee effect in breeding populations. *Ecol Modell.* doi:
 487 10.1016/j.ecolmodel.2007.11.009

488 Penteriani V, Otalora F, Sergio F, Ferrer M (2005b) Environmental stochasticity in
 489 dispersal areas can explain the “mysterious” disappearance of breeding
 490 populations. *Proc R Soc B Biol Sci* 272:1265–1269. doi: 10.1098/rspb.2005.3075

491 Penteriani V, Zarzo-Arias A, Bombieri G, et al (2018) Density and reproductive
 492 characteristics of female brown bears in the Cantabrian Mountains, NW Spain. *Eur*
 493 *Zool J* 85:313–321.

494 Penteriani V, Zarzo-Arias A, Delgado MM, et al in press. Female brown bears use areas
 495 with infanticide risk in a spatially confined population. *Ursus*.

496 Pérez T, Naves J, Vázquez JF, et al (2010) Evidence for improved connectivity between
 497 Cantabrian brown bear subpopulations. *Ursus* 21:104–108. doi:
 498 10.2192/09SC018.1

499 Pérez T, Naves J, Vázquez JF, et al (2014) Estimating the population size of the
 500 endangered Cantabrian brown bear through genetic sampling. *Wildlife Biol*
 501 20:300–309. doi: 10.2981/wlb.00069

502 QGIS Geographic Information System (2015) Open Source Geospatial Foundation
 503 Project <<http://qgis.osgeo.org>>.

504 R Foundation for Statistical Computing (2018) R: A Language and Environment for
 505 Statistical Computing.

506 Roces-Díaz J V., Jiménez-Alfaro B, Álvarez-Álvarez P, Álvarez-García MA (2014)
 507 Environmental niche and distribution of six deciduous tree species in the spanish
 508 atlantic region. *IForest* 8:214–221. doi: 10.3832/ifor1183-008

509 Roff D (1974a) Spatial heterogeneity and the persistence of populations. *Oecologia*
 510 15:245–258.

511 Roff D (1974b) The analysis of a population model demonstrating the importance of
 512 dispersal in a heterogeneous environment. *Oecologia* 15:259–275.

513 San Miguel A, Ballesteros F, Blanco JC, Palomero (eds.) G (2012) Manual de buenas
 514 prácticas para la gestión de corredores oseros en la Cordillera Cantábrica.
 515 Fundación Oso Pardo (FOA) and Ministerio de Agricultura, Alimentación y Medio
 516 Ambiente, Madrid, Spain

517 Schrag AM, Bunn AG, Graumlich LJ (2008) Influence of bioclimatic variables on tree-
 518 line conifer distribution in the Greater Yellowstone Ecosystem : implications for
 519 species of conservation concern. *J Biogeogr* 35:698–710. doi: 10.1111/j.1365-
 520 2699.2007.01815.x

521 Sikes RS, The Animal Care and Use Committee of the American Society of
 522 Mammalogists (2016) Guidelines of the American Society of Mammalogists for
 523 the use of wild mammals in research and education. *J Mammal* 97:663–688.

524 Wiegand T, Naves J, Stephan T, Fernandez A (1998) Assessing the Risk of Extinction
525 for the Brown Bear (*Ursus arctos*) in the Cordillera Cantabrica , Spain. Ecol
526 Monogr 68:539–570.

527 Zarzo-Arias A, Delgado M, Ordiz A, et al (2018) Brown bear behaviour in human-
528 modified landscapes: The case of the endangered Cantabrian population , NW
529 Spain. Glob Ecol Conserv 16:e00499. doi: 10.1016/j.gecco.2018.e00499

530

Table 1. Values of AICc, $\Delta AICc$ and Weighted AICc of the best models or sets of models ($\Delta AICc < 2$) for the four spatial scales at which potential differences of brown bear habitat use between the western and eastern sectors of the Cantabrian Mountains were analysed.

	<i>Selected models*</i>	<i>AICc</i>	<i>$\Delta AICc$</i>	<i>Weighted AICc</i>
500 m	1/2/3/4	4356.39	0	1
1000 m	1/2/3/4/5	3946.89	0	0.98
2000 m	1/3/4/5	3881.63	0	0.7
	1/2/3/4/5	3883.29	1.66	0.3
3000 m	1/2/3/4/5/6/7/8	3104.07	0	1

* 1 = altitude; 2 = ruggedness; 3 = % forest; 4 = % shrubland; 5 = river length; 6 = road length; 7 = trail length; 8 = rocky areas.

Table 2. Model-averaged coefficients and relative importance values (RIV) of each explanatory variable included in the best model or set of models ($\Delta\text{AICc} < 2$) describing the difference in habitat use between the western and eastern brown bear subpopulations in the Cantabrian Mountains. β = estimate; SE = standard error; CI = confidence interval. P-values were < 0.0001 for all variables at all scales. Values are shown at each of the spatial scales considered in the analyses (i.e., 500, 1000, 2000 and 3000 m).

Spatial scales	Explanatory variables	β	SE	CI	RIV
500 m	Intercept	3.05	0.07	2.90; 3.19	
	Altitude	-1.35	0.06	-1.46; -1.23	1
	Ruggedness	1.68	0.05	1.58; 1.78	1
	% forest	-0.29	0.05	-0.39; -0.19	1
	% shrubland	0.40	0.06	0.29; 0.51	1
Deviance = 38.0 %					
1000 m	Intercept	3.28	0.08	3.12; 3.44	
	Altitude	-1.60	0.07	-1.74; -1.45	1
	Ruggedness	1.59	0.05	1.49; 1.69	1
	River length	-0.37	0.04	-0.46; -0.29	1
	% forest	-0.16	0.06	-0.27; -0.06	0.98
	% shrubland	0.61	0.06	0.49; 0.72	1
Deviance = 43.9 %					
2000 m	Intercept	3.18	0.08	3.02; 3.33	
	Altitude	-1.51	0.07	-1.65; -1.37	1
	Ruggedness	1.26	0.05	1.17; 1.35	1
	River length	-0.48	0.04	-0.56; -0.39	1
	% shrubland	0.81	0.05	0.71; 0.91	1
Deviance = 44.9 %					
3000 m	Intercept	3.88	0.10	3.67; 4.09	
	Altitude	-1.12	0.09	-1.29; -0.94	1
	Ruggedness	1.99	0.07	1.85; 2.14	1
	River length	-0.78	0.05	-0.87; -0.68	1
	Road length	0.66	0.06	0.55; 0.78	1
	Trail length	1.18	0.07	1.04; 1.33	1
	% forest	-0.45	0.09	-0.61; -0.28	1
	% shrubland	0.66	0.08	0.51; 0.82	1
	% rocky areas	-0.38	0.05	-0.49; -0.28	1
Deviance = 58.4 %					

Table 3. Characterisation of the habitat structure of the western *vs.* eastern brown bear subpopulations in the Cantabrian Mountains. For both forest and shrubland cover, we took into consideration: Edge Density (m/ha, ED), Number of Patches (NP), Largest Patch Index (% , LPI), Mean Patch Size (ha, MPS) and Mean Nearest-Neighbour Distance (m, MNN). Shannon’s (SHDI) and Simpson’s (SIDI) indices were calculated for the whole landscape surface occupied by brown bears (see Methods).

Land cover classes	East <i>vs.</i> West	Landscape fragmentation indices				
		ED	NP	LPI	MPS	MNN
Forest	E	17.8	701	6.6	260.8	315.6
	W	20.3	1374	2.1	151.0	273.3
Shrubland	E	2.3	258	0.1	50.4	1132.1
	W	2.5	356	0.1	60.6	1184.9
		SHDI	SIDI			
Whole landscape	E	1.94	0.78			
	W	1.77	0.74			

Figure legends

Fig. 1 Distribution of the brown bear locations (western *vs.* eastern subpopulations) in the Cantabrian Mountains (north-western Spain) used in this study.

Fig. 2 Boxplots showing the median, mean (red dots) and range of the values of the habitat variables (see Table 1) that were included in the best models built to characterise the western (1) *vs.* eastern (0) brown bear habitat use in the Cantabrian Mountains at all scales. Habitat variables are reported for each of the four spatial scales considered (circular plots with radii of 500, 1000, 2000 and 3000 m).