FAR-INFRARED EXCITED HYDROXYL LINES FROM ORION KL OUTFLOWS¹

JAVIER R. GOICOECHEA,² JOSÉ CERNICHARO,³ MERCEDES R. LERATE,^{4,5} FABIEN DANIEL,³ MICHAEL J. BARLOW,⁵
BRUCE M. SWINYARD,⁴ TANYA L. LIM,⁴ SERENA VITI,⁵ AND JEREMY YATES⁵

Received 2006 January 6; accepted 2006 March 2; published 2006 March 22

ABSTRACT

As part of the first far-IR line survey toward Orion KL, we present the detection of seven new rotationally excited OH Λ -doublets (at ~ 48 , ~ 65 , ~ 71 , ~ 79 , ~ 98 , and $\sim 115 \mu\text{m}$). Observations were performed with the Long Wavelength Spectrometer Fabry-Pérot on board the *Infrared Space Observatory*. In total, more than 20 resolved OH rotational lines, with upper energy levels up to ~ 620 K, have been detected at angular and velocity resolutions of $\sim 80''$ and $\sim 33 \text{ km s}^{-1}$, respectively. The OH line profiles show a complex behavior evolving from pure absorption, P Cygni type, to pure emission. We also present a large-scale, $6'$ declination raster in the OH $^2\Pi_{3/2}$ $J = 5/2^+ - 3/2^-$ and $^2\Pi_{3/2}$ $J = 7/2^- - 5/2^+$ lines (at 119.441 and $84.597 \mu\text{m}$) revealing a decrease of excitation outside the core of the cloud. From the observed profiles, mean intrinsic line widths, and velocity offsets between emission and absorption line peaks, we conclude that most of the excited OH arises from Orion outflow(s), that is, the “plateau” spectral component. We determine an averaged OH abundance relative to H_2 of $\chi(\text{OH}) = (0.5 - 1.0) \times 10^{-6}$, a kinetic temperature of ≥ 100 K, and a density of $n(\text{H}_2) \approx 5 \times 10^5 \text{ cm}^{-3}$. Even with these conditions, the OH excitation is heavily coupled with the strong dust continuum emission from the inner “hot core” regions and from the expanding flow itself.

Subject headings: infrared: ISM — ISM: individual (Orion KL) — ISM: jets and outflows — ISM: molecules — radiative transfer

1. INTRODUCTION

The Kleinmann-Low (KL) infrared nebula ($L \sim 10^5 L_\odot$) in the Orion molecular cloud is the nearest (~ 450 pc) and probably the most studied massive star-forming region (Genzel & Stutzki 1989). Early studies soon determined that the large-scale distribution of gas and dust is heavily influenced by violent phenomena such as the interaction of compact and large-scale outflows with the quiescent gas, producing strong line and continuum emission. IRc2 was believed to be the main source of luminosity, heating, and dynamics in the region. However, great advances in near- and mid-IR subarcsecond resolution imaging and in (sub-) millimeter interferometric observations have dramatically changed our understanding of the region. First, the $8\text{--}12 \mu\text{m}$ emission peak of IRc2 is not coincident with the Orion SiO maser origin (related to the origin of the outflow or outflows), and second, its intrinsic IR luminosity ($L \sim 1000 L_\odot$) is only a fraction of the complex luminosity (Gezari et al. 1998). In addition, $3.6\text{--}22 \mu\text{m}$ images show that IRc2 is in fact resolved into four components that may even not be self-luminous. Therefore, the relevance of IRc2 as the powerful engine of Orion KL is no longer supported, and its nature is now even less clear (Dougados et al. 1993; Shuping et al. 2004; Greenhill et al. 2004). A new step forward was taken by Menten & Reid (1995) with the detection of the very embedded radio continuum source “I” (located $0''.5$ south of IRc2), which coincides with the centroid of the SiO maser

distribution. Menten & Reid also detected the radio continuum emission of IR source “n” and suggested that it could also contribute to the origin of some of the phenomena observed at larger scales. Thus, in addition to BN, the core of Orion KL contains at least two more compact H II regions (I and n) that seem to be running away from a common point, suggesting that BN, I, and n were originally part of a common massive stellar system that disintegrated ~ 500 years ago (Gómez et al. 2005). A causal relation between the dynamical decay of this stellar system and the observed large-scale molecular outflow(s), H_2 fingers, and bow shocks (Stolovy et al. 1998) has been proposed (Bally & Zinnecker 2005).

The different physical conditions and velocity fields along the line of sight result in a segregation of the gas and dust chemistry that complicates the interpretation of observations toward Orion, especially for low angular resolution molecular surveys. For this reason, it is common to distinguish between different spectral components: the “ridge” (extended quiescent molecular gas), the “hot core” (a collection of very dense and hot gas clumps in which sources I and n are embedded), and the “plateau” (a mixture of outflow[s], shocks, and interactions with the ambient cloud). Submillimeter aperture synthesis line surveys have finally provided the spatial location and extent of many molecular species (Blake et al. 1996; Wright et al. 1996; Liu et al. 2002; Beuther et al. 2005). As a result, the formation of complex oxygen-bearing species in the interaction region between the outflow(s) and the quiescent gas is now generally understood in the context of an oxygen-rich molecular outflow (Blake et al. 1987; Wright et al. 1996; Liu et al. 2002). However, the thermal lines from basic oxygen reservoirs (O, H_2O , and OH) lie in the far-IR, a spectral region blocked by Earth’s atmosphere. Besides, the wide range of excitation conditions provided by oxygen hydrides supply unique information about the energetics and dynamics of the region. In particular, far-IR OH line emission (first detected by Storey et al. 1981) is predicted to be a powerful diagnostic of the plateau gas (Draine & Roberge 1982). Observations with the Kuiper

¹ Based on observations with *ISO*, an ESA project with instruments funded by ESA member states (especially the PI countries: France, Germany, the Netherlands, and the UK) and with the participation of ISAS and NASA.

² Laboratoire d’Etude du Rayonnement et de la Matière en Astrophysique, UMR 8112, CNRS, Ecole Normale Supérieure and Observatoire de Paris, 24 rue Lhomond, F-75231 Paris Cedex 5, France; javier@lra.ens.fr.

³ Departamento de Astrofísica Molecular e Infrarroja, Instituto de Estructura de la Materia, CSIC, Calle Serrano 121, E-28006 Madrid, Spain.

⁴ Rutherford Appleton Laboratory, Chilton, Didcot OX11 0QX, UK.

⁵ Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK.

FIG. 1.—*Left*: Rotational energy diagram of ^{16}OH showing the lines detected by *ISO* (in microns). Newly detected transitions are shown in black, while previous KAO detections are shown in gray. The $2\Pi_{3/2}$ and $2\Pi_{1/2}$ rotational ladders produced by the spin-orbit interaction are shown. The Λ -doubling splitting of each rotational level has been enhanced for clarity. Hyperfine structure is not shown. *Right*: *ISO*-LWS-FP observations of ^{16}OH toward Orion. The vertical axes correspond to the continuum-normalized flux and the horizontal axes to the wavelength in microns. In each panel, the upper level energy (in kelvins) and the intrinsic line strength of the transition are also shown. The upper left panel corresponds to the unresolved OH $2\Pi_{3/2}-2\Pi_{1/2}$ $J = 3/2-5/2$ lines at $\sim 34\ \mu\text{m}$ detected by the *ISO* Short Wavelength Spectrometer (Wright et al. 2000).

Airborne Observatory (KAO) allowed the detection of several low-excitation OH rotational lines, as illustrated in Figure 1 (Watson et al. 1985; Viscuso et al. 1985; Betz & Boreiko 1989; Melnick et al. 1987, 1990). The specific contributions of the hot core, plateau, and ridge components and the physical conditions within the OH-emitting gas remain open to question.

2. OBSERVATIONS AND DATA REDUCTION

Most of the OH pure rotational lines appear in the wavelength range of the *Infrared Space Observatory* (*ISO*) Long Wavelength Spectrometer (LWS; Clegg et al. 1996). The size of the LWS circular aperture is $\sim 80''$. In its Fabry-Pérot (FP) mode, the instrumental response is close to a Lorentzian with a spectral resolution of $\sim 33\ \text{km s}^{-1}$. In this Letter, we present observations that are part of the first full far-IR line survey of Orion KL using the unprecedented *ISO*-LWS-FP wavelength coverage and velocity resolution (Lerate et al. 2006). OH spectra were obtained in the Astronomical Observation Template (AOT) L04 and L03 modes. Processing of the OH spectra from AOT L03 was carried out using the Off-Line Processing (OLP) pipeline and the LWS Interactive Analysis (LIA) package, version 10. AOT L04 spectra were analyzed using the *ISO* Spectral Analysis Package (ISAP).⁶ Typical routines include dark current optimization, deglitching and removal of the LWS grating profile, oversampling and averaging individual scans, baseline removal, and line flux measurements. A full description of the

complex data calibration and reduction process and associated Target Dedicated Time numbers (TDTs) of the survey are given in Lerate et al. (2006). The resulting OH lines are shown in Figure 1. Finally, we present a $\pm 180''$ declination raster (see Fig. 2) in the OH $2\Pi_{3/2}$ $J = 5/2^+-3/2^-$ and $2\Pi_{3/2}$ $J = 7/2^--5/2^+$ lines at 119.441 and $84.597\ \mu\text{m}$ (L04, TDT 70101216).

3. RESULTS AND DISCUSSION

The far-IR OH lines exhibit complicated behavior, evolving from pure absorption to pure emission profiles. The detection of P Cygni type profiles in $2\Pi_{3/2}$ OH excited rotational lines at ~ 65 and $\sim 84\ \mu\text{m}$ (Fig. 1) and in the ^{18}OH ground-state line (Fig. 2; also detected by KAO [Melnick et al. 1990]) confirms that most of the OH emission arises from an extended molecular outflow. Since its optical depth is small, the ^{18}OH emission helps to trace the dominant outflow origin of the OH emission. Compared with stellar wind profiles, the molecular far-IR P Cygni profiles are favored by the presence of far-IR (dust) continuum emission throughout the expanding flow (and not only from a central continuum source). Hence, the detailed balance between emission and absorption components critically depends on the dust and OH emissivities, opacities, and beam filling factors at each wavelength. Simple nonlocal non-LTE radiative transfer models for dust and OH easily reproduce these characteristics (see below). On the other hand, all observed OH lines within the $2\Pi_{1/2}$ ladder (~ 71 , ~ 98 , and $\sim 163\ \mu\text{m}$) appear in emission with a peak velocity of $v \gtrsim +15\ \text{km s}^{-1}$, showing less indication of self-absorption (the cloud rest velocity is $\sim 9\ \text{km s}^{-1}$; Scoville et al. 1983). Taking into account the high energy of the associated levels, effective

⁶ ISAP is a joint development by the LWS and SWS Instruments Teams and Data Centers. Contributing institutes are CESR, IAS, IPAC, MPE, RAL, and SRON.

FIG. 2.—Continuum-subtracted *ISO*-LWS-FP spectra taken at different declination offsets from Orion IRC2. Offsets in arcseconds are indicated. The left ($0'', 0''$) panel shows the different ground-state line profiles produced by ^{16}OH and ^{18}OH (also detected by KAO).

radiative pumping by the far-IR dust continuum emission must be playing a role in the OH excitation. This is clearly favored by the detection of several $2\Pi_{3/2}-2\Pi_{1/2}$ cross-ladder transitions in almost pure absorption (~ 34 , ~ 48 , and $\sim 53 \mu\text{m}$). Hence, temperatures and densities in the outflow can be significantly lower than those required to thermalize the OH emission lines. Other newly detected OH excited cross-ladder transitions (~ 96 and $\sim 115 \mu\text{m}$) are observed in pure emission, and they contribute to the radiative de-excitation of associated $2\Pi_{1/2}$ upper levels. Note that the absorption peak of pure absorption OH lines occurs at negative velocities, $v \lesssim -15 \text{ km s}^{-1}$. Similar conclusions apply for water lines (Cernicharo et al. 1999b). In particular, pure absorption mid-IR H_2O lines peak at $-8 \pm 3 \text{ km s}^{-1}$ (Wright et al. 2000). Taking into account the velocity resolution and wavelength calibration of the LWS/FP instrument, we find $25 \pm 5 \text{ km s}^{-1}$ as the most likely expansion velocity of the far-IR OH flow.

The inferred expansion velocity seems more consistent with the $18 \pm 2 \text{ km s}^{-1}$ *low-velocity outflow*, originally revealed by water maser motions (Genzel et al. 1981). In addition, it is known that OH 1.6 GHz masers toward the Orion KL region are detected over an area of $30''$ in diameter. The best kinematic model fitting the OH maser emission is found for a uniform expansion velocity of $21.0 \pm 3.5 \text{ km s}^{-1}$ away from a source (at or near IRC2) with a radial velocity of $9.0 \pm 0.5 \text{ km s}^{-1}$ (Cohen et al. 2006). Contribution from the extended *high-velocity outflow* (Martín-Pintado et al. 1990) could also be present (see predictions by Melnick et al. 1990). Only the ^{16}OH ground-state lines at $\sim 119 \mu\text{m}$ show high-velocity broad red wing emission above $\sim 100 \text{ km s}^{-1}$. This high-velocity emission is not observed in any other OH line at the *ISO* observations' sensitivity. The large-scale OH declination raster clearly shows that OH lines are weaker (at least by a factor of 5) outside the

FIG. 3.—Selected radiative transfer models for an expanding OH shell. Common parameters are $\chi(\text{OH}) = 5 \times 10^{-7}$ and $n(\text{H}_2) = 5 \times 10^5 \text{ cm}^{-3}$. Three different temperatures are considered: $T_k = 50, 100$, and 150 K . Each panel corresponds to a rotational Λ -doublet. The wavelength of each transition is shown. Line profiles have been convolved with a Gaussian beam of $80''$ and with a Lorentzian of 33 km s^{-1} .

central position (Fig. 2). This is likely due to lower excitation (continuum emission and densities decrease) in the extended cloud, but not necessarily to a steep decrease of OH abundance in the ridge. Unfortunately, it is impossible to infer the exact spatial distribution of the newly detected OH lines from the large *ISO* beam. KAO heterodyne observations of the $\sim 119.2 \mu\text{m}$ line (with a beam of $\sim 33''$) showed that the OH emission may be spatially compact. Betz & Boreiko (1989) deduced that the OH emission comes from a source less than $25''$ in FWHM diameter and, therefore, the same inner regions of the large-scale outflow revealed by H_2O masers at $\sim 325 \text{ GHz}$ (Cernicharo et al. 1999a), and HDO emission at ~ 893 and $\sim 850 \text{ GHz}$ (Pardo et al. 2001). Besides, OH may also arise from the regions where the outflow plunge into the quiescent cloud, producing dissociative shocks. According to recent $[\text{C I}]$ observations (with $10''$ angular resolution), molecular dissociation occurs in a shell of $\sim 40''$ diameter (Pardo et

al. 2005) and, therefore, within the LWS beam. Nevertheless, the OH emission measured by Betz & Boreiko (1989) toward the bright H_2 shocked regions “peak 1” and “peak 2” is significantly weaker than toward the outflow itself. More quantitative conclusions about the spatial distribution of the OH flow require much higher angular resolution observations.

In order to estimate the OH abundance and the physical conditions leading to the observed line profiles, we have modeled the first 20 OH rotational levels with the nonlocal code used in our OH study of Sgr B2 (Goicoechea & Cernicharo 2002). We have modeled a spherical shell with a $25''$ diameter expanding at 25 km s^{-1} that surrounds a uniform core, optically thick in the far-IR, with a $10''$ diameter and a color temperature of 150 K. These values agree with those obtained from high-dipole molecules toward the hot core. In all models, the dust emission reasonably fits the far-IR continuum levels measured by LWS. Obviously, this model simplifies the inhomogeneous and clumpy nature of the region, as *ISO* observations average all the substructure morphology and the different physical conditions found in the plateau. In our view, the far-IR OH excited emission may be spatially associated with the SO and SO_2 shell of low-velocity expanding gas, observed at $\sim 1''$ resolution, where the outflow shocks dense clumps of ambient material (Wright et al. 1996). A grid of models has been generated by varying $T_k = T_{\text{dust}}$ from 50 to 250 K, $n(\text{H}_2)$ from 10^5 to 10^7 cm^{-3} , and $\chi(\text{OH})$ from 10^{-8} to 10^{-5} in the outflow. The newly detected OH excited lines add some important constraints to the plateau physical conditions. Basically, the fact that excitation temperatures in several excited OH cross-ladder transitions have to be below T_{dust} constrains the maximum T_k and $n(\text{H}_2)$ allowed to produce absorption in these transitions. These absorptions can only be reproduced if $n(\text{H}_2) < 5 \times 10^6 \text{ cm}^{-3}$; otherwise, collisional excitation and reemission dominates. For this range of $n(\text{H}_2)$, the bulk of gas has to be above $T_k = 50 \text{ K}$, or else absorption will dominate the lower energy cross-ladder and $^3\Pi_{3/2}$ intraladder transitions. On the other hand, collisional excitation will dominate for $T_k > 200 \text{ K}$. Since conditions of high T_k and $n(\text{H}_2)$ result in a pure OH emission-line

spectrum, a dominant contribution from the hot core is not expected. In addition, the large far-IR line-plus-continuum opacity in the plateau itself will hide most of the hot core OH emission. The lack of spectral resolution makes a more accurate analysis of possible absorptions impossible.

Nevertheless, the detected OH self-absorptions and P Cygni type profiles allow one to constrain the physical parameters that lead to the observed balance between collisions and radiative pumping. The best single-component fit to the OH observations is obtained around $T_k \sim 100 \text{ K}$ and $n(\text{H}_2) \sim 5 \times 10^5 \text{ cm}^{-3}$ (see models in Fig. 3). In comparison with far-IR H_2O ground-state or CO high- J lines, OH cross-ladder transitions have both small spontaneous emission rates and small line strengths. In most applications, this indicates that the associated far-IR lines are optically thin. The lack of opacity broadening also implies that their line profiles are more sensitive to gas velocity fields and turbulence. A good agreement with observations is found for $\chi(\text{OH}) = (0.5\text{--}1.0) \times 10^{-6}$. Therefore, *ISO* observations clearly show that OH is very abundant in the plateau. These results provide additional insights into the fact that complex oxygen-rich molecules such as methanol (CH_3OH) (Wright et al. 1996) or formic acid (HCOOH) (Liu et al. 2002) are specifically enhanced (when resolved at much higher resolution) in the interaction surfaces between the outflow (OH-rich) material and the ambient cloud. Only the wide range of excitations provided by water lines and the input from chemical models, able to predict the chemistry evolution of the physically distinct regions in Orion, can extract more accurate information from far-IR observations.

We are grateful to the LWS instrument team and the *ISO* database team for the quality of the provided data. J. R. G. was supported by a Marie Curie Intra-European Individual Fellowship within the Sixth European Community Framework Programme, contract MEIF-CT-2005-515340. We acknowledge the Spanish DGES and PNIE grants ESP 2001-4516 and AYA 2003-2785.

REFERENCES

- Bally, J., & Zinnecker, H. 2005, *AJ*, 129, 2281
 Betz, A. L., & Boreiko, R. T. 1989, *ApJ*, 346, L101
 Beuther, H., et al. 2005, *ApJ*, 632, 355
 Blake, G. A., Sutton, E. C., Masson, C. R., & Phillips, T. G. 1987, *ApJ*, 315, 621
 Blake, G. A., Mundy, L. G., Carlstrom, J. E., Padin, S., Scott, S. L., Scoville, N. Z., & Woody, D. P. 1996, *ApJ*, 472, L49
 Cernicharo, J., Pardo, J. R., González-Alfonso, E., Serabyn, E., Phillips, T. G., Benford, D. J., & Mehinger, D. 1999a, *ApJ*, 520, L131
 Cernicharo, J., et al. 1999b, in *The Universe as Seen by ISO*, ed. P. Cox & M. F. Kessler (ESA-SP 427) (Noordwijk: ESA), 565
 Clegg, P. E., et al. 1996, *A&A*, 315, L38
 Cohen, R. J., Gasprong, N., Meaburn, J., & Graham, M. F. 2006, *MNRAS*, 367, 541
 Dougados, C., Léna, P., Ridgway, S. T., Christou, J. C., & Probst, R. G. 1993, *ApJ*, 406, 112
 Draine, B. T., & Roberge, W. G. 1982, *ApJ*, 259, L91
 Genzel, R., Reid, M. J., Moran, J. M., & Downes, D. 1981, *ApJ*, 244, 884
 Genzel, R., & Stutzki, J. 1989, *ARA&A*, 27, 41
 Gezari, D. Y., Backman, D. E., & Werner, M. W. 1998, *ApJ*, 509, 283
 Goicoechea, J. R., & Cernicharo, J. 2002, *ApJ*, 576, L77
 Gómez, L., Rodríguez, L. F., Loinard, L., Lizano, S., Poveda, A., & Allen, C. 2005, *ApJ*, 635, 1166
 Greenhill, L. J., Gezari, D. Y., Danchi, W. C., Najita, J., Monnier, J. D., & Tuthill, P. G. 2004, *ApJ*, 605, L57
 Lerate, M. R., et al. 2006, *MNRAS*, submitted
 Martín-Pintado, J., Rodríguez-Franco, A., & Bachiller, R. 1990, *ApJ*, 357, L49
 Melnick, G. J., Genzel, R., & Lugten, J. B. 1987, *ApJ*, 321, 530
 Melnick, G. J., Stacey, G. J., Genzel, R., Lugten, J. B., & Poglitsch, A. 1990, *ApJ*, 348, 161
 Menten, K. M., & Reid, M. J. 1995, *ApJ*, 445, L157
 Liu, S.-Y., Girart, J. M., Remijan, A., & Snyder, L. E. 2002, *ApJ*, 576, 255
 Pardo, J. R., Cernicharo, J., Herpin, F., Kawamura, J., Kooi, J., & Phillips, T. G. 2001, *ApJ*, 562, 799
 Pardo, J. R., Cernicharo, J., & Phillips, T. G. 2005, *ApJ*, 634, L61
 Scoville, N., Kleinmann, S. G., Hall, D. N. B., & Ridgway, S. T. 1983, *ApJ*, 275, 201
 Stolovy, S. R. et al. 1998, *ApJ*, 492, L151
 Storey, J. W. V., Watson, D. M., & Townes, C. H. 1981, *ApJ*, 244, L27
 Shuping, R. Y., Morris, M., & Bally, J. 2004, *AJ*, 128, 363
 Viscuso, P. J., Stacey, G. J., Harwit, M., Haas, M. R., Erickson, E. F., & Duffy, P. B. 1985, *ApJ*, 296, 149
 Watson, D. M., Genzel, R., Townes, C. H., & Storey, J. W. V. 1985, *ApJ*, 298, 316
 Wright, C. M., van Dishoeck, E. F., Black, J. H., Feuchtgruber, H., Cernicharo, J., González-Alfonso, E., & de Graauw, T. 2000, *A&A*, 358, 689
 Wright, M. C. H., Plambeck, R. L., & Wilner, D. J. 1996, *ApJ*, 469, 216