**Age, origins and extinctions of the avifauna of Macaronesia: a synthesis of phylogenetic
and fossil information**

Juan Carlos Illera^{1,2*}, Juan Carlos Rando³, David S. Richardson⁴ and Brent C. Emerson^{2,4}

¹Research Unit of Biodiversity (UO-CSIC-PA), Oviedo University, Campus of Mieres,
Research Building, 5th floor. C/ Gonzalo Gutiérrez Quirós , s/n, 33600 Mieres, Asturias,
Spain.

²Island Ecology and Evolution Research Group, IPNA, CSIC, 38206 La Laguna, Tenerife,
Canary Islands, Spain.

³Department of Animal Biology (Zoology), University of La Laguna, 38206 La Laguna,
Tenerife, Canary Islands, Spain.

⁴Centre for Ecology, Evolution and Conservation, School of Biological Sciences, University
of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

*Correspondence: Juan Carlos Illera. Research Unit of Biodiversity (UO-CSIC-PA), Oviedo
University, Campus of Mieres, Research Building, Fifth floor. C/ Gonzalo Gutiérrez Quirós
s/n, 33600 Mieres, Asturias, Spain. Tel.: + 34 985458132. E-mail: illerajuan@uniovi.es

Abstract

Understanding the age, origins and extinction of oceanic island biota has captivated the interest of evolutionary biologists since Darwin and Wallace. Because oceanic islands are discrete entities of small geographical size but with considerable habitat diversity, they provide ideal templates within which to study evolutionary processes. The peripheral North Atlantic islands of the Mediterranean basin, collectively referred to as Macaronesia, are considered a hot spot of biodiversity due to the fact that they contain a large proportion of endemic taxa (ca. 25%). Recent molecular studies are providing insight into the patterns of colonization and radiation within the extant avifauna, while paleontological studies have described many extinct avian species, sometimes identifying the causes and chronology of extinction. The aim of this review is to develop an understanding of the evolutionary and biogeographic history of the macaronesian avifauna, combining information from phylogenetic and paleontological studies. Phylogenetic analyses have confirmed the close relationships between endemic Macaronesian avifauna and the closest mainland areas (Europe and Africa), however, in contrast to other archipelagos of a similar age, we show that most extant birds appear to have colonized macaronesian archipelagos relatively recently, within the last four million years, despite some islands being approximately 30 million years old. Fossil records support the idea that higher species richness previously existed, with recent dating on bone collagen of selected extinct species suggesting that their extinction coincided with the arrival of aboriginal people ca. 2,500 years ago in the Canary Islands, or the arrival of Europeans across all the Macaronesian islands in the 14th century. It is plausible that these human mediated extinctions may have selectively acted upon older lineages, but there is little evidence available to evaluate this and other alternative hypotheses.

49 **Key words:** Macaronesia, oceanic islands, phylogeography, Quaternary fossil record,
50 diversification, speciation, extinction.
51

1. Introduction

Investigating the origin and diversification of oceanic island biotas has been a key focus of biogeography, evolutionary biology and conservation biology (MacArthur and Wilson, 1967; Filardi and Moyle, 2005; Emerson and Kolm 2005; Savolainen et al., 2006; Steadman, 2006; Grant and Grant, 2008; Ricklefs, 2010a). During the last two decades, the use of molecular markers has provided new insights into the evolutionary histories of a great many taxa, and the resulting conclusions have often challenged biogeographic and evolutionary ideas based on classical taxonomy (e.g. Raxworthy et al., 2002; Jordal et al., 2004; Glor et al., 2005; Macias-Hernández et al., 2008). Phylogeographic syntheses may provide a framework for a general understanding of colonization and diversification within the biotas of archipelagos with different physical and ecological conditions such as age, latitude, altitude and distance to mainland (Ricklefs, 2010a, 2010b). The reconstruction of ancestral distributions and hypotheses about the diversification of taxa have typically used data from extant species, as comprehensive paleontological information on a regional scale is often unavailable. However such analyses may result in biased interpretations if now extinct species are a non-random subset of species, and/or are informative with regard to complex ecological interactions and diversification within a community. Multidisciplinary approaches incorporating information from geology, archaeology, palaeontology and genetics could minimize these limitations and potential biases (Crisci et al., 2003).

Recently, taxa of the North Atlantic Macaronesian islands have become a focus for molecular studies of colonization and diversification of oceanic biotas. Many of these studies have focused on groups of organisms with high rates of diversification such as plants, invertebrates and reptiles (Juan et al., 2000; Emerson, 2002; Kim et al., 2008). Macaronesian birds exhibit a significant number of endemic species and subspecies (Starsfield et al., 1998;

Martín and Lorenzo, 2001), but have only more recently become the subject of phylogeographic study. These studies have revealed diverse evolutionary histories within species, or species complexes, including strong genetic differentiation between islands (Pestano et al., 2000; Dietzen et al., 2003; Kvist et al., 2005; Päckert et al., 2006), incipient differentiation and contemporary gene flow (Hille et al., 2003; Illera et al., 2007; Barrientos et al., 2008; Spurgin et al., 2011) and reverse colonization (Illera et al., 2011). However, the extent to which the overall macaronesian avifauna is the result of ancient or recent colonization and diversification events has not yet been addressed. In light of recent phylogenetic analyses of Macaronesian terrestrial taxa such as insects and plants, we may expect to find both old and young evolutionary lineages (e.g. Emerson, 2008; Aigo et al., 2009). Even though the avian radiation in the region is less spectacular than in other archipelagos such as those of the Pacific and Indian Oceans that does not necessarily mean that extant lineages are themselves young. In addition to molecular phylogenetic and phylogeographic data, fossil data may provide complementary insight into the evolutionary history and community composition of the avifauna of a region. Recent radiocarbon dating on bone collagen of extinct taxa is generating more rigorous approximations of species extinction dates and thus allowing us to better infer the causes of such extinctions. Fossil data can also contribute to a broader understanding of historical faunas and community composition, providing indirect inferences about past environments and ecological interactions.

The aim of this review is to develop a general understanding of the evolutionary and biogeographic history of the Macaronesian avifauna by synthesising information from phylogenetic, phylogeographic and paleontological studies that yield information on colonization, diversification and extinction within this group. We then compare patterns for

Macaronesia with those of other oceanic archipelagos to evaluate to what extent patterns may be generalised across regions, but also to identify knowledge gaps that should be the focus of future research.

2. Macaronesia

The Macaronesian region consists at 31 principal volcanic islands in the north eastern Atlantic Ocean with floral and faunal affinities to the African and European mainland (Juan et al., 2000; Gillespie and Clague, 2009). The islands are grouped into five archipelagos (Azores, Madeira, Selvagens, Canary Islands and Cape Verde) situated between 39°N 31°W and 15°N 23°W (Fig. 1). The broad range of distances between archipelagos and the mainland (<100 km for the Canary Islands and 1,365 km for Azores), latitude, variation of geological ages (0.25-26 million years) and altitudes (130-3,700 m, above sea level) on each island (Geldmacher et al., 2001; Azevedo and Ferreira, 2006; Hazevoet, 1995) result in a broad array of ecological conditions. The region is characterised by high habitat diversity with xerophytic shrublands, pine and laurel forests, lagoons and alpine habitats (Gillespie and Clague, 2009). This habitat diversity is considered to have contributed to the evolution of a rich endemic biota, with 23% of terrestrial taxa being endemic to the region (Izquierdo et al., 2004; Arechavaleta et al., 2005; Borges et al., 2008, 2010).

The degree of geographic isolation of each archipelago and the differing levels of vagility would seem to explain the geographical distribution of vertebrates across Macaronesia. Native bats and birds are distributed throughout the region. However, the colonization history of relatively poor dispersers, such as reptiles and non-volant mammals, appears to be determined by the isolation of each archipelago or island. For example, native

reptiles are absent from Azores, the most isolated archipelago, while non-volant mammals have colonized only the eastern and central Canary Islands, closest to mainland Africa.

3. Species origin: radiation *versus* multiple colonization

3.1. Geographic origins

The taxonomic affinities of native bird species suggest that the origin of the extant terrestrial Macaronesian avifauna is largely from the closest Palaearctic mainland areas, although the Cape Verde archipelago, due to its much lower latitude, contains species related to the semi-arid African mainland region (Hazevoet, 1995; Martín and Lorenzo, 2001; Clarke et al., 2006). Recent phylogenetic analyses have confirmed the close relationships between endemic Macaronesian avifauna and Palaearctic taxa. Although these affinities are similar for other terrestrial vertebrate groups such as reptiles and mammals (Carranza et al., 2002; 2008; Juste et al., 2004; Dubey et al., 2008; Cox, 2010), they contrast with for example, one Azorean native damselfly (*Ischnura hastata*), known only from North America and Caribbean Islands (Belle and van Tol, 1990; Cordero-Rivera et al., 2005), and some elements of the Macaronesian flora that exhibits some rare affinities with East Africa, India, South America and Australia (see Juan et al., 2000 for a review). That the avifauna is largely derived from the Palearctic is surprising given that every year the macaronesian archipelagos receive many North American, Asian and Sub-Saharan African migrants (Martín and Lorenzo, 2001; Clarke et al., 2006).

While it is clear that much of the macaronesian terrestrial avifauna has a Palaearctic origin, deducing the specific origins of species within the Palaearctic region is not easy. Such deductions are complicated by the often broad Palaearctic distributions that many of the species related to macaronesian taxa have, and the complex colonisation pathways, involving

more than one colonization event, reverse colonization, speciation and extinction, that may occur (Emerson, 2002).

3.2. Evolutionary history: isolation and gene flow

During the last two decades a plethora of molecular studies including Macaronesian avifauna has been published. Such studies have proven very useful for characterising the evolutionary history and, in some cases, deducing arrival dates, of Macaronesian taxa (e.g. Helbig et al., 1996; Marshall and Baker, 1998; Arnaiz-Villena et al., 1999; Voelker, 1999; Groombridge et al., 2002; Gómez-Díaz et al., 2006; Illera et al., 2008; Olsson et al., 2010). Phylogeographic approaches have tackled regional questions such as geographic origins, colonisation pathways, times of colonization, gene flow between islands, and diversification in Macaronesia (e.g. Marshall and Baker, 1999; Idaghdour et al., 2004; Päckert et al., 2006; Illera et al., 2007, 2011). Intriguingly, results have, at times, been counter-intuitive, with cases involving complex colonisation histories. For example populations of both *Regulus* spp. and *Erithacus rubecula* appear to represent multiple independent colonization events from the continent, rather than a single origin with subsequent colonisation of neighbouring islands (Päckert et al., 2006; Dietzen et al., 2003). There is little in the way of commonality across different bird species, with heterogeneous results including strong genetic differentiation among islands and archipelagos, but also species exhibiting more recent colonization events and/or high rates of contemporary gene flow. The role of Macaronesian palaeo-islands as stepping-stones and repositories of extant biotas has been evaluated recently (Fernández-Palacios et al., 2011), however, in contrast to taxa such as the planktonic gastropod *Tectarius striatus* (van den Broeck et al., 2008) and the endemic terrestrial moss *Hedenasiastrium percurrans* (Aigoïn et al., 2009) there is no avian lineage older than emerged islands.

The radiation of a single ancestral species into multiple descendant species is a complex process influenced by factors such as ecology, geology, genetics, and historical contingency (Ricklefs and Bermingham, 2007; Blount et al., 2008; Gavrillets and Losos, 2009). Despite their relative isolation and range of ages and habitats, the Macaronesia archipelagos harbour relatively few endemic birds (17% of species are endemic, and there is no endemic genus) (Stattersfield et al., 1998; Clarke et al., 2006). This contrasts with the higher levels of endemism observed in other oceanic archipelagos, such as the Lesser Antilles (43% and 27%, endemic species and genus respectively), islands of the Gulf of Guinea (36% and 9%), Galápagos (90% and 50%) and Hawaii (100% and 87%) (Table 1). The extant Macaronesian avifauna shows both a lower number of endemic species and lower radiation within lineages. This is despite the fact that some genera inhabiting Macaronesia, such as *Anthus*, *Saxicola* and *Turdus*, have diversified extensively outside of the region (Voelker, 1999; Illera et al., 2008; Nylander et al., 2008). A plausible explanation is that macaronesian species are the products of geologically recent colonisation events, or high rates of gene flow after colonization from continental or nearby islands have precluded diversification in allopatry (Ricklefs and Bermingham, 2007). However, although it could be the case for some endemic taxa such as *Anthus berthelotii* and *Serinus canarius*, which despite inhabiting three archipelagos did not experience radiation through (Dietzen et al. 2006; Illera et al., 2007), it does not occur in others as *Fringilla coelebs*, *Fringilla teydea* and *Cyanistes teneriffae*, which radiated among and within archipelagos (Marshall and Baker, 1999; Pestano et al., 2000; Illera et al., 2011). Alternatively, the rate of extinction of species or lineages across Macaronesia may have been higher than that of other oceanic archipelagos. If so, it would be predicted that there would be more extinct genera and species in the fossil record of Macaronesia than that seen in other archipelagos (see below).

4. Arrival dates

Island age limits the maximum time that a taxon can have inhabited an area. The range of geological ages of the Macaronesian islands is considerable (0.25-26 my) and so therefore is the range of possible colonization dates for the avifauna. However, recent studies estimating genetic divergence between island and continental lineages suggest a mainly a Pleistocene (0.01-2.6 my) colonization of Macaronesia (Voelker, 1999; Idaghdour et al., 2004; Päckert et al., 2006; Illera et al., 2008; Töpfer et al., 2011), but with some Pliocene (3-4 my) (Dietzen *et al.*, 2006; Illera et al., 2011) and Holocene (< 0.01 my; Pérez-Tris et al., 2004; Dietzen et al., 2008) elements (Figure 2; S-Table 1). These results contrast with the early to mid Miocene colonization dates (17-20 my) estimated for lizards (Cox *et al.*, 2010), and skinks (Carranza *et al.*, 2008), geckos (Carranza *et al.*, 2002) and invertebrates (e.g. Emerson and Oromí, 2005; Contreras-Díaz et al., 2007) respectively. A notable exception are the two endemic pigeons in the Canary Islands: González et al. (2009) estimated that the laurel pigeon (*C. junoniae*) colonised during early Miocene (20.8 ± 2.2 my), while Bolle's pigeon (*Columba bollii*) colonized during the late Miocene or early Pliocene (5.2 ± 1 my). However, lineage extinction and incomplete sampling of extant taxa are important when estimating colonization times from phylogenetic relationships (Emerson et al., 2000). Consequently, the colonization time estimates of González et al. (2009) may be overestimated, as they did not include all extant Macaronesian and neighbouring continental species.

The estimated recent arrival of birds to Macaronesia compared to reptiles (about 15 million years later) is surprising for such a vagile group. It is also perplexing given that the radiation of most bird lineages occurred during the Palaeogene in Gondwana (Cracraft, 2001; Brown et al., 2008), and that other, now submerged Macaronesian landmasses were emergent some 60 mya (Fernández-Palacios et al., 2011) providing a wide interval of time for the

colonisation of the region. The typically young age of the macaronesian avifauna raises the general question of whether this is representative of colonization times within the avifaunas of other archipelagos. Time of colonization is constrained by island age, thus to evaluate this question comparisons must be performed with volcanic islands or archipelagos of similar age. For instance, colonization of the Galápagos archipelago is necessarily constrained to four or five million years because that is the estimated age of the oldest islands (Cox, 1983). A candidate group of islands for such a comparison is the Lesser Antilles, where the oldest parts show similar ages (≈ 20 my) to the oldest Macaronesian islands and the nearest mainland (South America) is a similar distance away (150 km, Ricklefs and Bermingham, 2007). The last two decades have seen a phylogenetic focus on the avifauna of the Lesser Antilles (Ricklefs and Bermingham, 2002; Cadena et al., 2005; Ricklefs, 2010b) with studies showing some similarities, but also significant differences, compared to the patterns seen in the Macaronesian avifauna. In the Lesser Antilles 38 avian lineages have been analysed, with the most ancient of these being the Lesser Antillean tanager *Tangara cucullata* with a divergence time from its continental relative estimated to be between 9-12 my. Twenty lineages are estimated to have colonised the Lesser Antilles between 0.75-10 my, and the remaining 17 lineages are estimated to have colonized more recently during the mid-Pleistocene (0.55-0.75 my) (Ricklefs and Bermingham, 1999, 2002). The similarity between Macaronesian islands and the Lesser Antilles lies in the accumulation of colonists during the Pleistocene epoch. This may have been facilitated by climate oscillations and the changing mosaic of continental species distributions associated with them (Hewitt, 2000). The main difference is the absence of avian lineages older than 3.8 my in Macaronesia. This absence could reflect a mass extinction event in the region, but there is no evidence for such a catastrophe in other animal or plant groups, so the explanation remains unknown (see below).

5. Species extinction: patterns and processes

Recent paleontological studies in Madeira and the Canary Islands are now changing perceptions about the diversification, distribution, and composition of terrestrial avian communities on these islands (S-Table 2). Upper Pleistocene-Holocene fossil records reveal endemic species extinctions among bird families such as Procellariidae (shearwaters), Phasianidae (quails), Rallidae (rails), Strigidae (owls), Turdidae (thrushes) and Fringillidae (finches and buntings) (Pieper, 1985; Alcover and Florit, 1987; McMinn et al., 1990; Walker et al., 1990; Jaume et al., 1993; Rando et al., 1999, 2008; 2012a; Rando and Alcover, 2010). Recent ^{14}C accelerator mass spectrometry dating of bone collagen of certain species has made it possible to establish an approximate date for their extinction, shedding light on the possible causes of their extinctions. Radiocarbon dates show a correlation between the last occurrence for these extinct species and the colonisation of the regions they inhabited by humans, suggesting a negative causal effect of human arrival (Rando and Alcover, 2008; 2010). Similar conclusions have been reached concerning extinct endemic mammals and native plant species in the Canaries (Criado and Atoche, 2003; Rando et al., 2008; 2012b; de Nascimento et al., 2009). Hunting pressure, habitat loss, introduction of exotic mammals, parasites and diseases introduced as a result of the arrival of humans are thought to have caused biodiversity loss on various oceanic archipelagos (Brooks et al., 2002; Blackburn et al., 2004; Steadman, 2006; Wyatt et al., 2008). In the Canaries, these effects appear to have caused the extinction of some endemic and native species and resulted in the contraction of the once wider distributions of others (Rando, 1995a, 1995b, 2002, 2007).

5.1. The ghost of ancient ecological interactions

267 In addition to documenting and quantifying the number of extinct species, palaeozoological
268 data can also provide evidence for changes in biogeographic ranges, morphological traits,
269 genetic variability and ecological interactions between extinct and extant species (Lyman,
270 2012). Although competition between extinct and extant species is difficult to study,
271 multidisciplinary approaches can shed light on this phenomenon. To date the only study
272 integrating phylogenetic and fossil information comes from the Macaronesian finches. The
273 family Fringillidae is the classic group used by evolutionary biologists to explain bird
274 adaptive radiation on islands (Lerner et al., 2011), with Darwin's finches in Galápagos and
275 the Hawaiian honeycreepers being regarded as among the best examples (Prat, 2005; Grant
276 and Grant, 2008). In contrast the Macaronesia avifauna reveals much less diversification
277 within this family, with only three endemic species: the blue chaffinch (*Fringilla teydea*) in
278 Tenerife and Gran Canaria (Canary Islands), the Azores bullfinch (*Pyrrhula murina*) in São
279 Miguel (Azores), and the wild canary (*Serinus canarius*) in the Azores, Madeira and Canary
280 Islands. The chaffinch (*Fringilla coelebs*) is, perhaps, a notable exception. Five endemic
281 subspecies have been described in Macaronesia: *F. c. moreletti* in Azores, *F. c. maderensis* in
282 Madeira, and three other subspecies in the Canary Islands, *F. c. palmae* in La Palma, *F. c.*
283 *ombriosa* in El Hierro and *F. c. canariensis* in Tenerife, La Gomera and Gran Canaria (Martín
284 and Lorenzo, 2001). Phylogeographic analyses of mitochondrial and nuclear genes have
285 provided support for monophyly within the Atlantic island common chaffinch populations,
286 suggesting they are the result of a single colonization event from the Iberian Peninsula
287 (Marshall and Baker, 1999; Samarasin-Dissanayake, 2010). Interestingly, the picture of low
288 diversification within Macaronesian finches changes when extinct species are taken into
289 account. These extinct species include the Long-legged Bunting (*Emberiza alcoveri*), a
290 minimum of two *Carduelis* finches, which inhabited the woodlands of the Canary Islands and

another undescribed thick-billed finch species in the Madeira Island (S-Table 2). Furthermore, the coexistence of the Blue and Common Chaffinches on two islands (Tenerife and Gran Canaria), and the variation in beak morphology within extant chaffinches - the common chaffinches on the Azores have deeper and wider beaks than in Madeira and the Canary Islands - has intrigued evolutionary biologists for many years (Grant, 1979; Dennison and Baker, 1991). This issue has recently been addressed by combining the study of: (1) the morphology of the extinct greenfinch *Carduelis* spp., and the extant chaffinches, (2) molecular estimation of the colonization times of the extant finch species, and (3) the use of radiocarbon techniques to estimate the extinction date of the greenfinches (Rando et al., 2010). Results point to an ecological interaction between extinct and extant finches, as radiocarbon dating and colonization times reveal that the three forest finch species occurred in sympatry on Tenerife for approximately 1 million years. The differing bill sizes of the three species is thus most compellingly explained by resource competition between the extinct and extant sympatric species (Rando et al., 2010), suggesting a role for character displacement within the diversification of this group in Macaronesia.

5.2. Turnover in bird assemblages

Paleontological data are a useful complement to phylogenetic data to aid the quantification of speciation and extinction events within a region. Because of the volcanic origin of Macaronesia, there are few ancient paleontological sites, and fossils have been mainly collected in the absence of chronologies. The most detailed information so far comes from sites within the Canary Islands (S-Table 2). The oldest bird fossil recorded in Macaronesia corresponds to eggshells found in calcarenite rocks on Lanzarote, with an age estimate of between five and 12 million years ago (Sauer and Rothe, 1972). Based on egg morphology

and pore structure Sauer and Rothe (1972) suggested affinities with ostriches (genus *Struthio*) and an extinct aepyornithoid (genus *Aepyornis*). Because all extant ratite birds are flightless, and the Canary Islands have never been connected with the mainland, the presence of these eggshells remain intriguing, and the object of different interpretations (García-Talavera, 1990). However, despite the uncertainty of the dating of the eggshell it is clear that it predated the arrival of any currently extant species.

The most complete fossil information comes from Upper Pleistocene-Holocene paleontological and archaeological sites. The data show very different bird assemblages with the highest species richness present before the arrival of humans. Within these sites various species of now extinct birds have been identified. In addition, there have been historic documented extinctions of the endemic Canary Island Oystercatcher (*Haematopus meadewaldoi*) and the Red Kite (*Milvus milvus*) during the last century. Among these extinct taxa seven (58%) were endemic. With only five extant endemic species left this means that the Canary Islands have lost more than half of their endemic richness and approximately 13% of the native avifauna. The emerging picture of extinction of species places the Canary Islands above Galápagos and islands of the Gulf of Guinea, but at a lower level to other oceanic archipelagos, such as Hawaii and the Lesser Antilles (Table 1). In Hawaii approximately 58% of original avifauna went extinct after the arrival of Polynesians ca 1,600 yrs ago. In contrast, despite the fact that at least 14 separate bird populations have gone extinct in Galápagos during the last 150-300 yrs, no overall avian species extinctions have been recorded so far (Boyer, 2008; James, 1995; James and Olson, 1991; Olson and James, 1991; Steadman, 1991). In a similar way, no species in the islands of the Gulf of Guinea have gone extinct and there is no evidence for any bird species within the fossil record (Jones and Tye, 2006). In the Lesser Antilles, around 45% of the original avifauna has gone extinct

(Tyrberg, 2009), although the impact on the small land birds does not appear to have been dramatic.

Given that many of the current Neornithine lineages are estimated to be Late Cretaceous (Cracraft, 2001), and considering the Miocene age of Macaronesian islands there is little doubt that older lineages were, at some time, well distributed throughout the region. Because the extinct Macaronesian taxa have provided evidence that higher avian diversity existed before human arrival, a plausible explanation for the lack of older avian lineages is that these lineages have gone extinct. The taxon cycle theory predicts older lineages to have higher extinction probabilities due to evolutionary and ecological interactions among colonizing and native species (Wilson, 1961; Ricklefs and Bermingham, 2002). The absence of geochronologies at the paleontological sites limits inferences regarding colonization and diversification. Approaches for the sequencing of ancient DNA provide an opportunity address this issue (Bunce et al., 2005; Gilbert et al., 2005). However, the only species where this has been achieved to date is the extinct lava shearwater (*Puffinus olsoni*). Phylogenetic analyses performed on the genus *Puffinus* revealed low genetic differentiation between the lava shearwater and its extant sister species, the Manx shearwater (*P. puffinus*), suggesting incipient differentiation within the last million years, with incomplete lineage sorting between both taxa (Ramírez et al., 2010). This result suggests that extinct taxa are not necessarily older than extant lineages, but further studies are needed to shed light on this topic.

6. Considerations about the extinction of avian lineages and future directions

This review indicates that, despite the relatively old (Miocene: ca 5-23 mya) age of the majority of the Macaronesian islands, the extant avian lineages on these islands appear to have colonized the region relatively recently (less than four mya). Fossil records suggest that

greater species richness existed in the past, and that the arrival of aboriginal people from North Africa ($\approx 2,500$ yrs), and then the later colonisation of the islands by Europeans both caused significant extinctions. Nevertheless, it is striking that the extinction process has extirpated all ancient lineages from the current bird assemblages. Because there is not evidence for catastrophic volcanic events to have produced such result, other hypotheses need to be considered.

Introduction of small mammals to islands is known to produce terrible effects on native birds due to the lack of effective anti-predator behaviour and their reduced fecundity (Steadman, 2006; Boyer, 2008). The presence of mice and rats has been documented to occur with the human arrival in the Canary Islands (Rando et al., 2012b), and the introduction of these species had negative effects on the native birds (Martín and Lorenzo, 2001). This arrival, however, was not the first one. In the Canary Islands there is an endemic extant shrew (*Crocidura canariensis*) and three endemic extinct rodents (*Malpaisomys insularis*, *Canariomys bravoii* and *C. tamarani*) (Rando et al., 2012b). The two first ones are endemic to the eastern islands (Fuerteventura and Lanzarote) and *Canariomys* spp. was endemic to the central islands (Gran Canaria and Tenerife). The colonization of the former is estimated to have been approximately 2.14 mya (Dubey et al., 2008), whereas the arrival of *Malpaisomys insularis* is estimated to have been approximately 6 mya (Pagès et al., 2012), while there is no information regarding the estimated arrival date of *Canariomys*. The arrival time of the first land mammals predates the time of colonization of all extant bird lineages (Figure 2). Therefore, we can speculate that shrews and rodents may have produced a limited negative effect on bird species that can be considered recent colonists, while there may have been more dramatic consequences for the ancient endemic avifauna that had evolved in the absence of these mammals in the Canary Islands. In the remaining archipelagos with no native land

mammals, avian extinctions may have been more recent, driven by rodents introduced as a consequence of human colonisation. An additional and not mutually exclusive potential explanation for the observed pattern of young lineages is the possible negative consequences of novel avian pathogens arriving with more recent colonists. Avian malaria has been implicated in the decline and extinction of many endemic species in Hawaii (van Riper et al., 1986, 2002), and it is known that avian malaria and avian pox are pathogens present in Macaronesia, but apparently do not limit the distributions or population sizes of their avian host (Illera et al., 2008; Carrete et al., 2009; Spurgin et al., 2012). However, it is plausible that recent colonists could have introduced novel pathogen strains to which older established lineages of birds were naïve, which can potentially produce a fast and dramatic effect on their survival (Ricklefs and Bermingham, 2007). If the first arrival of mammals was the major driver of extirpation of older lineages, only continued efforts to obtain more fossils in combination with further geochronologies and dating of bone collagen of each new fossil found could shed light on this hypothesis. If so, we expect to find bird fossils in dated layers older than the estimated colonization times for these mammals, which might suggest a cause-effect relationship between both events. Interestingly, the oldest bird eggshells found in the Canary Islands are contained in rock layers dated within the range of mammal arrival to Lanzarote. In contrast, if the extinction of ancient lineages were due to a more recent process, a multidisciplinary approach would seem to be the best way forward to investigate this issue further. DNA amplification has been achieved in bones more than 50,000 years old (e.g. Shapiro et al., 2004; Willerslev and Cooper, 2005), and most bird extinctions in the Canaries have been dated at less than 15,000 years ago (Table 2). Consequently the successful extraction and analysis of ancient DNA from Canary Island avian fossils is theoretically within reach (de Bruyn et al., 2011). Next generation DNA sequencing platforms are very

promising technologies in ancient DNA research since they reduce the amount of starting DNA template necessary to yield a large number of short endogenous sequences (Knapp and Hofreiter, 2010; Orlando et al., 2011). Results obtained from ancient DNA may provide insights on phylogenetic relationships and understanding on dates of divergence for the most recent common ancestors, which can in turn be used to infer the time of colonization. In combination with morphological data (of both extinct and extant taxa) and palaeological data, this may help to improve our understanding of the adaptive and non-adaptive events that occurred among related species, facilitating inferences about the ecological processes that may have caused them and, ultimately, provide insights about the extinction process.

Acknowledgements

J.C. Illera was funded by a Spanish postdoctoral fellowship (Ramón y Cajal). Mario Díaz provided valuable comments on the manuscript.

References

- Agudo, R., Rico, C., Vilà, C., Hiraldo, F., Donázar, J.A., 2010. The role of humans in the diversification of a threatened island raptor. *BMC. Evol. Biol.* 10, 384.
- Aigoín, D.A., Devos, N., Huttunen, S., Ignatov, M.S., González-Mancebo, J.M., Vanderpoorten, A., 2009. And if Engler was not completely wrong? Evidence for multiple evolutionary origins in the moss flora of Macaronesia. *Evolution* 63, 3248-3257.
- Alcover, J.A., Florit, F., 1987. Una nueva especie de *Carduelis* (Fringillidae) de La Palma. *Vieraea* 17, 75-86.
- Arechavaleta, M., Zurita, N., Marrero, M.C., Martín, J.L., 2005. Lista preliminar de especies silvestres de Cabo Verde (hongos, plantas y animales terrestres). Consejería de Medio Ambiente y Ordenación Territorial. Gobierno de Canarias. 155 pp.
- Arnaiz-Villena, A., Álvarez-Tejado, M.A., Ruiz-del-Valle, V., García-de-la-Torre, C., Varela, P., Recio, M.J., Ferre, S., Martínez-Laso, J., 1999. Rapid Radiation of Canaries (Genus *Serinus*). *Mol. Biol. Evol.* 16, 2-11.
- Azevedo, J.M.M., Ferreira, M.P., 2006. The volcanotectonic evolution of Flores Island, Azores (Portugal). *J. Volc. Geo. Res.* 156, 90-102.
- Barrientos, R., Kvist, L., Barbosa, A., Valera, F., López-Iborra, G.M., Moreno, E., 2008. Colonization patterns and genetic structure of Peripherals populations of the trumpeter finch (*Bucanetes githagineus*) from Northwest Africa, the Canary Islands and the Iberian Peninsula. *J. Biog.* 36, 210-219.
- Belle, J., van Tol, J., 1990. *Anomalagrion hastatum* (Say), an American damselfly indigenous to the Azores (Odonata, Coenagrionidae). *Tijdschr. Entomol.* 133, 143-147.

448 Birdwatch News Team, 2010. Killdeer found nesting in Azores. Birdwatch,
 449 http://www.birdwatch.co.uk/channel/newsitem.asp?c=1&date=__9102
 450 Blackburn, T.M., Cassey, P., Duncan, R.P., Evans, K.L., Gaston, K.L., 2004. Avian
 451 extinction and mammalian introductions on oceanic islands. *Science* 305, 1955-1958.
 452 Blount, Z.D., Borland, C.Z., Lenski, R.E., 2008. Historical contingency and the evolution of a
 453 key innovation in an experimental population of *Escherichia coli*. *Proc. Natl. Acad.*
 454 *Sci. USA* 105, 7899-7906.
 455 Boessneck, J., Kinzelbach, R., 1993. Ein prähistorischer Brutplatz von Seevögeln auf der Insel
 456 Sal (Kapverden). *J. Ornithol.* 134, 245-271.
 457 Borges, P.A.V., Abreu, C., Aguiar, A.M.F., Carvalho, P., Jardim, R., Melo, I., Oliveira, P.,
 458 Sérgio, C., Serrano, A.R.M., Vieira, P., 2008. A list of the terrestrial fungi, flora and
 459 fauna of Madeira and Selvagens archipelagos. Direcção Regional do Ambiente da
 460 Madeira and Universidade dos Açores, Funchal and Angra do Heroísmo, 440 pp.
 461 Borges, P.A.V., Costa, A., Cunha, R., Gabriel, R., Gonçalves, V., Martins, A.F., Melo, I.,
 462 Parente, M., Raposeiro, P., Rodrigues, P., Santos, R.S., Silva, L., Vieira, P. and
 463 Vieira, V. 2010. A list of the terrestrial and marine biota from the Azores. *Princípios*,
 464 Cascais, 432 pp.
 465 Boyer, A.G., 2008. Extinction patterns in the avifauna of the Hawaiian Islands. *Divers.*
 466 *Distrib.* 14, 509-517.
 467 Brooks, T.M., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Rylands, A.B.,
 468 Konstant, W.R., Flick, P., Pilgrim, J., Oldfield, S., Magin, G., Hilton-Taylor, G., 2002.
 469 *Habitat Loss and Extinction in the Hotspots of Biodiversity. Cons. Biol.*, 16: 909-923.

470 Brown, J.W., Rest, J.S., García-Moreno, J., Sorenson, M.D., Mindell, D.P., 2008. Strong
 471 mitochondrial DNA support for a Cretaceous origin of modern avian lineages. BMC
 472 Biol. 6, 6.

473 Bunce, M., Szulkin, M., Lerner, H.R.L., Barnes, I., Shapiro, B., Cooper, A., Holaway, R.N.,
 474 2005. Ancient DNA Provides New Insights into the Evolutionary History of New
 475 Zealand's Extinct Giant Eagle. PLoS Biol 3, e9.

476 Cadena, C.D., Ricklefs, R.E., Jiménez, I., Bermingham, E., 2005. Is speciation driven by
 477 species diversity? Nature 434, 1015-1017.

478 Carracedo, J.C., Day, S., 2002. Canary Islands. Classic geology in Europe series. Terra
 479 Publishing. Hertfordshire, 294 pp.

480 Carranza, S., Arnold, E.N., Geniez, Ph., Roca, J., Mateo, J.A., 2008. Radiation, multiple
 481 dispersal and parallelism in the skinks, *Chalcides* and *Sphenops* (Squamata:
 482 Scincidae), with comments on *Scincus* and *Scincopus* and the age of the Sahara
 483 Desert. Mol. Phyl. Evol. 46, 1071-1094.

484 Carranza, S., Arnold, E.N. Mateo, J.A., Geniez, P., 2002. Relationships and evolution of the
 485 North African geckos, *Geckonia* and *Tarentola* (Reptilia: Gekkonidae), based on
 486 mitochondrial and nuclear DNA sequences. Mol. Phyl. Evol. 23, 244-256.

487 Carrete, M., Serrano, D., Illera, J.C., López, G., Vögeli, M., Delgado, A. & Tella, J.L., 2009.
 488 Goats, birds and emergent diseases: Apparent and hidden effects of an exotic species
 489 in an island environment. Ecol. Appl. 19, 840-853.

490 Clarke, T., Orgill, C., Disley, T., 2006. A Field Guide to the Birds of the Atlantic Islands:
 491 Canary Islands, Madeira, Azores, Cape Verde. Christopher Helm. London, 368 pp.

492 Contreras-Díaz, H.G., Moya, O., Oromí, P., Juan, C., 2007. Evolution and diversification of
 493 the forest and hypogean ground-beetle genus *Trechus* in the Canary Islands. Mol.
 494 Phyl. Evol. 42, 687-699.

495 Cordero Rivera A, Lorenzo Carballa, M.O., Utzeri, C., Vieira, V., 2005. Parthenogenetic
 496 *Ischnura hastata* (Say, 1839), widespread in the Azores (Zygoptera: Coenagrionidae).
 497 Odonatologica 34, 1-9.

498 Cox, A., 1983. Ages of the Galápagos Islands. In: Bowman, R.I., Berson, M., Leviton, A.E.
 499 (Eds.), Patterns of evolution in Galápagos organisms. American Association for the
 500 Advancement of Science, Pacific Division, San Francisco, pp. 11-23.

501 Cox, S.C., Carranza, S., Brown, R.P., 2010. Divergence times and colonization of the Canary
 502 Islands by *Gallotia* lizards. Mol. Phyl. Evol. 56, 747-757.

503 Cracraft, J., 2001. Avian evolution, Gondwana biogeography and the Cretaceous-Tertiary
 504 mass extinction event. Proc. R. Soc. Lond. B 268, 459-469.

505 Criado, C., Atoche, P., 2003. Estudio Geoarqueológico del yacimiento de El Bebedero (Siglos
 506 I a.C. a XIV d.C., Lanzarote, Islas Canarias). Rev. Cuater. Geomor. 17 (1-2), 91-104.

507 Crisci, J.V., Katinas, L., Posadas, P. 2003. Historical biogeography: an introduction. Harvard
 508 University Press, Cambridge, 250 pp.

509 De Bruyn, M., Hoelzel, R., Carvalho, G.R., Hofreiter, M., 2011. Faunal histories from
 510 Holocene ancient DNA. TREE 26, 405-413.

511 de Nascimento, L., Willis, K.J., Fernández-Palacios, J.M., Criado, C., Whittaker, R.J., 2009.
 512 The long-term ecology of the forest of La Laguna, Tenerife (Canary Islands). J. Biog.
 513 36, 499-514.

514 Dennison, M.D., Baker, A.J., 1991. Morphometric variability in continental and Atlantic
 515 island populations of Chaffinches (*Fringilla coelebs*). Evolution 45, 29-39.

516 Dietzen, C., Witt, H-H., Wink, M., 2003. The phylogeographic differentiation of the
 517 European robin, *Erithacus rubecula*, on the Canary Islands revealed by mitochondrial
 518 DNA sequence data and morphometrics: evidence for a new robin taxon on Gran
 519 Canaria? Avian Science 2-3, 115-131.

520 Dietzen, C., Garcia-del-Rey, E., Delgado, G., Wink, M., 2008. Phylogenetic differentiation of
 521 *Sylvia* species (Aves: Passeriformes) of the Atlantic islands (Macaronesia) based on
 522 mitochondrial DNA sequence data and morphometrics. Biol. J. Linn. Soc. 95, 157-
 523 174.

524 Dietzen, C., Voigt, C., Wink, M., Gahr, M., Leitner, S., 2006. Phylogeography of island
 525 canary (*Serinus canaria*) populations. J. Ornithol. 147, 485-494.

526 Dubey, S., Salamin, N., Ruedi, M., Barrière, P., Colyn, M., Vogel, P., 2008. Biogeography
 527 origin and radiation of the Old World crocidurine shrews (Mammalia: Soricidae)
 528 inferred from mitochondrial and nuclear genes. Mol. Phyl. Evol. 48, 953-963.

529 Dyhr, C.T., Holm, P.M., 2010. A volcanological and geochemical investigation of Boa Vista,
 530 Cape Verde Islands; ⁴⁰Ar/³⁹Ar geochronology and field constraints. J. Volc. Geo.
 531 Res. 189, 19-32.

532 Emerson, B.C., 2002. Evolution on oceanic islands: molecular phylogenetic approaches to
 533 understanding pattern and process. Mol. Ecol. 11, 951-966.

534 Emerson, B.C., 2003. Genes, geology and biodiversity: faunal and floral diversity on the
 535 island of Gran Canaria. ABC 26, 9-20.

536 Emerson, B.C. 2008. Speciation on islands: what are we learning? Biol. J. Linn. Soc. 95, 47-
 537 52.

538 Emerson, B.C., Kolm, N., 2005. Species diversity can drive speciation. Nature 434, 1015-
 539 1017.

540 Emerson, B.C., Oromí, P., 2005. Diversification of the forest beetle genus *Tarphius* on the
 541 Canary Islands, and the evolutionary origins of island endemics. *Evolution* 59, 586-
 542 598.

543 Emerson, B.C., Oromí, P., Hewitt, G.M., 2000. Colonization and diversification of the species
 544 *Brachyderes rugatus* (Coleoptera) on the Canary Islands: evidence from mitochondrial
 545 DNA COII gene sequences. *Evolution* 54, 911-923.

546 Fernández-Palacios, J.M., de Nascimento, L., Otto, R., Delgado, J.D., García-del-Rey, E.,
 547 Arévalo, J.R., Whittaker, R.J., 2011. A reconstruction of Palaeo-Macaronesia, with
 548 particular reference to the long-term biogeography of the Atlantic island laurel forests.
 549 *J. Biog.* 38, 226-246.

550 Filardi, C.E., Moyle, R.G., 2005. Single origin of a pan-Pacific bird group and upstream
 551 colonization of Australasia. *Nature* 438, 216-219.

552 Friesen, V.L., Smith, A.L., Gómez-Díaz, E., Bolton, M., Furness, R.W. González-Solís, J.,
 553 Monteiro, L.R., 2007. Sympatric speciation by allochrony in a seabird. *Proc. Natl.*
 554 *Acad. Sci. USA* 104, 18589-18594.

555 Garcia-del-Rey, E., Delgado, G., Gonzalez, J., Wink, M., 2007. Canary Island great spotted
 556 woodpecker (*Dendrocopos major*) has distinct mtDNA. *J. Ornithol.* 148, 531-536.

557 García-Talavera, F., 1990. Aves gigantes en el Mioceno de Famara (Lanzarote). *Rev. Acad.*
 558 *Canar. Cienc.* 2, 71-79.

559 Gavrillets, S., Losos, J.B., 2009. Adaptive radiation: contrasting theory with data. *Science* 323,
 560 732-737.

561 Geldmacher, J., Hoernle, K., Bogaard, Pvd., Zankl, G., Garbe-Schönberg, D., 2001. Earlier
 562 history of the ≥ 70 -Ma-old Canary hotspot based on the temporal and geochemical

563 evolution of the Selvagen Archipelago and neighboring seamounts in the eastern
 564 North Atlantic. J. Volc. Geo. Res. 111, 55-87.

565 Gilbert, M.T.P., Bandelt, H-J., Hofreiter, M., Barnes, I., 2005. Assessing ancient DNA
 566 studies. TREE 20, 541-544.

567 Gillespie, R.G., Clague, D.A., 2009. Encyclopedia of islands. University of California press.
 568 California, 1,074 pp.

569 Glor, R.E., Losos, J.B., Larson, A., 2005. Out of Cuba: overwater dispersal and speciation
 570 among lizards in the *Anolis carolinensis* subgroup. Mol. Ecol. 14, 2419-2432.

571 Gómez-Díaz, E., González-Solís, J., Peinado, M.A., Page, R.D.M., 2006. Phylogeography of
 572 the *Calonectris* shearwaters using molecular and morphometric data. Mol. Phy. Evol.
 573 41, 322-332.

574 González, J., Delgado Castro, G., García-del-Rey, E., Berger, C., Wink, M., 2009. Use of
 575 mitochondrial and nuclear genes to infer the origin of two endemic pigeons from the
 576 Canary Islands. J. Ornithol. 150, 357-367.

577 Grant, P.R., 1979. Evolution of the chaffinch, *Fringilla coelebs*, on the Atlantic Islands. Biol.
 578 J. Linn. Soc. 11, 301-332.

579 Grant, P.R., Grant, B.R., 2008. How and Why Species Multiply. The Radiation of Darwin's
 580 Finches. Princeton series in evolutionary biology, 218 pp.

581 Groombridge, J.J., Jones, C.G., Bayes, M.K., van Zyl, A.J., Carrillo, J., Nichols, R.A.,
 582 Bruford, M.W., 2002. A molecular phylogeny of African kestrels with reference to
 583 divergence across the Indian Ocean. Mol. Phyl. Evol. 25, 267-277.

584 Hazevoet, C.J., 1995. The birds of the Cape Verde Islands. B.O.U. Check-list N° 13. Tring,
 585 192 pp.

586 Hewitt, G., 2000. The genetic legacy of the Quaternary ice ages. Nature 405, 907-913.

587 Helbig, A.J., Martens, J., Seibold, I., Henning, F., Schottler, B., Wink, M., 1996. Phylogeny
 588 and species limits in the Palaearctic chiffchaff *Phylloscopus collybita* complex. Ibis
 589 138, 650-666.

590 Hille, S.M., Nesje, M., Segelbacher, G., 2003. Genetic structure of kestrel populations and
 591 colonization of the Cape Verde archipelago. Mol. Ecol. 12, 2145-2151.

592 Idaghdour, Y., Broderick, D., Korrida, A., Chbel, F., 2004. Mitochondrial control region
 593 diversity of the houbara Bustard *Chlamydotis undulata* complex and genetic structure
 594 along the Atlantic seaboard of North Africa. Mol. Ecol. 13, 43-54.

595 Illera, J.C., Emerson, B.C., Richardson, D.S., 2007. Population history of Berthelot's pipit:
 596 colonisation, gene flow and morphological divergence in Macaronesia. Mol. Ecol. 16,
 597 4599-4612.

598 Illera, J.C., Emerson, B.C., Richardson, D.S., 2008. Genetic characterization, distribution and
 599 prevalence of avian pox and avian malaria in the Berthelot's pipit (*Anthus berthelotii*)
 600 in Macaronesia. Parasitol. Res. 103, 1435-1443.

601 Illera, J.C., Koivula, K., Broggi, J., Päckert, M., Martens, J., Kvist, L., 2011. A multi-gene
 602 approach reveals a complex evolutionary history in the *Cyanistes* species group. Mol.
 603 Ecol. 20, 4123-4139.

604 Illera, J.C., Richardson, D.S., Helm, B., Atienza, J.C. Emerson, B.C., 2008. Phylogenetic
 605 relationships, biogeography and speciation in the avian genus *Saxicola*. Mol. Phy.
 606 Evol. 48, 1145-1154.

607 Iwanuit, A.N, Olson, S.L., James, H.F., 2009. Extraordinary cranial specialization in a new
 608 genus of extinct duck (Aves: anseriformes) from Kauai, Hawaiian Islands. Zootaxa
 609 2296, 47-67.

- 610 Izquierdo, I., Martín, J.L., Zurita, N., Arechavaleta, M., 2004. Lista de especies silvestres de
611 Canarias (hongos, plantas y animales terrestres) 2004. Consejería de Medio Ambiente
612 y Ordenación Territorial, Gobierno de Canarias, 500 pp.
- 613 James, H.F., 1995. Prehistoric extinctions and ecological changes on oceanic islands. *Ecol.*
614 *Stud.* 115, 87-102.
- 615 James, H.E., Olson, S.L., 1991. Descriptions of thirty-two new species of birds from the
616 Hawaiian Islands: Part II, Passeriformes. *Ornithol. Monographs* 45, 1-88.
- 617 Jaume, D., McMinn, M., Alcover, J.A., 1993. Fossil bird from the Bujero del Silo, La Gomera
618 (Canary Islands), with a description of a new species of Quail (Galliformes:
619 Phasianidae). *Bol. Mus. Mun. Funchal* 2, 147-165.
- 620 Jesus, J., Menezes, D., Gomes, S., Oliveira, P., Nogales, M., Brehm, A., 2009. Phylogenetic
621 relationships of gadfly petrels *Pterodroma* spp. From the northeastern Atlantic Ocean:
622 molecular evidence for specific status of Bugio and Cape Verde petrels and
623 implications for conservation. *Bird Conserv. Int.* 19, 199-214.
- 624 Jones, P., Tye, A., 2006. The birds of São Tomé and Príncipe with Annobón: islands of the
625 Gulf of Guinea. BOU Checklist Series No. 22. British Ornithologists' Union, Oxford,
626 172 pp.
- 627 Jordal, B.H., Kirkendall, L.R., Harkestad, K., 2004. Phylogeny of a Macaronesian radiation:
628 host-plant use and possible cryptic speciation in *Liparthrum* bark beetles. *Mol. Phyl.*
629 *Evol.* 31, 554-571.
- 630 Juan, C., Emerson, B.C., Oromi, P., Hewitt, G.M., 2000. Colonization and diversification:
631 towards a phylogeographic synthesis for the Canary Islands. *TREE* 15, 104-109.

632 Juste, J., Ibáñez, C., Muñoz, J., Trujillo, D., Benda, P., Karatas, A., Ruedi, M., 2004.
633 Mitochondrial phylogeography of the long-eared bats (*Plecotus*) in the Mediterranean
634 Palaeartic and Atlantic Islands. Mol. Phy. Evol. 31, 1114-1126.

635 Kim, S-C., McGowen, M.R., Lubinsky, P., Barber, J.C., Mort, M.E., Santos-Guerra, A., 2008.
636 Timing and Tempo of Early and Successive Adaptive Radiations in Macaronesia.
637 PLoS ONE 3(5): e2139.

638 Klassert, T.E., Hernández, M.A., Campos, F., Infante, O., Almeida, T., Suárez, N.M. Pestano,
639 J., Hernández, M., 2008. Mitochondrial DNA points to *Lanius meridionalis* as a
640 polyphyletic species. Mol. Phy. Evol. 47, 1227-1231.

641 Knapp, M., Hofreiter, M., 2010. Next generation sequencing of ancient DNA: requirements,
642 strategies and perspectives. Genes 1, 227-243.

643 Kvist , L., Broggi, J., Illera, J.C., Koivula, K., 2005. Colonisation and diversification of the
644 blue tits (*Parus caeruleus teneriffae*-group) in the Canary Islands. Mol. Phy. Evol. 34,
645 501-511.

646 Lee, D-C., Halliday, A.N., Fitton, J.G., Poli, G., 1994. Isotopic variations with distance and
647 time in the volcanic islands of the Cameroon line: evidence for a mantle plume origin.
648 Earth Planet. Sci. Lett. 123, 119-138.

649 Lerner, H.R.L., Meyer, M., James, H.F., Hofreiter, M., Fleischer, R.C., 2011. Multilocus
650 resolution of phylogeny and timescale in the extant adaptive radiation of Hawaiian
651 honeycreepers. Curr. Biol. 21, 1838-1844.

652 Lyman, R.L., 2012. A warrant for applied palaeozoology. Biol. Rev. doi: 10.1111/j.1469-
653 185X.2011.00207.x

654 MacArthur, R.H., Wilson, E.O., 1967. The theory of island biogeography. Princeton Univ.
655 Press, Princeton, 203 pp.

656 Macias-Hernandez, N., Oromi, P., Arnedo, M.A., 2008. Patterns of diversification on old
 657 volcanic islands as revealed by the woodlouse-hunter spider genus *Dysdera* (Araneae,
 658 Dysderidae) in the eastern Canary Islands. Biol. J. Linn. Soc. 94, 589-615.
 659 Marshall, H.D., Baker, A.J., 1998. Rates and Patterns of Mitochondrial DNA Sequence
 660 Evolution in Fringilline Finches (*Fringilla* spp.) and the Greenfinch (*Carduelis*
 661 *chloris*). Mol. Biol. Evol. 15, 638-646.
 662 Marshall, H.D., Baker, A.J., 1999. Colonization history of Atlantic island common
 663 chaffinches (*Fringilla coelebs*) revealed by mitochondrial DNA. Mol. Phyl. Evol. 11,
 664 201-212.
 665 Martín, A., 2011. Lista de los vertebrados terrestres de Canarias. Available at:
 666 <http://webpages.ull.es/users/aumartin/>
 667 Martín, A., Lorenzo, J.A., 2001. Aves del Archipiélago Canario. Francisco Lemus Editor. La
 668 Laguna, 787 pp.
 669 McMinn, M., Jaume, D., Alcover, J.A., 1990. *Puffinus olsoni* n. sp.: nova espècie de baldritja
 670 recentment extinguida provinent de depòsits espeleològics de Fuerteventura i
 671 Lanzarote (Illes Canàries, Atlàntic Oriental). Endins 16, 63-71.
 672 Moyle, R.G., Filardi, C.E., Smith, C.E., Diamond, J. (2009). Explosive Pleistocene
 673 diversification and hemispheric expansion of a “great speciator”. Proc. Natl. Acad.
 674 Sci. USA 106, 1863-1868.
 675 Nylander, J.A., Olsson, U., Alström, P., Sanmartín, I., 2008. Accounting for phylogenetic
 676 uncertainty in biogeography: a Bayesian approach to dispersal-vicariance analysis of
 677 the Thrushes (Aves: *Turdus*). Syst. Biol. 57, 257-268.
 678 Olson, S.L., den Hartog, J.C., 1990. Former breeding of *Sula dactylatra* in the Cape Verde
 679 Islands. Bull. B.O.C. 110, 10-12.

680 Olson, S.L., James, H.E., 1991. Descriptions of thirty-two new species of birds from the
 681 Hawaiian Islands: Part I, Non-Passeriformes. Ornithol. Monographs 45, 1-88.
 682 Olsson, U., Alström, P., Svensson, L., Aliabadian, M., Sundberg, P., 2010. The *Lanius*
 683 *excubitor* (Aves, Passeriformes) conundrum - taxonomic dilemma when molecular
 684 and non-molecular data tell different stories. Mol. Phyl. Evol. 55, 347-357.
 685 Orlando, L., Ginolhac, A., Raghavan, M., Vilstrup, J., Rasmussen, M., et al., 2011. True
 686 single-molecule DNA sequencing of a Pleistocene horse bone. Genome Res. 21, 1705-
 687 1719.
 688 Päckert, M., Dietzen, C., Martens, J., Wink, M., Kvist, L., 2006. Radiation of Atlantic
 689 goldcrests *Regulus regulus* spp.: evidence of a new taxon from the Canary Islands. J.
 690 Avian Biol. 37, 364-380.
 691 Pagès, M., Chevret, P., Gros-Balthazard, M., Hughes, S., Alcover, J.A., Hutterer, R., Rando
 692 J.C., Michaux, J., Hänni, C., 2012. Paleogenetic Analyses Reveal Unsuspected
 693 Phylogenetic Affinities between Mice and the Extinct *Malpaisomys insularis*, an
 694 Endemic Rodent of the Canaries. PLoSOne
 695 Pérez-Tris, J., Bensch, S., Carbonell, R., Helbig, A.J., Telleria, J.L., 2004. Historical
 696 diversification of migration patterns in a passerine bird. Evolution 58, 1819-1832.
 697 Pestano J., Brown, R.P., Rodríguez, F., Moreno, A., 2000. Mitochondrial DNA control region
 698 diversity in the endangered blue chaffinch, *Fringilla teydea*. Mol. Ecol. 9, 1421-1425.
 699 Pieper, H., 1985. The fossil land birds of Madeira and Porto Santo. Bocagiana 88, 1-6.
 700 Prat, H.D., 2005. The Hawaiian Honeycreepers. Oxford University Press, 342 pp.
 701 Ramírez, O., Illera, J.C., Rando, J.C., González-Solis, J., Alcover, J.A., Lalueza-Fox, C.,
 702 2010. Ancient DNA of the extinct lava shearwater (*Puffinus olsoni*) from the Canary

703 Islands reveals incipient differentiation within the *P. puffinus* complex. PLoS ONE
704 5(12), e16072.

705 Rando, J.C., 1995a. Presencia de restos de Pigargo (*Haliaeetus* sp.) (Aves: Accipitridae) en
706 yacimientos paleontológicos de Fuerteventura. Vieraeta 24, 65-69.

707 Rando, J.C., 1995b. Restos de hubara, *Chlamydotis undulata* (Jacquin, 1784) (Aves:
708 Otidae), en la Cueva del Viento (Tenerife, Islas Canarias). Vieraeta 24, 192.

709 Rando, J.C., 2002. New data of fossil birds from El Hierro (Canary Islands): Probable causes
710 of extinction and some biogeographical considerations. Ardeola 49, 39-49.

711 Rando, J.C., 2007. New fossil record of choughs genus *Pyrrhocorax* in the Canary Islands:
712 hypotheses to explain its extinction and current narrow distribution. Ardeola 54, 185-
713 195.

714 Rando, J.C., Alcover, J.A., 2008. Evidence for a second western Palaearctic seabird extinction
715 during the last Millennium: the Lava Shearwater *Puffinus olsoni*. Ibis 150, 188-192.

716 Rando, J.C., Alcover, J.A., 2010. On the extinction of dune shearwater (*Puffinus holeae*) from
717 the Canary Islands. J. Ornithol. 151, 365-369.

718 Rando, J.C., Alcover, J.A., Illera, J.C., 2010. Disentangling ancient interactions: a new extinct
719 passerine provides insights on character displacement among extinct and extant island
720 finches. PLoS ONE 5(9): e12956.

721 Rando, J.C., Alcover, J.A., Michaux, J., Hutterer, R., Navarro, J.F., 2012b. Late Holocene
722 asynchronous extinction of endemic mammals on the eastern Canary Islands. The
723 Holocene, doi:10.1177/0959683611430414.

724 Rando, J.C. Alcover, J.A., Navarro, J.F., García-Talavera, F., Hutterer, R., Michaux, J., 2008.
725 Chronology and causes of the extinction of the Lava Mouse, *Malpaisomys insularis*
726 (Rodentia: Muridae) from the Canary Islands. Quat. Res. 70, 141-148.

727 Rando, J.C., López, M., Seguí, B., 1999. A new species of extinct flightless passerine
728 (emberizidae: *Emberiza*) from the Canary Islands. *Condor* 101, 1-13.

729 Rando, J.C., Pieper, H., Alcover, J.A., Olson, S., 2012a. A new species of extinct fossil scops
730 owl (Aves: Strigiformes: Otus) from the Archipelago of Madeira (North Atlantic
731 Ocean). *Zootaxa*

732 Raxworthy, C.J., Forstner, M.R.J., Nussbaum, R.A., 2002. Chameleon radiation by oceanic
733 dispersal. *Nature* 415, 784-787.

734 Ricklefs, R.E., 2010a. Dynamics of colonization and extinction on islands. Insights from
735 Lesser Antillean birds. In: Losos, J.B., Ricklefs, R.E. (Eds.): *The theory of island
736 biogeography revisited*, Princeton University Press, Princeton, pp. 388-414.

737 Ricklefs, R.E., 2010b. Colonization of the Lesser Antilles by land birds. *Ecology* 91, 1811-
738 1821.

739 Ricklefs, R.E., Bermingham, E., 1999. Taxon cycles in the Lesser Antillean avifauna. *Ostrich*
740 70, 49-59.

741 Ricklefs, R.E., Bermingham, E., 2002. The concept of the taxon cycle in biogeography.
742 *Global Ecol. Biogeogr.* 11, 353-361.

743 Ricklefs, R.E., Bermingham, E., 2007. The causes of evolutionary radiations in archipelagoes:
744 passerine birds in the Lesser Antilles. *Amer. Natur.* 169, 285-297.

745 Samarasin-Dissanayake, P., 2010. Population differentiation, historical demography and
746 evolutionary relationships among widespread common chaffinch populations
747 (*Fringilla coelebs* ssp.). Ph.D. dissertation. University of Toronto.

748 Sauer, E.G.F., Rothe, P., 1972. Ratite Eggshells from Lanzarote, Canary Islands. *Science* 176,
749 43-45.

750 Savolainen, V., Anstett, M.C., Lexer, C., Hutton, I., Clarkson, J.J., Norup, M.V., Powell,
 751 M.P., Springate, D., Salamin, N., Baker, W.J., 2006. Sympatric speciation in palms on
 752 an oceanic island. *Nature* 441, 210-213.

753 Shapiro, B., Drummond, A.J., Rambaut, A., Wilson, M.C. et al. (2004). Rise and fall of the
 754 Beringian steppe bison. *Science* 306, 1561-1565.

755 Spurgin, L.G., Illera, J.C., Padilla, D.P., Richardson, D.S., 2012. Biogeographical patterns
 756 and co-occurrence of pathogenic infection across island populations of Berthelot's
 757 pipit (*Anthus berthelotii*). *Oecologia*, DOI 10.1007/s00442-011-2149-z.

758 Spurgin, L., Van Oosterhout, C., Illera, J.C., Bridgett, S., Gharbi, K., Emerson, B.C.,
 759 Richardson, D., 2011. Gene conversion rapidly generates MHC diversity in recently
 760 founded bird populations. *Mol. Ecol.* 20, 5213-5225.

761 Stattersfield, A.J., Crosby, M.J., Long, A.J., Wege, D.C., 1998. Endemic Bird Areas of the
 762 World. BirdLife Conservation, Series 7. BirdLife International, Cambridge, 860 pp.

763 Steadman, D.W., 1991. Chronology of Holocene Vertebrate Extinction in the Galápagos
 764 Islands. *Quat. Res.* 36, 126-133.

765 Steadman, D.W., 2006. Extinction and biogeography of tropical Pacific birds. London:
 766 University of Chicago Press, 594 pp.

767 Töpfer, T., Haring, E., Birkhead, T.R., Lopes, R.J., Severinghaus, L., Martens, J., Päckert, M.,
 768 2011. A molecular phylogeny of bullfinches *Pyrrula* Brisson, 1760 (Aves:
 769 Fringillidae). *Mol. Phy. Evol.* 58, 271-282.

770 Tyrberg, T., 2009. Holocene avian extinctions. In: Turvey, S.T. (Ed.), *Holocene extinctions*,
 771 Oxford University Press, pp. 63-106.

772 van den Broeck, H., Breugelmans, K., de Wolf, H., Backeljau, T., 2008. Completely disjunct
 773 mitochondrial DNA haplotype distribution without a phylogeographic break in a
 774 planktonic developing gastropod. *Marine Biology* 153, 421-429.
 775 van Riper, C.III., van Riper, S.G., Goff, M.L., Laird, M. 1986. The epizootiology and
 776 ecological significance of malaria in Hawaiian land birds. *Ecol Monogr.* 56, 327-344
 777 van Riper, C.III., van Riper, S.G., Hansen, W.R., 2002. Epizootiology and effect of avian pox
 778 on Hawaiian forest birds. *Auk* 119, 929-942.
 779 Voelker, G., 1999. Dispersal, vicariance, and clocks: historical biogeography and speciation
 780 in a cosmopolitan passerine genus (*Anthus*: Motacillidae). *Evolution* 53, 1536-1552.
 781 Walker, C.A., Wragg, G.M., Harrison, C.J.O., 1990. A new shearwater from the Pleistocene
 782 of the Canary Islands and its bearing on the evolution of certain *Puffinus* shearwaters.
 783 *Hist. Biol.* 3, 203-224.
 784 Willerslev, E., Cooper, A., 2005. Ancient DNA. *Proc. R. Soc. B.* 272, 3-16.
 785 Wilson, E.O., 1961. The nature of the taxon cycle in the Melanesian ant fauna. *Amer. Natur.*
 786 95, 169-193.
 787 Wyatt, K.B., Campos, P.F., Gilbert, M.T.P., Kolokotronis, S-O., Hynes, W.H., et al., 2008.
 788 Historical Mammal Extinction on Christmas Island (Indian Ocean) Correlates with
 789 Introduced Infectious Disease. *PLoS ONE* 3, e3602.
 790
 791
 792
 793

Table 1. Characteristics of the five Macaronesian and five other oceanic archipelagos. Number of genera and species (extant native and endemic, extinct native and endemic) per archipelago (and overall in Macaronesia) is shown, including only upper Pleistocene-Holocene and historical extinct birds. Age is the maximum oldest radiometric ages in millions of years (youngest-oldest islands). A: Africa; E: Europe; SA: South America; AU: Australia; NA: North America. G. Guinea: islands of the Gulf of Guinea (except Bioko). Geological data was obtained from Lee et al. (1994), Geldmacher et al. (2001). Carracedo and Day (2002), Azevedo et al. (2006), Ricklefs and Bermingham (2007) and Dyhr and Holm (2010). Physical features obtained from Stattersfield et al. (1998), Ricklefs and Bermingham (2007), Borges et al. (2008), Gillespie and Clague (2009). Bird information obtained from Pieper (1985), Olson and den Hartog (1990), Boessneck and Kinzelbach (1993), Martín and Lorenzo (2001); Arechavaleta et al. (2005); Jones and Tye (2006); Steadman (2006); Ricklefs and Bermingham (2007); Borges et al. (2008, 2010), Boyer (2008), Martín (2011), Tyrberg (2009), and Iwanuik et al. (2009). *: With human settlements.

Characteristics	Azores	Madeira	Selvagens	Canary Islands	Cape Verde	Macaronesia	Lesser Antilles	Vanuatu
N° Islands	9	2*	2	8*	10	31	21	6
Area (km²)	2,387	785	2.73	7,447	4,020	14,586	6,300	12,000
Age (my)	0.25-8.12	0.7-14	24.2-29.5	1.2-22	1-22	0.25-29.5	20	2
Maximum altitude (m)	2,381	1,810	0.153	3,718	2,829	3,718	1,500	1,500
Minimum dist. (km) continent	1,365 (E)	633 (A)	373 (A)	96 (A)	568 (A)	96 (A)	150 (SA)	1,150 (AU)
N° of extant genera (endemic)	29 (0)	30 (0)	8 (0)	60 (0)	33 (0)	75 (0)	30 (8)	18 (0)
N° of extinct genera (endemic)	0	7 (0)	0	10 (0)	3 (0)	13 (0)	9 (0)	4 (0)
N° of extant species (endemic)	36 (2)	37 (4)	9 (0)	77 (5)	37 (6)	109 (19)	37 (16)	24 (0)
N° of extinct species (endemic)	0	10 (8)	0	12 (7)	3 (0)	21 (14)	15 (13)	5 (0)

809 **Table 2.** Radiocarbon age (years before present) and 2 σ calibration intervals (cal. yrs. AD) from endemic extinct bird bones of the Canary
810 Islands. The 2 σ calibration intervals were obtained using the program OxCal. V4.0, the marine 04.14C calibration curve and $\Delta R = 275 \pm 67$, for
811 marine (*) bird bones, and intcal04.14C calibration curve, for the land species (see Rando and Alcover, 2008, 2010; Rando et al., 2010, for
812 further details). The lower value of each interval must be considered a minimum age estimate of extinction. AD: *Annus Domini*. BP: Before
813 Present. BC: Before Christ.

Laboratory number	Species	Radiocarbon age (yrs. BP)	2 σ calibration intervals
KIA-29159	<i>Puffinus olsoni</i> *	1,265 $\pm$ 25	1,270-1,475 cal. yrs. AD
KIA-36249	<i>Puffinus holeae</i> *	3,395 $\pm$ 30	1,159-790 cal. yrs. BC
KIA-30992	<i>Carduelis aurelioi</i>	11,460 $\pm$ 60	11,478-11,258 cal. yrs. BC

816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831

832 **S-Table 1.** Summary of divergence times and molecular markers and methods or software used to for their estimation colonization dates from the
833 most recent common ancestor in Macaronesian avifauna. Cyt-b: Cytochrome b. CR: Control region. ND2: NADH dehydrogenase 2. ND5:
834 NADH dehydrogenase 5. 16S: ribosomal RNA genes 16S. fib-7: intron 7 of b-fibrinogen. GAPDH-11: intron 11 of glyceraldehyde 3-phosphate
835 dehydrogenase.
836

Taxa / n° populations	Marker	Divergence times	Method	References
<i>Calonectris diomedea</i>	Cyt-b / CR	900,000-700,000 y	0.9% divergence per my	Gómez-Díaz et al. 2006
<i>Puffinus olsoni</i>	Cyt-b	900,000-400,000 y	0.9% divergence per my	Ramírez et al., 2010
<i>Puffinus puffinus</i>	Cyt-b	900,000-400,000 y	0.9% divergence per my	Ramírez et al., 2010
<i>Pterodroma madeira</i>	Cyt-b	2.5 my	0.78% divergence per my	Jesus et al., 2009
<i>Pterodroma deserta</i>	Cyt-b	1.75 my	0.78% divergence per my	Jesus et al., 2009
<i>Pterodroma feae</i>	Cyt-b	1.75 my	0.78% divergence per my	Jesus et al., 2009
<i>Oceanodroma castro</i>	CR	110,000-180,000 y	MDIV: 21% divergence per my	Friesen et al. 2007
<i>Neophron percnopterus</i>	Microsatellites	2,500 y	MSVAR 0.4 and 1.3	Agudo et al., 2010
<i>Falco tinnunculus</i>	Cyt-b	0.1-0.4 my	0.7-2% divergence per my	Groombridge et al. 2002
<i>Chlamydotis undulata</i>	CR	20-25,000 y	Using 2 substitution rates	Idaghdour et al. 2004
<i>Dendrocopus major</i>	Cyt-b / ND2	150,000 y	2% divergence per my	García del Rey et al. 2007
<i>Anthus berthelotii</i>	Cyt-b	2.5 my	2% divergence per my	Voelker 1999
<i>Erithacus rubecula</i>	Cyt-b	2.3 my	2% divergence per my	Dietzen et al., 2003
<i>Saxicola dacotiae</i>	Cyt-b	1.6 my	BEAST: 0.5-3.5% divergence per my	Illera et al. 2008
<i>Sylvia atricapilla</i>	CR	4,000-13,000 y	Using 3 substitution rates	Pérez-Tris et al. 2004
<i>Sylvia atricapilla</i>	Cyt-b	4,000-40,000 y	Using 3 substitution rates	Dietzen et al. 2008
<i>Sylvia melanocephala</i>	Cyt-b	6,500-65,000 y	Using 3 substitution rates	Dietzen et al. 2008
<i>Sylvia conspicillata</i>	Cyt-b	306,000 y	BEAST: 2% divergence per my	Illera et al. unpublished data
<i>Phylloscopus canariensis</i>	Cyt-b	1.0 my	BEAST: 2% divergence per my	Illera et al. unpublished data
<i>Cyanistes teneriffae</i>	CR / Cyt-b / fib-7	3.1 my	BEAST: using age constraints in two nodes	Illera et al. 2011
<i>Lanius meridionalis</i>	Cyt-b	0.8-0.9 my	1.9% divergence per my	Klassert et al. 2008

<i>Fringilla teydea</i>	Cyt-b	1.99 my	BEAST: 0.5-3.5% divergence per my	Rando et al. 2010
<i>Fringilla coelebs</i>	Cyt-b	1.3 my	BEAST: 0.5-3.5% divergence per my	Rando et al. 2010
<i>Serinus canarius</i>	Cyt-b	3.8 my	0.4% divergence per my	Arnaiz-Villena et al. 1999; Dietzen et al. 2006
<i>Regulus regulus</i>	ND5 / Cyt-b / CR	2.2-0.7 my	r8s: 1.68-3.6% divergence per my	Päckert et al. 2006
<i>Pyrrula murina</i>	Cyt-b / 16S / rRNA / fib-7 / GAPDH-11	1.5 my	BEAST: using age constraints in four nodes	Töpfer et al., 2011

837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859

860

861 **S-Table 2.** Native breeding bird species (extant and extinct) in Macaronesia. Endemic:

862 endemic to Macaronesia. † = extinct species. Only Upper Pleistocene-Holocene and

863 historical extinct species are included. Information obtained from Pieper (1985), Olson

864 and den Hartog (1990), Boessneck and Kinzelbach (1993), Arechavaleta et al. (2005),

865 Borges et al. (2005, 2008), Martín and Lorenzo (2001), Birdwatch News Team (2010),

866 Borges et al. (2010), Martín (2011), Rando et al. (2012a).

867

	Azores	Madeira	Salvagens	Canaries*	C. Verde	Endemic
Procellariiformes						
<i>Bulweria bulwerii</i>	✓	✓	✓	✓	✓	
<i>Calonectris diomedea</i>	✓	✓	✓	✓	-	
<i>Calonectris edwardsii</i>	-	-	-	-	✓	✓
<i>Puffinus holeae</i>	-	-	-	†	-	✓
<i>Puffinus olsoni</i>	-	-	-	†	-	✓
<i>Puffinus puffinus</i>	✓	✓	✓	✓	-	
<i>Puffinus baroli</i>	✓	✓	✓	✓	✓	
<i>Pterodroma sp.</i>	-	-	-	†	-	
<i>Pterodroma feae</i>	-	-	-	-	✓	✓
<i>Pterodroma deserta</i>	-	✓	-	-	-	✓
<i>Pterodroma madeira</i>	-	✓	-	-	-	✓
<i>Pelagodroma marina</i>	-	✓	-	✓	✓	
<i>Hydrobates pelagicus</i>	-	-	-	✓	-	
<i>Oceanodroma castro</i>	✓	✓	✓	✓	✓	
<i>Oceanodroma monteiroi</i>	✓	-	-	-	-	✓
Pelecaniformes						
<i>Fregata magnificens</i>	-	-	-	-	✓	
<i>Phaethon aethereus</i>	✓	-	-	✓	✓	
<i>Sula leucogaster</i>	-	-	-	-	✓	
<i>Sula dactylatra</i>	-	-	-	-	†	
Ciconiiformes						
<i>Ixobrychus minutus</i>	-	-	-	✓	-	
<i>Nycticorax nycticorax</i>	-	-	-	✓	✓	
<i>Bubulcus ibis</i>	-	-	-	✓	-	
<i>Egretta garzetta</i>	-	-	-	✓	✓	
<i>Ardea cinérea</i>	-	-	-	-	✓	
<i>Ardea purpurea</i>	-	-	-	-	✓	
Anseriformes						
<i>Tadorna ferruginea</i>	-	-	-	✓	-	
<i>Anas rubripes</i>	✓					

<i>Anas platyrhynchos</i>	✓	-	-	-	-	
<i>Marmaronetta angustirostris</i>	-	-	-	✓	-	
Accipitriformes						
<i>Milvus migrans</i>	-	-	-	✓	✓	
<i>Milvus milvus</i>	-	-	-	†	✓	
<i>Haliaeetus sp.</i>	-	-	-	†	-	
<i>Neophron percnopterus</i>	-	-	-	✓	✓	
<i>Accipiter gentilis</i>	-	-	-	†	-	
<i>Accipiter nisus</i>	-	✓	-	✓	-	
<i>Buteo buteo</i>	✓	✓	-	✓	✓	
<i>Pandion haliaetus</i>	-	-	-	✓	✓	
Falconiformes						
<i>Falco tinnunculus</i>	-	✓	-	✓	-	
<i>Falco eleonora</i>	-	-	-	✓	-	
<i>Falco peregrinus</i>	-	-	-	-	✓	
<i>Falco pelegrinoides</i>	-	-	-	✓	-	
Galliformes						
<i>Coturnix coturnix</i>	✓	✓	✓	✓	✓	
<i>Coturnix gomerae</i>	-	-	-	†	-	✓
<i>Coturnix n. spp.1</i>	-	†	-	-	-	✓
<i>Coturnix n. spp.2</i>	-	†	-	-	-	✓
Gruiformes						
<i>Rallidae n. spp. 1</i>	-	†	-	-	-	✓
<i>Rallidae n. spp. 2</i>	-	†	-	-	-	✓
<i>Rallidae n. spp. 3</i>	-	†	-	-	-	✓
<i>Gallinula chloropus</i>	✓	✓	-	✓	†	
<i>Fulica atra</i>	✓	-	-	✓	†	
<i>Chlamydotis undulada</i>	-	-	-	✓	-	
Charadriiformes						
<i>Haematopus meadewaldoi</i>	-	-	-	†	-	✓
<i>Himantopus himantopus</i>	-	-	-	✓	✓	
<i>Burhinus oedicephalus</i>	-	-	-	✓	-	
<i>Cursorius cursor</i>	-	-	-	✓	✓	
<i>Charadrius dubius</i>	-	-	-	✓	-	
<i>Charadrius alexandrinus</i>	✓	✓	-	✓	✓	
<i>Charadrius vociferus</i>	✓	-	-	-	-	
<i>Gallinago gallinago</i>	✓	-	-	-	-	
<i>Scolopax rusticola</i>	✓	✓	-	✓	-	
<i>Pinguinus impennis</i>	-	†	-	-	-	
Lariformes						
<i>Larus michahellis</i>	✓	✓	✓	✓	-	

<i>Larus fuscus</i>	-	-	-	✓	-	
<i>Sterna dougallii</i>	✓	-	-	✓	-	
<i>Sterna hirundo</i>	✓	✓	✓	✓	-	
<i>Sterna [Onychoprion] fuscata</i>	✓	-	-	-	-	
<i>Sterna [Onychoprion] anaethetus</i>	✓					
Pteroclidiformes						
<i>Pterocles orientalis</i>	-	-	-	✓	-	
Columbiformes						
<i>Columba livia</i>	✓	✓	-	✓	✓	
<i>Columba palumbus</i>	✓	-	-	-	-	
<i>Columba bollii</i>	-	-	-	✓	-	✓
<i>Columba junoniae</i>	-	-	-	✓	-	✓
<i>Columba trocaz</i>	-	✓	-	-	-	✓
<i>Streptopelia turtur</i>	-	-	-	✓	-	
<i>Streptopelia decaocto</i>	✓	-	-	✓	-	
Strigiformes						
<i>Tyto alba</i>	-	✓	-	✓	✓	
<i>Asio otus</i>	✓	-	-	✓	-	
<i>Otusn. spp.</i>	-	†	-	-	-	✓
Apodiformes						
<i>Apus unicolor</i>	-	✓	-	✓	-	✓
<i>Apus apus</i>	-	-	-	✓	-	
<i>Apus pallidus</i>	-	-	-	✓	-	
<i>Apus alexandri</i>	-	-	-	-	✓	✓
Coraciiformes						
<i>Halcyon leucocephala</i>	-	-	-	-	✓	
<i>Upupa epops</i>	-	✓	-	✓	-	
Piciformes						
<i>Dendrocopos major</i>	-	-	-	✓	-	
Passeriformes						
<i>Alaemon alaudipes</i>	-	-	-	-	✓	
<i>Alauda razae</i>	-	-	-	-	✓	✓
<i>Ammomanes cincturus</i>	-	-	-	-	✓	
<i>Eremopterix nigriceps</i>	-	-	-	-	✓	
<i>Calandrella rufescens</i>	-	-	-	✓	-	
<i>Hirundo rustica</i>	-	-	-	✓	-	
<i>Anthus berthelotii</i>	-	✓	✓	✓	-	✓
<i>Motacilla cinerea</i>	✓	✓	-	✓	-	
<i>Erithacus rubecula</i>	✓	✓	-	✓	-	
<i>Oenanthe oenanthe</i>	✓	-	-	-	-	-

<i>Saxicola dacotiae</i>	-	-	-	✓	-	✓
<i>Turdus merula</i>	✓	✓	-	✓	-	
<i>Turdus n. spp.</i>	-	†	-	-	-	✓
<i>Acrocephalus brevipennis</i>	-	-	-	-	✓	✓
<i>Sylvia conspicillata</i>	-	✓	-	✓	✓	
<i>Sylvia melanocephala</i>	-	-	-	✓	-	
<i>Sylvia atricapilla</i>	✓	✓	-	✓	✓	
<i>Phylloscopus canariensis</i>	-	-	-	✓	-	✓
<i>Regulus regulus</i>	✓	-	-	✓	-	
<i>Regulus madeirensis</i>	-	✓	-	-	-	✓
<i>Cyanistes teneriffae</i>	-	-	-	✓	-	
<i>Lanius meridionalis</i>	-	-	-	✓	-	
<i>Pyrrhonorax graculus</i>	-	-	-	†	-	
<i>Pyrrhonorax pyrrhonorax</i>	-	-	-	✓	-	
<i>Corvus corax</i>	-	-	-	✓	-	
<i>Corvus ruficollis</i>	-	-	-	-	✓	
<i>Sturnus vulgaris</i>	✓	-	-	✓	-	
<i>Passer hispaniolensis</i>	-	✓	-	✓	✓	
<i>Passer iagoensis</i>	-	-	-	-	✓	✓
<i>Petronia petronia</i>	-	✓	-	✓	-	
<i>Fringilla coelebs</i>	✓	✓	-	✓	-	
<i>Fringilla teydea</i>	-	-	-	✓	-	✓
<i>Serinus serinus</i>	-	-	-	✓	-	
<i>Serinus canarius</i>	✓	✓	-	✓	-	✓
<i>Carduelis triasi</i>	-	-	-	†	-	✓
<i>Carduelis aurelioi</i>	-	-	-	†	-	✓
<i>Carduelis chloris</i>	-	✓	-	✓	-	
<i>Carduelis carduelis</i>	-	✓	-	✓	-	
<i>Carduelis cannabina</i>	-	✓	-	✓	-	
<i>Carduelis spinus</i>	-	✓	-	-	-	
<i>Fringillidae n. spp.</i>	-	†	-	-	-	✓
<i>Bucanetes githagineus</i>	-	-	-	✓	-	
<i>Pyrrhula murina</i>	✓	-	-	-	-	✓
<i>Coccothraustes coccothraustes</i>	-	†	-	-	-	
<i>Emberiza alcoveri</i>	-	-	-	†	-	✓
<i>Miliaria calandra</i>	-	-	-	✓	-	

868

869

870 * Material stored in scientific collections was just considered.

871

872

873

874

Figure 1. Map of Macaronesia with archipelagos in boxes. The maximum geological age estimated per archipelago is presented below each box.

Figure 2. Accumulation of species through time (millions of years) in Macaronesia. Species were grouped into six age categories: 3-4 my; 2- <3 my; 1.5- <2.0 my; 1.0- <1.5 my; 0.5- <1.0 my; 0- <0.5 my.

Figure 2