

1 Application of mass spectrometry-based metabolomics approaches for food  
2 safety, quality and traceability.

3 María Castro-Puyana<sup>1</sup>, Raquel Pérez-Míguez<sup>1</sup>, Lidia Montero<sup>2</sup>, Miguel Herrero<sup>2\*</sup>

4

5

6 <sup>1</sup>Departamento de Química Analítica, Química Física e Ingeniería Química, Universidad  
7 de Alcalá, Ctra. Madrid-Barcelona, Km. 33.600, 28871 Alcalá de Henares, Madrid, Spain

8 <sup>2</sup>Laboratory of Foodomics, Institute of Food Science Research (CIAL-CSIC), Nicolás  
9 Cabrera 9, Campus UAM Cantoblanco, 28049 Madrid, Spain

10

11

12 Corresponding Author: M. Herrero

13 e-mail: m.herrero@csic.es

14 TEL: +34 910 017 946

15 FAX: +34 910 017 905

16

17 **ABSTRACT**

18 The always more-demanding fields of food safety, quality and traceability are  
19 continuously fostering the development of robust, efficient, sensitive and cost-effective  
20 analytical methodologies. Mass spectrometry-based metabolomics is a key tool nowadays  
21 with great potential in many analytical fields and has been demonstrated to be capable of  
22 facing some important challenges related to these areas within the food science domain.

23 The main aim of this review is to present a critical overview of the most recent  
24 applications of MS-based metabolomics approaches for food quality, safety and  
25 traceability assessment, covering the most relevant works published from 2014 to 2017.

26 Information about the different steps needed to develop a MS-metabolomics approach,  
27 i.e. sample treatment, analytical platform, and data processing, is also provided and  
28 discussed.

29

30 **Keywords:** chemometrics, contaminants, food analysis, food quality, food safety, food  
31 traceability, foodomics, GC-MS, hyphenated techniques, LC-MS, mass spectrometry,  
32 metabolomics.

33 **Abbreviations:** CE, capillary electrophoresis; CID, collision-induced dissociation;  
34 DBDI, dielectric barrier discharge ionization; EVOO, extra virgin olive oil; GC, gas  
35 chromatography; GC × GC, comprehensive two-dimensional GC; HILIC, hydrophilic  
36 interaction chromatography; HRMS, high resolution mass spectrometry; HS-SPME,  
37 headspace solid-phase micro-extraction; ICP, inductively coupled plasma; IT, ion trap;  
38 LAESI, laser ablation electrospray ionization; LC, liquid chromatography; LC × LC,  
39 comprehensive two-dimensional liquid chromatography; LLE, Liquid-liquid extraction;  
40 MRL, maximum residue limit; MRM, multiple reaction monitoring; MS, mass  
41 spectrometry; MVOCs, microbial volatile organic compounds; NMR, nuclear magnetic  
42 resonance; PCA, principal components analysis; PLE, pressurized liquid extraction; PTR,  
43 proton transfer reaction; Q, quadrupole; QqQ, triple quadrupole; QTOF, quadrupole-  
44 time-of-flight; SLE, Solid-liquid extraction; SPE, solid-phase extraction; SPME, solid-  
45 phase microextraction; SRM, selected reaction monitoring; TOF, time-of-flight; UAE-  
46 DLLME, ultrasound-assisted extraction in tandem with dispersive liquid-liquid  
47 microextraction; UHPLC, ultra-high pressure liquid chromatography.

48

49 **1. INTRODUCTION.**

50 Metabolomics is one of the main branches in the field of the -omics techniques, and  
51 together with genomics, transcriptomics and proteomics, is involved in the study of the  
52 food and nutrition domains through Foodomics approaches. As per definition,  
53 metabolomics includes the exhaustive study of the whole small metabolite composition  
54 of a particular system or organism, understanding by small metabolite typically those with  
55 a molecular weight below 1500 Da. In practice, this aim is difficult to achieve, due to the  
56 huge chemical variability of metabolites that is often found; this implies that a universal  
57 approach to analyze using a single method metabolites belonging to very different  
58 chemical classes (significantly different polarity) as well as present in a very wide  
59 dynamic range is not attainable. In this regard, the food metabolome is not an exception  
60 as quite diverse compounds, such as carbohydrates, lipids, proteins, amino acids, amines,  
61 steroids, phenolic compounds, carotenoids, alkaloids or volatile compounds, among  
62 others are frequently present. For this reason, the selection of more than one analytical  
63 approach, and their combination for results interpretation is often carried out.

64 The analytical procedures usually employed within metabolomics can be grouped in  
65 different categories. On the one hand, methods can be classified under fingerprinting  
66 approaches or under profiling methodologies. Fingerprinting is referred to the analysis of  
67 as many compounds as possible within a system, including their detection and the  
68 subsequent statistical treatment of the obtained results in order to look for sample patterns.  
69 Under this approach, the identification and quantification of the detected metabolites may  
70 not be a necessity. In opposition, profiling refers mainly to the analysis of closely related  
71 metabolites, often belonging to the same chemical class, which are most frequently  
72 identified and quantified. Similarly, metabolomics approaches can be also classified as  
73 non-targeted or targeted analysis; whereas non-targeted approaches look for maximum

74 coverage of metabolites that can be simultaneously identified in a particular system,  
75 targeted approaches are based on the determination and identification of a certain type of  
76 metabolites, that could either belong to the same chemical class or being involved in a  
77 particular pathway. In any case, as the complexity of the set of metabolites to be analyzed  
78 is quite high in both approaches, suitable analytical techniques are needed, as well as  
79 proper sample treatment methodologies. This latter subject is of great relevance in food  
80 analysis, as food are usually quite complex matrices full of potentially-disturbing  
81 components for the analysis of metabolites. Sample treatment may be relatively simple  
82 or involve multiple steps. However, it has always to be considered that sample treatment  
83 may include unintended bias towards the metabolites present, as a universal sample  
84 treatment directed to the extraction of the full metabolome of a particular sample will not  
85 exist in practice, and thus, some components may be lost during this phase.

86 Concerning the analytical tools employed, most attention has been paid to the detection  
87 technique. However, it is evident that a proper separation before detection can increase  
88 the quality of the obtained results. Although gas chromatography (GC) was perhaps the  
89 separation technique of choice in the initial metabolomics studies, the need for  
90 derivatization in order to increase the coverage of compounds that can be analyzed  
91 following this approach has driven to shift the primary technique to liquid  
92 chromatography (LC). In fact, LC can be operated in several separation modes, which  
93 increases its versatility towards the separation of a variety of different metabolites.  
94 Particularly, in the last years, methods based on the use of ultra-high performance liquid  
95 chromatography (UHPLC) have gained considerable popularity thanks to the advantages  
96 that this technique can provide with, including high efficiency, good resolution, relatively  
97 short analysis times and the use of flow rates fully compatible with mass spectrometry  
98 (MS) detection.

99 Likewise, concerning the detection of the metabolites, nuclear magnetic resonance  
100 (NMR) was the most-used technique in the first years of metabolomics development.  
101 However, MS has gradually substituted the use of NMR. Some of the reasons behind this  
102 move include that MS is by far more suitable for coupling with a separation technique, as  
103 well as the development and improved affordability of high resolution MS instruments.  
104 In this regard, the use of high resolution instruments, like time-of-flight (TOF) analyzers,  
105 or even hybrid instruments such as quadrupole-TOF (QTOF) or orbitrap, allows to obtain  
106 accurate mass determination, which is the key for their use in metabolomics approaches,  
107 as well as to resolve isomeric and isobaric species. Moreover, the possibility of running  
108 MS/MS experiments with some of these instruments, significantly enhances the  
109 capabilities for the identification of unknown metabolites.

110 As a direct consequence of the improvement on the available analytical tools, samples  
111 with higher complexity can be analyzed in which even thousands of features may be  
112 detected. Thus, the datasets generated after sample analyses in a typical metabolomics  
113 study is of extremely great complexity, including retention times, intensities,  $m/z$ , and  
114 even MS/MS spectra. Under these conditions, the manual interpretation and elaboration  
115 of all these data is impossible. For this reason, normalized procedures have been  
116 developed relying on bioinformatics tools in order to be able to properly extract the key  
117 information of all the huge amount of data available. Usually, data-processing involves  
118 peak detection, integration, peak alignment and normalization. After these steps, different  
119 chemometric tools can be used to statistically assess possible differences among samples.

120 To do that, multivariate analysis is often used, although the particular statistical approach  
121 to be used will largely depend on the objectives of the study. Principal components  
122 analysis (PCA) is frequently employed at first, as it allows to group samples as a function  
123 of different variables. However, the particular statistical analyses made are usually

124 different depending also on the topic of the study, i.e., food-health relationships,  
125 biomarker discovery, food quality, food safety or traceability, among others.

126 The aim of this review is to update the information provided in our previous article [1],  
127 including a critical revision of the latest research published in the field of MS-based  
128 metabolomics applied to food quality, food safety and traceability from 2014 to 2017. For  
129 the sake of clarity, each of these three topics are described and discussed in separate  
130 sections so that the basic particularities of the approaches involved in those subjects can  
131 be appropriately described.

132

## 133 **2. MS-BASED METABOLOMICS FOR FOOD SAFETY**

134 Food safety is one of the most-important topics within food analysis; although one may  
135 tend to consider that every sold and consumed foodstuff possess proper safety, the truth  
136 is that food control is constantly required to maintain an appropriate degree of security  
137 for consumers. Food safety involves many sub-fields, including the legislation  
138 enforcement regarding the presence of selected compounds in foods that may be present  
139 below certain limits (MRL, maximum residue limits), the detection of microbial-related  
140 spoilage, the determination of allergens, the detection of environmental contaminants as  
141 well as banned external compounds, or the assessment of the occurrence of natural toxins,  
142 for example. In this regard, the use of MS within metabolomics-based approaches has  
143 allowed significantly raising the level of the analytical determinations possible nowadays.  
144 In this section, the most-relevant published procedures to this aim are described and  
145 commented.

146

147 **2.1. Detection of chemical contaminants: food production-related controlled**  
148 **substances (veterinary drug and pesticide residues), environmental pollutants and**  
149 **food-contact materials**

150 Although there is a wealth of published material developing always better analytical  
151 methods for the detection of selected contaminants in foods, this section is focused to  
152 those methods that take advantage of metabolomics-based approaches to carry out those  
153 determinations, thus, targeting the detection of multiple components in just one run.

154 The first part of any MS-based metabolomics study for the detection of food contaminants  
155 is sample preparation. As foods may be considered as very complex matrices involving  
156 the presence of a broad array of very different components, suitable sample preparation  
157 steps are needed in order to allow a proper detection of contaminants which will surely  
158 be present in very low amounts. Some of the naturally present compounds in foods will  
159 negatively influence the analysis of the targeted compounds, and thus, different methods  
160 have been widely used to extract and/or concentrate those. Solid-liquid extraction (SLE)  
161 or liquid-liquid extraction (LLE), depending on the physical nature of the samples, using  
162 conventional solvents and solid-phase extraction (SPE) are, probably, the three sample  
163 preparation methods traditionally most-employed. However, following the latest trends  
164 regarding the application of “Green Chemistry” principles, other miniaturized protocols  
165 limiting the volumes of solvents employed have been also proposed and employed in the  
166 last years. Among them, solid-phase microextraction (SPME) [2], and most notably,  
167 QuEChERS methods are highlighted [3]. Nowadays, QuEChERS involves a widely  
168 accepted methodology for the recovery of target analytes from complex matrices, which  
169 is based on an initial extraction with acetonitrile followed by a clean-up using dispersive  
170 SPE [4]. From this basic methodology, multiple modifications have been presented so  
171 far; these are mainly related to an adaptation to the nature and fat content of the sample

172 extracted [3]. Other advanced extraction techniques, such as pressurized liquid extraction  
173 (PLE), have also been successfully employed. These environmentally green tools even  
174 allow the coupling with in-line clean-up steps using adsorbents. This strategy was  
175 followed for the extraction of pesticides from honey that were subsequently analyzed by  
176 GC-MS/MS [5]. Readers interested on gaining deeper insight on extraction methods and  
177 sample preparation for the analysis of contaminants in foods are referred to recent  
178 excellent review papers [2,6-12].

179 Methods directed to quantification of chemical contaminants in food are strongly  
180 influenced by current international legislation, which is generally directed to the  
181 establishment of MRLs on certain substances, and to specify the banned compounds that  
182 cannot be present at any concentration. MRLs for pesticides [13, 14], veterinary drugs  
183 [15, 16] and contaminants [17], are available.

184 The most frequent analytical approach to determine contaminants in foods relies on the  
185 use of tandem MS detection. This detection procedure allows the quantification of known  
186 compounds with great selectivity and sensitivity. Typically, triple quadrupole analyzers  
187 have been widely used to this aim, run under selected reaction monitoring (SRM), also  
188 called multiple reaction monitoring (MRM), mode. This way, each parent ion is  
189 fragmented by collision-induced dissociation (CID) and its two most-intense product ions  
190 are detected. The most-intense one is used for quantification whereas the second is  
191 employed for qualification purposes. This detection procedure allows complying with  
192 European legislation on banned and controlled substances in foods [18]. This regulation  
193 establishes the requirements that an analytical method must meet for an unequivocal  
194 identification and quantification of a controlled substance in a food sample, which means  
195 to gain, at least, four identification points. By using the mentioned approach, the  
196 legislation specifies that one identification point is gained by retention time confirmation



197 with a commercial standard, whereas additional 1.5 identification points are gained for  
198 each ion transition successfully confirmed. As a result, and thanks to the quite fast  
199 scanning speed of modern triple quads, different remarkable applications have been  
200 developed in this field. In Table 1, some recent examples of this methodology for the  
201 quantification of more than 50 contaminants in foods in just one run are summarized. As  
202 it can be observed, most applications are based on the coupling of MS with a separation  
203 technique. LC and GC-based methods are widely extended, although the use of  
204 multidimensional chromatography has also been explored with success.  
205 Multidimensional procedures allow increasing resolving power and separation which can  
206 be beneficial for subsequent MS-based detection, considering that the targeted  
207 compounds will reach the detector more separated in time. This is the case of  
208 comprehensive two-dimensional gas chromatography (GC × GC) that has been coupled  
209 to a TOF-MS analyzer to determine dioxin-related pollutants in complex food samples  
210 [54]. Satisfactory separation of more than 200 micropollutants was achieved, with low  
211 limits of detection. Figure 1 illustrates the good separation attainable using this approach.  
212 Although no practical application of comprehensive two-dimensional liquid  
213 chromatography (LC × LC) has been published so far for the quantification of a wide  
214 group of contaminants, the use of this technique retains a very good potential. In fact, a  
215 first application for the quantification of pesticides in complex food samples, such as  
216 wine, has recently been presented [62]. As can be deduced from the information presented  
217 in Table 1, during the period covered by the present review (2014-2017), the use of triple  
218 quadrupoles in MRM mode is still the most-extended approach. Satisfactory results have  
219 been attained in a variety of applications involving the use of these approaches, using  
220 targeted approaches and reaching the quantification of a significant amount of  
221 components in relatively short analysis times with high sensitivity. Although the basic

222 principles remain relatively constant, different modifications have pushed even forward  
223 the limits of these procedures. This is the case, for instance, of the use of high resolution  
224 MS (HRMS) analyzers instead of the commonly employed triple quads; in fact, the use  
225 of HRMS in the field of food safety is showing an increase. For instance, thanks to the  
226 use of nano-LC and HRMS coupled through the use of ambient dielectric barrier  
227 discharge ionization (DBDI) source, extremely low detection limits, as low as 10 pg mL<sup>-1</sup>,  
228 were achieved for the quantification of pesticide residues [63]. In fact, one of the  
229 possible advantages of using HRMS is the possibility of constructing databases for the  
230 sought compounds, when operating under targeted approaches. The use of these databases  
231 together with parallel reaction monitoring using a Q-Orbitrap analyzer has been shown to  
232 be effective for the appropriate screening and quantification of 157 residues of different  
233 nature in honey [42]. Similar approaches have involved an expansion on the studied  
234 compounds to more than 600 different contaminants, including pesticides, veterinary drug  
235 residues, contaminants, perfluoroalkyl substances, mycotoxins and nitrosamines [61]. In  
236 any case, each MS detection method has its highs and lows; comparative studies testing  
237 the performance of tandem MS versus HRMS to quantify polychlorinated dioxins and  
238 biphenyls in foods have concluded that although the use of GC-MS/MS allows meeting  
239 with the requirements laid by the European Commission, GC-HRMS may fit better for  
240 monitoring purposes as it was shown to produce less false positives [64].

241 In spite of the developed methods, the use of the above described targeted approach has  
242 important limitations, which are mainly related to the determination of unknown  
243 compounds as well as the need of reference commercial standards. For this reason, the  
244 use of similar approaches already developed in other fields for the non-targeted analysis  
245 of contaminants is increasingly proposed, taking advantage of the capabilities of HRMS  
246 modern analyzers [65]. An interesting example has recently been published in order to

247 investigate which compounds of potential concern were present in a pizza box, as a model  
248 of food packaging material [26]. This approach involved the coupling with proper *in-vitro*  
249 assays based on aryl hydrocarbon receptor activity to limit the number of fractions to be  
250 studied after extraction. The most-active fractions were analyzed by using GC-QTOF-  
251 MS and UHPLC-QTOF-MS. The workflow followed in this work is shown in Figure 2.  
252 Seventy-five substances were tentatively identified, among which seven commercially  
253 available could be further studied but could not explain a significant proportion of the  
254 aryl hydrocarbon receptor response in the extract. Thus, it could be concluded that other  
255 very active substances still remained unidentified in the food container [26]. Using  
256 another different non-targeted approach Zomer and Mol also showed the high potential  
257 of state-of-the-art HRMS instrumentation [50]. Using a hybrid HRMS analyzer, a new  
258 fully non-targeted approach for data acquisition combining full-scan and fragmentation  
259 was developed utilizing variable data-independent acquisition for the generation of  
260 fragment ions. Quantitative validation of the methodology using a mixture of 184  
261 pesticides in two food matrices showed that this approach was suitable for ca. 93 % of  
262 the assayed pesticide/matrix/concentration combinations studied in agreement with EU  
263 guidelines. Thus, this LC-full-scan HRMS method has been suggested as an alternative  
264 for triple quad MS-based methods. Moreover, the same data could be used to screen  
265 samples for a large number of compounds with lower probability of being present,  
266 reducing the chance for false-negatives compared to other previously used full-scan-  
267 based protocols [50].

268 The most interesting aspect related to the non-targeted methodology is based on the  
269 possibility of detecting substances not previously pre-selected, thus, increasing the chance  
270 for the proper detection of unknown and unexpected compounds. These metabolomics  
271 approaches may gain advantage of data mining tools initially developed in other fields. A

272 proof-of-concept study, demonstrating the ability of these tools to identify unknown  
273 chlorinated chemicals in honey samples has been reported [29]. However, the use of these  
274 diverse non-targeted methodologies is still somewhat limited compared to the targeted  
275 approach, as it is clearly illustrated in Table 1. Further developments on this field in the  
276 near future are expected.

277

## 278 **2.2. Detection of microbial contaminants (pathogens and toxins)**

279 Risks of natural origin for food safety are mainly related to the presence or activity of  
280 microorganisms. Thus, foods may be contaminated directly by the presence of pathogens,  
281 which could cause an infection to the consumer, or may be indirectly contaminated by  
282 toxins produced by a particular microorganism. Contamination of food with pathogens  
283 may imply very serious consequences on health, being the most extended diarrhea, and  
284 can occur at any point of the food production chain due to inadequate hygiene conditions.  
285 On the other hand, the presence of toxin producers within or near food related products  
286 can be a potential source of contamination. This is the case, for instance, of cereal  
287 products contaminated with mycotoxins, or shellfish contaminated with microalgal toxins  
288 that are bioaccumulated in those filter-feeding animals.

289 For the detection and quantification of toxins in foods, similar approaches to those already  
290 described for chemical contaminants are widely employed. The methodology to quantify  
291 those components by tandem MS is very much the same; however, in this case, the natural  
292 toxin variability potentially present in a particular food product mean that less compounds  
293 have to be analyzed, and thus, advanced metabolomics-based approaches are not required.  
294 Instead, proper sample preparation for toxins extraction and quantification by MRM using  
295 triple quads is the most common MS-based methodology applied [66-67]. Nuts [68],  
296 maize [69], shellfish [70], tomato [71], or beer [72], among others, are examples of food

297 products assayed following this approach. However, some modifications have been also  
298 introduced to this methodology in order to increase the performance of methods as well  
299 as to allow a very sensitive detection, as some of the natural toxins that might be  
300 potentially found in foods are very toxic (even lethal) at extremely low concentrations.  
301 For instance, the use of a multiple antibody immunoaffinity column for the selective  
302 extraction of 7 toxins before HPLC-MS/MS determination has been recently reported  
303 [73]. This method allowed extending the linear range of the determination as well as to  
304 decrease the detection limits to the low  $\mu\text{g kg}^{-1}$  level compared to previously developed  
305 methods. Other sample preparation-oriented improvements have been directed to the  
306 implementation of inexpensive graphitized carbon for SPE of paralytic shellfish toxins,  
307 showing excellent capabilities [74].

308 Other sensitive gains have been attained through the analytical tool employed prior MS.  
309 The ultrasensitive detection, with detection limits as low as 0.38 fmol of saxitoxin was  
310 achieved in seafood samples thanks to a reaction involving diethylenetriamine-  
311  $\text{N}_3\text{N}'_3\text{N}''_3\text{N}'''_3$ -pentaacetic acid. This compound can couple with saxitoxin and  
312 simultaneously chelate with  $\text{Eu}^{3+}$  to allow metallic labeling of this toxin, that may be  
313 quantified with extremely high sensitivity using capillary electrophoresis-inductively  
314 coupled plasma-MS detection (CE-ICP-MS) [75]. Direct determination of toxins may  
315 have the further advantage of increasing throughput in food safety laboratories. As  
316 already mentioned, some direct analysis MS techniques have been employed for the  
317 quantification of chemical contaminants (see Table 1). In the case of toxins, some direct  
318 methods have been also presented. Indeed, domoic acid has been quantified in mussel  
319 tissues directly by MS/MS using SRM mode without any sample extraction, clean-up or  
320 separation. This has been obtained using laser ablation electrospray ionization (LAESI),  
321 reaching limits of detection of  $1 \text{ mg kg}^{-1}$  for this compounds. This LOD is not particularly

322 low compared to other more conventional approaches based on extraction/separation and  
323 MS/MS detection, but it has to be considered that each analysis takes around just 10 s,  
324 thus, being very attractive for routine analysis [76]. Although these recent advances have  
325 enhanced in different manners the detection of toxins in food, any of them shows a purely  
326 metabolomics-based strategy. In this regard, this subfield of analysis should benefit in the  
327 future from applications already developed for contaminants analysis as those previously  
328 described in Section 2.1. In spite from this, some efforts have already been made, such as  
329 the development of an analytical micro HPLC-MS/MS method for the simultaneous  
330 quantification of 26 mycotoxins in maize with total run times of 9 min and reduced  
331 solvent consumption (below 0.3 mL) [77].

332 Other food safety-related methodologies are mostly focused on the detection of pathogen  
333 microorganisms that could be present in the food products posing a serious risk to  
334 consumers' health. Although different molecular techniques and proteomics-based  
335 approaches may be used to detect and identify the microorganisms present in a sample,  
336 in recent years much effort has been also focused on the determination of microbial  
337 volatile organic compounds (MVOCs) as markers of microbiological contamination [78].  
338 To that aim, the most-extended analytical MS-based approach is based on the use of GC-  
339 MS coupled to a proper sample preparation/extraction protocol, such as SPME or  
340 headspace (HS) sampling. After the determination of a group of volatiles as wide as  
341 possible, multivariate analysis of data is necessary to correlate the presence of specific  
342 compounds with the growth of particular pathogens. This approach has been employed  
343 to predict shelf-life, evaluating potential chemical spoilage indices of Atlantic salmon  
344 stored under aerobic conditions [79], sea bass stored under air and under modified  
345 atmosphere [80], sea bream depending on the storage conditions [81-82], as well as  
346 minced meat [83] or pork [84]. Another possibility gaining interest in recent times is the

347 determination of MVOCs by real time analysis through the application of proton-transfer-  
348 reaction-MS (PTR-MS). This technique is able to provide with fast on-line analyses that  
349 are very appropriate for determination of the real-time evolution of volatiles. Different  
350 applications have been recently published to determine MVOCs of microbial origin from  
351 selected strains [85] as well as in food products such as chicken meat [86] or milk [87-  
352 88]. To allow the continuous on-line monitoring, different set-ups have been developed,  
353 for instance, allowing the monitoring of four meat samples in parallel [86] (Figure 3A),  
354 or other more manually-operated set-ups for milk (Figure 3B) [87].

355

### 356 **3. MS-BASED METABOLOMICS TO ASSESS FOOD QUALITY**

357 Nowadays, food quality is one of the major concerns of the food industry. Its evaluation  
358 is a complex task due to the multiple aspects that may be considered to achieve an  
359 appropriate food quality. Food composition, aroma, flavor, or nutritional properties are  
360 among the most important aspects that may be evaluated in food quality assessments.  
361 Different types of analysis are clearly needed to evaluate all these aspects. Is at this point  
362 where MS-based metabolomics approaches are gaining attention due to their  
363 demonstrated capability to establish links between relevant food aspects and food quality  
364 perception.

365 Table 2 summarizes the most relevant applications of MS-based metabolomics strategies  
366 for food quality published during the period of time covered by this review (2014-2017).  
367 As can be observed, these works are mainly focused on the use of this kind of platform  
368 to establish the relationship between the chemical composition and food quality, to  
369 control food authentication and adulteration, or to differentiate food samples according  
370 to their variety. To achieve these aims, non-targeted approaches have usually been  
371 employed followed by data-processing and multivariate analysis to assess possible  
372 differences among samples. An interesting strategy is the combination of non-targeted

373 and targeted methods; its usefulness has recently been reported for the qualitative analysis  
374 of curcuminoids in turmeric [91]. This integrated strategy involves a non-targeted  
375 analysis by LC-QTOF-MS/MS and a targeted approach by LC-QTRAP-MS/MS. Figure  
376 4 depicts the workflow followed in this study. Ninety-six curcuminoids were fully  
377 characterized following this exclusive methodology. Anyhow, the ultimate goal of the  
378 researches developed to assess food quality is to determine relevant compounds that may  
379 be selected as quality markers. Afterwards, just a few studies have developed targeted  
380 methodologies for the routine analysis of those markers [89, 90]. However, this fact is  
381 interesting from an analytical point of view, since a targeted method requires less  
382 sophisticated instrumentation, is usually simpler and the data are more easily analyzed,  
383 being, therefore, more applicable for routine analysis.

384 One of the relevant points to assess food quality by MS-based metabolomics is, again, the  
385 choice of proper sample preparation procedures. This fact will depend not only on the  
386 analytical technique employed to perform the analysis but also on the particular aim of  
387 the study. Although nowadays the use of modern mass spectrometers enables to perform  
388 analysis with high sensitivity which may simplify sample preparation, the inherent  
389 complexity of food samples makes this step a critical factor in the determination of  
390 metabolites, as previously mentioned. In any case, to prevent any substantial loss of  
391 possible relevant metabolites, minimum sample preparation is preferable. Even though  
392 simple solvent-based extraction procedures have been the method of choice during the  
393 last years (see Table 2), certain GC-MS methodologies have required the use of other  
394 sample preparation techniques such as ultrasound-assisted extraction in tandem with  
395 dispersive liquid-liquid microextraction (UAE-DLLME) [98], solid-phase extraction  
396 (SPE) [101], static headspace extraction (HS) [108] or headspace solid-phase micro-



397 extraction (HS-SPME) [115], in order to improve the extraction of volatile compounds or  
398 to achieve a preconcentration effect, thus, increasing method sensitivity and efficiency.  
399 As can be deduced from the information shown in Table 2, the majority of applications  
400 of MS-based metabolomics approaches included the coupling LC-MS and/or GC-MS.  
401 Concerning LC-MS, the use of methods based on the UHPLC has increased considerably  
402 in the last years due to its capability to perform complex analysis with high efficiency and  
403 resolution in a short time. Different metabolomics studies have employed UHPLC  
404 technology for example to carry out the authentication and the evaluation of possible  
405 adulterations in fruits juices [89, 90] or saffron [99], demonstrating the feasibility of these  
406 methodology to face one of the most growing problems in the global market. Another  
407 point that should be highlighted regarding LC is that although C18 columns are by far the  
408 most utilized, methods based on the use of hydrophilic interaction chromatography  
409 (HILIC) have also successfully been applied to food quality. This allows profiling highly  
410 polar and hydrophilic compounds providing complementary metabolic information to  
411 reversed-phase LC. Even though there are some drawbacks associated with HILIC  
412 (variability in retention times, low peak efficiency, and long re-equilibration times after  
413 gradient elution), this methodology has been used for the assessment of contamination  
414 and degradation of infant formulas [97] or to identify biomarkers of meat quality [104,  
415 106].

416 Regarding GC-MS, in spite of the need to include a derivatization step in the sample  
417 treatment to increase the range of metabolites that can be analyzed, GC-MS metabolomics  
418 approaches have been broadly used to evaluate food quality as it can be observed in Table  
419 2. In these cases, GC has been hyphenated to a great variety of mass analyzers including  
420 simpler MS instruments, like quadruple (Q) working at electron ionization mode  
421 [98,102,103,112,115], or ion trap (IT) [114], as well as high resolution instruments

422 [93,95,105,109,110], and even hybrid analyzers [96,101,107]. An interesting work based  
423 on the use of GC coupled to TOF-MS has been employed to develop a non-targeted  
424 metabolomics approach capable to establish differences between wine grape cultivars  
425 [93]. To do that, two grape cultivars were profiled and 115 metabolites were identified  
426 and quantified. Among them, sugars and amino acids showed an opposite behavior in  
427 both cultivars. To carry out the biological interpretation of the data and to obtain an  
428 overview of the abundance of these compounds in the development of the cultivars, their  
429 behavior in the primary metabolism pathways was investigated. Figure 5 depicts the level  
430 of each metabolite within each cultivar during the grape development stage in different  
431 pathways (tricarboxylic acid cycle, glycolysis, amino acid synthesis, and sucrose  
432 synthesis). Other interesting strategies based on GC-MS metabolomics platforms have  
433 been applied, for instance, to investigate the effect of volatile compounds for the  
434 classification of saffron based on the concentration of biomarkers [98], to classify olive  
435 oils according to their quality parameters [101], or to detect milk or meat adulteration  
436 [103,107].

437 Although LC-MS and GC-MS have been the preferred platforms to assess food quality,  
438 GC × GC [108] and CE methods [104] coupled to TOF analyzers have also been applied  
439 with success. The first one has allowed to establish associations between volatile  
440 metabolites and perception of rice aroma, creating a panel of biomarkers of rice flavor  
441 quality [108]. These results are valuable for breeding programs since can be used to  
442 choose pleasant rice aromas. In the latter, the feasibility of using a polymer-coated-  
443 capillary for the separation of anionic metabolites both in orange juice and wine has been  
444 demonstrated [104]. It offers a complementary coverage of the metabolome of these  
445 samples to those provide by other analytical techniques. Due to the demonstrated  
446 capabilities of both GC × GC and CE, it is expected that future developments in this field

447 will gain advantage of those methods, since the full potential of these techniques in food  
448 metabolomics has not been reached.

449

#### 450 **4. MS-BASED METABOLOMICS FOR FOOD TRACEABILITY**

451 Food traceability is also a relevant topic within food analysis, whose main purpose is to  
452 provide a continuous monitoring of a food in the entire supply chain; this monitoring has  
453 been often defined as “from farm to fork”. Undoubtedly, food traceability is closely  
454 related to food quality, food safety and public health. This topic has a great importance  
455 not only to food industries but also to consumers who are increasingly demanding more  
456 information about each stage of the food that they consume. In this regard, MS-based  
457 metabolomics approaches are essential since they are capable to provide the level of  
458 accuracy needed for traceability management.

459 Bearing in mind that traceability involves knowing the composition and origin of a food,  
460 it is clear that the determination of the geographical origin may be considered the starting  
461 point for food traceability. Geographical origin assessments have not only relevant  
462 implications from an economical point of view but also they are a key parameter in terms  
463 of food quality. The most common metabolomics strategies developed to discriminate  
464 food samples according to their geographical origin are non-targeted approaches based  
465 on the use of LC (mainly UHPLC) coupled with HRMS. Using the most suitable sample  
466 preparation protocols according to the features of each food sample and the appropriate  
467 multivariate data analysis, these MS-based methodologies are able to point out different  
468 metabolites as potential markers of food origin. This kind of approaches has successfully  
469 been applied for the origin assessment of extra virgin olive oil (EVOO) [117] orange  
470 [118], hazelnuts [119] or cocoa beans [120].

471 Other relevant branch in food traceability is focused on monitoring changes in the food

472 metabolic profiles produced by food-processing. Production steps, including for instance,  
473 heat treatments, fermentation, and storage, among others, can alter nutritional and  
474 organoleptic properties of foods, as well as lead to a substantial loss of health-promoting  
475 compounds. This fact has been demonstrated by a recent and interesting non-targeted  
476 UHPLC-QTOF-MS method developed to evaluate the phenolic profiles of three different  
477 processed tomato products and tomato paste produced by three different treatments [121].  
478 The combination of the results obtained from the metabolomics analysis with total  
479 phenolic and lycopene content, and antioxidant capacity showed that processing affects  
480 the nutritional and health-promoting potential of tomato products. Besides, the  
481 metabolomics approach shows its high potential in traceability purposes since the  
482 treatment provide a characteristic phenolic profile.

483 Other non-targeted LC-HRMS platforms have also been applied with success to study the  
484 effect of storage conditions on the metabolic profile of red wine [122] or iceberg lettuce  
485 [123], as well as to compare the effects of thermal processing on *Brassica* vegetables  
486 [124]. After processing and carrying out the multivariate data analysis, the final purpose  
487 of this kind of studies is to find the relationship between the changes on the metabolite  
488 profile with a loss of food quality. Figure 6 shows an example of the data analysis  
489 procedure followed to explore the metabolome of lettuce in order to evaluate changes  
490 related to storage time and genetics. Fermentation and ripening are also relevant process  
491 which may change the food metabolome. Two interesting examples have been described  
492 in the literature to explore the changes that occur in the metabolic profile of cocoa beans  
493 [125] and cheese [126] as a consequence of fermentation and ripening process,  
494 respectively. Bearing in mind that these two processes are critical steps in the processing  
495 of high quality cocoa beans or in the formation of specific characteristics of cheese, the  
496 results obtained in these metabolomics assays are of high value for the food industry since

497 they shed new light into fermentation and ripening optimization.

498 Even though most applications developed for food traceability in the period of time  
499 covered by this review are based on the coupling of MS with LC, GC-MS methodologies  
500 have also been proposed. For instance, using headspace GC-MS non-targeted approach  
501 was possible to distinguish the effect of different process steps (including not only thermal  
502 processing but also blanching and high hydrostatic pressure) on the chemical composition  
503 of mango [127]. Once again, the results obtained clearly demonstrate the influence of  
504 these steps on the volatile profiles of processed products. GC-MS metabolomics approach  
505 has also proven to be an excellent tool to evaluate the modifications that may occur during  
506 the cooking of different types of pasta [128].

507 Another possibility gaining interest in recent times is the use of CE coupled to MS as  
508 analytical platform for traceability assays. For example, Sugimoto et al. developed two  
509 CE-TOF-MS methodologies for anionic and cationic metabolite analysis of dry-cured  
510 ham [129]. The results obtained enabled to establish a correlation between the metabolite  
511 profiles of twelve kinds produced in different countries and processed under different  
512 conditions and the ripening period and processing conditions. Even though CE-MS  
513 strategies are being mainly developed and applied for biological samples, nowadays, is  
514 possible to find some applications devoted to food analysis. Further progress in this field  
515 are expected in the near future.

516 Although non-targeted strategies have been the most-extended approach to evaluate  
517 changes in the metabolic profiles of food samples during food-processing, targeted  
518 analysis may also be very useful; this kind of approaches has been employed to evaluate  
519 the metabolic changes that take place in two starch potato genotypes in response to  
520 osmotic stress [130] or during avocado development and maturation [131].

521

## 522 **5. CONCLUSION AND FUTURE OUTLOOKS**

523 As it can be deduced from the update shown in this review paper, the use of MS-based  
524 approaches for food safety, food quality and traceability is still far from reaching its  
525 maximum potential. It is quite obvious that the use of MS, particularly high resolution  
526 MS, will still be dominant in studies on the mentioned fields in the years to come. In this  
527 regard, the continuous improvement of available instruments will be translated to  
528 enhanced capabilities of the developed methods. As MS is most frequently used  
529 hyphenated to other analytical tools, the improvement on robustness of couplings and  
530 available interfaces and ionization tools, including those employed in direct analysis, will  
531 positively influence the obtainable results. This way, new to-be-controlled substances  
532 appearing in the market as well as unknown ways to perform frauds during production of  
533 valuable food products could be discovered. Specifically, within the food safety field,  
534 new multi-residue and multi-targeted methods will surely continue appearing, ready to  
535 help on the food control area. However, more interestingly, the development of novel  
536 non-targeted metabolomics-based approaches will help to gain a holistic view of the food  
537 safety issue. Those procedures are clearly more capable of discovering new safety hazards  
538 beyond the use of the regulated compounds and contaminants. But those approaches  
539 could have even more potential if accompanied by proper in-vitro and in-vivo assays, so  
540 that the perspectives may be further opened, for instance, to the discovery of markers of  
541 toxicity.

542 Food quality will also benefit from the extension of metabolomics MS-based approaches  
543 to other studies. Within this field, the further application and development of these  
544 methodologies could help to increase the available knowledge on which compounds  
545 present in food that may have a still concealed importance for food quality perception.  
546 This is the case, for example, of the application of this kind of procedure to reveal the

547 whole sensory pattern of a food product, a concept already applied in flavoromics  
548 researches. Likewise, as metabolomics methods evolve in the future, new relationships  
549 between food components and particular characteristics related to food quality will be  
550 discovered.

551 Regarding traceability, much effort is expected to be focused on the development of new  
552 methodologies to assess food authentication and geographical origin of valuable food  
553 products. However, this field is intimately linked to food quality as some traceability  
554 aspects are related to quality. For instance, development of traceability potential will help  
555 to discover how production processes throughout the food production and  
556 commercialization chain may affect quality parameters. In this regard, the use of  
557 alternative analytical techniques to LC and GC, such as CE or multidimensional  
558 approaches (including LC × LC and GC × GC) could offer complementary selectivity  
559 and thus, information, that would help to increase the metabolite coverage of the studied  
560 system. This enhanced coverage could positively influence the applicability of MS-based  
561 metabolomics studies in the three different mentioned fields.

562 In summary, it is clear that although the interest of using MS-based metabolomics  
563 approaches in food safety, quality and traceability is already high, further developments  
564 in these methodologies will have a great influence on the mentioned fields in the near  
565 future.

566

567

## 568 **ACKNOWLEDGEMENTS**

569 The authors would like to thank Projects AGL2014-53609-P (MINECO, Spain) and  
570 S2013/ABI-3028 (AVANSECAL-CM, from Comunidad Autónoma of Madrid, Spain,  
571 and the European funding from FEDER program) for financial support. M.C.P. thanks

572 the Ministry of Economy and Competitiveness (Spain) for her “Ramón y Cajal” research  
573 contract (RYC-2013-12688). R.P.M. thanks the University of Alcalá for her pre-doctoral  
574 contract.



575 **REFERENCES**

- 576 [1] M. Castro-Puyana, M. Herrero, Metabolomics approaches based on mass  
577 spectrometry for food safety, quality and traceability, *TrAC, Trends Anal. Chem.* 52  
578 (2013) 74-87.
- 579 [2] E.A. Souza-Silva, E. Gionfriddo, J. Pawliszyn, A critical review of the state of the art  
580 of solid-phase microextraction of complex matrices II. Food analysis, *TrAC, Trends*  
581 *Anal. Chem.* 71 (2015) 236-248.
- 582 [3] M.A. González-Curbelo, B. Socas-Rodríguez, A.V. Herrera-Herrera, J. González-  
583 Sálamo, J. Hernández-Borges, M.A. Rodríguez-Delgado, Evolution and applications of  
584 the QuEChERS method, *TrAC, Trends Anal. Chem.* 71 (2015) 169-185.
- 585 [4] M.A. González-Curbelo, A.V. Herrera-Herrera, L.M. Ravelo-Perez, J. Hernández-  
586 Borges. Sample-preparation methods for pesticide-residue analysis in cereals and  
587 derivatives, *TrAC, Trends Anal. Chem.* 38 (2012) 32-51.
- 588 [5] L.M. Chiesa, G.F. Labella, S. Panseri, D. Britti, F. Galbiati, R. Villa, F. Arioli,  
589 Accelerated solvent extraction by using a “in-line” clean-up approach for multiresidue  
590 analysis of pesticides in organic honey, *Food Addit. Contam. Part A* (2017) in press DOI:  
591 10.1080/19440049.2017.1292558.
- 592 [6] V. Vazquez-Roig, Y. Picó, Pressurized liquid extraction of organic contaminants in  
593 environmental and food samples, *TrAC, Trends Anal. Chem.* 71 (2015) 55-64.
- 594 [7] P. Berton, N.B. Lana, J.M. Rios, J.F. Garcia-Reyes, J.C. Altamirano, State of the art  
595 of environmentally friendly sample preparation approaches for determination of PBDEs  
596 and metabolites in environment, *Anal. Chim. Acta* 905 (2016) 24-41.
- 597 [8] T. Rejczak, T. Tuzimski, Recent Trends in Sample Preparation and Liquid  
598 Chromatography/Mass Spectrometry for Pesticide Residue Analysis in Food and Related  
599 Matrixes, *J. AOAC Int.* 98 (2015) 1143-1162.
- 600 [9] C.H. Xu, G. S. Chen, Z.H. Xiong, X.X. Fang, X.C. Wang, Y. Liu, Applications of  
601 solid-phase microextraction in food analysis, *TrAC, Trends Anal. Chem.* 80 (2016) 12-  
602 29.
- 603 [10] M. Andraščíková, E. Matisová, S. Hrouzková, Liquid phase microextraction  
604 techniques as a sample preparation step for analysis of pesticide residues in food, *Sep.*  
605 *Pur. Rev.* 44 (2015) 1-18.
- 606 [11] R. Romero-Gonzalez, F.J.A. Liebanas, R.L. Pez-Ruiz, A.G. Frenich, Sample  
607 treatment in pesticide residue determination in food by high-resolution mass

608 spectrometry: Are generic extraction methods the end of the road?, J. AOAC Int. 99  
609 (2016) 1395-1402.

610 [12] A. Lawal, G.H. Tan, A.M.A. Alsharif, Recent advances in analysis of pesticides in  
611 food and drink samples using LPME techniques coupled to GC-MS and LC-MS: A  
612 review, J. AOAC Int. 99 (2016) 1383-1394.

613 [13] Commission Regulation (EC) no 396/2005 of the European parliament and of the  
614 council of 23 February 2005 on maximum residue levels of pesticides in or on food and  
615 feed of plant and animal origin and amending Council Directive 91/414/EEC, Off. J. Eur.  
616 Commun. L70, (2005) 1-16.

617 [14] U.S., Department of Agriculture, Foreign Agricultural Service, Maximum Residue  
618 Limit Database, 2014. Available from: [http://www.fas.usda.gov/maximum-residue-](http://www.fas.usda.gov/maximum-residue-limits-mrl-database)  
619 [limits-mrl-database](http://www.fas.usda.gov/maximum-residue-limits-mrl-database)

620 [15] Commission Regulation (EC) no 37/2010 of 22 December 2009, On  
621 pharmacologically active substances and their classification regarding maximum residue  
622 limits in foodstuffs of animal origin, Off. J. Eur. Union L15, (2009) 1-76.

623 [16] Tolerances for residues on new animal drugs in food. Code of Federal Regulations.  
624 Food and Drugs, Part 556, Title 21; U.S. Food Drug Administration.

625 [17] Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum  
626 levels for certain contaminants in foodstuffs, Off. J. Eur. Union L364, (2006) 1-68.

627 [18] European Union Commission Decision 2002/657/EC.

628 [19] M. Farré, Y. Picó, D. Barceló, Application of ultra-high pressure liquid  
629 chromatography linear ion-trap orbitrap to qualitative and quantitative assessment of  
630 pesticide residues, J. Chromatogr. A 1328 (2014) 66-79.

631 [20] H.G.J. Mol, M. Tienstra, P. Zomer, Evaluation of gas chromatography electron  
632 ionization full scan high resolution orbitrap mass spectrometry for pesticide residue  
633 analysis, Anal. Chim. Acta 935 (2016) 161-172.

634 [21] C. Cruzeiro, N. Rodrigues-Oliveira, S. Velhote, M. Ângelo Pardal, E. Rocha, M.  
635 João Rocha, Development and application of a QuEChERS-based extraction method for  
636 the analysis of 55 pesticides in the bivalve *Scrobicularia plana* by GC-MS/MS, Anal.  
637 Bioanal. Chem. 408 (2016) 3681-3698.

638 [22] G.C.R.M. Andrade, S.H. Monteiro, J.G. Francisco, L.A. Figueiredo, R.G. Botelho,  
639 V.L. Tornisielo, Liquid chromatography-electrospray ionization tandem mass

640 spectrometry and dynamic multiple reaction monitoring method for determining multiple  
641 pesticide residues in tomato, *Food Chem.* 175 (2015) 57-65.

642 [23] S. Moretti, G. Dusi, D. Giusepponi, S. Pellicciotti, R. Rossi, G. Saluti, G. Cruciani,  
643 R. Galarinia, Screening and confirmatory method for multiclass determination of 62  
644 antibiotics in meat, *J. Chromatogr. A* 1429 (2016) 175-188.

645 [24] A. Vavrou, L. Vapenka, J. Sosnovcova, K. Kejlova, K. Vrbík, D. Jírova, Method for  
646 analysis of 68 organic contaminants in food contact paper using gas and liquid  
647 chromatography coupled with tandem mass spectrometry, *Food Control* 60 (2016) 221-  
648 229.

649 [25] S.S. Shida, S. Nemoto, R. Matsuda, Simultaneous determination of acidic pesticides  
650 in vegetables and fruits by liquid chromatography-tandem mass spectrometry, *J. Environ.*  
651 *Sci. Health, Part B* 50 (2015) 151-162.

652 [26] L. Bengtström, A.K. Rosenmai, X. Trier, L.K. Jensen, K. Granby, A.M. Vinggaard,  
653 M. Driffield, J.H. Petersen, Non-targeted screening for contaminants in paper and board  
654 food-contact materials using effect-directed analysis and accurate mass spectrometry,  
655 *Food Addit. Contam., Part A* 33 (2016) 1080-1093.

656 [27] G.R. Chang, H.S. Chen, F.Y. Lin, Analysis of banned veterinary drugs and herbicide  
657 residues in shellfish by liquid chromatography-tandem mass spectrometry (LC/MS/MS)  
658 and gas chromatography-tandem mass spectrometry (GC/MS/MS), *Mar. Pollut. Bull.* 113  
659 (2016) 579-584.

660 [28] M.E. Dasenaki, C.S. Michali, N.S. Thomaidis, Analysis of 76 veterinary  
661 pharmaceuticals from 13 classes including aminoglycosides in bovine muscle by  
662 hydrophilic interaction liquid chromatography-tandem mass spectrometry, *J.*  
663 *Chromatogr. A* 1452 (2016) 67-80.

664 [29] J. Cotton, F. Leroux, S. Broudin, M. Marie, B. Corman, J.C. Tabet, C. Ducruix, C.  
665 Junot, High-resolution mass spectrometry associated with data mining tools for the  
666 detection of pollutants and chemical characterization of honey samples, *J. Agric. Food*  
667 *Chem.* 62 (2014) 11335-11345.

668 [30] S. Chawla, H.K. Patel, K.M. Vaghela, F. Khan Pathan, H.N. Gor, A.R. Patel, P.G.  
669 Shah, Development and validation of multi residue analytical method in cotton and  
670 groundnut oil for 87 pesticides using low temperature and dispersive cleanup on gas

671 chromatography and liquid chromatography-tandem mass spectrometry, *Anal. Bioanal.*  
672 *Chem.* 408 (2016) 983-997.

673 [31] D. Orso, L. Floriano, L.C. Ribeiro, N.M.G. Bandeira, O.D. Prestes, R. Zanella,  
674 Simultaneous determination of multiclass pesticides and antibiotics in honey samples  
675 based on ultra-high performance liquid chromatography-tandem mass spectrometry,  
676 *Food Anal. Methods* 9 (2016) 1638-1653.

677 [32] H. Zhang, J. Wang, L. Li, Y. Wang, Determination of 103 Pesticides and their main  
678 metabolites in animal origin food by QuEChERS and liquid chromatography-tandem  
679 mass spectrometry, *Food Anal. Methods* (2016) *in press* DOI 10.1007/s12161-016-0736-  
680 7.

681 [33] O. Golge, B. Kabak, Evaluation of QuEChERS sample preparation and liquid  
682 chromatography-triple-quadrupole mass spectrometry method for the determination of  
683 109 pesticide residues in tomatoes, *Food Chem.* 176 (2015) 319-332.

684 [34] J. Cho, J. Lee, C.U. Lim, J. Ahn, Quantification of pesticides in food crops using  
685 QuEChERS approaches and GC-MS/MS, *Food Addit. Contam., Part A* 33 (2016) 1803-  
686 1816.

687 [35] O. Golge, B. Kabak, Determination of 115 pesticide residues in oranges by high-  
688 performance liquid chromatography-triple-quadrupole mass spectrometry in  
689 combination with QuEChERS method, *J. Food Compos. Anal.* 41 (2015) 86-97.

690 [36] L. Han, J. Matarrita, Y. Sapozhnikova, S.J. Lehotay, Evaluation of a recent product  
691 to remove lipids and other matrix co-extractives in the analysis of pesticide residues and  
692 environmental contaminants in foods, *J. Chromatogr. A* 1449 (2016) 17-29.

693 [37] C. Rasche, B. Fournes, U. Dirks, K. Speer, Multi-residue pesticide analysis (gas  
694 chromatography-tandem mass spectrometry detection)-Improvement of the quick, easy,  
695 cheap, effective, rugged, and safe method for dried fruits and fat-rich cereals-Benefit and  
696 limit of a standardized apple purée calibration (screening), *J. Chromatogr. A* 1403 (2015)  
697 21-31.

698 [38] G. Ramadan, M. Al Jabir, N. Alabdulmalik, A. Mohammed, Validation of a method  
699 for the determination of 120 pesticide residues in apples and cucumbers by LC-MS/MS,  
700 *Drug Test. Anal.* 8 (2016) 498-510.

701 [39] D. Chen, J. Yu, Y. Tao, Y. Pan, S. Xie, L. Huang, D. Peng, X. Wang, Y. Wang, Z.  
702 Liu, Z. Yuan, Qualitative screening of veterinary anti-microbial agents in tissues, milk,  
703 and eggs of food-producing animals using liquid chromatography coupled with tandem  
704 mass spectrometry, *J. Chromatogr. B* 1017-1018 (2016) 82-88.

705 [40] M. Piatkowska, P. Jedziniak, J. Zmudzki, Multiresidue method for the simultaneous  
706 determination of veterinary medicinal products, feed additives and illegal dyes in eggs  
707 using liquid chromatography-tandem mass spectrometry, *Food Chem.* 197 (2016) 571-  
708 580.

709 [41] S. Pizzini, R. Piazza, G. Cozzi, C. Barbante, Simultaneous determination of  
710 halogenated contaminants and polycyclic aromatic hydrocarbons: a multi-analyte method  
711 applied to filter-feeding edible organisms, *Anal. Bioanal. Chem.* 408 (2016) 7991-7999.

712 [42] Y. Li, J. Zhang, Y. Jin, L. Wang, W. Zhao, W. Zhang, L. Zhai, Y. Zhang, Y. Zhang,  
713 J. Zhou, Hybrid quadrupole-orbitrap mass spectrometry analysis with accurate-mass  
714 database and parallel reaction monitoring for high-throughput screening and  
715 quantification of multi-xenobiotics in honey, *J. Chromatogr. A* 1429 (2016) 119-126.

716 [43] J. Náchér-Mestre, R. Serrano, T. Portolés, M.H.G. Berntssen, J. Pérez-Sánchez, F.  
717 Hernández, Screening of pesticides and polycyclic aromatic hydrocarbons in feeds and  
718 fish tissues by gas chromatography coupled to high-resolution mass spectrometry using  
719 atmospheric pressure chemical ionization, *J. Agric. Food Chem.* 62 (2014) 2165-2174.

720 [44] S. Saito-Shida, S. Nemoto, R. Teshima, Multiresidue determination of pesticides in  
721 tea by gas chromatography-tandem mass spectrometry, *J. Environ. Sci. Health, Part B* 50  
722 (2015) 760-776.

723 [45] S.E. Kern, L.A. Lin, F.L. Fricke, Accurate mass fragment library for rapid analysis  
724 of pesticides on produce using ambient pressure desorption ionization with high-  
725 resolution mass spectrometry, *J. Am. Soc. Mass Spectrom.* 25 (2014) 1482-1488.

726 [46] D.L. Christodoulou, P. Kanari, O. Kourouzidou, M. Constantinou, P. Hadjiloizou,  
727 K. Kika, P. Constantinou, Pesticide residues analysis in honey using ethyl acetate  
728 extraction method: validation and pilot survey in real samples, *Int. J. Environ. Anal.*  
729 *Chem.* 95 (2015) 894-910.

730 [47] D.L. Christodoulou, P. Kanari, P. Hadjiloizou, P. Constantinou, Pesticide residues  
731 analysis in wine by liquid chromatography-tandem mass spectrometry and using ethyl

732 acetate extraction method: validation and pilot survey in real samples, *J. Wine Res.* 26  
733 (2015) 81-98.

734 [48] A. Páleníková, G. Martínez-Domínguez, F.J. Arrebola, R. Romero-González, S.  
735 Hrouzková, A. Garrido Frenich, Multifamily determination of pesticide residues in soya-  
736 based nutraceutical products by GC/MS-MS, *Food Chem.* 173 (2015) 796-807.

737 [49] F. Hildmann, C. Gottert, T. Frenzel, G. Kempe, K. Speer, Pesticide residues in  
738 chicken eggs-A sample preparation methodology for analysis by gas and liquid  
739 chromatography/tandem mass spectrometry, *J. Chromatogr. A* 1403 (2015) 1-20.

740 [50] P. Zomer, H.G.J. Mol, Simultaneous quantitative determination, identification and  
741 qualitative screening of pesticides in fruits and vegetables using LC-Q-Orbitrap-MS,  
742 *Food Addit. Contam., Part A* 32 (2015) 1628-1636.

743 [51] V. Hanot, S. Gosciny, M. Deridder, A simple multi-residue method for the  
744 determination of pesticides in fruits and vegetables using a methanolic extraction and  
745 ultra-high-performance liquid chromatography-tandem mass spectrometry: Optimization  
746 and extension of scope, *J. Chromatogr. A* 1384 (2015) 53-66.

747 [52] H. Shendy, M.A. Al-Ghobashy, M.N. Mohammed, S.A. Gad Alla, H.M. Lotfy,  
748 Simultaneous determination of 200 pesticide residues in honey using gas  
749 chromatography-tandem mass spectrometry in conjunction with streamlined  
750 quantification approach, *J. Chromatogr. A* 1427 (2016) 142-160.

751 [53] S.B. Turnipseed, J.J. Lohne, J.M. Storey, W.C. Andersen, S.L. Young, J.R. Carr,  
752 M.R. Madson, Challenges in implementing a screening method for veterinary drugs in  
753 milk using liquid chromatography quadrupole time-of-flight mass spectrometry, *J. Agric.*  
754 *Food Chem.* 62 (2014) 3660-3674.

755 [54] C. Planche, J. Ratel, F. Mercier, P. Blinet, L. Debrauwer, E. Engel, Assessment of  
756 comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry  
757 based methods for investigating 206 dioxin-like micropollutants in animal-derived food  
758 matrices, *J. Chromatogr. A* 1392 (2015) 74-81.

759 [55] Z. He, L. Wang, Y. Peng, M. Luo, W. Wang, X. Liu, Multiresidue analysis of over  
760 200 pesticides in cereals using a QuEChERS and gas chromatography-tandem mass  
761 spectrometry-based method, *Food Chem.* 169 (2015) 372-380.

762 [56] M.A. Zhao, Y.N. Feng, Y.Z. Zhu, J.H. Kim, Multi-residue method for determination  
763 of 238 pesticides in chinese cabbage and cucumber by liquid chromatography-tandem  
764 mass spectrometry: comparison of different purification procedures, *J. Agric. Food*  
765 *Chem.* 62 (2014) 11449-11456.

766 [57] B.D. Morris, R.B. Schriener, Development of an automated column solid-phase  
767 extraction cleanup of QuEChERS extracts, using a zirconia-based sorbent, for pesticide  
768 residue analyses by LC-MS/MS, *J. Agric. Food Chem.* 63 (2015) 5107-5119.

769 [58] Z. Wang, Q. Chang, J. Kang, Y. Cao, N. Ge, C. Fan, G.F. Pang, Screening and  
770 identification strategy for 317 pesticides in fruits and vegetables by liquid  
771 chromatography-quadrupole time-of-flight high resolution mass spectrometry, *Anal.*  
772 *Methods* 7 (2015) 6385-6402.

773 [59] J. Wang, W. Chow, J. Chang, J.W. Wong, Ultrahigh-performance liquid  
774 chromatography electrospray ionization Q-Orbitrap mass spectrometry for the analysis of  
775 451 pesticide residues in fruits and vegetables: Method development and validation, *J.*  
776 *Agric. Food Chem.* 62 (2014) 10375-10391.

777 [60] V. Amelin, A. Korotov, A. Andoralov, Identification and determination of 492  
778 contaminants of different classes in food and feed by high-resolution mass spectrometry  
779 using the standard addition method, *J. AOAC Int.* 99 (2016) 1600-1609.

780 [61] P. Pérez-Ortega, F.J. Lara-Ortega, J.F. García-Reyes, B. Gilbert-López, M.  
781 Trojanowicz, A. Molina-Díaz, A feasibility study of UHPLC-HRMS accurate-mass  
782 screening methods for multiclass testing of organic contaminants in food, *Talanta* 160  
783 (2016) 704-712.

784 [62] P. Donato, F. Rigano, F. Cacciola, M. Schure, S. Farnetti, M. Russo, P. Dugo, L.  
785 Mondello, Comprehensive two-dimensional liquid chromatography-tandem mass  
786 spectrometry for the simultaneous determination of wine polyphenols and target  
787 contaminants, *J. Chromatogr. A* 1458 (2016) 54-62.

788 [63] M.F. Mirabelli, J.C. Wolf, R. Zenobi, Pesticide analysis at ppt concentration levels:  
789 coupling nano-liquid chromatography with dielectric barrier discharge ionization-mass  
790 spectrometry, *Anal. Bioanal. Chem.* 408 (2016) 3425-3434.

791 [64] G. ten Dam, I.C. Pussente, G. Scholl, G. Eppe, A. Schaechtele, S. van Leeuwen, The  
792 performance of atmospheric pressure gas chromatography-tandem mass spectrometry

793 compared to gas chromatography–high resolution mass spectrometry for the analysis of  
794 polychlorinated dioxins and polychlorinated biphenyls in food and feed samples, J.  
795 Chromatogr. A 1477 (2016) 76-90

796 [65] A.M. Knolhoff, T.R. Croley, Non-targeted screening approaches for contaminants  
797 and adulterants in food using liquid chromatography hyphenated to high resolution mass  
798 spectrometry, J. Chromatogr. A 1428 (2016) 86-96

799 [66] Y. Man, G. Liang, L. Pan, Analytical methods for the determination of *Alternaria*  
800 mycotoxins, Chromatographia 80 (2017) 9-22.

801 [67] N.W. Turner, H. Bramhmbhatt, M. Szabo-Vezse, A. Poma, R. Coker, S.A. Piletsky,  
802 Analytical methods for determination of mycotoxins: an update (2009-2014), Anal.  
803 Chim. Acta 901 (2015) 12-33.

804 [68] B. Skribic, J. Zivancev, M. Godula, Multimycotoxin analysis of crude extracts of  
805 nuts with ultra-high performance liquid chromatography/tandem mass spectrometry, J.  
806 Food Compos. Anal. 34 (2014) 171-177.

807 [69] M. Ludovici, C. Ialongo, M. Reverberi, M. Beccaccioli, M. Scarpari, V. Scala,  
808 Quantitative profiling of oxylipins through comprehensive LC-MS/MS analysis of  
809 *Fusarium verticillioides* and maize kernels, Food Addit. Contam., Part A 31 (2014) 2026-  
810 2033.

811 [70] M. García-Altare, A. Casanova, V. Bane, J. Diogene, A. Furey, P. de la Iglesia,  
812 Confirmation of pinnatoxins and spiroxins in shellfish and passive samplers from  
813 Catalonia (Spain) by liquid chromatography coupled with triple quadrupole and high-  
814 resolution hybrid tandem mass spectrometry, Mar. Drugs 12 (2014) 3706-3732.

815 [71] Y. Rodriguez-Carrasco, J. Mañez, H. Berrada, C. Juan, Development and validation  
816 of a LC-ESI-MS/MS method for the determination of *Alternaria* toxins alternariol,  
817 alternariol methyl-ether and rentoxin in tomato and tomato-based products, Toxins 8  
818 (2016) 328.

819 [72] Y. Rodriguez-Carrasco, M. Fattore, S. Albrizio, H. Berrada, J. Mañez, Occurrence  
820 of *Fusarium* mycotoxins and their dietary intake through beer consumption by the  
821 European population, Food Chem. 178 (2015) 149-155.



822 [73] Z. Zhang, X. Hu, Q. Zhang, P. Li, Determination for multiple mycotoxins in  
823 agricultural products using HPLC–MS/MS via a multiple antibody immunoaffinity  
824 column, *J. Chromatogr. B* 2021 (2016) 145-152.

825 [74] M.J. Boundy, A.I. Selwwood, D.T. Harwood, P.S. McNabb, A.D. Turner,  
826 Development of a sensitive and selective liquid chromatography–mass spectrometry  
827 method for high throughput analysis of paralytic shellfish toxins using graphitised carbon  
828 solid phase extraction, *J. Chromatogr. A*, 1387 (2015) 1-12.

829 [75] Y. He, F. Mo, D. Chen, L. Xu, Y. Wu, F. Fu, Capillary electrophoresis inductively  
830 coupled plasma mass spectrometry combined with metal tag for ultrasensitively  
831 determining trace saxitoxin in seafood, *Electrophoresis* 38 (2017) 469-476.

832 [76] D.G. Beach, C.M. Walsh, P. McCarron, High-throughput quantitative analysis of  
833 domoic acid directly from mussel tissue using Laser Ablation Electrospray Ionization -  
834 tandem mass spectrometry, *Toxicon* 92 (2014) 75-80.

835 [77] S. Hickert, J. Gerding, E. Ncube, F. Hübner, B. Flett, B. Cramer, H.U. Humpf, A  
836 new approach using micro HPLC-MS/MS for multi-mycotoxin analysis in maize  
837 samples, *Mycotoxin Res.* 31 (2015) 109-115.

838 [78] Y. Wang, Y. Li, J. Yang, J. Ruan, C. Sun. Microbial volatile organic compounds and  
839 their application in microorganism identification in foodstuff, *TRAC Trends Anal. Chem.*  
840 78 (2016) 1-16.

841 [79] M. Miks-Krajnik, Y. J. Yoon, D. O. Okuku, H.G. Yuk, Volatile chemical spoilage  
842 indexes of raw Atlantic salmon (*Salmo salar*) stored under aerobic condition in relation  
843 to microbiological and sensory shelf lives, *Food Microbiol.* 53 (2016) 182-191.

844 [80] F.F. Parlapani, S.A. Haroutounian, G.J.E. Nychas, I.S. Boziaris, Microbiological  
845 spoilage and volatiles production of gutted European sea bass stored under air and  
846 commercial modified atmosphere package at 2 °C, *Food Microbiol.* 50 (2015) 44-53.

847 [81] F.F. Parlapani, A. Mallouchos, S.A. Haroutounian, I.S. Boziaris, Microbiological  
848 spoilage and investigation of volatile profile during storage of sea bream fillets under  
849 various conditions, *Int. J. Food Microbiol.* 189 (2014) 153-163.

850 [82] F.F. Parlapani, A. Mallouchos, S.A. Haroutounian, I.S. Boziaris, Volatile organic  
851 compounds of microbial and non-microbial origin produced on model fish substrate un-

852 inoculated and inoculated with gilt-head sea bream spoilage bacteria, LWT - Food Sci.  
853 Technol. 78 (2017) 54-62.

854 [83] A.A. Argyri, A. Mallouchos, E.Z. Panagou, G.J.E. Nychas, The dynamics of the  
855 HS/SPME–GC/MS as a tool to assess the spoilage of minced beef stored under different  
856 packaging and temperature conditions, Int. J. Food Microbiol. 193 (2015) 51-58.

857 [84] T.T. Nieminen, P. Dalgaard, J. Björkroth, Volatile organic compounds and  
858 *Photobacterium phosphoreum* associated with spoilage of modified-atmosphere-  
859 packaged raw pork, Int. J. Food Microbiol. 218 (2016) 86-95.

860 [85] A. Romano, V. Capozzi, G. Spano, F. Biasiolo, Proton transfer reaction–mass  
861 spectrometry: online and rapid determination of volatile organic compounds of microbial  
862 origin, Appl. Microbiol. Biotechnol. 99 (2015) 3787-3795.

863 [86] C. Franke, J. Beauchamp, Real-time detection of volatiles released during meat  
864 spoilage: a case study of modified atmosphere-packaged chicken breast fillets inoculated  
865 with *Br. thermosphacta*, Food Anal. Methods 10 (2017) 310-319.

866 [87] P. Silcock, M. Alothman, W. Zardin, S. Heenan, C. Siefarth, P.J. Bremer, J.  
867 Beauchamp, Microbially induced changes in the volatile constituents of fresh chilled  
868 pasteurised milk during storage, Food packaging and shelf life 2 (2014) 81-90.

869 [88] M. Alothman, K.A. Lusk, P. Silcock, P.J. Bremer, Comparing PTR-MS profile of  
870 milk inoculated with pure or mixed cultures of spoilage bacteria, Food Microbiol. 64  
871 (2017) 155-163.

872 [89] Z. Jandric, D. Roberts, M.N. Rathor, A. Abraham, M. Islam, A. Cannavan,  
873 Assessment of fruit juice authenticity using UPLC–QToF MS: A metabolomics approach,  
874 Food Chem. 148 (2014) 7-17.

875 [90] Z. Jandrić, M. Islam, D.K. Singh, A. Cannavan, Authentication of Indian citrus  
876 fruit/fruit juices by untargeted and targeted metabolomics, Food Control 72 (2017) 181-  
877 188.

878 [91] S. Jin, C. Song, S. Jia, S. Li, Y. Zhang, C. Chen, Y. Feng, Y. Xu, C. Xiong, Y. Xiang,  
879 H. Jiang, An integrated strategy for establishment of curcuminoid profile in turmeric  
880 using two LC–MS/MS platforms, J. Pharm. Biomed. Anal. 132 (2017) 93-102.

881 [92] L. Millán, M.C. Sampedro, A. Sánchez, C. Delporte, P.V. Antwerpen, M. A.  
882 Goicolea, R.J. Barrio, Liquid chromatography–quadrupole time of flight tandem mass  
883 spectrometry–based targeted metabolomic study for varietal discrimination of grapes  
884 according to plant sterols content, J. Chromatogr. A. 1454 (2016) 67-77.

885 [93] A. Cuadros-Inostroza, S. Ruíz-Lara, E. González, A. Eckardt, L. Willmitzer, H.  
886 Peña-Cortés, GC-MS metabolic profiling of cabernet sauvignon and merlot cultivars  
887 during grapevine berry development and network analysis reveals a stage- and cultivar-  
888 dependent connectivity of primary metabolites, *Metabolomics* 12 (2016) 39.

889 [94] M. Arbulu, M.C. Sampedro, A. Gómez-Caballero, M.A. Goicolea, R.J. Barrio,  
890 Untargeted metabolomic analysis using liquid chromatography quadrupole time-of-flight  
891 mass spectrometry for non-volatile profiling of wines, *Anal. Chim. Acta* 858 (2015) 32-  
892 41.

893 [95] B. Khakimov, R.J. Mongi, K.M. Sørensen, B.K. Ndabikunze, B.E. Chove, S.B.  
894 Engelsen, A comprehensive and comparative GC-MS metabolomics study of non-  
895 volatiles in Tanzanian grown mango, pineapple, jackfruit, baobab and tamarind fruits,  
896 *Food Chem.* 213 (2016) 691-699.

897 [96] E.J. Gu, D.W. Kim, G.J. Jang, S.H. Song, J.I. Lee, S.B. Lee, B.M. Kim, Y. Cho, H.J.  
898 Lee, H.J. Kim, Mass-based metabolomic analysis of soybean sprouts during germination,  
899 *Food Chem.* 217 (2017) 311-319.

900 [97] K. Inoue, C. Tanada, T. Sakamoto, H. Tsutsui, T. Akiba, J.Z. Min, K. Todoroki, Y.  
901 Yamano, T. Toyoka, Metabolomics approach of infant formula for the evaluation of  
902 contamination and degradation using hydrophilic interaction liquid chromatography  
903 coupled with mass spectrometry, *Food Chem.* 181 (2015) 318-324.

904 [98] G. Aliakbarzadeh, H. Sereshti, H. Parastar, Pattern recognition analysis of  
905 chromatographic fingerprints of *Crocus sativus* L. secondary metabolites towards source  
906 identification and quality control, *Anal. Bioanal. Chem.* 408 (2016) 3295-3307.

907 [99] J. Rubert, O. Lacina, M. Zachariasova, J. Hajslova, Saffron authentication based on  
908 liquid chromatography high resolution tandem mass spectrometry and multivariate data  
909 analysis, *Food Chem.* 204 (2016) 201-209.

910 [100] M. Guijarro-Díez, L. Nozal, M.L. Marina, A.L. Crego, Metabolomic fingerprinting  
911 of saffron by LC/MS: novel authenticity markers, *Anal. Bioanal. Chem.* 407 (2015) 7197-  
912 7213.

913 [101] C. Sales, M.I. Cervera, R. Gil, T. Portolés, E. Pitarch, J. Beltran, Quality  
914 classification of Spanish olive oils by untargeted gas chromatography coupled to hybrid  
915 quadrupole-time of flight mass spectrometry with atmospheric pressure chemical  
916 ionization and metabolomics-based statistical approach, *Food Chem.* 216 (2017) 365-  
917 373.

918 [102] F.R. Pinu, S. de Carvalho-Silva, A.P. Trovatti Uetanabaro, S.G. Villas-Boas,  
919 Vinegar metabolomics: An explorative study of commercial balsamic vinegars using gas  
920 chromatography-mass spectrometry, *Metabolites* 6 (2016) 22.

921 [103] P. Scano, A. Murgia, F.M. Pirisi, P. Caboni, A gas chromatography-mass  
922 spectrometry-based metabolomic approach for the characterization of goat milk  
923 compared with cow milk, *J. Dairy Sci.* 97 (2014) 6057-6066.

924 [104] T. Acunha, C. Simó, C. Ibáñez, A. Gallardo, A. Cifuentes, Anionic metabolite  
925 profiling by capillary electrophoresis-mass spectrometry using a noncovalent polymeric  
926 coating. Orange juice and wine as case studies, *J. Chromatogr. A*, 1428 (2016) 326-335.

927 [105] J. Welzenbach, C. Neuhoff, C. Looft, K. Schellander, E. Tholen, C. Große-  
928 Brinkhaus, Different statistical approaches to investigate porcine muscle metabolome  
929 profiles to highlight new biomarkers for pork quality assessment, *PLoS One*, 2 (2016) 11.

930 [106] A.K. Subbaraj, Y.H. Brad-Kim, K. Fraser, M.M. Farouk, A hydrophilic interaction  
931 liquid chromatography–mass spectrometry (HILIC-MS) based metabolomics study on  
932 colour stability of ovine meat, *Meat Science* 117 (2016) 163-172.

933 [107] D.K. Trivedi, K.A. Hollywood, N.J.W. Rattray, H. Ward, J. Greenwood, D.I. Ellis,  
934 R. Goodacre, Meat, the metabolites: an integrated metabolite profiling and lipidomics  
935 approach for the detection of the adulteration of beef with pork, *Analyst* 141 (2016) 2155-  
936 2164.

937 [108] V.D. Daygon, S. Prakash, M. Calingacion, A. Riedel, B. Oviden, P. Snell, J.  
938 Mitchell, M. Fitzgerald, Understanding the jasmine phenotype of rice through metabolite  
939 profiling and sensory evaluation, *Metabolomics* 12 (2016) 63.

940 [109] G. Min-Lee, D. Ho-Suh, E. Sung-Jung, C. Hwan-Lee, Metabolomics provides  
941 quality characterization of commercial gochujang (fermented pepper paste), *Molecules*  
942 21 (2016) 921.

943 [110] D.E. Lee, G.R. Shin, S. Lee, E.S. Jang, H.W. Shin, B.S. Moon, C.H. Lee,  
944 Metabolomics reveal that amino acids are the main contributors to antioxidant activity in  
945 wheat and rice gochujangs (Korean fermented red pepper paste), *Food Res. Int.* 87 (2016)  
946 10-17.

947 [111] S. Jin, C. Song, S. Jia, S. Li, Y. Zhang, C. Chen, Y. Feng, Y. Xu, C. Xiong, Y.  
948 Xiang, H. Jiang, An integrated strategy for establishment of curcuminoid profile in  
949 turmeric using two LC–MS/MS platforms, *J. Pharm. Biomed. Anal.* 132 (2017) 93–102.

950 [112] L.L. Monti, C.A. Bustamante, S. Osorio, J. Gabilondo, J. Borsani, M.A. Lauxmann,  
951 E. Mauli3n, G. Valentini, C.O. Budde, A.R. Fernie, M.V. Lara, M.F. Drincovich,

952 Metabolic profiling of a range of peach fruit varieties reveals high metabolic diversity  
953 and commonalities and differences during ripening, *Food Chem.* 190 (2016) 879-888.

954 [113] A. Kårlund, U. Moor, G. McDougall, M. Lehtonen, R.O. Karjalainen, K.  
955 Hanhineva, Metabolic profiling discriminates between strawberry (*Fragaria × ananassa*  
956 Duch.) cultivars grown in Finland or Estonia, *Food Res. Int.* 89 (2016) 647-653.

957 [114] I. Akhatou, R. González-Domínguez, A. Fernández-Recamales, Investigation of the  
958 effect of genotype and agronomic conditions on metabolomic profiles of selected  
959 strawberry cultivars with different sensitivity to environmental stress, *Plant Physiol*  
960 *Biochem.* 101 (2016) 14-22.

961 [115] M.N.A. Khalil, M.I. Fekry, M.A. Farag, Metabolome based volatiles profiling in  
962 13 date palm fruit varieties from Egypt via SPME GC-MS and chemometrics, *Food*  
963 *Chem.* 217 (2017) 171-181.

964 [116] Z. Jandrić, S.A. Haughey, R.D. Frew, K. McComb, P. Galvin-King, C.T. Elliott, A.  
965 Cannavan, Discrimination of honey of different floral origins by a combination of various  
966 chemical parameters, *Food Chem.* 189 (2015) 52-59.

967 [117] R. Gil-Solsona, M. Raro, C. Sales, L. Lacalle, R. Díaz, M. Ibañez, J. Beltran, J.V.  
968 Sancho, F.J. Hernández, Metabolomic approach for extra virgin olive oil origin  
969 discrimination making use of ultra-high performance liquid chromatography-Quadrupole  
970 time-of-flight mass spectrometry, *Food Control* 70 (2016) 350-359.

971 [118] R. Díaz, O.J. Pozo, J.V. Sancho, F. Hernández, Metabolomic approaches for orange  
972 origin discrimination by ultra-high performance liquid chromatography coupled to  
973 quadrupole time-of-flight mass spectrometry, *Food Chem.* 157 (2014) 84-93.

974 [119] S. Klockmann, E. Reiner, R. Bachmann, T. Hackl, M. Fischer, Food fingerprinting:  
975 Metabolomic approaches for geographical origin discrimination of hazelnuts (*Corylus*  
976 *avellana*) by UHPLC-QTOF- MS, *J. Agric. Food Chem.* 64 (2016) 9253-9262.

977 [120] K. Hori, T. Kiriya, K. Tsumura, A liquid chromatography time-of-flight mass  
978 spectrometry-based metabolomics approach for the discrimination of cocoa beans from  
979 different growing regions, *Food Anal. Methods* 9 (2016) 738-743.

980 [121] L. Lucini, G. Rocchetti, D. Kane, M. Trevisan, Phenolic fingerprint allows  
981 discriminating processed tomato products and tracing different processing sites, *Food*  
982 *Control* 73 (2017) 696-703.

983 [122] P. Arapitsas, A. Della Corte, H. Gika, L. Narduzzi, F. Mattivi, G. Theodoridis,  
984 Studying the effect of storage conditions on the metabolite content of red wine using  
985 HILIC LC-MS based metabolomics, *Food Chem.* 197 (2016) 1331-1340.

986 [123] C.J. García, R. García-Villalba, Y. Garrido, M.I. Gil, F.A. Tomás-Barberán,  
987 Untargeted metabolomics approach using UPLC-ESI-QTOF-MS to explore the  
988 metabolome of fresh-cut iceberg lettuce, *Metabolomics* 12 (2016) 138.

989 [124] K. Hennig, R.C.H. de Vos, C. Maliepaard, M. Dekker, R. Verkerk, G. Bonnema, A  
990 metabolomics approach to identify factors influencing glucosinolate thermal degradation  
991 rates in Brassica vegetables, *Food Chem.* 155 (2014) 287-297.

992 [125] A.L. Mayorga-Gross, L.M. Quirós-Guerrero, G. Fourny, F. Vaillant, An untargeted  
993 metabolomic assessment of cocoa beans during fermentation, *Food Res. Int.* 89 (2016)  
994 901-909.

995 [126] C. Boucher, F. Courant, A.L. Royer, S. Jeanson, S. Lortal, G. Dervilly-Pinel, A.  
996 Thierry, B. Le Bizec, LC-HRMS fingerprinting as an efficient approach to highlight fine  
997 differences in cheese metabolome during ripening, *Metabolomics* 11 (2015) 1117-1130.

998 [127] F. Liu, T. Grauwet, B.T. Kebede, A.V. Loey, X. Liao, M. Hendrickx, Comparing  
999 the effects of high hydrostatic pressure and thermal processing on blanched and  
1000 unblanched mango (*Mangifera indica* L.) nectar: using headspace fingerprinting as an  
1001 untargeted approach, *Food Bioprocess Technol.* 7 (2014) 3000-3011.

1002 [128] R. Beleggia, V. Menga, C. Platani, F. Nigro, M. Fragasso, C. Fares, Metabolomic  
1003 analysis can detect the composition of pasta enriched with fibre after cooking, *J. Sci. Food*  
1004 *Agric.* 96 (2016) 3032-3041.

1005 [129] M. Sugimoto, S. Obiya, M. Kaneko, A. Enomoto, M. Honma, M. Wakayama, T.  
1006 Soga, M. Tomita, Metabolomic profiling as a possible reverse engineering tool for  
1007 estimating processing conditions of dry-cured hams, *J. Agric. Food Chem.* 65 (2017) 402-  
1008 410.

1009 [130] C. Bündig, C. Blume, C. Peterhänsel, T. Winkelmann, Changed composition of  
1010 metabolites in *Solanum tuberosum* subjected to osmotic stress in vitro: Is sorbitol taken  
1011 up?, *Plant Cell Tiss. Organ. Cult.* 127 (2016) 195-206.

1012 [131] E. Hurtado-Fernández, J.J. González-Fernández, J.I. Hormaza, A. Bajoub, A.  
1013 Fernández-Gutiérrez, A. Carrasco-Pancorbo, Targeted LC-MS approach to study the  
1014 evolution over the harvesting season of six important metabolites in fruits from different  
1015 avocado cultivars, *Food Anal. Methods* 9 (2016) 3479-3491.

1016

1017 **FIGURE CAPTIONS.**

1018 **Figure 1.** GC×GC-TOF/MS contour plot of the 209 PCBs and 17 PCDD/Fs with the Rtx-  
1019 Dioxin2/BP-X50 column set. Adapted with permission from [54].

1020 **Figure 2.** Workflow for the identification of compounds in fractions from pizza  
1021 packaging material analyzed by GC-EI-qTOF MS and UHPLC-ESI-qTOF MS.  
1022 Reproduced with permission from [26].

1023 **Figure 3.** Schematic set-ups for continuous on-line monitoring of microbial volatile  
1024 organic compounds by proton-transfer-reaction MS in A) the headspace of four meat  
1025 samples in parallel (adapted with permission from [86]) and, B) in the headspace of milk  
1026 samples (adapted with permission from [87]).

1027 **Figure 4.** Workflow for establishment of curcuminoid profile in turmeric by an integrated  
1028 strategy. Reproduced with permission from [101].

1029 **Figure 5.** Scheme of the primary metabolism pathways of metabolites in Cabernet  
1030 Sauvignon (CS) and Merlot (ME) cultivars during different grape development stage.  
1031 Pathways are simplified version of tricarboxylic acid cycle, glycolysis, amino acid and  
1032 sucrose synthesis. FLW, flowering; FS, fruit setting; PRV, pre-veraison; VR, veraison;  
1033 PSV, post-veraison; RP, ripening. Metabolite intensity is color coded. Reproduced with  
1034 permission from [93].

1035 **Figure 6.** Data analysis workflow to explore the metabolome of lettuce in order to  
1036 evaluate changes related to storage time and genetics. FB fast-browning cultivar, SB  
1037 slow-browning cultivar, d0 day 0, d5 day 5. Reproduced with permission from [123].

1038

1039 **Table 1.** Selected remarkable applications published during the period 2014-2017 dealing with the simultaneous identification and quantification  
 1040 of a large number of contaminants (> 50) in food samples.

Contaminants quantified	Food matrix	Sample preparation	MS-based approach	MS-based technique	Sensitivity		Reference
					LOD	LOQ	
Pesticides (54)	Fruits and fish	QuEChERS	Targeted	UHPLC-HRMS (Orbitrap)	< 2 ng mL <sup>-1</sup>		19
Pesticides (54)	Tomatoes, oranges	QuEChERS	Non-targeted	GC-EI-HRMS (Orbitrap)		10 µg kg <sup>-1</sup>	20
Pesticides (55)	Bivalves ( <i>Scrobicularia plana</i> )	QuEChERS	Targeted	GC-MS/MS (IT in SIM mode)		0.33-10.3 µg L <sup>-1</sup>	21
Pesticides (57)	Tomato	QuEChERS	Targeted	LC-MS/MS (QqQ in MRM mode)	< 5000 µg kg <sup>-1</sup>		22
Antibiotics (62)	Meat	Solvent-based extraction (ACN)	Targeted	LC-HRMS (Orbitrap)	1 µg kg <sup>-1</sup>	3.3 µg kg <sup>-1</sup>	23
Contaminants (68)	Food contact materials	QuEChERS (modified)	Targeted	LC-MS/MS (QqQ in MRM mode) GC-MS/MS (QqQ in MRM mode)		1.3 – 220 µg kg <sup>-1</sup>	24
Pesticides (73)	Fruits, vegetables	Solvent-based extraction (ACN)	Targeted	LC-MS/MS (QqQ in MRM mode)		< 10 µg kg <sup>-1</sup>	25
Contaminants (75)	Food contact materials	Soxhlet-based protocol	Non-targeted / Targeted	UPLC-HRMS (QTOF, database)	< 2 ng ml <sup>-1</sup>	< 20 ng ml <sup>-1</sup>	26
Herbicides (76) and veterinary drug residues	Shellfish	QuEChERS	Targeted	LC-MS/MS (QqQ in MRM mode) GC-MS/MS (QqQ in MRM mode)		0.25-0.50 µg kg <sup>-1</sup> veterinary residues 2-20 µg kg <sup>-1</sup> pesticides	27
Veterinary drug residues (76)	Meat	Solvent-based extraction (ACN)	Targeted	UHPLC-MS/MS (QqQ in SRM mode)		0.038- 74 µg kg <sup>-1</sup>	28
Pesticides and antibiotics (83)	Honey	Solvent-based extraction (ACN)	Targeted / Non-targeted	LC-HRMS (Orbitrap)	< MRLs		29
Pesticides (87)	Groundnut oil	QuEChERS	Targeted	LC-MS/MS		4 - 180 µg kg <sup>-1</sup>	30



Pesticides (79) and antibiotics (13)	Honey	Solvent-based extraction (ACN) and clean-up	Targeted	(QqQ MRM mode) UHPLC-MS/MS (QqQ in MRM mode)	0.03 to 1.51 $\mu\text{g kg}^{-1}$	0.1 to 5 $\mu\text{g kg}^{-1}$	31
Pesticides (103)	Chicken, fish	QuEChERS	Targeted	LC-MS/MS (QqQ in dynamic MRM mode)		1-10 $\mu\text{g kg}^{-1}$	32
Pesticides (109)	Tomatoes	QuEChERS	Targeted	LC-MS/MS (QqQ in MRM mode)	0.5-10.8 $\mu\text{g kg}^{-1}$	1.3-30.4 $\mu\text{g kg}^{-1}$	33
Pesticides (113)	Rice, red pepper, mandarin	QuEChERS	Targeted	GC-MS/MS (QqQ in MRM mode)		0.1–25 $\mu\text{g kg}^{-1}$	34
Pesticides (115)	Oranges	QuEChERS	Targeted	LC-MS/MS (QqQ in MRM mode)	1 – 11 $\mu\text{g kg}^{-1}$	2 – 30 $\mu\text{g kg}^{-1}$	35
Pesticides (65) and environmental contaminants (52)	Kale, salmon, pork, avocado	QuEChERS	Targeted	GC-MS/MS (QqQ in MRM mode)			36
Pesticides (120)	Fruits, cereals	QuEChERS	Targeted	GC-MS/MS (QqQ in MRM mode)	10 $\mu\text{g kg}^{-1}$		37
Pesticides (120)	Apples, cucumbers	QuEChERS	Targeted	LC-MS/MS (QqQ in SRM mode)	1.2 – 11 $\mu\text{g kg}^{-1}$	10 $\mu\text{g kg}^{-1}$	38
Veterinary drugs (120)	Meat, eggs, milk	Ultrasound-assisted extraction and SPE	Targeted	LC-MS/MS (QqQ in MRM mode)	0.5–3.0 $\mu\text{g kg}^{-1}$	1.5–10.0 $\mu\text{g kg}^{-1}$	39
Contaminants (120)	Eggs	Solvent-based extraction (ACN) and purification	Targeted	LC-MS/MS (QqQ in MRM mode)		2.04–1316 $\mu\text{g kg}^{-1}$ (CC $\beta$ )	40
PCBs (127), polychlorinated naphthalenes (6), PAHs (16)	Mussels, clams	PLE (100°C, dichloromethane:hexane)	Targeted	GC-MS (quadrupole, SIM)		0.2-15 pg	41
Pesticides (105), antibiotics (49) and steroids (3)	Honey	Solvent-based extraction (ACN)	Targeted	UHPLC-HRMS (Orbitrap, in PRM mode and database)		0.009 - 6.21 $\mu\text{g kg}^{-1}$ (CC $\beta$ )	42
Pesticides (133), PAHs (24)	Fish	QuEChERS	Targeted	GC-HRMS (QTOF)	10 $\mu\text{g kg}^{-1}$		43
Pesticides (162)	Tea	Solvent-based extraction (ACN) and purification	Targeted	GC-MS/MS (QqQ in MRM mode)	< 10 $\mu\text{g kg}^{-1}$		44

Pesticides (164)	Apples, broccoli, oranges	Polyurethane foam disks swabbing	Targeted	DART-HRMS (Orbitrap)	10 µg kg <sup>-1</sup>		45
Pesticides (167)	Honey	Solvent-based extraction (ethyl acetate)	Targeted	LC-MS/MS (QqQ in MRM mode)		10 – 100 µg kg <sup>-1</sup>	46
Pesticides (172)	Wines	Solvent-based extraction (ethyl acetate)	Targeted	LC-MS/MS (QqQ in MRM mode)		10 – 50 µg kg <sup>-1</sup>	47
Pesticides (177)	Soy-based products	QuEChERS	Targeted	GC-MS/MS (QqQ in MRM mode)	0.1 - 10 µg kg <sup>-1</sup>	0.5-20 µg kg <sup>-1</sup>	48
Pesticides (178)	Eggs	Matrix solid-phase dispersion	Targeted	LC-MS/MS (QqQ in MRM mode) GC-MS/MS (QqQ in MRM mode)		5 – 10 µg kg <sup>-1</sup>	49
Pesticides (184)	Lettuce, oranges	QuEChERS	Non-targeted	LC-HRMS (Orbitrap)	10 µg kg <sup>-1</sup> (SDL) for 134 compounds 50-200 µg kg <sup>-1</sup> (SDL) for 39 compounds		50
Pesticides (200)	Green lettuce, orange	Ultra-Turrax homogenization with methanol and dilution	Targeted	UHPLC-MS/MS (QqQ in MRM mode)		1.0 – 5.0 µg kg <sup>-1</sup>	51
Pesticides (200)	Honey	QuEChERS	Targeted	GC-MS/MS (QqQ in MRM mode)	1.00 to 3.00 ng mL <sup>-1</sup>		52
Veterinary drug residues (>200)	Milk	Solvent-based extraction (ACN)	Targeted	LC-HRMS (QTOF)	< 100 ng mL <sup>-1</sup> (for 72% of compounds)		53
Dioxin-like micropollutants (206)	Meat	PLE (100 °C, hexane)	Targeted	GC×GC-TOF/MS	0.050-0.100 µg kg <sup>-1</sup> PCBs 65-227 ng kg <sup>-1</sup> PCDD/Fs		54
Pesticides (219)	Cereals	QuEChERS	Targeted	GC-MS/MS (QqQ in MRM mode)		5 - 50 µg kg <sup>-1</sup>	55
Pesticides (238)	Cabbage, cucumber	QuEChERS	Targeted	LC-MS/MS (QqQ in MRM mode)	0.02 - 6.32 µg kg <sup>-1</sup>	0.06 – 21.06 µg kg <sup>-1</sup>	56

Pesticides (269)	Avocado, citrus	QuEChERS with automated zirconia-based SPE	Targeted	LC-MS/MS (QqQ MRM mode)	< MRLs		57
Pesticides (317)	Vegetables, fruits	SPE	Targeted	LC-HRMS (QTOF and database)	10 µg kg <sup>-1</sup> (84 %)		58
Pesticides (451)	Fruits, vegetables	QuEChERS	Non-targeted	LC-HRMS (Orbitrap)		< 5 µg kg <sup>-1</sup> (85% of compounds)	59
Contaminants (492)	Milk, meat, eggs, liver, kidney, fish	Solvent-based extraction	Targeted	HPLC-HRMS (TOF-MS)	0.0005–100 ng mL <sup>-1</sup>	0.003–250 ng mL <sup>-1</sup>	60
Multiclass contaminants (625)	Baby foods, oranges, tomato	Solvent-based extraction (ACN)	Targeted	UHPLC-HRMS (QTOF and database)		< MRLs (excepting ca. 10% analytes)	61

1041 ACN, acetonitrile; CCβ, detection capability; DART, direct analysis in real time; HRMS, high resolution mass spectrometry; IT, ion trap; MRL, maximum residue limit; MRM, multiple reaction monitoring; PAH, polycyclic aromatic hydrocarbons; PCBs, polychlorinated biphenyls; PCDD, polychlorinated dibenzo-p-dioxins; PLE, pressurized liquid extraction; PRM, parallel reaction monitoring; QqQ, triple quadrupole; SDL, screening detection limit; SIM, selected ion monitoring; SPE, solid phase extraction; SRM, selected reaction monitoring; TOF: time-of-flight.

**Table 2.** The most remarkable MS-based metabolomics approaches devoted to food quality published during the period 2014-2017.

Food matrix	Metabolites	Sample preparation	MS-based approach	MS-based technique	Application	References
Pineapple, orange, apple, clementine, pomelo, and grapefruit juices	Flavonoids and limonoid glucosides	Centrifugation and filtering	Non-targeted / Targeted	UHPLC-HRMS (QTOF)	Detection of fruit juice adulteration	89
Citrus fruits, Jaffa, Mosambi orange and Red blush grapefruit	Flavonoids and limonoid glucosides	Centrifugation and filtering	Non-targeted / Targeted	UHPLC-HRMS (QTOF) for non-targeted LC-MS/MS (QqQ in MRM mode) for targeted	Discrimination of authentic and adulterated citrus fruits/fruit juices	90
Tumeric	Curcuminoids	Solvent-based extraction (using mixtures methanol:water)	Non-targeted/ Targeted	LC-HRMS (QTOF) for non-targeted LC-QTRAP-MS/MS (MRM mode) for targeted	Quality evaluation of raw turmeric from different regions	91
Grapes	Phytosterols	Solvent-based extraction (chloroform: methanol 1:1 (v/v))	Targeted	LC-HRMS (QTOF)	Discrimination of grapes according to plant sterols content	92
	Amino acids, fatty acids, acids (aromatic acids, hydroxy acids, dicarboxylic acids, phenylpropanoic acids), flavonoid, and sugars	Solvent-based extraction (water: methanol:chloroform (1:2.5:1, (v/v/v))	Non-targeted	GC-HRMS (TOF)	Differentiation of cultivars through their metabolite profile	93
Graciano <i>Vitis vinifera</i> wine	Non-volatiles/ semivolatile metabolites (sugars, amino acids, higher alcohol, biogenic amines, organic acids and phenolic compounds)	Centrifugation and filtering	Non-targeted	LC-HRMS (QTOF)	Analysis of the metabolome of the Graciano <i>Vitis vinifera</i> wine variety	94

Tropical fruits (Mango, pineapple, jackfruit, baobab, tamarind)	Non-volatiles metabolites (carbohydrates, organic acids, amino acids, and fatty acids).	Solvent-based extraction (water), acid hydrolysis and derivatization with trimethylsilyl cyanide	Non-Targeted	GC-HRMS (TOF)	Comparison of non-volatile metabolites of tropical fruits	95
Soybean sprouts	Amino acids, organic acids, lipids, sugars, phytosterol, isoflavones, and soyasaponins.	Solvent-based extraction (50 % methanol for UHPLC; 50 % methanol followed by methoxylation, and derivatization with BSTFA for GC analysis)	Non-targeted	GC-MS/MS (QqQ in MRM mode), and UHPLC-HRMS (QTOF)	Evaluation of the relationship between germination and nutritional quality	96
Infant formulas	Low-molecular-weight compounds (nicotinic acid and nicotinamide were identified)	Solvent-based extraction (water) and ultrafiltration	Non-targeted	HILIC-HRMS (QTOF)	Assessment of contamination and degradation of infant formulas	97
Saffron	Volatile metabolites	UASE-DLLME	Non-targeted	GC-MS (Q with EI)	Investigation of the effect of volatile components on the saffron's classification	98
	Glycerophospholipids and their oxidized lipids	Solvent-based extraction (ethanol:water 70:30 v/v) with sonication	Non-targeted	UHPLC-HRMS (QTOF)	Authentication of saffron	99
	Mainly flavonols and anthocyanins	Solvent-based extraction (ethanol:borate buffer at pH 9.0, 50:50 v/v) with sonication	Non-targeted	LC-HRMS (QTOF)	Investigation of the quality and authenticity of saffron	100
Olive oil	Volatile organic compounds	SPE	Non-targeted	GC-HRMS (QTOF)	Classification of olive oils according to their quality	101
Vinegar	Amino acids, carboxylic acids, sugars, sugar alcohols, fatty acids, vitamin, peptides and aroma compounds	MCF derivatization/TMS derivatization/ or extraction with diethyl ether	Non-targeted	GC-MS (Q with EI)	Comprehensive metabolite profile of vinegar	102
Milk	Short-chain hydroxylated carboxylic acids, long-chain stearic and palmitic acids, free amino acids, and sugars	Solvent-based extraction (methanol:chloroform) and derivatization with pyridine	Non-targeted	GC-MS (Q with EI)	Discrimination between milk typologies and detection of milk fraud	103

Orange juice and red wine	Mainly sugars, amino acids, and organic acids	Filtering	Non-targeted	CE-HRMS (TOF)	Comprehensive anionic metabolite profile of orange juice and red wine	104
Meat	Organic acids, amino acids, sugars, sugar alcohols, phosphorylated intermediates and lipophilic compounds	Solvent-based extraction (methanol:water 80:20 (v/v)) Derivatization with MSTFA for GC analysis	Non-targeted	GC-HRMS (TOF)/HILIC-HRMS (QTOF)	Identify biomarkers of meat quality traits	105
	Amino acids, sugars, nucleosides, nucleosides, and organic acids	Solvent-based extraction (methanol:water 80:20 (v/v)) followed by chloroform:water 67:33 (v/v))	Non-targeted	HILIC-HRMS (Orbitrap)	Study of colour stability of ovine meat	106
Rice (Jasmine phenotype)	Amino acids, organic acids, alkane hydrocarbon, and sugar alcohols,	Solvent-based extraction (chloroform:methanol:water) and derivatization with MSTFA	Non-targeted	GC-HRMS (TOF)	Detection of the adulteration of beef meat	107
	Volatile organic compounds	Static HS extraction	Non-targeted	GC×GC-TOF/MS	Determination of the metabolites that define the 'Jasmine' quality of rice	108
Gochujang (fermented pepper paste)	Amino acids, organic acids, fatty acids, sugars, sugar alcohols, flavonoids, capsaicinoids, capsinoids, lipids	Solvent-based extraction (80 % methanol) Derivatization with MSTFA for GC analysis	Non-targeted	GC-HRMS (TOF)/UHPLC-IT-MS	Quality characterization	109
	Mainly amino acids, organic acids, and sugars	Solvent-based extraction (80 % methanol) Derivatization with MSTFA for GC analysis	Non-targeted	GC-HRMS (TOF)/UHPLC-HRMS (QTOF)	Evaluation of the metabolite differences according to the raw material used in the production of gochujangs	110
Green tea	Mainly catechins, amino acids, caffeine	Solvent-based extraction (hot water)	Non-targeted	UHPLC-HRMS (QTOF)	Study of the chemical composition of green tea to assess its quality	111
Peach fruit	Sugars, organic acids, and amino acids	Solvent-based extraction (methanol) and derivatization with MSTFA	Non-targeted	GC-MS (Q with EI)	Explore the chemical composition which defines fruit quality	112

Strawberry	Phenolic acids, flavonoids, flavan-3-ol derivatives, terpenes, and many types of glycosidically bound aroma and flavour precursors	Solvent-based extraction (80 % methanol)	Non-targeted	LC-HRMS (QTOF)	Separation and identification of major metabolites showing significant variation between strawberry cultivars	113
	Sugars, organic acids, and amino acids	Solvent-based extraction (methanol:water 1:1 (v/v)) and derivatization with MSTFA	Non-targeted	GC-MS (IT)	Differentiation of strawberry cultivars and assessment of the influence of agronomic conditions	114
Date palm fruit	Volatile metabolites (lipid-derived volatiles, phenylpropanoid derivatives, amino acid derived volatiles, and sugar derived volatiles)	HS-SPME	Non-targeted	GC-MS (Q with EI)	Differentiation among date varieties	115
Honey	Not described	Solvent-based extraction (methanol:water 1:1 (v/v) containing 1 % formic acid)	Non-targeted	UHPLC-HRMS (QTOF)	Discrimination of honeys according to their floral origin	116

1046

1047 BSTFA, bis(trimethylsilyl)trifluoroacetamide; DLLME, dispersive liquid-liquid microextraction; EI, electron ionization; HILIC, hydrophilic interaction liquid chromatography;  
 1048 HRMS, high resolution mass spectrometry; HS-SPME, headspace solid-phase micro-extraction; IT, ion trap, MCF, Methylchloroformate; MRM, multiple reaction monitoring;  
 1049 MSD, mass selective detector; MSTFA, N-Methyl-N-(trimethylsilyl) trifluoroacetamide; Q, quadrupole; QqQ, triple quadrupole; QTOF, quadrupole-time-of-flight; QTRAP,  
 1050 hybrid triple-quadrupole linear ion trap; SPE, solid-phase extraction; TMS, trimethyl Silyl; TOF, time-of-flight; UASE-DLLME, ultrasound-assisted solvent extraction in  
 1051 tandem with dispersive liquid-liquid microextraction.

1052

1053