Fine-scale determinants of vertebrate roadkills across a biodiversity hotspot in Southern Spain

David Canal^{1,2*}, Carlos Camacho³, Beatriz Martín⁴, Manuela de Lucas¹ and Miguel Ferrer¹

¹ Applied Ecology Group, Estación Biológica de Doñana (EBD-CSIC), Av. Américo Vespucio s/n, 41092 Seville, Spain.

² Centre for the Study and Conservation of Birds of Prey of Argentina (CECARA-UNLPam) & Institute of Earth and Environmental Sciences of La Pampa (INCITAP), National Scientific and Technical Research Council (CONICET), Av. Uruguay 151, 6300 Santa Rosa, La Pampa, Argentina.

³ Department of Evolutionary Ecology, Estación Biológica de Doñana (EBD-CSIC), c/Américo Vespucio s/n, 41092 Seville, Spain.

⁴ Fundación Migres, Complejo Huerta Grande, Crta. N 340 km 96.7, 11390 Pelayo, Algeciras, Spain.

*Corresponding author: davidcanal@ebd.csic.es

Running head: Fine-scale determinants of vertebrate roadkills

Abstract

An increasing number of wildlife-vehicle collisions occur each year worldwide, which involves extensive economic costs and constitutes one of the main anthropogenic causes of animal mortality. Because of this, there is an urgent need to identify the factors leading to collision hotspots and thus implementing effective mitigation measures. By using a stratified random sampling survey, we investigated the fine-scale determinants of roadkill probability in small and medium-sized birds and mammal across a country-size region of Southern Spain, Andalusia (87000 km²), located within a global biodiversity hotspot. During two consecutive seasons, we regularly surveyed 45 road transects of 10 km each and characterized the site-specific attributes of both roadkill and random points, including traffic density, road design (embankments, medians, fences, roadside vegetation and distance to curves), and adjacent landscape matrix. Based on this information, we investigated variation in collision risk according to landscape and road features, and the life history of the affected taxa. Mortality rates of mammals and birds increased with traffic density, and were also significantly affected by the distance to the nearest curve, slope of embankments, height of roadside vegetation, and land use adjacent to roads. Road mortality of both birds and mammals was related to the presence and typology of fences and center medians, so more densely vegetated medians and smaller mesh sizes reduced roadkill probability. Overall, our results indicate that roadkill risk may vary at exceedingly small spatial scales. The information provided by this extensive survey may be used to identify taxa-specific factors associated to roadkill risk and priority points for action. Our findings will therefore be relevant for the design of safer roads for both drivers and wildlife through the application of effective mitigation measures.

43 **Key words:** Wildlife vehicle collisions; road mortality; collision hotspots; roadkill
44 predictors; predictive models; mitigation measures.

45

46

Introduction

Roads have multiple ecological impacts, as they can act as barriers by limiting connectivity among populations, contaminate adjacent ecosystems, alter animal behavior, and facilitate dispersal of exotic species, among others (reviewed in (Sandberg et al. 1998; Trombulak and Frissell 2000; Forman et al. 2003; Coffin 2007; van der Ree et al. 2015b). However, traffic-related mortality due to wildlife-vehicles collisions (WVCs, hereafter) appear to be the most important ecological impact of roads (Trombulak and Frissell 2000; Coffin 2007).

WVCs are an important traffic safety issue that involves significant monetary costs, primarily due to human injury and material damage (Bissonette et al. 2008; Huijser et al. 2009), but also high environmental costs (Forman et al. 2003; Erritzøe et al. 2003; Bissonette et al. 2008). Traffic related mortality is considered one of the most important sources of non-natural mortality in wildlife populations (Forman et al. 2003; Erritzøe et al. 2003; Colino-Rabanal and Lizana 2012). For instance, considering only birds, 27 and 80 millions of fatalities are estimated to occur annually in Europe (Erritzøe et al. 2003) and the United States (Erickson et al. 2005), respectively, although actual numbers may be higher. Traffic related mortality may dramatically affect population dynamics (e.g. through differential incidence into a gender or age class; Madsen et al. 2002; Colino-Rabanal and Lizana 2012) and constitutes a major threat for endangered species (Mumme et al. 2000; Gibbs and Shriver 2002). Importantly, because of the expansion of the road network and the increase in traffic volume, the ecological impact of roads on wildlife is expected to increase over the next decades in both developed and developing countries (Fulton and Eads 2004; Meyer et al. 2012; van der Ree et al. 2015c). Thus, quantifying the impact of roads on wildlife and developing

effective mitigation measurements is urgently needed to balance future development requirements and biodiversity conservation (van der Ree et al. 2015c).

Road characteristics have long been recognized as a crucial determinant of roadkills. Factors such as traffic density and velocity, road sinuosity, and the presence of road crosses and elevation changes are frequently associated with collision risk (e.g. (Trombulak and Frissell 2000; Forman et al. 2003; Clevenger et al. 2003; Malo et al. 2004; Seiler 2005; Gomes et al. 2009; Langen et al. 2012; Zuberogoitia et al. 2014; D'Amico et al. 2015). Further, roadside strips of vegetation and land-use adjacent to roads may influence roadkill risk by determining the presence and movements of animals (e.g. Forman et al. 2003; Clevenger et al. 2003; Malo et al. 2004; Grilo et al. 2009; Gunson et al. 2011).

Mitigation measures commonly used to reduce animal roadkill include, among others, wildlife crossing structures (e.g. underpasses and overpasses), warning signs, animal detection systems and a variety of fences (reviewed in Glista et al. 2009; van der Grift et al. 2013). However, none of these measures has been fully effective in preventing WVCs, since their effectiveness strongly depends on the interplay between particular life history traits (e.g. foraging strategy, dispersal or migratory movements) of the species affected by roadkills and environmental factors influencing collision risk. Given that the implantation of mitigation measures along the entire road network is economically and logistically unfeasible, and that WVCs are typically clustered (Gunson et al. 2011), identifying the factors that increase the risk of collision is essential to implement effective mitigation measures (Gunson et al. 2011). Furthermore, recent calls have highlighted the need to conduct additional research that broadens the taxonomic, spatial, and temporal scale of roadkill data sets to optimize the implementation of the mitigation measures (van der Ree et al. 2015c).

In this study, we investigated the fine-scale determinants of roadkill probability in small and medium-sized birds and mammal at a large spatio-temporal scale. Over 22 months, we regularly surveyed 45 road sections of 10km each distributed across Andalusia (South Spain), an extensive Mediterranean region (87,268 Km²) located in a biodiversity hotspot (Myers et al. 2000). During the surveys, we characterized site-specific attributes at both WVC and control points (i.e. randomly-generated points without casualties recorded), using a landscape-level and road level approach (Table 1). This information was used to investigate the WVC risk in relation to the species' biology, adjacent land use, and road design based on the predictions presented in Table 1.

Material and methods

The study was conducted during two consecutive periods (July 2009-June 2010 -except September and October- and April 2011-March 2012) in the Autonomous Community of Andalusia, Spain, a region that stretches from the southeast to the southwest of the Iberian Peninsula (Fig 1). The ecosystems in Andalusia, characterized by an extraordinary diversity of species and landscapes, are considered to be highly sensitive to global-change drivers and are thus predicted to experience dramatic biodiversity changes in the next decades (Myers et al. 2000).

Briefly, the climate in Andalusia is Mediterranean, but due to a marked interannual variation in rainfall (it may varies from 170 mm/year to more than 1800 mm/year) and a wide elevation range (from sea level to approximately 3500 m.a.s.l.), there is a high diversity in vegetation and landscape conditions (including semiarid zones, forest, mountains or marshland). To capture such environmental diversity, the study region was divided into five ecoregions, defined as areas characterized by similar

landscape characteristics and environmental conditions (GIASA et al. 2006). For a detailed description of characteristics and environmental conditions on these ecoregions see Canal et al. (2018) and (GIASA et al. 2006).

Besides environmental conditions, the selection of the sampling roads aimed at representing the road network in Andalusia. Thus, according to the physical characteristics of the roads (number of lanes, speed limit or traffic density), we grouped the surveyed roads into three categories: 1) Highways characterized by a dual carriageway and 120 km/h speed limit; 2) National roads, including all roads belonging to the State Network, Regional and Interregional network except highways and 3) Local roads, all roads belonging to Complementary Regional Network and the Provincial Councils. Both national and local roads are characterized by a single carriageway and a 90 km/h speed limit.

Nine road sections (three replicates per road category) were randomly selected from the road network crossing each of the five ecoregions (3 replicates x 3 road categories x 5 ecoregions = 45 road sections). For each road, a random number was generated to set the starting point (kilometric point) of the sampling stretch. Overall, we monthly surveyed 450 km along 45 road segments crossing all ecoregions included in Andalusia (Fig. 1 and Table 1 in Supplementary material).

Monthly surveys were carried out by two experienced observers by driving a vehicle at low speed (~ 25-30 km/h) along the shoulder of the road with the emergency lights flashing. The sampling order of the surveyed sections was set at random from month to month and survey session. Roadkilled animals encountered on the paved road or the road verge were identified at the species level (whenever possible) and its location was recorded using a GPS. All carcasses were removed from roads to avoid duplicating records during posterior surveys. At each point, we recorded site-specific

attributes of roads and their immediate vicinity, including structures potentially influencing animal accessibility to roads (see Table 1 for a description of the measured variables and its predicted effect on WVC). Traffic density of the surveyed roads, defined as the average number of vehicles per day, was obtained from official data at http://www.fomento.es/MFOM/LANG_CASTELLANO/DIRECCIONES_GENERALES/CARRETERAS/TRAFFICO_VELOCIDADES/MAPAS/ and <http://www.juntadeandalucia.es/fomentoyvivienda/portal-web/web/areas/carreteras/aforos>. Twenty control points were randomly selected as outlined above (without previous knowledge of roadkill points) within each of the sampling sections (20 x 45 road sections = 900 control points) and characterized following the same procedure as for the collision points.

Statistical analyses

We used Generalized Linear Mixed Models (GLMM) to model the probability of WVC in relation to landscape and road features. Separate GLMMs were fitted for birds and mammals (see below) according to the noticeable differences between their life-history strategies (e.g. spatial ecology or locomotor capacity). Even though the classification of roadkill species at the class level might appear simplistic due to major species-specific differences in life history traits, such a classification may broaden the applicability of the mitigation measures derived from our survey. Further support for the use of a coarse-grained approach comes from the common implementation of similar mitigation measurements for different animal groups. For instance, similar measures are applied for ground birds and large terrestrial mammals, whereas the same applies for bats and flying birds (Abbott et al. 2015; Kociolek et al. 2015). Finally, an analysis of the probability of WVC at a lower taxonomic level (higher functional similarity) including

passerines, carnivores and lagomorphs, the groups most affected by roadkills, showed similar results to those found at the class level (see Table 3 in Supplementary material for details).

GLMMs analyzing the probability of WVC separately in birds and mammals included the presence (1) or absence (0, control points) of collision as the dependent variable (binomial distribution and logit link function) and seven explanatory variables as descriptors of each point: road type, distance to the nearest curve and its quadratic term, maximum height of roadside vegetation and its quadratic term, adjacent land use, and type of embankment. Road ID, nested within the ecoregion, was fit as a random factor.

Based on exploratory analyses, the multiple levels of road embankments was reduced to two classes: roads with embankments in any of the road sides instigating birds to fly high above the road (e.g., steep, buried sections) and those allowing animals to fly close to the road surface (e.g. roads sites at ground level). Similarly, for mammals, we reduced the type of road embankments to four classes: roads at ground level, raised, buried and roads with opposing types of embankment at each side (e. g. buried on the right side and raised on the left side).

Note that traffic density was not included in the two models above as the values for roadkill and control points within a given road section would be the same. Thus, to test if the accumulated number of roadkills in a road section was related to traffic density, we used a Pearson correlation.

Special considerations for fences and center medians

Because all highways in Spain are fenced, the effect of fences on vertebrate roadkills could only be investigated using data from secondary and local roads. The latter roads,

however, lack median centers and, thus, the influence of this structure on WVC was explored using exclusively data from highways.

Road points were categorized into four classes according to the presence and type of structures preventing the access of wildlife to roads (unfenced points, presence of barbed fences, wire mesh fences and walls; Table 1). The presence and type of barrier (fence/wall) was recorded at both roadsides and the difficulty of accessing roads was then determined according to the roadside having the less restrictive type of barrier. Exploratory analyses revealed no differences between mesh fences and walls in roadkill likelihood, and thus the difficulty of wildlife to access roads was assessed using a three-level variable: easy (unfenced point), medium (barber fenced) and high (mesh fences and walls) difficulty of access.

Median strips were initially categorized as Jersey barriers and structures with absent, medium or much vegetation (Table 1). However, frequency diagrams of the types of median strips revealed that the number of points with Jersey barriers as medians was very small. These points were therefore excluded from the models to avoid a disproportionate influence of rare categories on model outputs and, consequently, only medians varying in the amount of vegetation cover were analyzed.

The influence of barriers (fence/walls; except for highways) and medians (only in highways) on roadkill likelihood was analyzed separately for mammals and birds using GLMMs with the same structure as described above.

Model selection

In total, we fitted three GLMMs per vertebrate class analyzing the probability of roadkill in relation to i) the landscape and road attributes, ii) the type of barriers and ii) the amount of vegetation cover in the median strips.

Selection of the final models -i.e. containing only statistically significant terms- was carried out by sequentially dropping non-significant terms from fully saturated models (containing all main effects and interactions) in a hierarchical way, starting with the least significant order terms. To confirm whether the inclusion of a predictor was significantly informative, we compared the models including and excluding the focal term using chi-square likelihood ratio tests (through maximum likelihood estimations).

We systematically performed model diagnostics statistics to avoid misleading conclusions based on statistical artifacts. Accordingly, we visually checked assumptions about the distribution of residuals through diagnostics plots, and examined collinearity and the existence of influential data points. To meet statistical assumptions, the distances to the nearest curve and traffic intensity were log10-transformed. After these transformations, diagnostics analyses did not show obvious deviation from GLMM assumptions.

Our dataset was unbalanced since twenty control points were systematically recorded per road section, whereas the number of recorded roadkill varied among roads (Fig. 2). Even though the accuracy of binomial models is robust to unbalanced sampling (Crone and Finlay 2012), we repeated the analyses above after creating a balanced dataset (roadkill and control were points randomly selected) to check for consistency between the results based on balanced and unbalanced samples. Because the results obtained using the balanced and raw datasets were similar, we present along the paper the models using the whole dataset to make full use of the available data as suggested by Crone and Finlay (2012). During our surveys, we found a small fraction of domestic mammals (mostly dogs; see Canal et al. 2018) killed by vehicles. Results from the analyses excluding and including domestic animals were qualitatively similar. For this reason, the results of the analyses including the whole dataset are presented here.

Statistical analyses were carried out in R 3.1.2 (R Development Core Team 2015). For running the GLMMs, we used the packages lme4 (Bates et al. 2014), lmerTest (Kuznetsova et al. 2017) and Rcpp (Eddelbuettel and François 2011). For a part of the model diagnostics, we used the package DHARMA (Hartig 2016) and the VIF function from the package car (Fox & Weisberg, 2011).

Results

A total of 835 mammals and 555 birds belonging to 19 and 70 species, respectively, were recorded as killed by vehicles during the two study seasons (Table 1 and 2 in Supplementary material; Canal et al. 2018). 2.8% of roadkills could not be identified at the species level due to severe damage and/or poor conservation status.

Road mortality analyses revealed common factors associated with the occurrence of roadkills in birds and mammals (Table 2 and 3). In both groups, roadkills were related to traffic density (mammals: $R = 0.57$, $P < 0.001$; birds: $R = 0.39$, $P = 0.01$; Fig. 2), the distance to the nearest curve, and the height of the roadside vegetation. For the latter two factors, roadkill probability showed an inverted-U shape, increasing until a maximum distance and height, and decreasing afterwards (Table 2, 3 and Fig. 3). Also, in both groups, roadkill risk was affected by the adjacent land use type and the slope of road embankments. Roads crossing forests showed the highest probability of roadkill in birds (Fig. 4), whereas, in mammals, forests and farmlands were the habitats with highest mortality rates (Table 2, 3 and Fig. 4). The presence and type of road embankments also affected roadkill risk. In mammals, the probability of roadkill was lowest in elevated road sites, whereas roads with embankments boosting "high-altitude" flights (e.g. buried roads sections) reduced roadkill probability in birds (Table 2 and 3).

The presence and type of physical structures (fences and walls) preventing access of wildlife to roads reduced the roadkill likelihood in mammals and birds. In both taxa, the number of casualties decreased as the difficulty of accessing roads increased from unfenced points, followed by barbed fences and points having mesh wire fences or walls (mammals: estimate (SE) -0.57 (0.225), $Z = -2.53$, $P = 0.011$; birds: estimate (SE) -0.587 (0.185), $Z = -3.16$, $P = 0.002$; Fig. 5). Roadkill risk also decreased as vegetation cover in the median strips increased (mammals: estimate (SE) -0.330 (0.184), $Z = -1.79$, $P = 0.07$; birds: estimate (SE) -0.57 (0.17), $Z = -3.345$, $P < 0.001$; Fig. 5).

Discussion

Based on a large-scale survey and accurate description of the sampling sites, we have shown that roadkill risk in small and medium-sized birds and mammals may vary at exceedingly small spatial scales and that collision risk is group-specific. A fine-scale description of the road attributes at both roadkill and random points allowed us to unravel the road characteristics (e.g. steep embankments at roadsides and fences) determining the risk of WCV in birds and mammals. Other factors like the adjacent landscape matrix, the roadside vegetation, and vegetation density in center medians also contributed to determine roadkill probability.

Road related features

Traffic density is one of the most important predictors of roadkills (e.g. Clevenger et al. 2003; Seiler 2005; Barrientos and Bolonio 2009; Zuberogoitia et al. 2014; Gagné et al. 2015), although its influence on mortality is often non-linear; i.e. mortality peaks occur at intermediate levels of traffic density because animals are reluctant to cross highly

transited roads (Madsen et al. 2002; Seiler 2005; Zuberogitia et al. 2014). In our survey, the number of roadkills was associated with traffic density, but we did not found the expected reduction in mortality at high traffic density, even when we surveyed roads with enormous levels of traffic. At least two factors might explain the lack of a non-linear relationship between WVCs and traffic density. First, there might be a mismatch between the levels of traffic density and animal activity, since traffic density on the surveyed roads may be high only during the day, and many roadkilled species, especially mammals (see Table 2 in the Supplementary material), are most active during the night. Second, although often having a deterrent effect, traffic noise or lighting may also attract some bird species to roadsides increasing their mortality rates (Blackwell et al. 2015; Kociolek et al. 2015).

The influence of road topography in the WVC risk was in agreement with our predictions. For birds, collision risk decreased in road sections with steep buried or elevated roadside embankments as opposed to those at the ground level or with soft slopes. Possibly, flat roads enable birds to fly close to the ground while crossing, thereby increasing collision risk, whereas the reverse is likely true if steep embankments are present (Clevenger et al. 2003; Kociolek et al. 2015). Note, however, that the effect of topography (as well as that of road characteristics; see below) may be species-specific and/or conditional on other factors. For example, car lights may dazzle nocturnal birds and increase their susceptibility to WVC or predators (Blackwell et al. 2015; Kociolek et al. 2015). Further, scavengers (e.g. raptors) attracted to roads for foraging on roadkilled animals or species typically showing low-flight behaviors (e.g. owls; Massemin and Zorn 1998) may be particularly vulnerable to traffic mortality, and such susceptibility may in turn be increased or diminished by the type of road embankments. For mammals, raised road points and those at the ground level showed,

respectively, the highest and the lowest roadkill rates. These results suggest that roads with steep slopes at the roadside may discourage mammals from crossing (Alexander and Waters 2000; Clevenger et al. 2003; Malo et al. 2004; Gunson et al. 2011). Elevating roads may therefore be a good option to mitigate roadkills of small- and medium-sized mammals, especially when combined with other elements such as fences or crossing structures (Clevenger et al. 2003; Malo et al. 2004; Glista et al. 2009; Gunson et al. 2011).

The distance to the nearest curve, as determining the trade-off between improved visibility (reduced WVC risk) and increased velocity (increased WVC risk), was another important predictor of roadkill in mammals and birds. Given that vehicles must decelerate as approaching a curve, the quadratic effect of the distance to the curve on roadkill probability found in our study can be reasonably expected. Non-linear relationship between proximity to the nearest curve and roadkill risk has been previously reported in other studies (Table 1), although the distance with the highest risk of roadkill varies widely among them, possibly due to a number of additional factors (e.g. presence of dense roadside vegetation, type of road and focal species) influencing the likelihood of roadkill (Malo et al. 2004; Ramp et al. 2005; Zuberogoitia et al. 2006; Grilo et al. 2009; Gunson et al. 2011).

The presence of roadside barriers (fences with varying mesh sizes and walls) also shaped mortality risk in birds and mammal; so the presence of walls or fences with small mesh size that difficult the access to roads minimized roadkill risk. Our findings are in agreement with previous works suggesting that, overall, these mitigation measures may be effective in reducing roadkills (Gunson et al. 2011; van der Grift et al. 2013; van der Ree et al. 2015a), but at least two considerations should be taken into account. First, in the case of terrestrial vertebrates (mammals, amphibians and reptiles)

fences may act as barriers hampering wildlife (pre-breeding and/or dispersal) movements, thus reducing connectivity between populations (Trombulak and Frissell 2000; Forman et al. 2003; Coffin 2007). Second, as reiterated in the literature (Glista et al. 2009; van der Grift et al. 2013; D'Amico et al. 2015), the use of barriers as a mitigate measure to prevent wildlife access to roads should ideally be combined with other measures, such as underpasses and scape structures, to keep permeability between populations and thus avoid the fatal consequences of trap-effects (Colino-Rabanal et al. 2011; Cserksz et al. 2013; Zuberogoitia et al. 2014; van der Ree et al. 2015a).

The influence of median strips on roadkill risk has been scarcely assessed (Bellis and Graves 1971; Clevenger et al. 2003; Clevenger and Kociolek 2013), even when these structures may have a critical effect on WVC (reviewed in Clevenger & Kociolek, 2013). Medians are usually covered by dense vegetation that may provide relatively undisturbed breeding habitat, food resources (depending on the vegetation composition; Kociolek et al. 2015), and concealment from predators and can therefore attract many animals (Adams 1984; Clevenger and Kociolek 2013). Medians may thus increase roadkill risk by increasing wildlife presence and movements around roads (Bellis and Graves 1971; Clevenger et al. 2003; Clevenger and Kociolek 2013). By contrast, we have found that the roadkill rate of birds and mammals (except lagomorphs; see Supplementary material) decreased as the amount of vegetation cover in the medians increased. Several factors may explain these results. Perhaps, in our study area, the composition and/or structure of vegetation in the median strips are not suitable as a foraging or breeding site. Densely vegetated medians might also function as an obstacle for crossing birds, encouraging them to fly high (see above) and thus avoid potential collisions (Kociolek et al. 2015). In addition, the specific requirements of the species affected and the synergistic effect of medians and microhabitat attributes might explain

the apparent discrepancies between studies. For example, rabbits (the lagomorph most frequently found roadkilled during surveys) predominantly use the roadside vegetation and embankments as a refuge (Planillo and Malo 2013, 2018), which might explain the lack of relationship between vegetation cover in the medians and the probability of roadkill in this group. Regardless of the determinants of collision risk, our findings provide invaluable information about the effects of medians on WVC given the limited knowledge on this topic (Clevenger and Kociolek 2013). Further research (e.g. testing the impact of continuous and discontinuous strips of median cover on different vertebrate groups and/or their effect if combined with other crossing structures) is needed to better understand the effect of these linear developments on animal movement and mortality (Clevenger and Kociolek 2013).

Landscape features

Bird roadkills were more likely to occur on roads with adjacent wooded areas, perhaps because wooded sites offer lower visibility in relation to more open habitats, as scrubland and farmlands (Clevenger et al. 2003). Dense tree cover may at the same time increase bird abundance around roads, as they often use trees as foraging and nesting sites. Indeed, the abundance of a species in the road surroundings was likely a major determinant of their roadkill rate since, although no data on local bird abundances are available (see below), top ranked species recorded in our study are among the most common species in Andalusia (e.g. *Passer domesticus*, *Turdus merula*, *Sylvia atricapilla* or *Erithacus rubecula*). In mammals, the highest rates of fatalities occurred in forested areas, but also in points adjacent to farm areas. These findings are in agreement with other reports showing that mammal casualties increased in forested areas (Clevenger et al. 2003; Malo et al. 2004; Ramp et al. 2005; Seiler 2005; Grilo et

al. 2009; Gunson et al. 2011). Moreover, the influence of landscape on the roadkill risk may depend on species-specific habitat preferences (Gunson et al. 2011) and, in our survey, mammal mortality was dominated by wild rabbits and European hares (see Canal et al. 2018), typically associated to open and/or farm areas.

For birds and mammals, roadkill risk increased when the roadside vegetation was either very tall or very short. Short roadside vegetation -or lack thereof- may reduce WVC by increasing the reaction capacity of drivers and animals to dodge the collisions. On the contrary, by providing protection or food, medium-sized vegetation at the roadside such as small trees and shrubs may attract individuals to roads and, subsequently, influence the probability of roadkill (Barrientos and Bolonio 2009). Small trees and shrubs may also increase collision rates by, for example, favoring low-to-ground-level flights while crossing roads (Clevenger et al. 2003; Ramp et al. 2006), especially in narrow roads (personal observation) or when central median with scarce or no vegetation are present (see above). Furthermore, the roadside cover in the study area consists of lush plants, such as *Pistacia lentiscus*, *Rubus ulmifolius*, *Arbutus unedo*, which contribute to reduce visibility and, consequently, increases roadkill probabilities. In fact, roadside management (e.g. regular cutting and removal of dense vegetation) has proven an effective mitigation measure in diverse carnivores (Trombulak and Frissell 2000; Grilo et al. 2009) and birds (Kociolek et al. 2015). For non-flying mammals, it is not surprising that the influence of vegetation height on collision risk decreased, after a threshold. In birds, however, tall vegetation should encourage high flights to cross roads, thereby reducing the probability of vehicle collision as vegetation height increases (Clevenger et al. 2003; Ramp et al. 2006). Not surprisingly, adding fences/walls adjacent to dense vegetation sites has proven an effective mitigation measures in birds and bats (Kociolek et al. 2015). However, as discussed above (see

“road related features”), it is important to ensure that those barriers do not entail additional, undesirable impacts on wildlife, such as collision (as may occur if walls are made of clear glass) or insurmountable barriers to movement (Kociolek et al. 2015).

Potential limitations of the study

Field effort in terms of road distance covered and sampling frequency can strongly influence the accuracy of roadkill counts, because roadkills may be clustered in time and space and several biotic (scavengers) and abiotic (rainfall) factors may affect carcass persistence (Guinard et al. 2012; Teixeira et al. 2013; Barrientos et al. 2018). Thus, it is possible that the total number of casualties for some species is underestimated by monthly sampling (Teixeira et al. 2013). However, this should not be an issue because an accurate estimate of the number of road casualties is not the primary aim of this study; rather, our goal was to investigate the landscape and road features underlying the probability of roadkill. Our conclusions concerning the determinants of roadkill probability are unlikely to be biased by the sampling strategy, since we randomly alternated the road surveys (i.e. were randomly conducted in relation to the ecoregion, weather, type of road, and their fine-scale characteristics) and, therefore, there is no reason to think that roadkills passed systematically unnoticed at the most risky points, and viceversa.

During the first year of study, survey effort during autumn months was comparatively reduced due to logistic issues. This might have affected the roadkill estimates, for example, by decreasing the detection probability of those species that are most active or abundant during autumn, such as migratory birds. However, despite this potential inaccuracy, autumn peaks of mortality for birds and mammals clearly emerged from our survey and, importantly, the composition and temporal distribution of roadkills

(see Canal et al. 2018) are in line with those found in surveys conducted at a shorter sampling periodicity (weekly or fortnightly) in the Iberian Peninsula (e.g. Frias, 1999; Grilo et al., 2009; Garriga et al., 2012; Zuberogoitia et al., 2014; D’Amico et al., 2015). Thus, we are confident that our results were not qualitatively affected by the lesser survey effort performed during the autumn months of the first year of study.

Finally, due to the large spatial scale and range of taxa covered by our survey, local estimates of animal abundance and movements could not be obtained for the entire study region, and their potential effects could not be accounted for as suggested by many authors (Fahrig and Rytwinski 2009; Gunson et al. 2011; van der Ree et al. 2015b; D’Amico et al. 2015). Nonetheless, to partially control for this limitation, the characteristics of the landscape (e.g. land use) adjacent to roads were considered in our analyses, as they often influence animal distribution, abundance, and movements (D’Amico et al. 2015). Future confirmatory studies should explicitly account for these variables when developing models on WVC risk (van der Ree et al. 2015b).

Conclusions

Data on wildlife roadkills were collected at an unusually large temporal (22 months) and spatial (regional) scale, providing stronger inferences of the patterns detected. Fine-grained characterization of road and adjacent landscape characteristics allowed the identification of important factors determining collision risk in small-to medium-sized mammals and birds. Given that roadkill risk may vary at very small spatial scales, we highlight the importance of assessing collision risk based on site-specific attributes and not uniquely on geographic information systems. Overall, reduced traffic density, steep roadside embankments, and structures hampering road access substantially reduced roadkills. By contrast, the effect of other predictors, such as land use adjacent to roads

or the presence of curves, varied between vertebrate groups. It was also evident from our analyses that roadkill risk actually reflects the interplay between different variables. Hence, we suggest that future studies should focus on assessing the effect of particular predictors in road sections with no or little variation in other influential factors e.g. by assessing the effect of different median designs at sites showing similar roadside and landscape characteristics. Further research addressing the impact of medians on wildlife movement and mortality is urgently required because, despite their widespread use, the actual conservation impact of medians remains unclear.

Acknowledgements

We thank Miguel A. Sanza for his dedicated assistance during the road surveys. Two anonymous reviewers provided numerous comments and suggestions that substantially improved the manuscript. This project was funded by Consejería Medio Ambiente (Junta de Andalucía). CC was supported by the Spanish Ministry of Economy and Competitiveness (SVP-2013-067686).

References

- Abbott I, Berthinussen A, Stone M, et al (2015) Bats and roads. In: van der Ree R, Smith D, Grilo C (eds) Handbook of road ecology. John Wiley & Sons, Oxford, pp 290–299
- Adams LW (1984) Small Mammal use of an Interstate Highway Median Strip. *J Appl Ecol* 21:175 . doi: 10.2307/2403045
- Alexander SM, Waters NM (2000) The effects of highway transportation corridors on wildlife: a case study of Banff National Park. *Transp Res Part C Emerg Technol*

494 8:307–320 . doi: 10.1016/S0968-090X(00)00014-0

495 Barrientos R, Bolonio L (2009) The presence of rabbits adjacent to roads increases
 496 polecat road mortality. *Biodivers Conserv* 18:405–418 . doi: 10.1007/s10531-008-
 497 9499-9

498 Barrientos R, Martins RC, Ascensão F, et al (2018) A review of searcher efficiency and
 499 carcass persistence in infrastructure-driven mortality assessment studies. *Biol*
 500 *Conserv* 222:146–153 . doi: 10.1016/j.biocon.2018.04.014

501 Bates D, Mächler M, Bolker B, Walker S (2014) Fitting Linear Mixed-Effects Models
 502 using lme4. *J Stat Softw* 67: . doi: 10.18637/jss.v067.i01

503 Bellis E, Graves H (1971) Deer Mortality on a Pennsylvania Interstate Highway. *J*
 504 *Wildl Manage* 35:232–237

505 Bissonette JA, Kassar CA, Cook LJ (2008) Assessment of costs associated with deer-
 506 vehicle collisions: human death and injury, vehicle damage, and deer loss. *Human-*
 507 *Wildlife Interact* 2:17–27

508 Blackwell B, DeVault T, Seamans T (2015) Understanding and mitigating the negative
 509 effects of road lighting on ecosystems. In: van der Ree R, Smith D, Grilo C (eds)
 510 *Handbook of road ecology*. John Wiley & Sons, Oxford, pp 143–150

511 Canal D, Camacho C, Martín B, et al (2018) Magnitude, composition and
 512 spatiotemporal patterns of vertebrate roadkill at regional scales: a study in southern
 513 Spain. *Anim Biodivers Conserv* 2:281–300

514 Clevenger AP, Chruszcz B, Gunson KE (2003) Spatial patterns and factors influencing
 515 small vertebrate fauna road-kill aggregations. *Biol Conserv* 109:15–26 . doi:
 516 10.1016/S0006-3207(02)00127-1

517 Clevenger AP, Kociolek A V. (2013) Potential Impacts of Highway Median Barriers on
 518 Wildlife: State of the Practice and Gap Analysis. *Environ Manage* 52:1299–1312 .

doi: 10.1007/s00267-013-0155-0

Coffin AW (2007) From roadkill to road ecology: A review of the ecological effects of roads. *J Transp Geogr* 15:396–406 . doi: 10.1016/j.jtrangeo.2006.11.006

Colino-Rabanal VJ, Lizana M (2012) Herpetofauna and roads: A review. *Basic Appl Herpetol* 26:5–31

Colino-Rabanal VJ, Lizana M, Peris SJ (2011) Factors influencing wolf *Canis lupus* roadkills in Northwest Spain. *Eur J Wildl Res* 57:399–409 . doi: 10.1007/s10344-010-0446-1

Crone SF, Finlay S (2012) Instance sampling in credit scoring: An empirical study of sample size and balancing. *Int J Forecast* 28:224–238 . doi: 10.1016/j.ijforecast.2011.07.006

Cserkész T, Ottlecz B, Cserkész-Nagy Á, Farkas J (2013) Interchange as the main factor determining wildlife–vehicle collision hotspots on the fenced highways: spatial analysis and applications. *Eur J Wildl Res* 59:587–597 . doi: 10.1007/s10344-013-0710-2

D’Amico M, Román J, de los Reyes L, Revilla E (2015) Vertebrate road-kill patterns in Mediterranean habitats: Who, when and where. *Biol Conserv* 191:234–242 . doi: 10.1016/j.biocon.2015.06.010

Eddelbuettel D, François R (2011) Rcpp : Seamless R and C++ Integration. *J Stat Softw* 40. doi: 10.18637/jss.v040.i08

Erickson W, Johnson G, Young DJ (2005) A summary and comparison of bird mortality from anthropogenic causes with an emphasis on collisions. In: Ralph, C. John; Rich TD (ed) *Bird Conservation Implementation and Integration in the Americas: Proceedings of the Third International Partners in Flight Conference*. 2002 March 20-24; Asilomar, California, Volume 2 Gen. Tech. Rep. PSW-GTR-191. Albany,

544 CA: U.S. Dept. of Agriculture, Fo

545 Erritzøe J, Mazgajski TD, Rejt Ł (2003) Bird Casualties on European Roads — A

546 Review. *Acta Ornithol* 38:77–93 . doi: 10.3161/068.038.0204

547 Fahrig L, Rytwinski T (2009) Effects of roads on animal abundance: an emperical

548 review and synthesis. *Ecol Soc* 14:21–41 . doi: Artn 21

549 Forman RTT, Sperling D, Bissonette JA, et al (2003) Road ecology. Island Press,

550 Washington DC

551 Frías O (1999) Estacionalidad de los atropellos de aves en el centro de España: número

552 y edad de los individuos y riqueza y diversidad de especies. *Ardeola* 46:23–30

553 Fulton L, Eads G (2004) IEA/SMP Model Documentation and Reference Case

554 Projection; International Energy Agency (IEA)/World Business Council for

555 Sustainable Development (WBCSD): Geneva, Switzerland

556 Gagné SA, Bates JL, Bierregaard RO (2015) The effects of road and landscape

557 characteristics on the likelihood of a Barred Owl (*Strix varia*)-vehicle collision.

558 *Urban Ecosyst* 18:1007–1020 . doi: 10.1007/s11252-015-0465-5

559 Garriga N, Santos X, Montori A, et al (2012) Are protected areas truly protected? The

560 impact of road traffic on vertebrate fauna. *Biodivers Conserv* 21:2761–2774 . doi:

561 10.1007/s10531-012-0332-0

562 GIASA et al. (2006) Libro blanco de las carreteras y los espacios naturales protegidos

563 de Andalucía. Consejería de Obras Públicas y Transportes, Junta de Andalucia

564 Gibbs JP, Shriver WG (2002) Estimating the Effects of Road Mortality on Turtle

565 Populations. *Conserv Biol* 16:1647–1652 . doi: 10.1046/j.1523-1739.2002.01215.x

566 Glista DJ, DeVault TL, DeWoody JA (2009) A review of mitigation measures for

567 reducing wildlife mortality on roadways. *Landsc Urban Plan* 91:1–7 . doi:

568 10.1016/j.landurbplan.2008.11.001

569 Gomes L, Grilo C, Silva C, Mira A (2009) Identification methods and deterministic
570 factors of owl roadkill hotspot locations in Mediterranean landscapes. *Ecol Res*
571 24:355–370 . doi: 10.1007/s11284-008-0515-z

572 Grilo C, Bissonette JA, Santos-Reis M (2009) Spatial–temporal patterns in
573 Mediterranean carnivore road casualties: Consequences for mitigation. *Biol*
574 *Conserv* 142:301–313 . doi: 10.1016/j.biocon.2008.10.026

575 Guinard É, Julliard R, Barbraud C (2012) Motorways and bird traffic casualties:
576 Carcasses surveys and scavenging bias. *Biol Conserv* 147:40–51 . doi:
577 10.1016/j.biocon.2012.01.019

578 Gunson KE, Mountrakis G, Quackenbush LJ (2011) Spatial wildlife-vehicle collision
579 models: A review of current work and its application to transportation mitigation
580 projects. *J Environ Manage* 92:1074–1082 . doi: 10.1016/j.jenvman.2010.11.027

581 Hartig F (2016) DHARMA: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)
582 Regression Models. In: R Packag. <https://cran.r-project.org/package=DHARMA>

583 Huijser MP, Duffield JW, Clevenger AP, et al (2009) Cost-benefit analyses of
584 mitigation measures aimed at reducing collisions with large ungulates in the united
585 states and canada: A decision support tool. *Ecol Soc* 14:15 . doi:
586 10.1016/j.contraception.2009.11.002

587 Kociolek A, Grilo C, Jacobson S (2015) Flight doesn't solve everything: mitigation of
588 road impacts on birds. In: van der Ree R, Smith D, Grilo C (eds) *Handbook of road*
589 *ecology*. John Wiley & Sons, Oxford, pp 281–289

590 Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in
591 Linear Mixed Effects Models. *J Stat Softw* 82: . doi: 10.18637/jss.v082.i13

592 Langen T a., Gunson KE, Scheiner C a., Boulerice JT (2012) Road mortality in
593 freshwater turtles: Identifying causes of spatial patterns to optimize road planning

594 and mitigation. *Biodivers Conserv* 21:3017–3034 . doi: 10.1007/s10531-012-0352-
595 9

596 Madsen A, Strandgaard H, Prang A (2002) Factors causing traffic killings of roe deer
597 *Capreolus capreolus* in Denmark. *Wildlife Biol* 8:55–61

598 Malo JE, Suárez F, Díez A (2004) Can we mitigate animal–vehicle accidents using
599 predictive models? *J Appl Ecol* 41:701–710

600 Massemin S, Zorn T (1998) Highway Mortality of Barn Owls in Northeastern France. *J*
601 *Raptor Res* 32:229–232

602 Meyer I, Kaniovski S, Scheffran J (2012) Scenarios for regional passenger car fleets and
603 their CO₂ emissions. *Energy Policy* 41:66–74 . doi: 10.1016/j.enpol.2011.01.043

604 Mumme RL, Schoech SJ, Woolfenden GE, Fitzpatrick JW (2000) Life and death in the
605 fast lane: Demographic consequences of road mortality in the Florida Scrub-Jay.
606 *Conserv Biol* 14:501–512 . doi: 10.1046/j.1523-1739.2000.98370.x

607 Myers N, Mittermeier RA, Mittermeier CG, et al (2000) Biodiversity hotspots for
608 conservation priorities. *Nature* 403:853–858 . doi: 10.1038/35002501

609 Planillo A, Malo JE (2013) Motorway verges: Paradise for prey species? A case study
610 with the European rabbit. *Mamm Biol* 78:187–192 . doi:
611 10.1016/j.mambio.2012.11.001

612 Planillo A, Malo JE (2018) Infrastructure features outperform environmental variables
613 explaining rabbit abundance around motorways. *Ecol Evol* 8:942–952 . doi:
614 10.1002/ece3.3709

615 Ramp D, Caldwell J, Edwards K a., et al (2005) Modelling of wildlife fatality hotspots
616 along the Snowy Mountain Highway in New South Wales, Australia. *Biol Conserv*
617 126:474–490 . doi: 10.1016/j.biocon.2005.07.001

618 Ramp D, Wilson VK, Croft DB (2006) Assessing the impacts of roads in peri-urban

619 reserves: Road-based fatalities and road usage by wildlife in the Royal National
 620 Park, New South Wales, Australia. *Biol Conserv* 129:348–359 . doi:
 621 10.1016/j.biocon.2005.11.002
 622 Reale D, Garant D, Humphries MMM, et al (2010) Personality and the emergence of
 623 the pace-of-life syndrome concept at the population level. *Philos Trans R Soc Lond*
 624 *B Biol Sci* 365:4051–4063 . doi: 10.1098/rstb.2010.0208
 625 Sandberg M, Eriksson L, Jonsson J, et al (1998) New chemical descriptors relevant for
 626 the design of biologically active peptides. A multivariate characterization of 87
 627 amino acids. *J Med Chem* 41:2481–91 . doi: 10.1021/jm9700575
 628 Seiler A (2005) Predicting locations of moose-vehicle collisions in Sweden. *J Appl Ecol*
 629 42:371–382 . doi: 10.1111/j.1365-2664.2005.01013.x
 630 Teixeira FZ, Coelho AVP, Esperandio IB, Kindel A (2013) Vertebrate road mortality
 631 estimates: Effects of sampling methods and carcass removal. *Biol Conserv*
 632 157:317–323 . doi: 10.1016/j.biocon.2012.09.006
 633 Trombulak SC, Frissell CA (2000) Review of ecological effects of roads on terrestrial
 634 and aquatic communities. *Conserv Biol* 14:18–30
 635 van der Grift EA, van der Ree R, Fahrig L, et al (2013) Evaluating the effectiveness of
 636 road mitigation measures. *Biodivers Conserv* 22:425–448 . doi: 10.1007/s10531-
 637 012-0421-0
 638 van der Ree R, Gagnon J, Smith D (2015a) Fencing: A valuable tool for reducing
 639 wildlife-vehicle collisions and funnelling fauna to crossing structures. In: van der
 640 Ree R, Smith D, Grilo C (eds) *Handbook of road ecology*. John Wiley & Sons,
 641 Oxford, pp 159–171
 642 van der Ree R, Smith D, Grilo C (2015b) *Handbook of Road Ecology*. John Wiley &
 643 Sons

644 van der Ree R, Smith D, Grilo C (2015c) The ecological effects of linear infrastructure
 645 and traffic: challenges and opportunities of rapid global growth. In: van der Ree R,
 646 Smith D, Grilo C (eds) Handbook of road ecology. John Wiley & Sons, Oxford, pp
 647 1–9

648 Zuberogoitia I, del Real J, Torres JJ, et al (2014) Ungulate Vehicle Collisions in a Peri-
 649 Urban Environment: Consequences of Transportation Infrastructures Planned
 650 Assuming the Absence of Ungulates. PLoS One 9:e107713 . doi:
 651 10.1371/journal.pone.0107713

652 Zuberogoitia I, Martínez JE, Martínez JA, et al (2006) Influence of management
 653 practices on nest site habitat selection, breeding and diet of the common buzzard
 654 *Buteo buteo* in two different areas of Spain. *Ardeola* 53:83–98
 655

Figure 1. Situation of the road sections and main ecological units (ecoregions) surveyed during the study period. Surveyed roads (roadkill and control points) are highlighted in black.

Figure 2. Number of recorded bird (left) and mammal (right) roadkills in relation to traffic density (average number of vehicles per day) of the roads.

Figure 3. Probability of vehicle collision in birds (left) and mammals (right) according to height of roadside vegetation (upper figures) and distance to the nearest curve (lower figures). Grey dots are predicted values, the solid line denotes the fitted response of GLMMS and dashed lines show the 95% confidence intervals. Distances to the nearest curve were log transformed to meet normality assumptions.

Figure 4. Probability of vehicle collision in birds (upper) and mammals (bottom) according to type of habitat surrounding the road. Boxplots show the extreme of the lower whisker, the lower hinge, the median, the upper hinge, and the extreme of the upper whisker. Dots are data points that lie beyond the extremes of the whiskers. Only significant P-values (< 0.05) from post-hoc Tukey tests are shown.

Figure 5. Probability of vehicle collision in birds (left) and mammals (right) according to type of fence (upper figures) and amount of vegetation cover in the median strips (lower figures). Boxplots show the extreme of the lower whisker, the lower hinge, the median, the upper hinge, and the extreme of the upper whisker. Dots are data points that lie beyond the extremes of the whiskers.

Table 1. Description of the variables measured both at collision and control points and expected influence (increase (+), decrease (-) or quadratic) that each factor has on wildlife-vehicle collisions according to the taxonomic group and existing literature. Note that only some studies among those available in the literature are reported (for a review, see e.g. Gunson et al. 2011). For further details about the influence of each predictor on WVC we refer to the original source.

Quantitative variable	Description	Birds	Mammals	Ref.
Max_Vegetation ⁺	Maximum height (m) of the vegetation; measured at both sides of the road	+/- Quadratic (concave)	+/- Quadratic (concave)	[1], [2], [3]
Dist_Curve	Distance (m) to the nearest curve estimated from a geographic information system	+/- Quadratic (concave)	+/- Quadratic (concave)	[4], [5], [6], [7]
Traffic density	Averaged number of vehicles per day	+/- Quadratic (concave)	+/- Quadratic (concave)	[1], [2], [7], [8]
Qualitative variable*				
Fence	Unfenced (1), wire mesh fence (2), barbed wire fence (3) and wall	- as the difficulty to access the road increases	-/+ due to trap effects	[7], [9], [10], [11]
Median	Absent (1), with medium (2) or much (3) vegetation and Jersey barriers (4)	+ vegetation in medians will attract animals	+ vegetation in medians will attract animals	[1], [9], [12]
Land use	Dominant use of adjacent habitat around 100 m of the focal point: Urban (1), scrubland (2), farmland (3) and forest (4)	Depends on local population density	Depends on local population density. Generally, + in forest	[1], [4], [5], [6], [8]
Embankments [†]	Combined both road sides each point was classified as: at the road level (1), buried (2), raised (3), buried-raised (4), ground level-buried (5) and ground level-raised (6)	- as the steep in the embankments increases	- as the steep in the embankments increases	[1], [4], [5], [8]

⁺ Vegetation height may be considered as a proxy of vegetation cover since the correlation between height and width in the most common Mediterranean scrubs is very high ($r = 0.96$; Camacho 2014)

*Class variables (land use, presence and type of road embankments, presence and type of fence and median centers) were visually estimated.

[†] Roadside topography was initially classified at each side as: at the ground level (1), gentle buried (2), steep buried (3), gentle raised (4), steep raised (5).

References: [1] Clevenger et al. 2003, [2] Barrientos and Bolonio 2009, [3] Ramp et al. 2006, [4] Malo et al. 2004, [5] Ramp et al. 2005, Grilo et al. 2009, [7] Zuberogitia et al. 2006, [8] Seiler 2005, [9] Bellis and Graves 1971, [10] Colino-Rabanal et al. 2011, [11] Van der Ree et al. 2015 [12], Clevenger and Kociolek 2013.

Table 2. Relationship between the occurrence of roadkills in birds and the characteristics of the road and adjacent land use.

Fixed effects	Estimate	SE	Z	P
Intercept	-3.854	1.014	-3.800	<0.001
Road category:				
National	-0.381	0.204	-1.869	0.062
Local	-0.957	0.229	-4.172	<0.001
Dist_Curve	1.309	0.388	3.379	0.001
Dist_Curve^2	-0.121	0.038	-3.226	0.001
Vegetation	0.503	0.090	5.602	<0.001
Vegetation^2	-0.056	0.012	-4.560	<0.001
Land use:				
Farmland	-0.552	0.167	-3.312	<0.001
Scrubland	-0.873	0.215	-4.069	<0.001
Urban	-0.556	0.264	-2.107	0.035
Embankments*	-0.472	0.189	-2.490	0.013
Random effects	Variance	SD		
RoadID:Region	0.150	0.387		
Region	0.090	0.300		

* Simplified to two levels according road topography while fitting the model: roads hampering birds to fly at low altitude and roads facilitating birds to fly close to the road surface.

Table 3. Relationship between the occurrence of roadkills in mammals and the characteristics of the road and adjacent land use.

Fixed effects	Estimate	SE	Z	P
Intercept	-4.539	1.105	-4.107	<0.001
Road category:				
National	-0.669	0.253	-2.643	0.008
Local	-1.499	0.288	-5.205	<0.001
Dist_Curve	1.062	0.389	2.728	0.006
Dist_Curve^2	-0.097	0.038	-2.572	0.010
Vegetation	0.604	0.091	6.616	<0.001
Vegetation^2	-0.062	0.012	-5.119	<0.001
Land use:				
Farmland	0.079	0.167	0.475	0.635
Scrubland	-1.244	0.259	-4.806	<0.001
Urban	-0.797	0.277	-2.880	0.004
Embankments*:				
Buried roads	1.342	0.227	5.908	<0.001
Roads at level	2.402	0.217	11.090	<0.001
Roads with mixed embankments	1.719	0.164	10.468	<0.001
Random effect	Variance	SD		
RoadID:Region	0.285	0.534	-	-
Region	0.745	0.863	-	-

* Simplified to four levels according road topography while fitting the model: road at ground level, buried, raised and road with mixed embankments (involving roads part buried, part raised and buried-raised)

719

SUPPLEMENTARY FILES.720 **SUP MAT. TABLE 1.** Ecoregion, road type and situation of the surveyed roads. For further information about the number of casualties

721 registered in each road section and taxonomic group, see Canal et al. (2018)

Ecoregion	Type	ID	Province	Km origin	Km end	Mammals	Birds
Lowlands and green fields of Guadalquivir river	Highways	A-92	Seville	42	52	66	19
		A-4	Seville	497	507	51	19
		A-49	Huelva	23	33	39	11
	National roads	A-380	Seville	0	10	51	21
		A-407/A-456	Seville	8	38*	57	16
		A-364	Seville	5	15	47	8
	Local roads	JA-6108	Jaen	0	10	4	6
		SE-7200	Seville	0	10	23	8
		SE-8105	Seville	0	10	17	6
Atlantic and Mediterranean Coastline	Highways	A-49	Huelva	107	117	23	23
		A-48	Cadiz	4	14	67	24
		A-7	Almeria	483	493	15	13
	National roads	A-494	Huelva	42	52	2	2
		A-405	Cadiz	23	33	8	24
		A-377	Malaga	5	15	3	12
	Local roads	A-2227	Cadiz	0	10	33	18
		A-2101	Cadiz	0	10	17	14
		AL-3106	Almeria	13	23	4	8

722

Ecoregion	Type	ID	Province	Km origin	Km end	Mammals	Birds
Atlantic and Continental bio-geographic regions of Sierra Morena	Highways	A-66	Huelva	755	765	5	13
		A-66	Seville	766	776	13	13
		A-66	Seville	782	792	21	31
	National roads	A-461	Huelva	2	12	1	3
		N-433	Huelva	75	85	10	19
		A-424	Córdoba	1	11	4	2
	Local roads	HU-9116/SE-6405	Huelva/Seville	1	15*	0	0
		CO-6103	Córdoba	1	11	2	4
		A-3200	Córdoba	9	19	1	1
Medium and high mountain areas of the Baetic system	Highways	A-381	Cadiz	43	53	49	21
		A-92M	Malaga	1	11	24	21
		A-92	Malaga	161	171	58	41
	National roads	A-308	Granada	23	33	13	8
		A-406	Seville	12	22	13	25
		A-333/A-328	Malaga/Córdoba	59	69*	18	24
	Local roads	CA-5102	Cadiz	8	18	2	8
		SE-8205	Seville	0	10	2	8
		MA-5102	Malaga	0	10	11	3

723

724

725

Ecoregion	Type	ID	Province	Km origin	Km end	Mammals	Birds	Reptiles	Amphibians
Arid zones in southeastern Andalusia	Highways	A-92N	Granada	10	20	5	5	1	0
		A-92N	Granada	50	60	15	9	1	0
		A-92	Almeria	365	375	9	9	3	2
	National roads	A-334	Granada	11	21	13	7	0	0
		A-330	Granada	4	14	7	8	1	0
		A-349	Almeria	2	12	8	9	8	0
	Local roads	GR-7100	Granada	0	10	3	3	0	0
		GR-9109	Granada	2	12	0	0	1	0
		AL-3102	Almeria	2	12	1	8	4	5

726

727

728

***10-km long sections including the same road designated with two different names, so that the official assignation of km also varies.**

729 **SUP MAT. TABLE 2.** Number of individuals of the mammal and birds species found killed by traffic along Andalusian roads between the
 730 2009-2010 and 2011-2012 surveys.

	Records
Mammals	835
<i>Oryctolagus cuniculus</i>	298
<i>Rattus norvegicus</i>	103
<i>Canis lupus familiaris</i>	93
<i>Erinaceus europaeus</i>	76
<i>Felis silvestris catus</i>	63
<i>Lepus sp</i>	54
<i>Lepus europaeus</i>	46
<i>Vulpes vulpes</i>	30
<i>Lepus granatensis</i>	15
<i>Apodemus sylvaticus</i>	11
<i>Genetta genetta</i>	8

731

732

Species	Records
Mammals	835
<i>Herpestes ichneumon</i>	8
<i>Pipistrellus pipistrellus</i>	7
<i>Martes foina</i>	6
<i>Mustela putorius</i>	4
<i>Eptesicus serotinus</i>	3
<i>Meles meles</i>	3
<i>Eliomys quercinus</i>	2
<i>Mustela nivalis</i>	1
<i>Undetermined mammals</i>	4
Birds	555
<i>Passer domesticus</i>	70
<i>Sylvia melanocephala</i>	48
<i>Athene noctua</i>	42
<i>Alectoris rufa</i>	41
<i>Turdus merula</i>	30
<i>Sylvia atricapilla</i>	29
<i>Erithacus rubecula</i>	28
<i>Columba livia</i>	12
<i>Fringilla coelebs</i>	12
<i>Galerida cristata</i>	12
<i>Saxicola torquata</i>	11
<i>Tyto alba</i>	11
<i>Caprimulgus ruficollis</i>	8

Species	Records
Birds	555
<i>Miliaria calandra</i>	8
<i>Pica pica</i>	8
<i>Melanocorypha calandra</i>	7
<i>Turdus philomelos</i>	7
<i>Hirundo rustica</i>	8
<i>Alauda arvensis</i>	6
<i>Serinus serinus</i>	6
<i>Cisticola juncidis</i>	5
<i>Hirundo daurica</i>	5
<i>Phylloscopus collybita</i>	5
<i>Bubo bubo</i>	4
<i>Calandrella brachydactyla</i>	4
<i>Carduelis carduelis</i>	4
<i>Jynx torquilla</i>	4
<i>Merops apiaster</i>	4
<i>Motacilla alba</i>	4
<i>Passer montanus</i>	4
<i>Picus viridis</i>	4
<i>Streptopelia decaocto</i>	4
<i>Asio otus</i>	3
<i>Bubulcus ibis</i>	3
<i>Cuculus canorus</i>	3
<i>Lanius senator</i>	3
<i>Sylvia conspicillata</i>	3

Species	Records
Birds	555
<i>Sylvia undata</i>	3
<i>Upupa epops</i>	3
<i>Anas platyrhynchos</i>	2
<i>Cettia cetti</i>	2
<i>Cyanopica cooki</i>	2
<i>Gallus gallus domesticus</i>	2
<i>Lanius meridionalis</i>	2
<i>Larus argentatus</i>	2
<i>Otus scops</i>	2
<i>Cyanistes caeruleus</i>	2
<i>Phoenicurus ochruros</i>	2
<i>Strix aluco</i>	2
<i>Sturnus unicolor</i>	2
<i>Sturnus vulgaris</i>	2
<i>Sylvia cantillans</i>	2
<i>Accipiter nisus</i>	1
<i>Asio flammeus</i>	1
<i>Carduelis cannabina</i>	1
<i>Carduelis chloris</i>	1
<i>Circus cyaneus</i>	1
<i>Carduelis chloris</i>	1
<i>Circus cyaneus</i>	1
<i>Corvus monedula</i>	1
<i>Coturnix coturnix</i>	1

Species	Records
Birds	555
<i>Delichon urbicum</i>	1
<i>Falco tinnunculus</i>	1
<i>Galerida theklae</i>	1
<i>Gallinula chloropus</i>	1
<i>Garrulus glandarius</i>	1
<i>Larus michahellis</i>	1
<i>Milvus milvus</i>	1
<i>Parus major</i>	1
<i>Petronia petronia</i>	1
<i>Phylloscopus trochilus</i>	1
<i>Streptopelia turtur</i>	1
<i>Undetermined birds</i>	35

733

SUP MAT. TABLE 3. Roadkill probability in (a) passerines, (b) carnivores and (c) lagomorphs, the orders most affected by roadkills during the study period (see Canal et al 2018), in relation to the characteristics of the road and adjacent land use.

A)

Fixed effects	Estimate	SE	Z	P
Intercept	-5.359	1.256	-4.270	<0.001
Road category:				
National	0.005	0.263	0.020	0.986
Local	-0.498	0.291	-1.710	0.087
Dist_Curve	1.526	0.477	3.200	0.001
Dist_Curve^2	-0.138	0.046	-2.990	0.003
Vegetation	0.712	0.119	5.980	<0.001
Vegetation^2	-0.095	0.018	-5.240	<0.001
Land use:				
Farmland	-0.644	0.197	-3.270	0.001
Scrubland	-1.210	0.271	-4.460	<0.001
Urban	-0.332	0.293	-1.130	0.258
Embankments*	-0.488	0.231	-2.110	0.035
Random effect	Variance	SD		
RoadID:Region	0.282	0.531	-	-
Region	0.134	0.366	-	-

* Simplified to two levels according road topography while fitting the model: roads hampering birds to fly at low altitude and roads facilitating birds to fly close to the road surface.

742 B)

Fixed effects	Estimate	SE	Z	P
Intercept	-7.975	1.685	-4.730	<0.001
Road category:				
National	0.081	0.247	0.330	0.743
Local	-1.203	0.315	-3.820	<0.001
Dist_Curve	1.782	0.617	2.890	0.004
Dist_Curve^2	-0.164	0.058	-2.830	0.005
Vegetation	0.801	0.145	5.510	<0.001
Vegetation^2	-0.090	0.021	-4.390	<0.001
Land use:				
Farmland	-0.105	0.252	-0.420	0.678
Scrubland	-0.902	0.338	-2.670	0.008
Urban	-0.507	0.392	-1.290	0.196
Embankments*:				
Buried roads	1.406	0.328	4.280	<0.001
Roads at level	2.565	0.307	8.370	<0.001
Roads with mixed embankments	1.625	0.248	6.570	<0.001
Random effect	Variance	SD		
RoadID:Region	0.101	0.317	-	-
Region	0.384	0.62	-	-

743

744 * Simplified to four levels according road topography while fitting the model: road at ground level,
745 buried, raised and road with mixed embankments (involving roads part buried, part raised and
746 buried-raised)
747

748

749 C)

Fixed effects	Estimate	SE	Z	P
Intercept	-4.182	1.555	-2.688	0.007
Road category:				
National	-1.628	0.416	-3.915	<0.001
Local	-2.105	0.456	-4.611	<0.001
Dist_Curve	0.597	0.556	1.073	0.283
Dist_Curve^2	-0.056	0.054	-1.030	0.303
Vegetation	0.539	0.123	4.385	0.000
Vegetation^2	-0.053	0.017	-3.159	0.002
Land use:				
Farmland	0.445	0.219	2.034	0.042
Scrubland	-0.925	0.382	-2.421	0.015
Urban	-1.313	0.423	-3.100	0.002
Embankments*:				
Buried roads	1.425	0.303	4.707	<0.001
Roads at level	2.564	0.299	8.575	<0.001
Roads with mixed Embankments	1.829	0.218	8.403	<0.001
Random effect	Variance	SD		
RoadID:Region	0.698	0.836	-	-
Region	1.503	1.226	-	-

750 ***Simplified to four levels according road topography while fitting the model: road at ground level,**
751 **buried, raised and road with mixed embankments (involving roads part buried, part raised and**
752 **buried-raised)**
753

754 The number of casualties increased with traffic density in carnivores ($R = 0.48$, P
755 <0.001), lagomorphs ($R = 0.42$, $P = 0.004$) and passerines ($R = 0.23$, $P = 0.12$), the latter
756 being marginally significant. In carnivores and passerines, the number of casualties
757 decreased as the difficulty of accessing roads increased from unfenced points, followed
758 by barber fences, up to points having mesh wire fences or walls (passerines: estimate
759 (SE) -0.703 (0.209), $Z = -3.37$, $P < 0.001$; carnivores: estimate (SE) -0.75 (0.27), $Z = -$
760 2.78, $P = 0.005$), whereas this factor did not affected the probability of roadkill in
761 lagomorphs (estimate (SE) -0.15 (0.38), $Z = -0.41$, $P = 0.68$). The risk of roadkill in
762 carnivores and passerines also decreased as vegetation cover in the median strips

763 increased (passerines: estimate (SE) -0.38 (0.2), $Z = -1.8$, $P = 0.06$; carnivores: estimate
764 (SE) -0.616 (0.31), $Z = -2.05$, $P = 0.04$), whereas in lagomorphs, the probability of
765 roadkill was unrelated to the vegetation cover in the median strips (estimate (SE) -0.019
766 (0.21), $Z = -0.92$, $P = 0.35$).

767

768