1 **Neuroprotective protein hydrolysates from hemp (*Cannabis sativa* L.)**
2 **seeds**

3
4 Noelia M **Rodriguez-Martin**^{a,b}, Rocio **Toscano**^{a,b}, Alvaro **Villanueva**^b,
5 Justo **Pedroche**^b, Francisco **Millan**^b, Sergio **Montserrat-de la Paz**^{a,*},
6 Maria C **Millan-Linares**^c

7
8 ^a Department of Medical Biochemistry, Molecular Biology, and Immunology,
9 School of Medicine, Universidad de Sevilla. Av. Dr. Fedriani 3, 41071 Seville,
10 Spain.

11 ^b Department of Food & Health. Instituto de la grasa, CSIC. Ctra. Utrera Km 1,
12 41013 Seville, Spain.

13 ^c Cell Biology Unit, Instituto de la grasa, CSIC. Ctra. Utrera Km 1, 41013 Seville,
14 Spain.

15
16 *** Corresponding author:**

17 Prof. Sergio Montserrat-de la Paz

18 Department of Medical Biochemistry, Molecular Biology, and Immunology,
19 School of Medicine, Universidad de Sevilla. Av. Dr. Fedriani 3, 41071 Seville,
20 Spain.

21 Tel.: +34-954-559-850

22 Email address: delapaz@us.es

23
24
25

ABSTRACT

View Article Online
DOI: 10.1039/C9FO01904A

Hemp (*Cannabis sativa* L.) seeds are well known for their potential use as source of nutrients, fiber, and bioactive compounds. Hemp protein isolate, prepared from defatted hemp flour, was hydrolyzed by Alcalase and Flavourzyme under specific conditions. The resulting hydrolysates were evaluated to the selection of potentially bioactive hemp protein hydrolysates (HPHs) by its DPPH scavenging and ferric reducing antioxidant power activity. *In vitro* cell-free experiments led to the identification of two bioactive HPHs, HPH20A and HPH60A+15AF, which were used at 50 and 100 µg/mL on BV-2 microglial cells in order to evaluate the anti-neuroinflammatory activities. Our results showed that HPH20A and HPH60A+15AF down-regulated *TNF-α*, *IL-1β*, and *IL-6* mRNA transcriptional levels in LPS-stimulated BV-2 microglial cells. In addition, HPH20A and HPH60A+15AF up-regulated gene expression of anti-inflammatory cytokine IL-10. This study suggests for the first time that HPHs may improve neuroinflammatory and inflammatory states, supporting the nutraceutical value of hemp seeds.

Keywords: Hemp; *Cannabis sativa*; protein hydrolysate; microglia; neuroinflammation.

1 **Introduction**

2 Non-drug varieties of *Cannabis sativa* L. (Cannabaceae), collectively known as
3 “hemp”, are a well-known plant of industrial value as an important source of
4 nutrients, fiber, and bioactive compounds.¹ In the last years, industrial hemp
5 global market is rapidly increasing, since low- Δ^9 -tetrahydrocannabinol (THC)
6 cultivars are legal and available.² Hemp seeds are raising great interest as
7 valuable crop, not requiring fertilizers, herbicides, and pesticides.³ Currently,
8 the great attention for hemp seed is related to its nutritional content: 25-35% oil,
9 20–30% carbohydrates, 20-25% proteins, 27.6% total fiber, vitamins, and
10 minerals.^{4,5}

11 Up-to-now, there is increasing attention for hemp seed proteins owing to their
12 digestibility, satisfactory essential amino acid composition, and techno-
13 functional properties.^{6,7} Nevertheless, the poor water solubility of hemp protein
14 isolate (HPI) hinders its application as a functional ingredient in food industry.
15 Recent research has shown that hydrolysates obtained by treating HPI with
16 different enzymes, such as pepsin, alcalase, papain, pancreatin, or other
17 proteases, containing from 60 % to 80 % of protein content.⁸ Induced
18 proteolysis not only modifies HPI hydrophobicity, but also changes protein
19 molecular weight and conformation, obtaining high availability of bioactive
20 peptides.¹

21 In this context, hemp protein hydrolysates (HPHs) contain a mixture of
22 polypeptides, oligopeptides, and amino acids that can be used as a key factor in
23 the development of applications in nutraceuticals and functional foods.⁹ To date,
24 the characterization of the bioactive HPHs remains largely unstudied. In

particular, few studies have showed that HPHs obtained from enzymatic hydrolysis demonstrated antihypertensive, hypocholesterolemic, antioxidant, antithrombotic, and immunomodulatory effects.^{8–14} However, to our knowledge, neuroprotective HPHs have not been described in the literature.

Microglia are phagocytic macrophages present in all regions of the adulthood brain and spinal cord, and represent between 10–15% of the total cells in the central nervous system (CNS).¹⁵ The principal function of microglia is to destroy and to clear foreign materials through phagocytic and cytotoxic mechanisms.¹⁶ Microglia cells also contribute to inflammation and homeostatic mechanisms by secreting cytokines and other signaling molecules, and they act as antigen presenting cells.¹⁷ When local inflammation in the CNS and brain is chronically active (neuroinflammation), it can produce a wide variety of neurodegenerative disorders including Alzheimer's disease, Parkinson's disease, Huntington's disease, tauopathies, age-related macular degeneration, and diseases of autoimmune origin such as multiple sclerosis.¹⁸

On the basis of these considerations, the overall objective of the present study was to investigate the effect of different HPHs from hemp seed on neuroinflammation in LPS-activated BV-2 microglial cells. In particular, to achieve this goal, the specific targets were (i) the chemical characterization of hemp products, (ii) the evaluation of the biological activity of HPHs on *in vitro* cell-free experiments, and (iii) the identification of neuroprotective effect of the two HPHs selected on BV-2 microglial cells.

1 **Materials and methods**

2 **Chemicals and sampling**

3 Seeds of *C. sativa* L. were provided by Sensi Seeds Bank. Alcalase 2.4 L (2.4
4 AU/g) and Flavourzyme (1000 L) were a gift from Novozymes (Bagsvaerd,
5 Denmark). The murine BV-2 cell line was gently provided by Cell Biology Unit
6 (Institute of Fats, CSIC, Spain). Primers were purchased from Eurofins Biolab
7 S.L.U (Barcelona, Spain). Total RNA was isolated using Trisure (Bioline,
8 Meridian Life Science, Inc. Memphis, USA) and reverse transcribed to cDNA
9 using a synthesis kit of Bio-rad (California, EE. UU). Quantitative PCR was
10 performed by iTaq™ Universal SYBR® Green Supermix of Bio-rad. All
11 chemicals (reagents and solvents) were of analytical grade and provided by
12 Sigma Chemical Co. (St. Louis, MO, USA), Bachem AG and Gibco.

13 **Preparation of hemp protein isolate (HPI)**

14 The HPI was obtained using the method of Lqari et al.¹⁹ Briefly, hemp defatted
15 flour (HDF) was extracted with 0.25% Na₂SO₃ (w/v) at pH 10.5 for 1 h. After
16 centrifuging the extract at 7500 rpm for 15 min, the supernatant was recovered
17 and the pellet was extracted again. Both supernatants were adjusted to the
18 isoelectric point of hemp proteins (pH 4.3) and the precipitate was washed with
19 distilled water adjusted to pH 4.3 and centrifuged to remove residual salts and
20 other non-protein compounds. Finally, the precipitated proteins were lyophilized
21 and stored at room temperature.

22

23

Hydrolysis of hemp protein isolate

Hydrolysis of HPI was performed in a bioreactor under continuous stirring at a controlled pH and temperature. HPI was suspended in distilled water (10% w/v) and temperature was adjusted to 50°C. Alcalase 2.4 L (2.4 AU/g, a serine endopeptidase which was kindly donated by Novozymes, Bagsvaerd, Denmark) was added at enzyme/substrate = 0.3 AU/g protein at pH 8 for 3 h. After 60 min of hydrolysis with Alcalase, Flavourzyme (1000 L, a leucine amino-peptidase which was kindly donated by Novozymes) was added at enzyme/substrate = 60 LAPU/g protein at pH 7 for 2 h. During peptic hydrolysis, the pH was constantly kept by titration with NaOH solution in appropriate concentration. Each hydrolytic time-point sample was stopped by holding at 85 °C for 15 min to ensure a complete inactivation of residual enzyme activity. The supernatant obtained by centrifugation at 7500 rpm for 15 min constituted the HPH. The HPHs obtained using only Alcalase were designated as 10A, 20A, 30A, 45A, and 60A. The HPHs obtained using Alcalase followed by Flavourzyme were designated as 60A+15AF, 60A+30AF, 60A+60AF, 60A+90AF, and 60A+120AF. The number indicates the time of hydrolysis in minutes.

Hemp product compositional analysis

The protein concentration was determined by elemental microanalysis as % nitrogen content x 6.25 using a Leco CHNS932 analyser (St. Joseph, MI, USA). Total dietary fiber was determined by the gravimetric method.²⁰ Ash content was determined according to the direct ignition method (550 °C for 36 h). Polyphenols and soluble sugars were measured using standard curves of chlorogenic acid and glucose, respectively.^{21,22} Amino acid quantification by

HPLC was done according to Alaiz et al.²³ Finally, tryptophan content was analyzed according to the method of Yust et al.²⁴

Evaluation of hydrolysis degree (HD)

The HD, defined as the percentage of peptide bonds cleaved, was calculated by the TNBS method which determine free amino acid groups.²⁵ The total number of amino groups was determined in a sample that had been 100% hydrolyzed in 6 N HCl at 110 °C for 24 h.

Molecular profile by Fast Protein Liquid Chromatography (FPLC)

The molecular profile of the samples was determined using FPLC Akta purifier 10 (GE Healthcare Bio-sciences AB, Uppsala, Switzerland). Superose 12 HR 10/300 GL, a prepacked column for high-performance size exclusion chromatography with a broad fractionation range for molecules with molecular weights between 1 and 300 kDa, was used. The column was calibrated with proteins of known molecular weights. From the logarithms of the molecular weights of these proteins and their elution volumes, the calibration line was made. The elution was carried out with 50 mL of 0.05 M sodium phosphate buffer, 0.5 M sodium chloride and 0.02% sodium azide (w/v) at pH 7.5, with a flow of 1 mL/min. The injected volume of the samples and its concentrations were of 500 µL and 1 mg/mL of protein, respectively. The elution of the proteins was recorded by measuring their absorbance at 280 nm.

DPPH radical scavenging activity

The scavenging effect of the samples on DPPH free radical was measured according to Wu et al.²⁶ Briefly, 1.5 mL of each sample (at concentration of 10

mg of protein/mL) was added to 1.5 mL of 0.1 mmol/L DPPH in ethanol. The mixture was shaken and left for 30 min at room temperature, and absorbance was measured at 517 nm. The control was conducted in the same manner but distilled water was used instead of sample. Butylated hydroxytoluene (BHT) at 0.08 mg/mL was used as positive standard.

Ferric reducing antioxidant power (FRAP)

The ability of the samples to reduce iron (III) was measured according to Oyaizu.²⁷ Briefly, 0.1 mL of sample (at concentration of 10 mg of protein/mL) was added to 0.25 mL of 0.2 mol/L sodium phosphate buffer pH 6.6 and 0.25 mL of 0.03 mol/L potassium ferricyanide. The mixture was incubated at 50 °C for 20 min. Then 0.25 mL of 0.6 mol/L trichloroacetic acid was added and the mixture was centrifuged at 1300 x g for 10 min. 500 µL of the upper layer were mixed with 0.5 mL of distilled water and 0.1 mL of 3.7 mmol/L ferric chloride. After 10 min, absorbance of the resulting solutions was measured at 700 nm. Increased absorbance of the reaction mixture indicated increased reducing power. BHT at 0.08 mg/mL was used as standard.

BV-2 Cell Culture and Treatments

BV-2 is a murine cell line used extensively in research related to neurodegenerative disorders.^{28,29} BV-2 microglial cells were obtained from the Cell Biology Unit at the Instituto de la Grasa (Seville, Spain). The cells were maintained in high glucose Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% heat-inactivated foetal bovine serum and 1% penicillin/streptomycin. They were maintained in 5% CO₂ at 37 °C in a CO₂ incubator (Thermo Con Electron Corporation, Waltham, MA, USA). Experiments

1 were done in twelve-well plates. BV-2 microglial cells were seeded at a density
2 of 5×10^5 cells/well. Different conditions were used: Untreated cells (negative
3 control), cells exposed to LPS at 100 ng/mL (positive control), and cells
4 exposed to LPS at 100 ng/mL + different treatments (HPI, HPH20A, and
5 HPH60A+15AF) at 50 or 100 μ g/mL. LPS (*E. coli* 055: B5) (Sigma-Aldrich) was
6 added for 1 h and then the hemp product. After incubation for an additional
7 period of 24 h, cells were collected for RNA extraction.

8 **Cell Viability Assay**

9 BV-2 microglial cells were incubated with different concentrations of hemp
10 products in 96-well plates (1×10^5 cells/well) during 24 h. Concentrations used
11 were the following: 10, 25, 50, 100, and 200 μ g/mL. Afterwards, the MTT
12 solution (Sigma-Aldrich) was incubated in the well until a purple precipitate was
13 visible. MTT-formazan crystals were solubilized with DMSO (Sigma-Aldrich),
14 and then measured with a microplate reader at 570 nm corrected to 650 nm.³⁰
15 Cell survival was expressed as the percentage of absorbance compared with
16 that obtained in control, non-treated cells.

17 **RNA Isolation and Real-Time quantitative PCR Analysis**

18 RNA from BV-2 microglial cells was isolated to quantify gene expression by RT-
19 qPCR. Total RNA was extracted by using TRIsure Reagent (Bioline). RNA
20 quality was assessed by A260/A280 ratio in a NanoDrop ND-1000
21 Spectrophotometer (ThermoFisher Scientific, Madrid, Spain). Briefly, RNA (250
22 ng) was subjected to reverse transcription (iScript, Bio-Rad). An amount of 40
23 ng of the resulting cDNA was used as template for qRT-PCR amplifications.
24 The mRNA levels for specific genes were determined in a CFX96 system (Bio-

Rad). For each PCR reaction, cDNA template was added to Brilliant SYBR green QPCR Supermix (Bio-Rad) containing the primer pairs for either gene or for glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and hypoxanthine phosphoribosyltransferase (HPRT) as housekeeping genes. All amplification reactions were performed in triplicate and average threshold cycle (Ct) numbers of the triplicates were used to calculate the relative mRNA expression of candidate genes. The magnitude of change of mRNA expression for candidate genes was calculated by using the standard $2^{-(\Delta\Delta Ct)}$ method. All data were normalized to endogenous reference (HPRT) gene content and expressed as relative fold-change of control. The sequences of the designed oligonucleotides are shown in **Supplementary Table 1**.

Statistical analysis

All values are expressed as arithmetic means $\pm$ standard deviations (SD). Data were evaluated with Graph Pad Prism Version 5.01 software (San Diego, CA, USA). The statistical significance of any difference in each parameter among the groups was evaluated by one-way analysis of variance (ANOVA), following Tukey multiple comparisons test as post hoc test. P values less than 0,05 were considered statistically significant.

Results and discussion

Chemical characterization of hemp products

Hemp has been cultivated since ancient times to produce fiber and seeds. Nowadays, seeds are essentially used for extracting high ω -3 fatty acids-containing oil.² The HDF, a by-product of hemp industry from oil extraction,

contains high amount of fibers and proteins. Hemp seeds were defatted obtaining HDF with a protein content of 33.3%, as reported in **Table 1**. HPI was separated in 13.8% yield from defatted hemp seed flour by basic extraction followed by acidic precipitation and the obtained HPI had 96.5% protein content. The protein content in our study was higher than previously described for HDF and HPI.¹

Amino acidic composition of hemp products

Amino acid profile might influence to peptide activity in hemp products. Hence, this study analyzed the amino acid profile in HDF, HPI, and two HPHs: HPH20A and HPH60A+15AF (**Table 2**). All these hemp products had similar amino acid profile and have been identified as a great source of essential amino acids. Briefly, hemp products were rich in negatively charged amino acid like Asp + Asn (11-12%) and Glu + Gln (18%), and also rich in Arg (12-13%). These amino acid profiles have some similarities with the amino acid profile of edestin, the main globulin protein in hemp seed, reported in the literature.³¹ The importance of these amino acids, Asp and Glu, are that they are involved in electron donation mechanism between reactive oxygen species and Arg for nitric oxide (NO) production,³² which may have contributed to regulate the hemp protein products bioactivity. Among the differences between hemp products and edestin amino acid profiles it turns out to appear a higher percentage of apolar amino acids in edestin than in HPI and HPHs, as a result, it could indicate that HPI and HPHs arises better solubility properties.^{1,7,10,}

Analysis of hydrolysis degree and molecular profile of hemp products

[View Article Online](#)

DOI: 10.1039/C9FO01904A

Extent of hydrolysis can be estimated from the DH values, which can be used an indication of peptide chain length, higher and lower values indicate mean shorter and longer lengths, respectively. Previous results indicated that the use of a single enzyme, Alcalase or Flavourzyme, led to the production of protein hydrolysates with HD < 30%, where peptides of 3–20 kDa are predominant.³³ However, most bioactive peptides described up to date have a molecular weight under 1 kDa. In this context, the extensive hydrolysis of HPI was performed using both enzymes in a sequential way as depicted in **Figure 1**. Herein, HPI was hydrolyzed for 1 h with Alcalase, which facilitated the subsequent action of Flavourzyme. The rate of hydrolysis was fast during the first 30 min of hydrolysis with Alcalase and remained almost stable in the following 30 min (approximately 30% of HD). The addition of Flavourzyme led to a new increase of HD up to 60% after 75 min with Alcalase plus 15 min with Flavourzyme (HPH60A+15AF).

As shown in **Figure 2**, the peptide sizes in hemp products were estimated by FPLC and the results showed distribution into five major peaks with approx. MW ranging from 0.3 (peak e) to 300 kDa (peak a). These bigger peptide peaks (peaks a-c) were most remarkable for HPI. The hydrolysis with Alcalase and Flavourzyme contained a minor peak with size 1.45 kDa (peak d), which indicates a polypeptide that is resistant to digestion by these enzymes. In addition, sequential hydrolysis with Alcalase and Flavourzyme released an oligopeptide with approx. MW ranging from 300–400 Da. The interrelationship between the MWs of hydrolysates and their bioactivities in human health application is very crucial in functional foods and nutraceutical formulations.

Thus, the presence of low molecular weight peptides enhances the potential amino acid absorption and their ability to have bioactive effects during *in vivo* tests.¹⁰

Cell-free *in vitro* evaluation of hemp product biological activity

Biological activity of peptides is related to chain length, HD, amino acid composition, sequence, and conformation.³⁴ Some literature evidence indicates that the most effective hemp products are ACE-inhibitory peptides identified up-to-now contain 2–20 amino acids and must have a good hydrophobicity/hydrophilicity balance and some particular structural characteristic related to its sequence.⁸⁻¹¹ Herein, we evaluated the potential antioxidant activity of HPI, five HPHs with Alcalase, and five HPHs with Alcalase and Flavourzyme (**Figure 3**) in cell-free *in vitro* experiments. The DPPH radicals have been widely used to investigate the scavenging activity of protein hydrolysates. As shown in **Figure 3A**, all HPHs tested showed DPPH radical scavenging activity. Furthermore, all hydrolysates had higher activity than HPI, so it maybe concluded that hydrolysis generated peptides that could react with free radicals to convert them to more stable products and complete the radical chain reaction. Hydrolysis with Alcalase raised up DPPH scavenging effect to 21-36%. The subsequent hydrolysis with Alcalase and Flavourzyme slightly increased the DPPH scavenging activity even more, obtaining values of approximately 40%. Similar results were founded in HPHs obtained by different proteases (AFP, HT, ProG, actinidin and zingibain) by Teh et al.³⁵ A previous study documented that the smaller size peptide fraction (<1 kDa) of the pepsin hydrolysate – followed by hydrolysis with pancreatin, had a higher DPPH

scavenging activity (25%) than a larger sized peptide fraction (5–10 kDa) (20%), while the unfractionated HPI had less than 5% DPPH scavenging activity.¹⁴ As compared to our study here, HPHs hydrolyzed with Alcalase and Flavourzyme had higher DPPH scavenging activity (40%). The FRAP was used in order to found a direct correlation between antioxidant activity and the donate-electron potential of bioactive compounds into HPI and HPHs as other studies have reported in similar seeds hydrolysates.³⁶ For FRAP value (**Figure 3B**), significant differences were observed for the HPHs with Alcalase compared to those hydrolyzed with Alcalase and Flavourzyme. HPHs hydrolyzed with Alcalase and Flavourzyme had higher FRAP value compared to those hydrolyzed with alcalase. All peptide fractions had significantly lower ($p < 0.05$) absorption values when compared to BHT action. Similar results were founded in HPHs obtained by reverse-phase HPLC by Girgih et al.¹³

Neuroprotective properties of hemp protein hydrolysates

Oxidative stress has been associated with chronic inflammation in neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis, which increasing numbers of elderly individuals suffer.¹⁸ The brain is particularly vulnerable to oxidative stress and damage, because of its high oxygen consumption, low antioxidant defenses, and high content of polyunsaturated fats very prone to be oxidized.³⁷ Thus, it is not surprising the importance of protecting systems, including antioxidant molecules, to maintain brain homeostasis. After analyzing the previous results, we decided to select two hydrolysates considering the HD, hydrolysis time, and antioxidant activity. The HPHs were HPH20A and HPH60A+15AF. At first, to evaluate the cytotoxic effects of hemp products on BV-2 microglia cell viability,

1 the cells were treated with HPI, HPH20A, and HPH60A+15AF up to 200 µg/mL
2 for 24 h. The cell viability evaluated by the MTT method was not affected by any
3 hemp product concentration (data not shown). In order to evaluate the
4 neuroprotective properties of HPHs, gene expression of cytokines TNF-α, IL-1β,
5 IL-6, and IL-10 was measured by RT-qPCR in BV-2 microglial cells (**Figure 4**).
6 As expected, LPS increased mRNA levels of TNF-α, IL-1β, and IL-6.
7 Meanwhile, these LPS-induced changes were blocked in the presence of
8 HPH20A and HPH60A+15AF at 50 or 100 µg/mL. HPI were also effective to
9 down-regulate the expression levels of TNF-α and IL-6. Hemp products also
10 showed a statistically significant dose-dependent effect on IL-6 gene
11 expression. Furthermore, HPH20A and HPH60A+15AF induced a increase in
12 mRNA levels of anti-inflammatory IL-10 gene at 100 µg/mL. In the process of
13 developing new strategies to modulate the inflammatory response in
14 neurodegenerative diseases, a number of phytochemicals, mainly phenolic
15 compounds, have been shown to target neuroinflammation in Alzheimer's
16 disease due to their anti-oxidant and anti-inflammatory activities,³⁸ however,
17 some of the benefits of these natural products were further dependent on their
18 poor bioavailability or activity when metabolically transformed into conjugates.
19 One promising attempt for an improved and safety therapy of
20 neurodegenerative diseases by naturally occurring compounds has been
21 recently reported with the soy-derived peptide lunasin.³⁹ Inspired in part by
22 these studies, we envisioned that HPHs might be an ideal candidate to test its
23 efficacy in mitigating a neuroinflammatory environment. To date there is little if
24 any studies on the impact of HPHs on microglial activation. The present study
25 contributes to establishing novel data for a hemp seed global database of

1 bioactive constituents. Based on previous studies, hemp seeds contain a highly
2 amount of bioactive chemicals as canabisin-F or grossamide (lignanamides),
3 which improve neuroinflammatory states in LPS-stimulated BV-2 microglial cells
4 through down-regulation of TLR-4 mediated NF- κ B pathway.^{40,41} In particular,
5 this study showed for the first time that the treatment with hemp protein
6 products obtained by enzymatic digestion decreased inflammatory state in LPS-
7 stimulated BV-2 microglial cells. However, the study had limitations. First, the
8 responsible bioactive peptides were not identified in this work and second, the
9 study on activity-structure relation is missing. These two limitations of this study
10 can be future research directions.

11 To conclude, the initial HPI has limited bioactive properties as measured by
12 DPPH scavenging and FRAP activities. The bioactivity of the HPI was
13 increased upon hydrolysis by Alcalase and Flavourzyme, obtaining HPHs with
14 small peptides that exert bioactivities. Our findings advocate to considerate that
15 HPHs may inhibit oxidative stress and inflammatory response in activated
16 microglia. HPHs should be the focus of future analysis of the bioactive peptides
17 generated in the prevention or treatment of neurodegenerative diseases,
18 confirming that hemp seeds are a valuable source of bioactive compounds.

19 **Acknowledgements**

20 This work was supported by grant CYTED-2019 119RT0567 from the Spanish
21 Ministry of Science, Innovation, and Universities. Authors thank Cell Biology
22 Unit of Instituto de la Grasa for its assistance during the fulfilment of this work.
23 SMP acknowledges financial support from “V Own Research Plan” (University
24 of Seville).

1 **Conflicts of interest**

[View Article Online](#)
DOI: 10.1039/C9FO01904A

- 2 All the authors declare that they do not have conflicts of interest.

References

View Article Online
DOI: 10.1039/C9FO01904A

1. G. Mamone, G. Picariello, A. Ramondo, M. A. Nicolai, et al., Production, digestibility and allergenicity of hemp (*Cannabis sativa* L.) protein isolates, *Food Res. Int.*, 2019, **115**, 562–571.
2. B. D. Oomah, M. Busson, D. V. Godfrey and J. C. G. Drover, Characteristics of Hemp (*Cannabis sativa* L.) seed oil, *Food Chem.*, 2002, **76**, 33–43.
3. E. Vonapartis, M. P. Aubin, P. Seguin, A. F. Mustafa, et al., Seed composition of ten industrial hemp cultivars approved for production in Canada, *J. Food Compos. Anal.*, 2015, **39**, 8–12.
4. G. Aiello, E. Fasoli, G. Boschini, C. Lammi, et al., Proteomic characterization of hempseed (*Cannabis sativa* L.), *J. Proteomics*, 2016, **147**, 187–196.
5. S. Montserrat-de la Paz, F. Marín-Aguilar, M. D. García-Giménez and M. A. Fernández-Arche, Hemp (*Cannabis sativa* L.) seed oil: analytical and phytochemical characterization of the unsaponifiable fraction, *J. Agric. Food Chem.*, 2014, **62**, 1105–1110.
6. S. A. Malomo, R. He and R. E. Aluko, Structural and functional properties of hemp seed protein products, *J. Food Sci.*, 2014, **79**, C1512–C1521.
7. C. H. Tang, Z. Ten, X. S. Wang and X. Q. Yang, Physicochemical and functional properties of hemp (*Cannabis sativa* L.) protein isolate, *J. Agric. Food Chem.*, 2006, **54**, 8945–8950.
8. L. P. Orio, G. Boschini, T. Recca, C. F. Morelli, et al., New ACE-inhibitory peptides from hemp seed (*Cannabis sativa* L.) proteins, *J. Agric. Food Chem.*,

- 1 2017, **65**, 10482–10488.
- 2 9. G. Aiello, C. Lammi, G. Boschini, C. Zanoni, et al., Exploration of potentially
3 bioactive peptides generated from the enzymatic hydrolysis of hempseed
4 proteins, *J. Agric. Food Chem.*, 2017, **65**, 10174–10184.
- 5 10. S. A. Malomo, J. O. Onuh, A. T. Girgih and R. E. Aluko, Structural and
6 antihypertensive properties of enzymatic hemp seed protein hydrolysates,
7 *Nutrients*, 2015, **7**, 7616–7632.
- 8 11. A. T. Girgih, A. Alashi, R. He, S. Malomo, et al., Preventive and treatment
9 effects of a hemp seed (*Cannabis sativa* L.) meal protein hydrolysate against
10 high blood pressure in spontaneously hypertensive rats, *Eur. J. Nutr.*, 2014, **53**,
11 1237–1246.
- 12 12. C. Zanoni, G. Aiello, A. Arnoldi and C. Lammi, Hempseed peptides exert
13 hypocholesterolemic effects with a statin-like mechanism, *J. Agric. Food Chem.*,
14 2017, **65**, 8829–8838.
- 15 13. A. T. Girgih, C. C. Udenigwe and R. E. Aluko, Reverse-phase HPLC
16 separation of hemp seed (*Cannabis sativa* L.) protein hydrolysate produced
17 peptide fractions with enhanced antioxidant capacity, *Plant Foods Hum. Nutr.*,
18 2013, **68**, 39–46.
- 19 14. A. Girgih, A. Alashi, R. He, S. Malomo, et al., A Novel hemp seed meal
20 protein hydrolysate reduces oxidative stress factors in spontaneously
21 hypertensive rats, *Nutrients*, 2014, **6**, 5652–5666.
- 22 15. J. A. Kabba, Y. Xu, H. Christian, W. Ruan, et al., Microglia: Housekeeper of
23 the Central Nervous System, *Cell. Mol. Neurobiol.*, 2018, **38**, 53–71.

- 1 16. Z. Shen, X. Bao and R. Wang, Clinical PET Imaging of Microglial Activation, *Front. Aging*
2 Implications for Microglial Therapeutics in Alzheimer's Disease, *Front. Aging*
3 *Neurosci.*, 2018, **10**, 314. View Article Online
DOI: 10.1039/C9FO01904A
- 4 17. Z. Cai, M. D. Hussain and L. J. Yan, Microglia, neuroinflammation, and
5 beta-amyloid protein in Alzheimer's disease, *Int. J. Neurosci.*, 2014, **124**, 307-
6 321.
- 7 18. R. M. Ransohoff, How neuroinflammation contributes to neurodegeneration,
8 *Science*, 2016, **353**, 777-783.
- 9 19. H. Lqari, J. Vioque, J. Pedroche and F. Millán, *Lupinus angustifolius* protein
10 isolates: Chemical composition, functional properties and protein
11 characterization, *Food Chem.*, 2002, **76**, 349–356.
- 12 20. L. Prosky, N. G. Asp, T. F. Schweizer, J. W. DeVries, et al., Determination
13 of insoluble, soluble, and total dietary fiber in foods and food products:
14 Interlaboratory study, *J. Assoc. Off. Anal. Chem.*, 1988, **71**, 1017–1023.
- 15 21. R. G. Moores, D. L. McDermott and T. R. Wood, Determination of
16 chlorogenic acid in coffee, *Anal. Chem.*, 1948, **20**, 620–624.
- 17 22. M. DuBois, K. A. Gilles, J. K. Hamilton, P. A. Rebers, et al., Colorimetric
18 method for determination of sugars and related substances, *Anal. Chem.*, 1956,
19 **28**, 350–356.
- 20 23. M. Alaiz, J. L. Navarro, J. Girón and E. Vioque, Amino acid analysis by high-
21 performance liquid chromatography after derivatization with diethyl
22 ethoxymethylenemalonate, *J. Chromatogr. A.*, 1992, **591**, 181–186.

24. M. M. Yust, J. Pedroche, J. Girón-Calle, J. Vioque, et al., Determination of tryptophan by high-performance liquid chromatography of alkaline hydrolysates with spectrophotometric detection, *Food Chem.*, 2004, **85**, 317–320.
25. J. Adler-Nissen, Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzenesulfonic acid, *J. Agric. Food Chem.*, 1979, **27**, 1256–1262.
26. H. C. Wu, H. M. Chen and C. Y. Shiau, Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber australasicus*), *Food Res. Int.*, 2003, **36**, 949–957.
27. M. Oyaizu, Studies on Products of Browning Reaction. Antioxidative activities of products of browning reaction prepared from glucosamine *Japanese, J. Nutr. Diet.*, 1986, **44**, 307–315.
28. T. Masuda and M. Prinz, Microglia: A Unique Versatile Cell in the Central Nervous System, *ACS Chem. Neurosci.*, 2016, **7**, 428-434.
29. N. Lannes, E. Eppler, S. Etemad, P. Yotovski, et al., Microglia at center stage: a comprehensive review about the versatile and unique residential macrophages of the central nervous system, *Oncotarget*, 2017, **8**, 114393-114413.
30. S. Montserrat-de la Paz, M. C. Naranjo, S. Lopez, R. Abia, et al., Niacin and its metabolites as master regulators of macrophage activation, *J. Nutr. Biochem.*, 2017, **39**, 40-47.
31. T. Docimo, I. Caruso, E. Ponzoni, M. Mattana, et al., Plant physiology and biochemistry molecular characterization of Edestin gene family in *Cannabis*

- 1 *sativa* L., *Plant Physiol. Biochem.*, 2014, **84**, 142–148.
- 2 32. J. Moon, O. Y. Kim, G. Jo, and M. J. Shin, Alterations in Circulating Amino
3 Acid Metabolite Ratio Associated with Arginase Activity Are Potential Indicators
4 of Metabolic Syndrome: The Korean Genome and Epidemiology Study,
5 *Nutrients*, 2017, **9**, 740.
- 6 33. A. Clemente, J. Vioque, R. Sanchez-Vioque, J. Pedroche, et al., Production
7 of extensive chickpea (*Cicer arietinum* L.) protein hydrolysates with reduced
8 antigenic activity, *J. Agric. Food Chem.*, 1999, **47**, 3776–3781.
- 9 34. C. Daskaya-Dikmen, A. Yucetepe, F. Karbancioglu-Guler, B. Ozcelik, et al.,
10 Angiotensin-I-Converting Enzyme (ACE)-Inhibitory peptides from plants,
11 *Nutrients*, 2017, **9**, 316.
- 12 35. S. S. The, A. L. D. A. Bekhit, A. Carne and J. Birch, Antioxidant and ACE-
13 inhibitory activities of hemp (*Cannabis sativa* L.) protein hydrolysates produced
14 by the proteases AFP, HT, Pro-G, actinidin and zingibain, *Food Chem.*, 2016,
15 **203**, 199-206.
- 16 36. A. Connolly, M. Cermeño, D. Crowley, Y. O'Callaghan, et al.,
17 Characterisation of the in vitro bioactive properties of alkaline and enzyme
18 extracted brewers' spent grain protein hydrolysates, *Food Res. Int.*, 2019, **121**,
19 524-532.
- 20 37. G. Cenini, A. Lloret and R. Cascella, Oxidative Stress in Neurodegenerative
21 Diseases: From a Mitochondrial Point of View, *Oxid. Med. Cell Longev.*, 2019,
22 **2019**, 2105607.
- 23 38. Y. Sawikr, N. S. Yarla, I. Peluso, M. A. Kamal, et al., Neuroinflammation in

- 1 Alzheimer's Disease: The Preventive and Therapeutic Potential of Polyphenolic
2 Nutraceuticals, *Protein Chem. Struct. Biol.*, 2017, **108**, 33-57.
- 3 39. A. Sarkar, N. Gogia, N. Glenn, A. Singh, et al., A soy protein Lunasin can
4 ameliorate amyloid-beta 42 mediated neurodegeneration in *Drosophila* eye, *Sci.*
5 *Rep.*, 2018, **8**, 13545.
- 6 40. Q. Luo, X. Yan, L. Bobrovskaya, M. Ji, et al., Anti-neuroinflammatory effects
7 of grossamide from hemp seed via suppression of TLR-4-mediated NF-KB
8 signaling pathways in lipopolysaccharide-stimulated BV2 microglia cells, *Mol.*
9 *Cell. Biochem.*, 2017, **428**, 129–137.
- 10 41. S. Wang, Q. Luo and P. Fan, Cannabisin F from hemp (*Cannabis sativa*)
11 seed suppresses lipopolysaccharide-induced inflammatory responses in BV2
12 microglia as SIRT1 modulator, *Int. J. Mol. Sci.*, 2019, **20**, 507.
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21

Figure legend

View Article Online
DOI: 10.1039/C9FO01904A

Figure 1. Time-course of hydrolysis degree of HPI during enzymatic hydrolysis with Alcalase at 10 (HPH10A), 20 (HPH20A), 30 (HPH30A), 45 (HPH45A), and 60 (HPH60A) min, and with Flavourzyme at 15 (HPH60A+15AF), 30 (HPH60A+30AF), 60 (HPH60A+60AF), 90 (HPH60A+90AF), and 120 (HPH60A+120AF) min. Data, expressed as the percentage of peptide bonds cleaved, are mean $\pm$ standard deviation of three determinations.

Figure 2. Molecular weight (MW) profiles by size-exclusion FPLC of HPI, HPH20A, and HPH60A+15AF. MW ranging from 0.3 (peak e) to 300 kDa (peak a). The bigger peptide peaks (peaks a-c) were most remarkable for HPI.

Figure 3. DPPH radical scavenging activity (**A**) and ferric reducing antioxidant power (FRAP) (**B**) by HPI and HPHs obtained using Alcalase and Flavourzyme in cell-free *in vitro* experiments. Values are presented as means $\pm$ SD ($n = 3$) and those marked with different letters are significantly different ($p < 0.05$).

Figure 4. Gene expression of TNF α (**A**), IL-1 β (**B**), IL-6 (**C**), and IL-10 (**D**) in BV-2 microglial cells after 24 h incubation with or without LPS (100 ng/mL) and HPI, HPH20A, and HPH60A+15AF at 50 and 100 μ g/mL. Values are presented as means $\pm$ SD ($n = 3$) and those marked with different letters are significantly different ($p < 0.05$).

Table 1. Chemical composition of HDF and HPI. Data, expressed as percentage in dry basis, are mean $\pm$ standard deviation of three determinations.

Fat (seed) (%)	32.35 $\pm$ 0.57	
	HDF	HPI
Proteins (%)	33.28 $\pm$ 1.39	96.46 $\pm$ 0.93
Moisture (%)	6.18 $\pm$ 0.02	2.19 $\pm$ 0.01
Ash (%)	7.01 $\pm$ 0.11	1.06 $\pm$ 0.10
Fiber (%)	40.90 $\pm$ 1.98	0.00 $\pm$ 0.00
Sugar (%)	5.24 $\pm$ 1.05	0.00 $\pm$ 0.00
Polyphenols (%)	0.48 $\pm$ 0.03	0.04 $\pm$ 0.00
Fat (%)	---	0.25 $\pm$ 0.00

- 1 **Table 2.** Amino acid composition of hemp products. Data, expressed as
 2 percentage of amino acids on total amino acid content, are mean $\pm$ standard
 3 deviation of three determinations.

Aa (%)	HDF	HPI	HPH20A	HPH 60A+15AF	FAO/ WHO
Asp + Asn	11.34 $\pm$ 0.37	12.35 $\pm$ 0.34	12.40 $\pm$ 0.29	12.08 $\pm$ 0.51	
Glu + Gln	18.41 $\pm$ 0.13	18.86 $\pm$ 0.35	18.83 $\pm$ 0.25	18.58 $\pm$ 0.12	
Ser	5.57 $\pm$ 0.11	6.31 $\pm$ 0.21	6.36 $\pm$ 0.19	6.31 $\pm$ 0.22	
His	3.07 $\pm$ 0.01	2.85 $\pm$ 0.05	2.80 $\pm$ 0.02	2.79 $\pm$ 0.03	1.9
Gly	4.74 $\pm$ 0.07	4.79 $\pm$ 0.23	4.64 $\pm$ 0.11	4.64 $\pm$ 0.12	
Thr	3.79 $\pm$ 0.08	3.79 $\pm$ 0.10	3.89 $\pm$ 0.06	4.02 $\pm$ 0.07	3.4
Arg	12.68 $\pm$ 0.21	13.44 $\pm$ 0.03	13.34 $\pm$ 0.24	13.78 $\pm$ 0.23	
Ala	4.89 $\pm$ 0.10	5.01 $\pm$ 0.14	4.99 $\pm$ 0.10	5.00 $\pm$ 0.11	
Pro	2.66 $\pm$ 0.69	3.02 $\pm$ 0.00	2.83 $\pm$ 0.05	1.29 $\pm$ 0.19	
Tyr	3.57 $\pm$ 0.06	4.08 $\pm$ 0.06	3.98 $\pm$ 0.03	4.06 $\pm$ 0.12	
Val	5.27 $\pm$ 0.11	3.77 $\pm$ 0.40	4.05 $\pm$ 0.42	4.71 $\pm$ 0.19	3.5
Met	2.14 $\pm$ 0.02	0.89 $\pm$ 0.34	0.60 $\pm$ 0.03	0.64 $\pm$ 0.04	2.5 ^a
Cys	1.22 $\pm$ 0.02	1.98 $\pm$ 0.08	1.81 $\pm$ 0.18	1.65 $\pm$ 0.21	
Ile	4.23 $\pm$ 0.08	3.02 $\pm$ 0.40	3.57 $\pm$ 0.15	3.85 $\pm$ 0.20	2.8
Trp	0.76 $\pm$ 0.05	0.55 $\pm$ 0.02	0.66 $\pm$ 0.01	0.82 $\pm$ 0.03	1.1
Leu	7.12 $\pm$ 0.05	7.13 $\pm$ 0.11	7.12 $\pm$ 0.13	7.34 $\pm$ 0.07	6.6
Phe	4.76 $\pm$ 0.09	5.12 $\pm$ 0.02	5.04 $\pm$ 0.05	5.19 $\pm$ 0.08	6.3 ^b
Lys	3.78 $\pm$ 0.09	3.04 $\pm$ 0.08	3.10 $\pm$ 0.06	3.23 $\pm$ 0.04	5.8

4 ^aMet + Cys, ^bPhe + Tyr.

5

Figure 1

View Article Online
DOI: 10.1039/C9FO01904A

Figure 2

View Article Online
DOI: 10.1039/C9FO01904A

Figure 3

View Article Online
DOI: 10.1039/C9FO01904A

Figure 4

View Article Online
DOI: 10.1039/C9FO01904A