Dear Author,

Please correct your galley proofs carefully and return them no more than four days after the page proofs have been received.

Please limit corrections to errors already in the text; cost incurred for any further changes or additions will be charged to the author, unless such changes have been agreed upon by the editor.

The editors reserve the right to publish your article without your corrections if the proofs do not arrive in time.

Note that the author is liable for damages arising from incorrect statements, including misprints.

Please note any queries that require your attention. These are indicated with a Q in the PDF and a question at the end of the document.

Reprints may be ordered by filling out the accompanying form.

Return the reprint order form by fax or by e-mail with the corrected proofs, to Wiley-VCH : advhealthmat@wiley.com

Corrections should be made directly in the PDF file using the PDF annotation tools. If you have questions about this, please contact the editorial office. The corrected PDF and any accompanying files should be uploaded to the journal's Editorial Manager site.

To avoid commonly occurring errors, **please ensure that the following important items are correct** in your proofs (please note that once your article is published online, no further corrections can be made):

- **Names** of all authors present and spelled correctly
- **Titles** of authors correct (Prof. or Dr. only: please note, Prof. Dr. is not used in the journals)
- **Addresses** and **postcodes** correct
- **E-mail address** of corresponding author correct (current email address)
- **Funding bodies** included and grant numbers accurate
- **Title** of article OK
- All **figures** included
- **Equations** correct (symbols and sub/superscripts)

Author Query Form

WILEY

Journal ADHM

Article adhm201800335

Dear Author,

During the copyediting of your manuscript the following queries arose.

Please refer to the query reference callout numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools.

Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

Query No.	Description	Remarks
Q1	Please provide TOC keyword.	
Q2	Please confirm that forenames/given names (blue) and surnames/family names (vermilion) have been identified correctly.	
Q3	Please provide the highest academic title (either Dr. or Prof.) for all authors, where applicable.	
Q4	Please provide the zip code for all affiliations.	
Q5	Please shorten Table of Contents text to a maximum of 60 words. All abbreviations should be defined.	
Q6	Refs. have been renumbered as experimental section has been moved at the end of section. Please check for correctness.	
Q7	Experimental section has been moved at the end of section as per journal style and subsequent heading level has been renumbered. Please check for correctness.	
Q8	Please provide the significance of * as given in the table body.	
Q9	Refs. (2 and 63, 24 and 64, and 62 and 65) were identical, the later instances have been deleted and renumbered the remaining. Please validate.	
Q10	Please provide the book title, publisher and its location in ref. (29).	
Q11	Book title has been updated in ref. (49). Please check for correctness.	
Q12	Please provide the publisher location in ref. (57).	

Author: Please confirm that Funding Information has been identified correctly.

Please confirm that the funding sponsor list below was correctly extracted from your article: that it includes all funders and that the text has been matched to the correct FundRef Registry organization names. If a name was not found in the FundRef registry, it may not be the canonical name form, it may be a program name rather than an organization name, or it may be an organization not yet included in FundRef Registry. If you know of another name form or a parent organization name for a "not found" item on this list below, please share that information.

FundRef Name	FundRef Organization Name
MINECO/FEDER	
Instituto de Salud Carlos III	Instituto de Salud Carlos III
VI National R&D&I Plan	
Instituto de Salud Carlos III	Instituto de Salud Carlos III
MINECO	Secretaría de Estado de Investigación, Desarrollo e Innovación
Agència de Gestió d'Ajuts Universitaris i de Recerca	

XXXX

C. Fornaguera, M. Guerra-Rebollo,
M. Ángel Lázaro, C. Castells,
O. Meca-Cortés, V. Ramos,
A. Cascante, N. Rubio, J. Blanco,
S. Borrós* 1800335

mRNA Delivery System for Targeting Antigen-Presenting Cells In Vivo

Oligopeptide end-modified poly (β -amino esters) (OM-PBAEs) are complexed with mRNA and form discrete nanoparticles. It is demonstrated that the selection of the appropriate end-oligopeptide modifications enables the specific targeting and major transfection of antigen-presenting cells (APC) in vivo. Therefore, it can be confirmed that OM-PBAE are appropriate for the design of mRNA-based immunotherapy approaches aimed to in vivo transfect APCs.

Q5

mRNA Delivery System for Targeting Antigen-Presenting Cells In Vivo

Cristina Fornaguera, Marta Guerra-Rebollo, Miguel Ángel Lázaro, Cristina Castells, Oscar Meca-Cortés, Victor Ramos, Anna Cascante, Núria Rubio, Jerónimo Blanco, and Salvador Borrós*

The encapsulation of mRNA in nanosystems as gene vaccines for immunotherapy purposes has experienced an exponential increase in the recent years. Despite the many advantages envisaged within these approaches, their application in clinical treatments is still limited due to safety issues. These issues can be attributed, in part, to liver accumulation of most of designed nanosystems and to the inability to transfect immune cells after an intravenous administration. In this context, this study takes advantage of the known versatile properties of the oligopeptide end-modified poly (β -amino esters) (OM-PBAEs) to complex mRNA and form discrete nanoparticles. Importantly, it is demonstrated that the selection of the appropriate end-oligopeptide modifications enables the specific targeting and major transfection of antigen-presenting cells (APC) in vivo, after intravenous administered, thus enabling their use for immunotherapy strategies. Therefore, with this study, it can be confirmed that OM-PBAE are appropriate systems for the design of mRNA-based immunotherapy approaches aimed to in vivo transfect APCs and trigger immune responses to fight either tumors or infectious diseases.

1. Introduction

Nanomedicines can be defined as any entity showing at least one dimension at the nanoscale, developed for medical purposes: diagnosis and/or treatment.^[1,2] They appeared some decades

ago with the aim to become novel therapies for unmet needs and, since then, they have experienced an exponential increase. Among the multiple applications of nanomedicine, its use for gene therapy purposes has been in the forefront of research laboratories. Gene therapy has the potential to treat current diseases in need of improved therapies; it aims to treat inheritable diseases such as hemophilia and cystic fibrosis, but also acquired diseases such as cancer.^[3–5] Specifically, their use as vaccines to become novel immunotherapies for cancer or infectious diseases is of utmost relevance. The first attempts of cancer vaccination consisted on the use of viral vectors, due to virus evolutionary design to transfer their genetic information to cells, thus showing a high efficiency in cell transduction.^[3] However, important drawbacks appeared soon. Apart from the limited encapsulating capacity, problems concerning safety issues aroused

together with an unspecific activation of the immune system, which ends up to the elimination of the exogenous elements and does not enable repeated administration. These problems, together with others associated with insertional mutagenesis, limit the use of viral vectors in humans.^[3–8] For these reasons, the use of nonviral synthetic vectors was presented as a safer alternative.^[5] Problems related with undesired adaptive immune response are not expected, so these vectors could be readministered, and they can be produced using more cost-effective and easy to scale-up methods, compared to viral vectors.^[3–5,9–15] Although lipid carriers have been the most widely studied so far, they aroused some toxicity problems related, mostly, with liver accumulation.^[7,12,16] In contrast, polymeric nanoparticles represent a promising choice for many reasons, such as the protection and controlled release of the gene material, the easy, versatile and tailored production and their biocompatibility and biodegradability.^[7,9,17] Poly(β -amino esters) (PBAE) are a class of biocompatible and biodegradable polymers, composed of ester bonds, easily degradable in physiological conditions. They were previously demonstrated to self-assemble in aqueous media forming nanometric discrete structures that enhance the transfection of encapsulated genetic material.^[7,12,15,18,19] In addition, they can be designed to increase their buffering capacity, and their ability to compact nucleic acids improving their capacity to promote cellular internalization.

C. Fornaguera, M. Ángel Lázaro, C. Castells, A. Cascante, S. Borrós
Sagetis Biotech SL
Barcelona, Spain
E-mail: salvador.borros@iqs.url.edu

C. Fornaguera, V. Ramos, A. Cascante, S. Borrós
Grup d'Enginyeria de Materials (GEMAT)
Institut Químic de Sarrià (IQS)
Universitat Ramon Llull (URL)
Barcelona, Spain

M. Guerra-Rebollo, O. Meca-Cortés, N. Rubio, J. Blanco, S. Borrós
CIBER of Biomaterials
Bioengineering and Nanomedicine (CIBER-BBN)
Barcelona, Spain

M. Guerra-Rebollo, N. Rubio, J. Blanco
Institute of Advanced Chemistry of Catalonia (IQAC-CSIC)
Barcelona, Spain

The ORCID identification number(s) for the author(s) of this article can be found under <https://doi.org/10.1002/adhm.201800335>.

DOI: 10.1002/adhm.201800335

Once inside the endosomes, thanks to the proton sponge effect, endosomal escape, required for the expression of the encoded gene, is also enhanced.^[19] PBAEs can be synthesized varying their structural components (e.g., backbone, side chain, and terminal groups), tailored to achieve different features. Despite their proven capacity to encapsulate gene material and transfect cells, some authors described their difficulty to efficiently encapsulate gene material.^[19] For this reason, and in order to improve the gene material complexation, in our group, libraries of oligopeptide end-modified PBAEs (OM-PBAEs) were recently designed with the aim to have more hydrophilic and cationic structures that could easily encapsulate genetic material by electrostatic interactions, but at the same time, maintaining the capacity to release the genetic material in the cell cytoplasm.^[10,12,20,21] A combination of two kinds of oligopeptide end modifications was previously demonstrated to be more efficient in gene transfection. While lysine (K)-modified PBAE have demonstrated enhanced interaction to cell membranes,^[22] histidine (H)-modified PBAEs are used due to histidine buffering capacity at pH $\approx 5-6$,^[23,24] which is required for the endosomal escape and penetration to the cytoplasm, where mRNA can be translated to the protein it codifies for.

OM-PBAEs can encapsulate any type of genetic material, from plasmid DNA to siRNA and miRNA. Recently, mRNA delivery has gained a lot of attention for vaccination purposes because of its multiple advantages when compared to DNA. In particular, it has a superior safety profile, since it is easily degraded in biological environment, thus having only a transient expression and avoiding genome integration. It is able to transfect nondividing, slowly proliferating and quiescent cells. Moreover, nucleus penetration, which is a specially difficult step, is not required, as the mRNA translation takes place in the cytoplasm.^[3,11,12,18,23,25] In addition, mRNA can be produced easily under good manufacturing practices.^[7] For all these reasons, mRNA is the gene material of choice for immunotherapies, where the transfection of antigen-presenting cells (APC) is envisaged. These cells, once transfected, express the encoded protein, that can trigger specific activation of the immune system for the elimination of tumor or infected cells, together with the inflammation enhancement produced by the detection of the mRNA as an exogenous element.^[26]

However, robust, safe, and efficient delivery vectors, able to intracellularly deliver their cargos inside APC are still missing.^[27] Although much bibliography is available regarding mRNA vaccines, in most cases, the *in vivo* biodistribution represents a drawback, since most nanosystems end up in liver accumulation, thus not reaching the intended target organ and not enabling the genetic cargo to perform the therapeutic activity.^[1,2] For this reason, the main goal of this work is to design OM-PBAEs nanoparticles able to efficiently transfect APCs in the spleen after intravenous administration, while decreasing their accumulation in the liver. The achievement of this aim opens the possibility to develop novel genetic vaccines for immunotherapy approaches.^[28]

2. Results

2.1. Physicochemical Characterization of the Complexes

We have previously demonstrated that the OM-PBAE from our library are cell-specific depending on their terminal

oligopeptide composition.^[12] Bearing this in mind, preliminary *in vitro* studies were performed to determine which formulation was the most appropriate for the transfection of APCs (see Figure S1, Supporting Information). The so-called KH formulation, which consists on 60% C6CK3 plus 40% C6CH3, was selected, according to previous experiments, as the most appropriate end-oligopeptide modification to target APCs in the spleen. Freshly prepared nanoparticles were characterized by means of size and concentration (Figure 1A,B). The average hydrodynamic diameter obtained from DLS measurements was ≈ 140 nm (Figure 1A,B). Accordingly, when measured by NTA, KH nanoparticles size was similar ≈ 120 nm^[29,30] (see Figure S2, Supporting Information, for further details on NTA measures). Regarding the data obtained, it is remarkable that nanoparticles were low polydisperse samples, since PDIs were around to or lower than 0.2.^[30]

Interestingly, after freeze-drying, nanoparticles only showed a slight size increase ($\approx 10-20$ nm), while maintaining their low polydispersity (Figure 1A,B; see Figure S3, Supporting Information, for further details); therefore, they can be lyophilized to enable a long-term storage time, in terms of nanoparticle physicochemical properties. These size properties make them adequate for an intravenous administration.^[31,32]

After freeze-drying, surface charge and encapsulation efficiency were also measured to ensure that they were appropriate for intravenous administration. Surface charges were positive, as expected, due to the presence of cationic oligopeptides at the end termini of the polymers (Figure 1A,C). Nevertheless, the surface charge was not as high as other reported nanosystems in absolute value.^[6,33] Concerning mRNA entrapment efficiency, it was especially high for KH nanoparticles ($\approx 80\%$). Interestingly, for this formulation, it did not decrease significantly after lyophilization.

It is important to remark that all nanoparticle properties have been tested in, at least, three independent batches, resulting in very similar values. The reproducibility of the preparation method, an important property when thinking on translating to medical application, thus, was confirmed.

2.2. In Vitro Biocompatibility and Transfection Efficiency

The next step in the characterization of KH nanoparticles was the *in vitro* testing of their safety and efficacy transfecting different cell lines. Four different cell lines were selected as models of more (HeLa) and less permissive (ARPE-19) cell cultures. Moreover, and taking into account the aim to transfect APC, a dendritic and a macrophage cell line was also included (JAWSII and RAW264.7, respectively). First, cell viability was studied by means of *in vitro* MTT analysis of different cell culture models presenting high viability ($>80\%$) in all cases 24 h after transfection (Figure 2A), as compared to negative controls (nontreated cells; 100% viability). This result was in accordance with previous studies performed with OM-PBAEs, in which higher polymer concentrations were incubated during longer periods of time, without observing any sign of toxicity *in vitro*.^[10,12]

Concerning cell transfection efficiency, as shown in Figure 2B,C, it was dependent on the cell line, as previously stated.^[10] HeLa cells, the most permissive cell line tested, showed similar trans-

Figure 1. Nanoparticles' physicochemical characterization: A) Table with all numerical parameters analyzed; B) Hydrodynamic diameter (bars) by DLS and NTA of fresh and lyophilized NPs. PDI values are presented in dots, of fresh and lyophilized NPs; C) Surface charge of lyophilized NPs; and D) Encapsulation efficiency (in %) of NPs before and after the lyophilization process. $n = 3$. No significant differences were found in any data of B and D plots.

fection for both commercial vectors and OM-PBAE nanoparticles. However, OM-PBAE presented significant higher transfection in ARPE-19 cells (70%) when compared with both commercial vectors (20%). Most importantly, KH nanoparticles enable transfection efficiencies significantly higher than positive controls; ≈ 55 –60% for RAW264.7 and JAWSII (macrophages and dendritic cells, respectively), the two most restrictive cell lines analyzed, and previously described as nearly impossible to transfect with nonviral vectors.^[34] It is worth noting that mRNA used was of 0.3 μ g per well in our nanoparticles, while it was one third for the positive controls. This is due to the fact that it was not possible to increase the concentration of the positive controls due to cytotoxicity problems, which also makes them more problematic than our nanoparticles.

2.3. In Vivo Organ Biodistribution Study

Next, and after showing that KH nanoparticles could safely and very efficiently transfect APC in vitro (JAWS and RAW264.7 cell lines tested), we decided to study their biodistribution after an intravenous injection, in order to determine if they were able to reach this type of cells in vivo after a systemic administration. A total of five animals were intravenously administered with 50 μ g of mRNA codifying for firefly luciferase (Fluc), encapsulated in KH nanoparticles. BLI was used to monitor their biodistribution. Negative controls consisting in nontreated and naked mRNA treated animals were also included ($n = 2$). Plots of total light events (PHCs/area) were recorded 6 and 24 h after the administration of nanoparticles, just 15 min after luciferine substrate injection (Figure 3A,B). For the negative controls, all results were negative, no BLI signal was observed in any case. Remarkably, high quantity of the injected OM-PBAE were

localized on the spleen, and, regarding other organs, only the liver shows the presence of signal, but at a significant much lower level. The highest expression of spleen signal appeared 6 h after injection and it was followed by a significant notable signal decrease up to 24 h (Figure 3A,B). For this reason, 6 h was chosen as the maximum expression level for further in vivo studies. Concerning ex vivo measurements, they were performed at 30 h after nanoparticle administration. According to in vivo measurements, a meaningful accumulation of nanoparticles in the spleen was observed after this time, as compared to liver accumulation (Figure 3A,C).

2.4. In Vivo Cell Targeting Study

To further understand the biodistribution of nanoparticles, and in order to determine if they were able to specifically transfect APCs in vivo, we studied the specific cell types transfected by the nanoparticles after their intravenous administration. In this case, the polymer was labeled with Cyanine 5 (Cy5), in order to study cell uptake, while the mRNA codifying for eGFP was used to determine cell transfection. Animals were injected with 50 μ g of mRNA, intravenously and after 6 h administration, the previously determined highest level of gene expression, animals were sacrificed and spleens were digested. Further, the mixture of cells obtained was stained with the appropriate antibodies, labeled with fluorescent dyes, and the colocalization of nanoparticles dyes with those concerning specific cells was performed by FACS analysis. Negative controls were used to determine the background signal (nontreated and naked mRNA treated animals; negative results; $n = 2$; data not shown). Two panel combinations of antibodies, selected based on many previous publications,^[35–40] were designed as described in

Figure 2. A) Viability (in %) of different cell lines after 24 h incubation with different formulations; B) Transfection efficiency (in %) of mRNA eGFP in vitro, in different cell lines; and C) Examples of the cell fluorescence after 24 h transfection with mRNA eGFP, as a function of cell line and sample incubated (Scale bar = 100 μm). $n = 3$. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. Negative control (C-) showed significant differences with any other sample ($p < 0.0001$).

Figure 4A to study colocalization of NPs signals (both polymer and mRNA) with different cell types markers.

It is interesting to see that nanoparticle uptake (Cy5+ cells) is high, reaching more than the 50% of the spleen cells (Figure 4B). On the other hand, the transfection itself (GFP+ cells) results to be around a 0.2% of the total cells. This result leads us to think that only a fraction of the particles uptaken are able to properly release and express the genetic material.

Going deep into spleen transfection, cell markers were used to determine which cell type internalized nanoparticles and which of these cells were transfected. In general, lymphocytes transfection was almost null (Figure 4E). Previous studies remarked the difficulty in transfecting lymphocytes (namely murine splenic lymphocytes). These cells are refractory to be transfected, especially when using nonviral vectors, due to their low efficiency and high toxicity.^[41]

Six hours after intravenous administration, lymphocytes did not internalize nanoparticles, unlike macrophages, neutrophils and dendritic cells, in which NPs uptake was similar among the different cell types (Figure 4D). Regarding transfection, the highest percentage was found for macrophages, followed by neutrophils and dendritic cells (no lymphocytes transfection was observed) (Figure 4E). Macrophages transfection was higher compared to the other two cell types, reaching values $\approx 6\%$, while it was ≈ 3 and 1.5% for neutrophils and dendritic cells, respectively. These levels of transfection are very similar to the ones obtained in other studies, in which lipid nanoparticles encapsulating mRNA were used for the design of novel mRNA immunotherapies.^[16,26,28] The transfection of macrophages could be useful due to their role as APC and also, it could be desired for reprogramming their inflammation response in many diseases, such as Parkinson's disease.^[42,43]

Figure 3. A) Bioluminescent images of a representative animal, at the different measurement times; B) Quantification of the BLI signal of the in vivo images (PHC/area); and C) Ex vivo images BLI signal quantification after 30 h administration (PHC/area). $n = 5$. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

In addition, neutrophils transfection, although it has been less studied, could be envisaged to design treatments for infectious diseases or even as cancer immunotherapeutics.^[44,45]

3. Discussion

With the aim to design novel nanoformulations able to be used as vaccines for immunotherapies in a near future, oligopeptide end-modified PBAE (OM-PBAE) nanoparticles have been used to encapsulate genetic material. In particular, mRNA has been selected as the gene material to be encapsulated and delivered due to its safety profile, amenability to be fabricated under good manufacturing procedures and because, unlike DNA, it does not require crossing the nuclear membrane to be expressed.^[3,11,12,46] Therefore, mRNA can be transfected efficiently to both dividing and quiescent cells.^[25] In addition, a modified mRNA was selected due to its higher performance and its decreased unspecific immune system activation.^[46] Concerning the polymer selection, it is worth remarking the advantages of using OM-PBAEs, instead of more simple PBAEs, since they are able to self-aggregate negatively charged gene material and spontaneously form small discrete nanoparticles.^[19]

Previous studies demonstrated the possibility to tailor OM-PBAE nanoparticles to target specific cell types by modifying the end-oligopeptide modifications.^[10,12] In this study, KH polymer combination was chosen to achieve efficient transfection of APCs in vivo.

Prior to in vivo studies, physicochemical properties of the nanoparticles were fully characterized, since a wide characterization and knowledge of nanoparticle properties has been described to enhance their performance in vivo.^[11,47] Cationic low polydisperse complexes with a diameter ≈ 150 nm were prepared, with a notable mRNA encapsulation efficiency ($>75\%$), higher than previously designed nanosystems.^[46,48] It is noteworthy that properties of these nanoparticles are dependent on the end-oligopeptides, as expected,^[12] making OM-PBAEs a very tunable system to be used for many biomedical applications. In this work, based on differential physicochemical properties and in vitro transfection efficiency found in previous work^[10,12] (see Figure S1, Supporting Information), KH were selected for further experimentation. It is worth remarking that the high encapsulation efficiency is a key point in terms of industrial production by pharmaceutical companies; a high efficient procedure is always more cost-effective.^[49–51] In addition, these complexes have a positive surface charge, which makes them stable against colloidal destabilization processes^[52] and enables a high interaction with cell membranes thanks to electrostatic interactions.^[53,54]

With the aim to prepare a ready to use product, maintaining OM-PBAEs properties, the lyophilization process has been carefully adjusted paying special attention to size, ζ potential and encapsulation efficiency. Moreover, it has been proven the maintenance of their transfection efficacy, both in vitro and in vivo; even after long-term storage of nanoparticles (see Figure S4, Supporting Information). The possibility to successfully lyophilize these NPs is of utmost importance, since, to our best knowledge, only two recent studies demonstrated the possibility to freeze-dry PBAE nanoparticles and redisperse them in an aqueous buffer maintaining their properties.^[7,15] Interestingly, the current work successfully lyophilized OM-PBAE nanoparticles for the first time. This process is a key step regarding nanoparticles long-term storage and production at larger scales,^[7,55] a milestone for the development of future therapies, since nanoparticle dispersions suffer from destabilization processes like sedimentation.^[56] Accordingly, after freeze-drying, nanoparticle showed an appropriate osmolality (around 300 mOsm kg^{-1}) and pH (7.4); size, surface charge, monodispersity, and encapsulation efficiency to be administered intravenously.^[57] Moreover, these nanoparticles are smaller and less polydisperse than most of previously published nanosystems, which is a key parameter, also to increase cell penetration and transfection.^[11,37,48,55] It is

Figure 4. A) Schematic representation of a cell lineage and the cell surface markers used to detect each cell type; B) % Cy5+ cells and C) % GFP+ cells within homogenized spleens of animals administered with KH nanoparticles, after 6 h administration; D) % Cy5+ cells; and E) % GFP+ cells in the Spleen, as a function of the cell type (same animals and same experimental time). Epithelial and endothelial cells are not plotted because their uptake and transfection was null. $n = 5$. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

worth noting that it is the first time that OM-PBAE nanoparticles have been synthesized with these small sizes, lower than 170 nm.^[12] In addition, lower N/P ratios (25/1 as compared to 200/1 from previous studies^[12]) were used, which represents lower polymer concentrations, being advantageous in terms of safety issues. The use of less polymer has also enabled lower cationic surface charges, as compared with many current polymeric nanoparticles, which is advantageous in terms of safety (lower phagocytosis and lower interaction with cell membranes) when administered intravenously.^[17,57] Highly charged nanosystems, in absolute value, were widely reported to strongly interact with blood proteins and cells, in an unspecific way, thus resulting in an activation of the innate immune system and nanoparticle elimination, hindering them to reach the target organ and produce their therapeutic action.^[9,27,31]

Concerning the interaction of these nanoparticles with cells, no cytotoxic effects were found in vitro at the mRNA doses tested, higher for nanoparticles than for controls (Figure 2A). Therefore, these OM-PBAE polymers are confirmed once more as non-toxic at these concentrations.^[5,9,10,12] Two commercial vectors were chosen as positive transfection controls: Lipofectamine 2000 and jetMESSENGER. While the former has a lipid nature, the latter was selected due to its polymeric nature, which makes it more similar to our OM-PBAE nanoparticles. Both positive controls showed very similar results in all cell

lines tested. In general, it was confirmed that OM-PBAE nanoparticles could transfect efficiently all the cell lines analyzed. KH nanoparticles showed transfection efficiencies comparable to commercial vectors when they were assayed in HeLa cells. Interestingly, these nanoparticles were able to triplicate the transfection efficiency in another model, ARPE-19 cells, which are not as permissive as HeLa cells to commercial transfection reagents. Concerning macrophages and dendritic cells, which are usually refractory to any kind of transfection,^[25] their transfection was improved by the use of KH nanoparticles, which showed significantly higher levels of transfection in both cell lines (around 60% GFP+ cells), compared to commercial vectors and most of the nanoformulations previously used.^[6,25,46]

Traditionally, most nanoformulations intended for biomedical purposes were composed of lipids instead of polymers.^[18,48] This was due to the belief that polymeric carriers performed very poorly in the context of in vivo mRNA delivery, which was attributed to an ineffective endosomal escape of the gene material due to its stringent binding with the cationic polymer.^[46] However, some studies demonstrated concerns about lipid carriers safety, as some cases of toxicity after administration, together with inflammation and cytokines activation were found.^[12,33] In addition, the use of polymers instead of lipids enables the possibility to better control the material nature and composition and tune its properties.^[25] In this study, OM-PBAEs are presented

as promising gene encapsulating polymers, since previous work already demonstrated PBAE capability of endosomal escape and cell transfection both in vitro and in vivo, without showing toxicity at the concentrations required.^[10,12,19,20,58] In this context, it is remarkable the relationship between encapsulation efficiency and endosomal escape. In previous studies, it was confirmed that, although high encapsulation was desired to protect genetic material from nucleases,^[11] a high encapsulation efficiency represented a drawback for the gene expression since high stability of the complex may hinder endosomal escape.^[25] However, in this study, we demonstrated a high and stable mRNA encapsulation. In parallel, endosomal escape was also confirmed, since it was possible to observe a high expression of the encapsulated reporter gene, both in vitro and in vivo. The release of the mRNA in the endosome could be attributed to the hydrophilic character of the peptide fraction of the polymer,^[21] as well as to the proton sponge effect of the polymer formulated with lysines.^[10,12]

Typically, nanomedicines, both lipid and polymeric, are accumulated in the liver when they are administered without an active targeting moiety, especially due to their sequestration by the mononuclear phagocyte system (MPS).^[2,24,48,59] For example, McCaskill et al.^[48] encapsulated siRNA in cationic liposomal nanostructures and found most of them in the liver (60% of the total) after intravenous administration, although their aim was lung transfection. Liver accumulation could be advantageous for the treatment of hepatic diseases, but in the rest of diseases, this could be a problem mainly due to the phagocytic clearance that the liver itself produces, which does not enable nanoparticles to reach their target organs and to produce their therapeutic effect^[24] and also due to hepatocellular toxicity by an excessive dose.^[2] In this sense, it is important to remark that our OM-PBAE nanoparticles, which do not include an active targeting moiety, are not predominantly accumulated in the liver. In contrast, they are mainly accumulated in the spleen (80% of PHC, see Figure S5, Supporting Information for a comparison spleen—liver; see also Figure 3); therefore, a significant detargeting of the liver is confirmed, thus overcoming the drawback of previous studies, which claimed only 1% of the dose reached tissues other than liver.^[59]

Most in vivo studies are only focused on the organ biodistribution and do not analyze the different cell populations inside the organ.^[2] In this work, a deeper in vivo study was performed to determine the cell specificity of these nanoparticles, since we tried to find an explanation for the spleen accumulation without the use of an active targeting moiety. For vaccination purposes, the main goal of the current work, the transfection of antigen presenting cells (APCs), such as dendritic cells, is envisaged. These cells will trigger the specific immune response against the encoded antigen, thus producing an immune attack against cells expressing such antigen, that should end up with their elimination and the reduction of tumor size, in the case of cancer, for example.^[16,26,36] Moreover, a labeling of the polymer (Cy5) has been also included, in order to study, not only transfection, but also cell penetration. Although all transfected cells must be previously penetrated by nanoparticles, it is possible that nanoparticles penetrate some other cells, which are not able to be transfected.^[7]

These data show that cell uptake was very high in general, in most of immune cells studied (higher than 20%) and higher than previous studies.^[2] As stated above, in this study, the transfection of APCs is desired for the design of vaccines. APCs define a group of different kinds of cells, all of them with the common characteristic of antigen presentation when transfected with exogenous gene material.^[16,23,26,37] Among them, dendritic cells are the most efficient APCs, although macrophages and neutrophils are also of interest; all of them can amplify or suppress an adaptive immune response and therefore, they play a key role in DNA/mRNA vaccination. Consequently, the penetration and further transfection of dendritic cells in the spleen, the organ with the highest immune cell concentration of the body, has been envisaged as a very useful approach for the design of gene vaccines.^[21,23,26,46,60] KH nanoparticles were designed with physicochemical properties that make them easily uptaken by dendritic cells (e.g., size of ≈ 150 nm).^[36] Concerning the uptake, it is worth noting that high uptake levels were found for APCs. This high uptake in APCs was previously reported by Hamdy et al.,^[28] who claimed that it was due to the phagocytosis of the nanoparticles. Therefore, the phagocytosis would be the reason why nontargeted nanoparticles are accumulated in the spleen; because it is the resident organ of phagocytic cells. However, their nanoparticles were much bigger than ours, comparable to pathogens, which enhance recognition and phagocytosis. Related with this study, Moffatt and Cristiano^[21] also produced microparticles with the objective to induce their phagocytosis by the dendritic cells, usually enhanced in the micrometer range. These authors claimed that this route was also useful for cell vaccination purposes, although this was not the desired route, because it is not antigen—specific. This statement is in good agreement with our results, where the transfection was also higher. Although it could be thought that transfecting $\approx 2\%$ of splenic dendritic cells is a low value, it is in the order of magnitude of previous studies, which have proven that these reduced percentages are able to confirm that the tested nanoparticles can modify the gene expression patterns of these cells.^[26] In addition, transfection of this low percentage APCs in the spleen could represent a high number, in absolute value, since the spleen is the organ where more APC are accumulated; so the total cellular count transfected could be more than enough to use these systems as effective mRNA vaccines that initiate the specific required immune response. Consequently, knowing that transfection takes place, the uptake cannot be completely attributed to phagocytosis: a combination between this route and another alternative route such as endocytosis is hypothesized.^[33,46]

Currently, many therapies aiming to transfect APCs perform cell transfection ex vivo and then readminister the modified cells. However, the migration of these cells to the lymph nodes is inefficient and the antigen presentation does not take place efficiently. In addition, ex vivo therapies have the added disadvantage of being tailored for each individual and therefore, they are time-consuming and expensive. In this context, our approach represents an advantage since APC transfection takes place in vivo.^[28] Furthermore, this in vivo immune response could be not only directed against tumoral cells, thus being novel cancer immunotherapies but also to treat infectious diseases, if desired.^[26,38]

Therefore, taking into account all our data, it could be concluded that the KH end-modification of the PBAE nanoparticles enables a specific in vivo targeting to the APCs in the spleen, thus representing promising vectors for vaccination to treat tumors or infectious diseases.

4. Conclusions

In this work, OM-PBAEs have been selected for the design of novel nanoparticles, encapsulating mRNA, for vaccination purposes. A wide physicochemical characterization study confirmed the suitable characteristics of the KH polymer combination to be administered intravenously, even after their lyophilization, process that has been applied with success for the first time to OM-PBAE nanoparticles. In vitro studies have demonstrated their safety and efficacy transfecting different model cell lines, among them APCs such as macrophages and dendritic cells. Furthermore, in vivo studies have demonstrated the capability of these nanoparticles to efficiently penetrate and transfect these cells after their intravenous administration. KH nanoparticles clearly accumulate in the spleen. Therefore, the main aim of the study has been achieved: the capacity to design OM-PBAE nanoparticles tailored to specifically target APCs has been demonstrated. Consequently, these OM-PBAE nanoparticles represent a promising alternative as novel vectors for the design of vaccination strategies against tumors and infectious diseases.

5. Experimental Section

Materials: Reagents and solvents used for synthesis were purchased from Sigma Aldrich and Panreac. CK3 (NH₂-Cys-Lys-Lys-Lys-COOH) and CH3 (NH₂-Cys-His-His-His-COOH) peptides was obtained from GL Biochem (Shanghai) Ltd with a purity of at least 98%. Antibodies used in flow cytometry (FACS) were directly conjugated monoclonal fluorescently labeled primary antibodies from BD Bioscience: PE-Cy7 Hamster Anti-Mouse CD3e, PerCP-Cy5.5 Rat Anti-Mouse CD11b, BV421 Hamster Anti-Mouse CD11c, BV786 Rat Anti-Mouse CD19, APC-Cy7 Rat Anti-Mouse CD45, PerCP-Cy5.5 Rat Anti-Mouse CD146, PE Rat Anti-Mouse CD326 and PE Rat Anti-Mouse F4/80; together with the Fixable Viability Stain 775 V. mRNAs used in this study were eGFP (CleanCap Enhanced Green Fluorescent Protein mRNA 5-methoxyuridine) for cell culture experiments and Fluc (CleanCap Fluc mRNA 5-methoxyuridine), for all other experiments; obtained from TriLink.

Cell Lines: HeLa (ATCC CCL-2) and RAW 264.7 (ATCC TIB-71) cell lines were maintained in DMEM supplemented with 10% (v/v) heat inactivated fetal bovine serum (FBS), 100 units mL⁻¹ penicillin G, 100 µg mL⁻¹ streptomycin, and 2 mmol L⁻¹ L-glutamine. ARPE-19 (ATCC CRL-2302) cell line was maintained in DMEM-F12 supplemented with 10% (v/v) FBS, 100 units mL⁻¹ penicillin G, 100 µg mL⁻¹ streptomycin and 2 mmol L⁻¹ L-glutamine. JAWS2 (ATCC CRL-11904) cell line was maintained in EMEM supplemented with 20% (v/v) FBS; 1 mmol L⁻¹ sodium pyruvate, 4 mmol L⁻¹ L-glutamine AND 5 ng mL⁻¹ of GM-CSF. All cells were cultured at 37 °C, under a 5% CO₂/95% air atmosphere until 90% confluence before starting transfections.

Animals: Adult 6–8 weeks old Balb/c mice were purchased (Envigo) and kept under pathogen-free conditions in laminar flow boxes. Animal maintenance and experiments were performed in accordance with established guidelines of the Catalan Government and following protocol number 8856, approved by the *Direcció General del Medi Natural*.

Synthesis of PBAE Polymers: Poly(β-amino ester)s (PBAEs) were synthesized as described in Dosta et al., (submitted to Molecular

Systems Design and Engineering). Briefly, addition reaction of primary amines to an excess of diacrylates was used to synthesize an acrylate-terminated polymer. Then, this polymer was end-capped with oligopeptide composed of Cys + 3 Lys (CKKK or CK3) or Cys + 3His (CHHH or CH3), in DMSO. Synthesized structures were confirmed by ¹H NMR, recorded in a 400 MHz Varian (NMR Instruments, Claredon Hills, IL) and methanol-d₄ used as a solvent. Molecular weight (MW) relative to polystyrene standard was determined by HPLC (HPLC Elite LaChrom system of VWR-Hitech equipped with a GPC Shodex KF-603 column and THF as mobile phase).

Synthesis and Lyophilization of PBAE Nanoparticles: OM-PBAE nanoparticles (also named as complexes) were prepared following a modification of the Dosta et al., protocol.^[12] In brief, they were prepared at a 25/1 OM-PBAE/mRNA ratio, by mixing equal volumes of mRNA at 0.5 mg mL⁻¹ with the polymer (being the polymer a mixture of 60% C6CK3 i 40% C6CH3) at 12.5 mg mL⁻¹, in 12.5 × 10⁻³ M AcONa buffer solution. The mRNA was added over the polymer solution and mixed by pipetting, followed by 30 min incubation at 25 °C. For the formation of discrete structures, this mixture (volume = Vi) was nanoprecipitated in a same volume (Vi) of DEPC water. Thereafter, the same volume (Vi) of a HEPES 20 × 10⁻³ M + 4 wt% sucrose solution was also added as cryo and lioprotectors. Then, nanoparticles were freeze-dried. The lyophilized powder was kept at -20 °C until use, when it was reconstituted with DEPC water (Vi).

Physicochemical Characterization of PBAE Nanoparticles: Hydrodynamic diameter (nm), polydispersity index (PDI), and surface charge of nanoparticles were measured at 25 °C, 633 nm laser wavelength and 173° signal detector; using a ZetaSizer Nano ZS (Malvern Instruments Ltd, Malvern, UK). For size measurements, nanoparticles prepared at 0.25 mg mL⁻¹ of mRNA were used, while a 1/10 dilution (final concentration 0.025 mg mL⁻¹) in water was performed to the lyophilized nanoparticles to measure surface charge. Three measurements of each nanoparticle batch were performed with 20 runs per measurement, taking into account the intensity approximation. Results correspond to the mean ± standard deviation of at least five independent nanoparticle batches.

Hydrodynamic size was also measured by nanoparticle tracking analysis (NTA), using a NanoSight LM10 from Malvern, equipped with a fluorescent laser (λ = 488 nm). Results correspond to the mean ± standard deviation of at least three independent nanoparticle batches.

Encapsulation efficiency was measured by the Quanti-It RiboGreen RNA assay kit (ThermoFisher Scientific), following manufacturer instructions.

In Vitro Cellular Cytotoxicity and Transfection: Cells were seeded in 96-well and 48-well (only for RAW264.7 cells) plates at 90% confluence, 24 h before starting the experiment. Cells were incubated with 0.3 µg mL⁻¹ of mRNA encapsulated within OM-PBAE or 0.1 µg mL⁻¹ of encapsulated mRNA for Lipofectamine2000 (Thermo Fisher Scientific; Spain) and jetMESSENGER (Polyplus transfection; France) as positive controls (concentration recommended by the manufacturers' guidelines). mRNA concentration was selected based on previous studies and corresponded to the minimum concentration required to achieve the maximum transfection. Nontreated cells were used as negative controls (results correspond to 100% viability, used to normalize absorbance of the tested samples). After 24 h incubation in the presence of FBS, cell viability and transfection efficiency were measured. MTT assay (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)^[61] was used to determine cell viability. 0.5 mg mL⁻¹ of MTT were added at each well, and incubated at 37 °C for 24 h. After this time, media was removed and DMSO was added to dissolve the formazan crystals. Absorbance was quantified at 570 nm, using a plate reader (SpectraMax M5 luminometer, Molecular Devices). Results were expressed as relative viability percentage to control (nontreated) cells. Transfection efficiency was measured by flow cytometry (FACS; in a BD Fortessa cell analyzer). Cells were trypsinized (except RAW264.7, which were scrapped) and fixed with 1% paraformaldehyd. At least 2000 cells were analyzed per sample; and fluorescence gating was determined using positive and negative controls. Data were analyzed with BD LSRFortessa cell

Table 1. Antibody panels used for the cellular distribution studies to label cells and detect them by FACS.

Panel	Laser	Violet laser			Blue laser		Green laser		Red laser	
		V780	V560	V450	B695	B515	G780	G575	R780	R660
	Dye	BV785	BV570	BV421	PERCP	PERCP	PECY7	PE	APCCY7	APCH7
					PERCP	PERCP	PECY7	PE	APCCY7	APCH7
1			FVS570v*		CD146	GFP	CD326		CD45	CyS
2		CD19	FVS570v*	CD11c	CD11b	GFP	CD3	F4/80	CD45	CyS

analyzer software. Results correspond to the mean $\pm$ standard deviation of at least three independent transfections, each one consisting on experiment triplicates.

Biodistribution Studies by In Vivo Bioluminescence: Mice ($n = 5$) were injected intravenously with 50 μ g (200 μ L) of mRNA encoding for firefly luciferase complexed with OM-PBAE selected polymer. Radiance was measured by bioluminescence imaging (BLI). In vivo BLI mice images were taken 6 h and 24 h postinjection, performed as described previously.^[62] Briefly, mice were anesthetized and injected intraperitoneally with 150 μ L of luciferin (Regis Technologies) (16.7 mg mL⁻¹ in PBS). Animals were placed in a high efficiency ImagEM X2 C9100-23BEM-CCD Imaging System (Hamamatsu Photonics) and images were acquired from the ventral direction. A second image of the animal was obtained using a white-light source, to register the position of the luminescence signal.

Ex vivo BLI images were taken 30 h after NPs injection. Mice were anesthetized, injected with luciferin, and euthanized. Spleen was removed immediately and images of the organs were taken as described previously.

Quantification and analysis of photons recorded in images was done using the Wasabi image analysis software (Hamamatsu Photonics) and the results were expressed as photons counts (PHC) / area, after subtracting the background.

In Vivo Cellular Distribution by FACS: Mice ($n = 5$) were injected with 50 μ g (200 μ L) of mRNA encoding for eGFP complexed with OM-PBAE polymer; 6 h postinjection animals were euthanized, liver and spleen were removed and disaggregated as described below.

Spleens were directly excised and digested following the protocol from Pujol-Autonell et al.^[63] In brief, they were removed and placed in 2 wt% FBS in PBS media, at 4 °C. They were mechanically digested by manual tissue mashing in around 5 mL of media using a syringe plunger. The liquid was filtered through 70 μ m nylon cell strainers to remove big fragments (conjunctival and stromal tissue). The filtered cell suspension was centrifuged for 10 min at 400 g, at 25 °C. The supernatant was removed and the pellet incubated with 5 mL hemolysis buffer (1.029 g Tris-HCl and 3.753 g ammonium chloride in 500 mL of milliQ water) for a maximum time of 5 min. To stop the hemolytic activity, 5 mL of the media were added and the cells were recentrifuged. The centrifugation was repeated twice to clean cells. Finally, cell density was counted and cells were frozen with FBS/DMSO 90/10 for their further use.

Spleen cellular dispersions were thaw and redispersed in media. The living cell number was counted using standard beads, by FACS. Then, cells were stained with FVS570 viability dye and fluorescently labeled with specific antibodies (see Table 1). This mixture was incubated for 30 min at 4 °C. After staining, cells were washed with PBS and analyzed fresh at BD Fortessa FACS (BD Bioscience). FACS data were processed using BD FACSDiva software (BD Biosciences). To identify myeloid populations, after the exclusion of debris and doublets, immune cells were identified and quantified by the pan-hematopoietic CD45 marker, for the labeling of lymphocytes. Then, CD45-cells were tested as epithelial (CD146+) or endothelial (CD326+) cells. Instead, CD45+ cells were classified in B lymphocytes (CD19+), T lymphocytes (CD3+) or myeloid cells (DCs: CD11b+/–, CD11c+, F4/80–; macrophages: CD11b+, CD11c–, F4/80+; or neutrophils: CD11b+, CD11c–, F4/80–).^[35] Results from the FACS experiments correspond to means $\pm$ standard deviation of at least five independent cell suspensions, coming from five independent animals.

Statistical Analysis: Data from the physicochemical characterization of PBAE nanoparticles (hydrodynamic diameter, PDI, surface charge and encapsulation efficiency) were obtained directly from each measurement, without any preprocessing. Given values correspond to the mean $\pm$ standard deviation of a sample size of $n = 3$ (exception: DLS data correspond to statistics of $n = 5$). GraphPad Prism software was used for the statistical analysis. A t-test has been applied to find out statistical differences between groups. p values < 0.05 were considered statistically significant.

In vitro viability data were obtained from the normalization of raw absorbance data to the negative controls (nontreated cells). Results correspond to mean $\pm$ standard deviation of a sample size of $n = 3$. GraphPad Prism software was used for the statistical analysis. A paired-sample t-test has been applied to find out statistical differences. p values < 0.05 were considered statistically significant.

Concerning fluorescent signal obtained from FACS analysis, it was used for in vitro and in vivo data. Results correspond to mean $\pm$ standard deviation of a sample size of $n = 3$ for in vitro data, while $n = 5$ for in vivo studies. GraphPad Prism software was used for the statistical analysis. A paired-sample t-test has been applied to find out statistical differences for in vitro data; while one-way analysis of variance (ANOVA) or Student's t-test was performed to determine statistical significance among multiple groups of the in vivo data. p values < 0.05 were considered statistically significant. In vivo luminescence data were normalized by the area (the background was subtracted) and the time of the measure, resulting in photon counts/area. Results correspond to mean $\pm$ standard deviation of a sample size of $n = 5$. GraphPad Prism software was used for the statistical analysis. A t-test has been applied to find out statistical differences. p values < 0.05 were considered statistically significant.

Supporting Information

Supporting Information is available from the Wiley Online Library or from the author.

Acknowledgements

Financial support from MINECO/FEDER (grants RTC-2015-3751-1, SAF2015-64927-C2-1-R and SAF2015-64927-C2-2-R) and Instituto de Salud Carlos III (Red Temática de Investigación Cooperativa en Terapia Celular-TERCEL) are acknowledged. CIBER-BBN is an initiative funded by the VI National R&D&I Plan2008–2011, Iniciativa Ingenio 2010, Consolider Program, CIBER Actions and financed by the Instituto de Salud Carlos III with assistance from the European Regional Development Fund. C.F. is grateful to MINECO for their Postdoctoral Fellowship (grant Torres Quevedo 2015). The Support of Agència de Gestió d'Ajuts Universitaris i de Recerca (AGAUR) from Generalitat de Catalunya for their support through SGR 2014 1170 grant. Authors acknowledge the kind support of Marta Vives group on the digestion of spleens; Marco A. Fernández and Gerard Requena for the FACS measures and analysis and Irene Porcar and Elena García-Ollé for their kind support in experiments performance.

Conflict of Interest

The authors declare no conflict of interest.

Keywords

antigen-presenting cells transfection, immunotherapy, mRNA encapsulation and delivery, oligopeptide-modified poly-(β -aminoester) polyplexes, spleen

Received: March 30, 2018

Revised: May 6, 2018

Published online:

- [1] J. K. Park, T. Utsumi, Y. E. Seo, Y. Deng, A. Satoh, W. M. Saltzman, Y. Iwakiri, *Biol. Med.* **2016**, 12, 1365.
- [2] Y. N. Zhang, W. Poon, A. J. Tavares, I. D. McGilvray, W. C. W. Chan, *J. Controlled Release* **2016**, 240, 332.
- [3] R. M. Levine, C. M. Scott, E. Kokkoli, *Soft Matter* **2013**, 9, 985.
- [4] T. J. Burney, J. C. Davies, *Appl. Clin. Genet.* **2012**, 5, 29.
- [5] C. J. Bishop, R. L. Majewski, T. R. M. Guiriba, D. R. Wilson, N. S. Bhise, A. Quiñones-Hinojosa, J. J. Green, *Acta Biomater.* **2016**, 37, 120.
- [6] H. Akita, S. Ishii, N. Miura, S. M. Shaheen, Y. Hayashi, T. Nakamura, N. Kaji, Y. Baba, H. Harashima, *Biomaterials* **2013**, 34, 8979.
- [7] H. Guerrero-Cázares, S. Y. Tzeng, N. P. Young, A. O. Abutaleb, A. Quiñones-Hinojosa, J. J. Green, *ACS Nano* **2014**, 8, 5141.
- [8] C. J. Bishop, S. Y. Tzeng, J. J. Green, *Acta Biomater.* **2015**, 11, 393.
- [9] H. Tian, Z. Tang, X. Zhuang, X. Chen, X. Jing, *Prog. Polym. Sci.* **2012**, 37, 237.
- [10] N. Segovia, P. Dosta, A. Cascante, V. Ramos, S. Borrós, *Acta Biomater.* **2014**, 10, 2147.
- [11] K. K. L. Phua, K. W. Leong, S. K. Nair, *J. Controlled Release* **2013**, 166, 227.
- [12] P. Dosta, N. Segovia, A. Cascante, V. Ramos, S. Borrós, *Acta Biomater.* **2015**, 20, 82.
- [13] E. W. F. W. Alton, A. C. Boyd, S. H. Cheng, J. C. Davies, L. A. Davies, A. Dayan, D. R. Gill, U. Griesenbach, T. Higgins, S. C. Hyde, J. A. Innes, G. McLachlan, D. Porteous, I. Pringle, R. K. Scheule, S. Sumner-Jones, *Gene Ther.* **2014**, 21, 89.
- [14] M. Elsbahy, K. L. Wooley, *Chem. Soc. Rev.* **2013**, 42, 5552.
- [15] D. R. Wilson, A. Mossenia, M. Suprenant, R. Upadhy, D. Routkevitch, R. Meyer, A. Quiñones-Hinojosa, J. J. Green, *J. Biomed. Mater. Res., Part A* **2017**, 105, 1813.
- [16] M. A. Oberli, A. M. Reichmuth, J. R. Dorkin, M. J. Mitchell, O. S. Fenton, A. Jaklenec, D. G. Anderson, R. Langer, D. Blankschtein, *Nano Lett.* **2017**, 17, 1326.
- [17] M. A. Dobrovolskaia, M. Shurin, A. A. Shvedova, *Toxicol. Appl. Pharmacol.* **2016**, 299, 78.
- [18] C. Pollard, J. Rejman, W. De Haes, B. Verrier, E. Van Gulck, T. Naessens, S. De Smedt, P. Bogaert, J. Grooten, G. Vanham, S. De Koker, *Mol. Ther.* **2013**, 21, 251.
- [19] K. L. Kozielski, S. Y. Tzeng, B. A. Hurtado De Mendoza, J. J. Green, *ACS Nano* **2014**, 8, 3232.
- [20] N. Segovia, M. Pont, N. Oliva, V. Ramos, S. Borrós, N. Artzi, *Adv. Healthcare Mater.* **2015**, 4, 271.
- [21] S. Moffatt, R. J. Cristiano, *Int. J. Pharm.* **2006**, 321, 143.
- [22] R. Duncan, R. Gaspar, *Mol. Pharm.* **2011**, 8, 2101.
- [23] F. Perche, T. Benvegnu, M. Berchel, L. Lebegue, C. Pichon, P. A. Jaffrès, P. Midoux, *Biol. Med.* **2011**, 7, 445.
- [24] E. Blanco, H. Shen, M. Ferrari, *Nat. Biotechnol.* **2015**, 33, 941.
- [25] T. Bettinger, R. C. Carlisle, M. L. Read, M. Ogris, L. W. Seymour, *Nucleic Acids Res.* **2001**, 29, 3882.
- [26] L. M. Kranz, M. Diken, H. Haas, S. Kreiter, C. Loquai, K. C. Reuter, M. Meng, D. Fritz, F. Vascotto, H. Hefesha, C. Grunwitz, M. Vormehr, Y. Husemann, A. Selmi, A. N. Kuhn, J. Buck, E. Derhovanessian, R. Rae, S. Attig, J. Diekmann, R. A. Jabulowsky, S. Heesch, J. Hassel, P. Langguth, S. Grabbe, C. Huber, Ö. Türeci, U. Sahin, *Nature* **2016**, 534, 396.
- [27] N.-H. Cho, T.-C. Cheong, J. H. Min, J. H. Wu, S. J. Lee, D. Kim, J.-S. Yang, S. Kim, Y. K. Kim, S.-Y. Seong, *Nat. Nanotechnol.* **2011**, 6, 675.
- [28] S. Hamdy, A. Haddadi, R. W. Hung, A. Lavasanifar, *Adv. Drug Delivery Rev.* **2011**, 63, 943.
- [29] C. Fornaguera, **2015**.
- [30] W. Schärfl, *Light Scattering from Polymer Solutions and Nanoparticle Dispersions*, Springer L, Springer, Mainz, Germany **2007**.
- [31] M. A. Dobrovolskaia, S. E. McNeil, *J. Controlled Release* **2013**, 172, 456.
- [32] C. Fornaguera, G. Calderó, C. Solans, C. Vauthier, *Eur. J. Pharm. Biopharm.* **2015**, 94, 284.
- [33] H. Akita, K. Kogure, R. Moriguchi, Y. Nakamura, T. Higashi, T. Nakamura, S. Serada, M. Fujimoto, T. Naka, S. Futaki, H. Harashima, *J. Controlled Release* **2011**, 149, 58.
- [34] S. M. Shaheen, H. Akita, T. Nakamura, S. Takayama, S. Futaki, A. Yamashita, R. Katoono, N. Yui, H. Harashima, *Biomaterials* **2011**, 32, 6342.
- [35] A. V. Misharin, L. Morales-Nebreda, G. M. Mutlu, G. R. S. Budinger, H. Perlman, *Am. J. Respir. Cell Mol. Biol.* **2013**, 49, 503.
- [36] V. Sokolova, T. Knuschke, A. Kovtun, J. Buer, M. Eppe, A. M. Westendorf, *Biomaterials* **2010**, 31, 5627.
- [37] K. J. Radford, K. M. Tullett, M. H. Lahoud, *Curr. Opin. Immunol.* **2014**, 27, 26.
- [38] A. Gardner, B. Ruffell, *Trends Immunol.* **2016**, 37, 855.
- [39] M. Merad, P. Sathe, J. Helft, J. Miller, A. Mortha, *Annu. Rev. Immunol.* **2013**, 31, 563.
- [40] P. Guha, J. W. Morgan, G. Mostoslavsky, N. P. Rodrigues, A. S. Boyd, *Cell Stem Cell* **2013**, 12, 407.
- [41] O. S. Fenton, K. J. Kauffman, J. C. Kaczmarek, R. L. McClellan, S. Jhunjunwala, M. W. Tibbitt, M. D. Zeng, E. A. Appel, J. R. Dorkin, F. F. Mir, J. H. Yang, M. A. Oberli, M. W. Heartlein, F. DeRosa, R. Langer, D. G. Anderson, *Adv. Mater.* **2017**, 29, 1.
- [42] Y. Zhao, M. J. Haney, R. Gupta, J. P. Bohnsack, Z. He, A. V. Kabanov, E. V. Batrakova, *PLoS One* **2014**, 9, 1.
- [43] X. Su, Y. Yu, Y. Zhong, E. Giannopoulou, X. Hu, H. Liu, J. R. Cross, G. Rättsch, C. M. Rice, L. B. Ivashkiv, *Nat. Immunol.* **2015**, 16, 838.
- [44] P. Xia, S. Wang, B. Ye, Y. Du, G. Huang, P. Zhu, Z. Fan, *Nat. Immunol.* **2015**, 16, 366.
- [45] K. Murata, H. Yoshitomi, M. Furu, M. Ishikawa, H. Shibuya, H. Ito, S. Matsuda, *Arthritis Rheumatol.* **2014**, 66, 549.
- [46] V. K. Udhayakumar, A. De Beuckelaer, J. Mccaffrey, C. M. Mccradden, J. L. Kirschman, D. Vanover, L. Van Hoecke, K. Roose, K. Deswarte, B. G. De Geest, S. Lienenklaus, P. J. Santangelo, J. Grooten, H. O. McCarthy, S. De Koker, *Adv. Healthcare Mater.* **2017**, 1601412, 1.
- [47] T. J. Anchordoquy, Y. Barenholz, D. Boraschi, M. Chorny, P. Decuzzi, M. A. Dobrovolskaia, Z. S. Farhangrazi, D. Farrell, A. Gabizon, H. Ghandehari, B. Godin, N. M. La-Beck, J. Ljubimova, S. M. Moghimi, L. Pagliaro, J. H. Park, D. Peer, E. Ruoslahti, N. J. Serkova, D. Simberg, *ACS Nano* **2017**, 11, 12.
- [48] J. McCaskill, R. Singhania, M. Burgess, R. Allavena, S. Wu, A. Blumenthal, N. A. McMillan, *Mol. Ther.–Nucleic Acids* **2013**, 2, e96.
- [49] Y. Ge, S. Li, S. Wang, R. Moore, *Nanomedicine: Principles and Perspectives*, Springer, Ottawa **2014**.
- [50] C. Fornaguera, M. García-Celma, *J. Pers. Med.* **2017**, 7, 12.
- [51] R. Bosetti, *Nanomedicine* **2015**, 10, 1851.

- [52] C. Solans, I. Solé, *Curr. Opin. Colloid Interface Sci.* **2012**, 17, 246.
- [53] A. Lesniak, A. Salvati, M. J. Santos-Martinez, M. W. Radomski, K. A. Dawson, C. Åberg, *J. Am. Chem. Soc.* **2013**, 135, 1438.
- [54] V. Forest, J. Pourchez, *Mater. Sci. Eng. C* **2016**, 70, 889.
- [55] J. Beirowski, S. Inghelbrecht, A. Arien, H. Giseler, *J. Pharm. Sci.* **2011**, 100, 1958.
- [56] C. Fornaguera, M. Llinàs, C. Solans, G. Calderó, *Colloids Surf., B* **2015**, 125, 58.
- [57] M. A. Dobrovolskaia, S. E. McNeil, *Handbook of Immunological Properties of Engineered Nanoparticles*, SAIC-Frederick Inc., **2013**.
- [58] K. L. Kozielski, Y. Rui, J. J. Green, *Expert Opin. Drug Delivery* **2016**, 13, 1475.
- [59] E. Samuelsson, H. Shen, E. Blanco, M. Ferrari, J. Wolfram, *Colloids Surf., B* **2017**, 158, 356.
- [60] S. M. Garland, S. K. Kjaer, N. Mu?oz, S. L. Block, D. R. Brown, M. J. Dinubile, B. R. Lindsay, B. J. Kuter, G. Perez, G. Dominiak-Felden, A. J. Saah, R. Drury, R. Das, C. Velicer, *Clin. Infect. Dis.* **2016**, 63, 519.
- [61] D. Putnam, C. A. Gentry, D. W. Pack, R. Langer, *Proc. Natl. Acad. Sci. USA* **2001**, 98, 1200.
- [62] M. Vilalta, I. R. Dégano, J. Bagó, E. Aguilar, S. S. Gambhir, N. Rubio, J. Blanco, *Gene Ther.* **2009**, 16, 547.
- [63] I. Pujol-Autonell, M.-J. Mansilla, S. Rodriguez-Fernandez, M. Cano-Sarabia, J. Navarro-Barriuso, R.-M. Ampudia, A. Rius, S. Garcia-Jimeno, D. Perna-Barrull, E. Martinez-Caceres, D. Maspoch, M. Vives-Pi, *Nanomedicine* **2017**, 12, 1231.

Reprint Order Form 2018 - please return with your proofs -

Editorial Office:

Wiley-VCH Verlag, Boschstrasse 12
69469 Weinheim, Germany

Tel.: (+49) 6201 606 235

Fax: (+49) 6201 606 500

E-mail: advhealthmat@wiley-vch.de
<http://www.advhealthmat.de>

Manuscript No. _____

Please send me and bill me for

no. of reprints via ☐ airmail (+ 25 Euro)
☐ surface mail

☐ high-resolution PDF file (330 Euro).

My e-mail address:

Please note: It is not permitted to present the PDF file
on the internet or on company homepages

★**Special Offer**★ If you order 200 or more reprints you
will get a PDF file for half price.

Information regarding VAT

Please note that from German sales tax point of view, the charge for
Reprints, Issues or Posters is considered as "supply of goods" and
therefore, in general, such delivery is a subject to German sales tax.
However, this regulation has no impact on customers located outside of the
European Union. Deliveries to customers outside the Community are
automatically tax-exempt. Deliveries within the Community to institutional
customers outside of Germany are exempted from the German tax (VAT)
only if the customer provides the supplier with his/her VAT number.
The VAT number (value added tax identification number) is a tax
registration number used in the countries of the European Union to identify
corporate entities doing business there. It starts with a country code (e.g.
FR for France, GB for Great Britain) and follows by numbers.

Cover Posters

Posters are available of all the published covers and
frontispieces in two sizes

☐ DIN A2 42 x 60 cm/ 17 x 24in (one copy: **39 Euro**)

☐ DIN A1 60 x 84 cm/ 24 x 33in (one copy: **49 Euro**)

Postage for shipping posters overseas by airmail:
+ 25 Euro

Postage for shipping posters within Europe by surface mail:
+ 15 Euro

Mail reprints / cover posters to:

Invoice address:

Date, Signature

Stamp

VAT no.: _____
(institutes / companies in EU countries only)

Purchase Order No.: _____

Credit Card Payment

VISA, MasterCard, AMERICAN EXPRESS

Please use the Credit Card Token Generator located at the
website below to create a token for secure payment. The token
will be used instead of your credit card number.

Credit Card Token Generator:

https://www.wiley-vch.de/editorial_production/index.php

Please transfer your token number to the space below.

Credit Card Token Number:

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Price list for reprints (The prices include mailing and handling charges. All Wiley-VCH prices are exclusive of VAT)

No. of pages	50 copies	100 copies	150 copies	200 copies	300 copies	500 copies
1-4	345	395	425	445	548	752
5-8	490	573	608	636	784	1077
9-12	640	739	786	824	1016	1396
13-16	780	900	958	1004	1237	1701
17-20	930	1070	1138	1196	1489	2022
for every additional 4 pages	147	169	175	188	231	315

★ **Special Offer** ★ If you order 200 or more reprints you will get a PDF file for half price.