This is the peer reviewed version of the following article: Allergy development is associated with consumption of breastmilk with a reduced microbial richness in the first month of life Pediatric Allergy And Immunology DOI:10.1111/pai.13176 (2019) which has been published in final form at <https://doi.org/10.1111/pai.13176>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions

Relationship between breastmilk IgA-coated microbiota, probiotic supplementation and allergy development in childhood

Majda Dzidic^{a,b}, MSc, Alex Mira^b, PhD, Alejandro Artacho^b, BSc, Thomas R. Abrahamsson^c, MD, PhD, Maria C. Jenmalm^{d,Ψ}, PhD, M. Carmen Collado^{1Ψ*}, PhD

Short title: Breastmilk microbiota and infant's allergy development

Affiliations:

a. Institute of Agrochemistry and Food Technology (IATA-CSIC), Department of Biotechnology, Unit of Lactic Acid Bacteria and Probiotics, Valencia, Spain

b. Department of Health and Genomics, Center for Advanced Research in Public Health, FISABIO, Valencia, Spain; and CIBER-ESP, Madrid; Spain

c. Department of Clinical and Experimental Medicine and Department of Pediatrics, Linköping University, Linköping, Sweden

d. Department of Clinical and Experimental Medicine, Division of Autoimmunity and Immune Regulation, Linköping University, Linköping, Sweden

Correspondence:

*To whom correspondence may be addressed:

Address: Institute of Agrochemistry and Food Technology (IATA-CSIC), Department of Biotechnology, Unit of Lactic Acid Bacteria and Probiotics, Valencia, Spain; mcolam@iata.csic.es

^ΨShared senior authors

Funding: Alex Mira: Spanish Ministry of Economy and Competitiveness (grant no. BIO2015-68711-R). Maria C. Jenmalm: The Swedish Research Council (2016-01698); the Swedish Heart and Lung Foundation (20140321 and 20170365); the Cancer and Allergy Foundation. M. Carmen Collado: European Research Council (ERC-starting grant 639226).

40
41 **Conflict of interest:** Maria C. Jenmalm has received funding for a clinical trial and honoraria
42 for lectures from BioGaia AB, as well as consultant fees and travel support from
43 Nutricia/Danone.

44

ABSTRACT

Background: Early colonization with a diverse microbiota seems to play a crucial role for appropriate immune maturation during childhood, and breastmilk microbiota is one important source of microbes for the infant, transferred together with maternal IgA antibodies. We previously observed that allergy development during childhood was associated with aberrant IgA responses to the gut microbiota already at 1 month of age, when the IgA antibodies are predominantly maternally derived in breastfed infants.

Objective: To determine the microbial composition and IgA-coated bacteria in breastmilk in relation to allergy development in children participating in an intervention trial with pre- and postnatal *Lactobacillus reuteri* supplementation.

Methods: A combination of flow cytometric cell sorting and 16S rRNA gene sequencing was used to characterize the bacterial recognition patterns by IgA in breastmilk samples collected one-month *post partum* from 40 mothers whose children did or did not develop allergic and asthmatic symptoms during the first 7 years of age.

Results: The milk fed to children developing allergic manifestations had significantly lower bacterial richness, when compared to the milk given to children that remained healthy. Probiotic treatment influenced the breastmilk microbiota composition. The proportions of IgA-bound bacteria and the total bacterial load were similar in breastmilk between mothers of healthy children and those developing allergies, but the pattern of IgA-coated bacteria tended to appear different.

Conclusion: Consumption of breastmilk with a reduced microbial richness in the first month of life may play an important role in allergy development during childhood.

Keywords: Allergy, Breastmilk, IgA, microbiota, mother-infant transfer.

INTRODUCTION

Human breastmilk is considered to be an optimal nutritional source for the immature immune system of the infant.¹ Among the bioactive factors, breastmilk contains immunoglobulins, that can be transferred to the offspring through breastfeeding.² Although all immunoglobulin isotypes can be encountered in breastmilk, secretory IgA (SIgA) is the dominating isotype and considered most important due to its anti-inflammatory properties and important role in defending the mucous membranes, thus regulating the binding and invasion of commensals and pathogenic microorganisms.³ This passive immunization through breastfeeding is crucial as early production of secretory IgA in newborns is limited.⁴

The increasing prevalence of allergic diseases in affluent societies is hypothesized to be caused by reduced intensity and diversity of microbial stimulation.^{5,6} In support of this theory, the gut microbiota differs in composition and diversity during the first months of life in children who later do or do not develop allergic disease.⁵ Breastmilk hosts a diverse array of microbiota and potential probiotic bacteria, transferred together with maternal IgA antibodies⁷, likely influencing the infant's developing mucosal immune system.⁸

We have previously observed that allergy development during childhood was associated with divergent patterns of IgA recognized bacteria in the gut already at 1 month of age, when the IgA antibodies are predominantly maternally derived in breast-fed children.⁹ However, the identities of the bacterial taxa targeted by IgA in the breastmilk and what role they may play in immune and allergy development are unknown. In this study, we aimed to characterize the composition and IgA coating pattern of the breastmilk microbiota from mothers whose children developed allergic symptoms during early childhood or stayed healthy.

98

99 **METHODS**

100 For detailed methods, experimental protocols and statistical analyses, see the Methods section
101 in this article's Online Repository.

102 **Study design**

103 The subjects included in this study were part of a larger randomized double-blind trial in
104 Sweden, recruiting participants between 2001 and 2003, where the potential allergy
105 preventive effect of probiotic *Lactobacillus reuteri* ATCC 55730 in the infants with family
106 history of allergic disease was evaluated.^{10,11} Among the 184 mothers of children that
107 completed the 7-year follow up in the original study, breastmilk samples, at one month *post*
108 *partum*, from 24 mothers whose children did not and from 26 mothers whose children did
109 develop allergic disease during early childhood, were randomly selected for flow cytometry
110 based-sorting of IgA-coated bacteria in the current study. From these flow cytometry-sorted
111 samples, subsequent 16S rRNA gene characterization was performed on the IgA-coated and
112 IgA-free fractions of breastmilk bacteria from 20 mothers whose children did not and from 20
113 mothers whose children did develop allergic manifestations as well as from total, non-sorted
114 breastmilk samples from the same mothers. Selection of the samples used for 16S rRNA
115 sequencing in this study was based on the sample availability and required sample volume for
116 Illumina sequencing, and a clear allergy diagnosis (based on proven symptoms to allergy
117 provocation) of the child. Allergic disease included eczema, gastrointestinal allergy, asthma,
118 allergic rhinoconjunctivitis and allergic urticaria. The criteria of these diagnoses are described
119 in detail in ^{10,11} and in the supplementary information of this manuscript. All the children
120 included in the current study were exclusively breastfed during the first month of life.

There were no differences regarding potential confounders, such as sex, mode of delivery, birth order, maternal atopy, breastfeeding, antibiotics, and probiotic supplementation, between the infants who did or did not have allergic manifestations (Table I).

Total IgA levels were measured by ELISA in a study by Böttcher *et al.*¹² The studies were approved by the Regional Ethics Committee for Human Research in Linköping, Sweden (Dnr 99323, M122-31 and M171-07, respectively).

Sample preparation and flow cytometry-based sorting

The breastmilk samples were stained with goat anti-mouse IgA labelled with fluorescein isothiocyanate (FITC), used as an isotype control corresponding to unspecific binding (Sigma; reference SLBD9273), or with goat anti-human IgA labelled with FITC (Life Technologies; reference A18782), according to the manufacturer's instructions. The sorting of the bacterial cells according to whether they were IgA-coated or IgA-free was performed with the MoFlo XDP Cell Sorter (Beckman Coulter, Brea, Calif), according to the procedures of Simon-Soro *et al.*¹³

DNA Extraction

DNA from sorted breastmilk bacteria, both IgA+ and IgA-, as well as the total milk sample was isolated by using the MasterPure complete DNA and RNA Purification Kit (Epicentre Biotechnologies, Madison, Wis), according to the manufacturer's instructions.

16S rRNA gene amplification and sequencing

DNA from sorted bacterial fractions (in total 80) together with total non-sorted breastmilk

samples (in total 40) was used for PCR amplification and Illumina sequencing to describe the bacterial composition of breastmilk. Sequences supporting the conclusions of this article are publicly available at European Nucleotide Archive database (ENA) with accession number PRJEB30065.

Sequence analysis

16S rRNA gene reads from the total milk samples were used in order to perform an accurate filtering of the flow cytometry IgA-sorted fractions that, due to low bacterial yield, were more susceptible to sequencing contaminations. This was done by eliminating OTUs in IgA sorted fractions that were absent in corresponding total non-sorted milk samples.

For analyzing IgA coating patterns, the IgA index score (calculated according the formula $\log(IgA+/IgA-)$) was used to describe the degree of mucosal immune responsiveness to the microbiota.

Statistical analyses were performed in R version 3.2.2 and GraphPad Prism 6 (GraphPad Software, San Diego, CA, USA, Version 6.1f), where $p < 0.05$ was considered significant.

RESULTS

IgA proportions in breastmilk

On average, approximately 40% of bacteria in breastmilk appeared to be IgA-coated. Proportions of IgA-coated bacteria were similar in breastmilk samples of mothers whose children did or did not develop allergic (Fig. 1A) and asthmatic symptoms (Fig. 1B) during the first 7 years of life. Furthermore, IgA proportions observed did not seem to be influenced by the total IgA levels in breastmilk samples of these mothers ($n=29$; Spearman correlation test $r=0.32$, $p=0.095$). The proportions of IgA-coated bacteria were lower in breastmilk from

mothers supplemented with probiotics during the last month of pregnancy, as compared with placebo ($p=0.04$; Fig. S1A in this article's Online Repository), and particularly in breastmilk of probiotic supplemented mothers whose children stayed healthy ($p=0.02$; Fig. S1B).

Bacterial diversity, richness and density in total non-sorted milk samples and IgA-coated fractions

The overall species richness (as determined by Chao1 index) in total non-sorted breastmilk samples was significantly higher ($p=0.02$, Fig. 2A) in mothers with healthy children, although the bacterial load (Fig. S2) and species diversity (Shannon index, Fig. 2B) were similar in breastmilk samples of mothers whose children did/did not develop allergic manifestations. The total species richness also tended to be higher in breastmilk from mothers whose children stayed healthy than from mothers whose children developed asthmatic symptoms ($p=0.066$, Fig. 2A). However, no significant differences between mothers of healthy and allergic subjects were observed upon comparing the richness and diversity of the IgA-bound fractions in breastmilk (Table SI). Additionally, no differences in bacterial load, species diversity or richness was observed between the total non-sorted breastmilk from the mothers treated with probiotics and placebo (data not shown).

Bacterial composition in allergy development and probiotic supplementation

Bacterial 16S rRNA gene sequencing, of the total non-sorted milk and IgA-coated/uncoated fractions, was performed in order to determine the milk microbial composition and to assess IgA responses towards specific bacteria. After quality filtering and removal of chimeric sequences, 40 total non-sorted breastmilk samples and 80 IgA separated fractions remained with 2,000,107 and 1,525,770 high quality reads, respectively. Sequencing of total non-sorted breastmilk samples resulted in an average of $48,987 \pm 2725$ (SEM) reads per sample, while the

IgA separated breastmilk fractions had an average of $20,322 \pm 1660$ (SEM) sequence reads per sample.

The relative abundance of genera in total breastmilk, i.e. non-sorted samples, is presented in Fig. 3A and 3B. Infants developing allergies during the first 7 years of life tended to have higher abundance of the genera *Enterococcus* ($p=0.01$; adj. p -value=0.18; Fig. S3A) and *Pseudomonas* ($p=0.01$; adj. p -value=0.18; Fig.S3B). The genus *Enterococcus* was found in significantly higher abundance ($p=0.001$; adj. p -value= 0.02) in breastmilk given to children that developed asthmatic symptoms, when compared to breastmilk given to the children that remained healthy (Fig. 3B, Fig. S3A). No effect of the confounding factors, such as probiotic supplementation during pregnancy, maternal atopy, sex and the delivery mode influenced the microbial abundance between the groups, as determined with the MaAslin multivariate statistical logarithm.

Microbiota composition patterns of total non-sorted breastmilk, were significantly different ($p=0.04$ Adonis testing) between mothers that received, or not, probiotic supplementation (Fig. 4A) with significantly increased relative abundance ($p=0.03$, Wilcoxon rank-sum test adjusted p -value) of the genus *Rothia* in mothers treated with placebo. However, upon comparing IgA pattern recognitions of bacteria in mothers who were treated, or not, with probiotics supplementation during the last month of the pregnancy, no statistically significantly different IgA responses could be observed for bacterial genera presented (Fig. 4B).

IgA responses towards milk microbiota in allergy development

Although the analysis of the relative abundance of dominant bacterial genera in breastmilk, and the composition patterns of sorted IgA fractions (Fig. S4) were generally similar between the IgA-bound and IgA-free fractions, upon considering the health status of the children,

some differences at genus level were observed when analyzing the bacterial targets of IgA responses, represented as the IgA index here. The value of the IgA index can range from positive values, reflecting genera found dominantly in the IgA positive fraction, to negative values (genera found dominantly in the IgA negative fraction). However, no statistically significant differences could be observed in IgA-coating patterns of breastmilk bacteria given to children that did and did not developed allergic and/or asthmatic symptoms (Fig. 5A-B).

DISCUSSION

The data presented in the current study demonstrate that total non-sorted breastmilk from mothers whose children developed allergic symptoms during early childhood had lower bacterial richness, when compared to milk fed to children staying healthy. Moreover, probiotic supplementation during pregnancy modifies the breastmilk microbiota composition and the proportion of IgA coated bacteria.

The influence of breastmilk composition on later allergy development appears to be linked to higher richness of bacterial species and not to the relative abundance of specific bacteria. Previously, low total diversity of the gut and oral microbiota have been associated with atopy and asthma development during early childhood.^{14–17} Moreover, aberrant IgA immune responses towards gut microbiota were observed as early as 1 month *post partum*, in exclusively breastfed children who subsequently developed allergies.⁹ At this time point, the IgA antibodies are predominantly maternally derived in exclusively breastfed children, as the levels of endogenously produced IgA in the baby during this period is limited.¹⁸ Therefore, any divergent responses observed at this time suggest that immunological interactions between mother and infant may play an important role in subsequent immune development.

Streptococcus, *Acinetobacter*, *Staphylococcus* and *Veillonella* were the most commonly found bacterial genera in milk samples used in this study. Furthermore, lactic acid bacteria *Lactococcus*, *Lactobacillus* and *Enterococcus* as well as oral inhabitant *Gemella* were also detected, in agreement with previous reports.¹⁹ Various studies have demonstrated that there is a mother-to-infant transfer of bacterial genera including *Lactobacillus*, *Staphylococcus*, *Enterococcus* and *Bifidobacterium* through breastfeeding.^{20–23} The constant intake during lactation of breastmilk bacteria leads to the establishment of an intestinal microbiota that deeply impacts on the newborn's immune maturation.²⁴ An

interesting finding is the significantly higher levels of the lactic acid bacteria *Enterococcus* in breastmilk fed to children developing asthma. *Enterococcus* is a common inhabitant of breastmilk microbiota and one of the first microbes to colonize the infant gut after birth^{15,25}, being more abundant in the gut of atopic infants, from the same cohort, at 12 months of age.¹⁵ However, whether our finding here reflects an overgrowth of *Enterococcus* due to the absence of competing species, or whether *Enterococcus* is suppressing the growth of allergy protecting bacteria remains to be addressed.

An important function of SIgA is immune exclusion, a mechanism where this antibody binds to commensals, through its recognition of multiple antigenic epitopes on the surface, encountered in gut lumen and prevents their attachment to mucosal barrier.²⁶ Moreover, this antibody can also promote bacterial adhesion to the mucosa, as shown *in vitro* and *in vivo*^{27,28}, thus enriching for the growth of particular microbial strains.²⁹ In this study, we were not able to observe significant differences in the patterns of IgA-binding in breastmilk fed to children staying healthy and children developing allergies but the possibility that the true differences are at the lower bacterial resolution (for instance species and even different strains), should not be excluded. The fact that the majority of the mothers included in this study, and also in the original study of this cohort, suffered from allergic disease, should be considered as it also may affect both the IgA-coating patterns and the microbiota transferred to the child.

IgA-coating proportions of breastmilk bacteria were higher in mothers treated with placebo, compared to those treated with probiotic *L. reuteri*. As the mothers were treated with probiotics until delivery, likely the changes in the microbial composition were more significant in colostrum than in breastmilk samples extracted at one month *post partum*, as in the present study. We have previously shown that *Lactobacilli* colonization was significantly increased in colostrum of the mothers treated with *L. reuteri*, but not in samples obtained one

month *post partum*.³⁰ This could suggest that part of the mother's IgA responses, while treated with probiotics, were directed towards *L. reuteri*, but that the ending of supplementation was reflected in diminished targeting of this species and perhaps decreased proportions of IgA-bound microbiota. Additionally, microbiota patterns of total non-sorted milk were also significantly different between mothers treated with probiotics and those with placebo, with the genus *Rothia* (a common oral inhabitant associated with good oral health³¹) significantly increased in the placebo group. However, the importance of these results needs to be further confirmed and evaluated in larger cohorts.

Together with the maternal intestinal and vaginal microorganisms that are ingested by the neonate during the passage through the birth canal, breastmilk microbiota appears to contribute to the initial microbial colonization in infants, thus having a pivotal role in modulating and influencing the newborns' immune system. The bacterial species that initially colonize the mucosal surfaces likely define the ecosystem conditions which in turn will affect the establishment of further co-colonization patterns. Also, milk microbes are transferred together with maternal IgA antibodies, that may enable maintenance of a mutually beneficial relationship with a diverse set of commensals, while protecting against pathogens.

In conclusion, consumption of breastmilk with a reduced microbial richness in the first month of life correlates with the risk for allergy development during childhood. In addition, our data show that probiotic supplementation during pregnancy alters the breastmilk microbiota composition and the proportion of IgA coated bacteria. This opens the possibility of modulating breastmilk microbiota and its interaction with antibodies as a strategy to promote a healthy microbial colonization with the purpose of reducing allergy risk.

REFERENCES

1. Walker A. Breast Milk as the Gold Standard for Protective Nutrients. *J Pediatr*. 2010 Feb;156(2):S3–7.
2. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*. 2013 Feb;60(1):49–74.
3. Lepage P, Van de Perre P. The Immune System of Breast Milk: Antimicrobial and Anti-inflammatory Properties. In: *Advances in experimental medicine and biology*. 2012. p. 121–37.
4. Newburg DS, Walker WA. Protection of the Neonate by the Innate Immune System of Developing Gut and of Human Milk. *Pediatr Res*. 2007 Jan;61(1):2–8.
5. West CE, Jenmalm MC, Prescott SL. The gut microbiota and its role in the development of allergic disease: a wider perspective. *Clin Exp Allergy*. 2015 Jan;45(1):43–53.
6. West CE, Dzidic M, Prescott SL, Jenmalm MC. Bugging allergy; role of pre-, pro- and synbiotics in allergy prevention. *Allergol Int*. 2017;66(4).
7. Jenmalm MC. The mother–offspring dyad: microbial transmission, immune interactions and allergy development. *J Intern Med*. 2017 Jul;282(6):484–95.
8. Cacho NT, Lawrence RM. Innate Immunity and Breast Milk. *Innate Immun Breast Milk Front Immunol*. 2017;8(584):e-publication.
9. Dzidic M, Abrahamsson TRTR, Artacho A, Björkstén B, Collado MCMC, Mira A, et al. Aberrant IgA responses to the gut microbiota during infancy precede asthma and allergy development. *J Allergy Clin Immunol*. 2017 Mar;139(3):1017-1025.e14.
10. Abrahamsson TR, Jakobsson T, Böttcher MF, Fredrikson M, Jenmalm MC, Björkstén B, et al. Probiotics in prevention of IgE-associated eczema: A double-blind, randomized, placebo-controlled trial. *J Allergy Clin Immunol*. 2007 May;119(5):1174–

- 326 80.
- 327 11. Abrahamsson TR, Jakobsson T, Björkstén B, Oldaeus G, Jenmalm MC. No effect of
328 probiotics on respiratory allergies: a seven-year follow-up of a randomized controlled
329 trial in infancy. *Pediatr Allergy Immunol*. 2013 Sep;24(6):556–61.
- 330 12. Böttcher MF, Abrahamsson TR, Fredriksson M, Jakobsson T, Björkstén B. Low breast
331 milk TGF- β 2 is induced by *Lactobacillus reuteri* supplementation and associates with
332 reduced risk of sensitization during infancy. *Pediatr Allergy Immunol*. 2008
333 Aug;19(6):497–504.
- 334 13. Simón-Soro Á, D'Auria G, Collado MC, Džunková M, Culshaw S, Mira A. Revealing
335 microbial recognition by specific antibodies. *BMC Microbiol*. 2015 Jul;15(1):132.
- 336 14. Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm
337 MC. Low gut microbiota diversity in early infancy precedes asthma at school age. *Clin*
338 *Exp Allergy*. 2014 Jun;44(6):842–50.
- 339 15. Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm
340 MC. Low diversity of the gut microbiota in infants with atopic eczema. *J Allergy Clin*
341 *Immunol*. 2012 Mar;129(2):434–40.
- 342 16. Dzidic M, Abrahamsson T, Artacho A, Collado MC, Mira A, Jenmalm MC. Oral
343 microbiota maturation during the first 7 years of life in relation to allergy development.
344 *Allergy*. 2018 Mar;0(ja).
- 345 17. Azad MB, Konya T, Guttman DS, Field CJ, Sears MR, HayGlass KT, et al. Infant gut
346 microbiota and food sensitization: associations in the first year of life. *Clin Exp*
347 *Allergy*. 2015 Mar;45(3):632–43.
- 348 18. Tourneur E, Chassin C. Neonatal immune adaptation of the gut and its role during
349 infections. *Clin Dev Immunol*. 2013 May;2013:270301.
- 350 19. Fitzstevens JL, Smith KC, Hagadorn JI, Caimano MJ, Matson AP, Brownell EA.

- 351 Systematic Review of the Human Milk Microbiota. *Nutr Clin Pract*. 2016 Sep;32:354–
352 64.
- 353 20. Martín R, Langa S, Reviriego C, Jiménez E, Marín ML, Xaus J, et al. Human milk is a
354 source of lactic acid bacteria for the infant gut. *J Pediatr* [Internet]. 2003;143(6):754–8.
- 355 21. Soto A, Martín V, Jiménez E, Mader I, Rodríguez JM, Fernández L. Lactobacilli and
356 Bifidobacteria in Human Breast Milk: Influence of Antibiotherapy and Other Host and
357 Clinical Factors. *J Pediatr Gastroenterol Nutr*. 2014 Jul;59(1):78–88.
- 358 22. Martín V, Maldonado-Barragán A, Moles L, Rodriguez-Baños M, Campo R del,
359 Fernández L, et al. Sharing of Bacterial Strains Between Breast Milk and Infant Feces.
360 *J Hum Lact*. 2012 Jan;28(1):36–44.
- 361 23. Benito D, Lozano C, Jiménez E, Albújar M, Gómez A, Rodríguez JM, et al.
362 Characterization of *Staphylococcus aureus* strains isolated from faeces of healthy
363 neonates and potential mother-to-infant microbial transmission through breastfeeding.
364 *FEMS Microbiol Ecol*. 2015 Mar;91(3):fiv007–fiv007.
- 365 24. Houghteling PD, Walker WA. Why Is Initial Bacterial Colonization of the Intestine
366 Important to Infants' and Children's Health? *J Pediatr Gastroenterol Nutr*. 2015
367 Mar;60(3):294–307.
- 368 25. Mshvildadze M, Neu J. The infant intestinal microbiome: friend or foe? *Early Hum*
369 *Dev*. 2010;86.
- 370 26. Hurley WL, Theil PK. Perspectives on immunoglobulins in colostrum and milk.
371 *Nutrients*. 2011;3(4):442–74.
- 372 27. Donaldson GP, Ladinsky MS, Yu KB, Sanders JG, Yoo BB, Chou WC, et al. Gut
373 microbiota utilize immunoglobulin A for mucosal colonization. *Science* (80-)
374 [Internet]. 2018;360(6390):795–800.
- 375 28. McLoughlin K, Schluter J, Rakoff-Nahoum S, Smith AL, Foster KR. Host Selection of

Microbiota via Differential Adhesion. *Cell Host Microbe*. 2016;19(4):550–9.

29. Rogier EW, Frantz AL, Bruno MEC, Wedlund L, Cohen DA, Stromberg AJ, et al.

Secretory antibodies in breast milk promote long-term intestinal homeostasis by regulating the gut microbiota and host gene expression. *Proc Natl Acad Sci*. 2014 Feb;111(8):3074–9.

30. Abrahamsson TR, Sinkiewicz G, Jakobsson T, Fredrikson M, Björkstén B. Probiotic Lactobacilli in Breast Milk and Infant Stool in Relation to Oral Intake During the First Year of Life. *J Pediatr Gastroenterol Nutr*. 2009;49(3).

31. Lundberg JO, Carlström M, Weitzberg E. Metabolic Effects of Dietary Nitrate in Health and Disease. *Cell Metab*. 2018;28(1):9–22.

388

Table I. Descriptive data of children compare in this study.

Children	Healthy (% [no.] n=20	Developing allergic disease (% [no.] n=20	P value*	Developing asthma (% [no.] n=10	P value*
Girls	50(10)	60(12)	0.53	60(6)	0.71
Older siblings	40(8)	35(7)	1.00	30(3)	0.70
Caesarean delivery	20(4)	20(4)	1.00	20(2)	1.00
Furred pets	15(3)	10(2)	1.00	10(1)	1.00
Breastfeeding					
1 month exclusive	90(18)	95(19)	1.00	90(9)	1.00
3 months exclusive	85(17)	70(14)	0.45	50(5)	0.08
12 months partial	20(4)	25(5)	1.00	0(0)	0.27
Maternal atopy	85(17)	80(16)	1.00	70(7)	0.37
Antibiotic treatment (1-12 m)	35(7)	30(6)	1.00	50(5)	0.46
Day care (1-12 m)	10(2)	5(1)	1.00	10(1)	1.00
Probiotics-<i>L. reuteri</i>	45(9)	55(11)	0.53	60(6)	0.70

404

*The x2 test was used to detect potential differences in frequencies between groups, except when the expected frequency for any cell was less than 5, in which case the Fisher exact test was used.

FIGURE LEGENDS

Figure 1. Proportions of IgA-coated breastmilk bacteria collected one-month *post partum*. A) Proportion of breastmilk IgA-bound bacteria in mothers whose children stayed healthy (n=24, circles) or developed allergic symptoms (n=26, triangles) during the first 7 years of life, as determined by flow cytometry based-cell sorting. B) Proportion of breastmilk bacteria bound to IgA at one month of age in children staying healthy (n=24) or developing asthmatic symptoms (n=13) during the first 7 years of life. Medians and interquartile ranges are indicated.

Figure 2. Bacterial richness (A) and diversity (B) at 16S rRNA OTU's species level, as described with Chao1 and Shannon indices, respectively, of the total non-sorted breastmilk samples from mothers whose children stayed healthy up to 7 years of age (n=20), mothers whose children developed allergic manifestations (n=20) and/or asthmatic symptoms (n=10). Medians and interquartile ranges are indicated. * p-value <0.05, Mann-Whitney U test.

Figure 3. Microbiota composition of most dominant bacterial genera in total non-sorted milk samples. A) The relative abundance (>0.5 % of the total) of dominant bacterial genera in 20 mothers whose children stayed healthy and in 20 mothers whose children developed allergic manifestation during their first 7 years of life. B) The relative abundance of dominant bacterial genera in breastmilk samples in 20 mothers whose children stayed healthy and 10 mothers whose children developed asthmatic manifestation.

Figure 4. Breastmilk microbiota composition and IgA-coating patterns of the dominant genera (>0.5% of total), in probiotic supplementation. A) Constrained correspondence analyses (CCA) based on microbiota composition patterns (determined by 16S rRNA

sequencing) in breastmilk of mothers treated with probiotics *L. reuteri* or placebo. The percentage of variation explained by constrained correspondence components is indicated on the axes. *p* value for CCA plots were determined by Adonis ($p=0.04$) and indicate if the factor provided (in this case *probiotics*) can significantly explain data variability. B) Plots represent IgA coating patterns (defined by IgA index, reflecting the ratio in IgA bound and IgA free breastmilk microbiota) to dominant genera in breastmilk samples collected at one month *post partum* from mothers whose were treated with probiotic *L. reuteri* or placebo. For a given genera, the value of the IgA index can range from positive values, reflecting genera found dominantly in the IgA bound fraction, to negative values (genera found dominantly in the IgA free fraction). $n_{\text{Probiotics}}=20$, $n_{\text{Placebo}}=20$.

Figure 5. IgA responses towards breastmilk microbiota. Plots represent IgA-coating patterns to dominant genera (>0.5 % of total) from mothers whose children stay healthy ($n=20$) compared to children that develop allergic (A, $n=20$) or asthmatic (B, $n=10$) symptoms. For a given genera, the value of the IgA index can range from positive values, reflecting genera found dominantly in the IgA-bound fraction, to negative values (genera found dominantly in the IgA-free fraction). Means with SEs are indicated. Wilcoxon rank-sum test with FDR correction was performed.

SUPPLEMENTARY INFORMATION for - *Relationship between breastmilk IgA-coated microbiota, probiotic supplementation and allergy development in childhood*

METHODS

Study design

The subjects included in this study were part of a larger randomized double-blind trial in South-eastern Sweden, recruiting participants between 2001 and 2003, where the potential allergy preventive effect of probiotic *Lactobacillus reuteri* ATCC 55730 in the infants with family history of allergic disease was evaluated.^{1,2} The mothers were supplemented with *L. reuteri* during pregnancy from postmenstrual week 36+0 to delivery and the infants from day 1-2 of life until 12 months of age. Among the 184 mothers, our of 232 recruited, that completed the 7-year follow up in the original study, breastmilk samples from 24 mothers whose children did not and from 26 mothers whose children did develop allergic disease during early childhood, were used for flow cytometry based-sorting of IgA-coated bacteria in the current study. Subsequently, 16S rRNA gene characterization was performed on the IgA-coated and IgA-free fractions of breastmilk bacteria from 20 mothers whose children did not and from 20 mothers whose children did develop allergic manifestations as well as from total, non-sorted breastmilk samples from the same mothers. The selection of the samples used for 16S rRNA sequencing in this study, was based on the sample availability and a clear allergy diagnosis (based on proven symptoms to allergy provocation) of the child. There were no differences regarding potential confounders, such as sex, mode of delivery, birth order, maternal atopy, breastfeeding, antibiotics, and probiotic supplementation, between the infants who did or did not have allergic manifestations. Allergic disease included eczema, gastrointestinal allergy, asthma, allergic rhinoconjunctivitis and allergic urticaria. The criteria of these diagnoses are described in detail in^{1,2}. All children with allergic disease in the current study were also sensitized (*i.e.* they had at least one positive skin prick test (evaluated at 6, 12 and 24 months and 7 years of age) and/or detectable circulating allergen specific-IgE antibodies (assessed at 6, 12 and 24 months), while

the healthy children were non-sensitized.

Breastmilk samples were collected one-month *post partum* by the mother at home. They were immediately placed in the freezer and brought to the hospital and stored at -70°C within 3 days. All the children included in the current study were exclusively breastfed during the first month of life. Total IgA levels were measured by ELISA in a study by Böttcher *et al.*³

The studies were approved by the Regional Ethics Committee for Human Research in Linköping, Sweden (Dnr 99323, M122-31 and M171-07, respectively). An informed consent was obtained from both parents before inclusion in the study.

Sample preparation and flow cytometry-based sorting

After thawing, the breastmilk fatty layer and whey were removed by centrifugations. The resulting sample fraction was further suspended in sterile saline solution (autoclaved H_2O ; NaCl Sodium Chloride 99.5% PA-ACS-ISO; Panreac, Barcelona, Spain; reference 131689.1211) with 5% BSA (Sigma-Aldrich, St Louis, Mo; reference A7030-100gr) to prevent nonspecific antibody binding. The samples were stained with goat anti-mouse IgA labelled with fluorescein isothiocyanate (FITC), used as an isotype control corresponding to unspecific binding (Sigma; reference SLBD9273), or with goat anti-human IgA labelled with FITC (Life Technologies; reference A18782), according to the manufacturer's instructions. The sorting of the bacterial cells according to whether they were IgA-coated or IgA-free was performed with the MoFlo XDP Cell Sorter (Beckman Coulter, Brea, Calif), according to the procedures of Simon- Soro *et al.*⁴

DNA Extraction

DNA from sorted breastmilk bacteria, both IgA+ and IgA-, as well as the total milk sample (2 ml) was isolated by using the MasterPure complete DNA and RNA Purification Kit (Epicentre

Biotechnologies, Madison, Wis), according to the manufacturer's instructions, with a previous glass bead beating (0.17 mm in diameter) and an additional enzymatic lysis step with lysozyme (20 mg/mL, 37°C, 30 minutes; Thermo-mixer comfort, Eppendorf, Hamburg, Germany), lysostaphin (2000 units/mg protein, 37 °C, 60 min; Sigma-Aldrich, Madrid, Spain) and mutanolysin (4000 units/mg protein, 37 °C, 60 min; Sigma-Aldrich).

16S rDNA gene amplification and sequencing

DNA from sorted bacterial fractions (in total 80) together with total non-sorted breastmilk samples (in total 40) was used for PCR amplification and Illumina sequencing to describe the bacterial composition of breastmilk. Universal bacterial degenerate primers 8F—5'-AGAGTTTGATCMTG GCTCAG-3' and 926R—5'-CCGTCAATTCMTTTRAGT- 3', which encompass the hypervariable regions V1–V5 of 16S ribosomal RNA (rRNA) gene were used for an initial amplification in order to increase the bacterial yield. Purification of PCR products was completed using Nucleofast 96 PCR filter plates (Macherey-Nagel, Düren, Germany).

An Illumina amplicon library was performed following the 16S rRNA gene Metagenomic Sequencing Library Preparation Illumina protocol (Part #15044223 Rev. A). The gene-specific primer sequences used in this protocol were selected from Klindworth et al.⁵, and target the 16S rRNA gene V3 and V4 regions, resulting in a single amplicon of approximately 460 bp. After 16S rRNA gene amplification, the DNA was sequenced on a MiSeq Sequencer according to manufacturer's instructions (Illumina) using the 2 × 300 bp paired-end protocol. Sequences supporting the conclusions of this article are publicly available at European Nucleotide Archive database (ENA) with accession number PRJEB30065.

Total bacterial load

Total bacterial load (number of bacterial cells per ml breastmilk) was measured by quantitative PCR. Amplifications were performed in duplicates on a LightCycler 480 Real-Time PCR System (Roche Technologies) by using annealing temperatures of 60 °C. Each reaction mixture

of 10mL was composed of SYBR Green PCR Master Mix (Roche), 0.5 mL of the specific primer (concentration 10 mmol/L), and 2 mL of DNA template. The universal forward and reverse primers were 5'-CGTGCCAGCAGCCGCGG-3' and 5'-TGGACTACCAGGGTATCTAATCCTG-3', targeting a 293bp long region of the bacterial 16S rRNA gene. The obtained Ct values were transformed in bacterial cell numbers by a calibrated standard curve.⁶

Bioinformatics

The PRINSEQ program was used for a sequence quality assessment.⁷ Sequences of <250 nucleotides in length were discarded; sequence end-trimming was performed by cutting out nucleotides with a mean quality of <30 in 20-bp windows. Chimeric 16S sequences were filtered out using USEARCH program.⁸ OTUs were built at 97% of identity by using vsearch program and Qiime modules (version qiime2-2017.12) were used for taxonomic annotation.⁹ In order to assign taxonomy up to species level to each OTU's centroid, we classified them using a naïve bayes classifier model previously fitted against the Green Genes database version 13.5.

16S rRNA gene reads from the total milk samples were used in order to perform an accurate filtering of the flow cytometry IgA-sorted fractions that, due to low bacterial yield, were more susceptible to sequencing contaminations. This was done by eliminating OTUs in IgA sorted fractions that were absent in corresponding total non-sorted milk samples. Moreover, OTUs were also removed in cases where either IgA positive or IgA negative fractions presented an abundance of less than five reads, compared to its corresponding OTU in non-sorted milk sample. In addition, we filtered out low signal OTUs that were presented in less than five sequences through the total set of samples.

α -diversity analyses (presented here as Shannon and Chao1 indices), were utilized to estimate

the samples' diversity and richness at the 97% OTU level using the R-package Vegan.¹⁰ Constrained correspondence analysis (CCA) was used here to emphasize variation and bring out strong patterns in the dataset. This analysis was performed by R software ade4 package together with permutational multivariate analysis (Adonis) determining the differences in variance between groups.

For analyzing IgA coating patterns, the threshold used for including the genera was 0.5 % or greater in relative abundance in either the IgA positive or IgA negative fractions. A pseudocount that was equal to 0.001 was added to every genus dedicated in both the IgA positive or IgA negative fractions, thus avoiding the fractions with a value of zero. The abundance proportions of a given genera were log-transformed before calculating the ratio between the IgA positive or IgA negative fractions, resulting in the IgA index.^{11,12} The IgA index (calculated according the formula $\log(IgA+/IgA-)$) score reflects the degree of mucosal immune responsiveness to the microbiota, where the positive values represent the genera predominantly found IgA-coated while the negative values the bacterial genera predominantly found IgA uncoated.

The MaAslin multivariate statistical framework was used in this study in order to evaluate if the confounding factors, including probiotics supplementation during pregnancy and maternal atopy, could influence microbial community abundance.¹³ Statistical analyses were performed in R version 3.2.2 and GraphPad Prism 6 (GraphPad Software, San Diego, CA, USA, Version 6.1f), where $p < 0.05$ was considered significant. Specific statistical tests (including Mann–Whitney U-test/ Wilcoxon rank-sum test for nonparametric comparisons together with false discovery rate control giving the adjusted p-value) are stated in figure legends.

125
126
127

RESULTS

Table S1. Chao1 and Shannon indices describing the species richness and diversity, respectively, in total milk samples, in the IgA bound and IgA free fractions. Mean and standard error of the mean (SEM) are indicated.

HEALTHY non-sorted (n=20)		ALLERGIC non-sorted (n=20)		ASTHMA non-sorted (n=10)		p-value
Chao1	SEM	Chao1	SEM	Chao1	SEM	H vs Allergic / H vs Asthmatic
542.26	32.96	430.30	40.74	424.88	71.38	0.018*/ 0.065
HEALTHY IgA-bound (n=40)		ALLERGIC IgA-bound (n=40)		ASTHMA IgA-bound (n=20)		p-value
Chao1	SEM	Chao1	SEM	Chao1	SEM	H vs Allergic / H vs Asthmatic
62.67	6.26	65.88	9.39	70.33	17.05	0.733/0.820
HEALTHY IgA-free (n=40)		ALLERGIC IgA-free (n=40)		ASTHMA IgA-free (n=20)		p-value
Chao1	SEM	Chao1	SEM	Chao1	SEM	H vs Allergic / H vs Asthmatic
57.09	5.45	58.14	6.75	58.42	11.78	0.898/ 0.820
HEALTHY non-sorted (n=20)		ALLERGIC non-sorted (n=20)		ASTHMA non-sorted (n=10)		p-value
Shannon	SEM	Shannon	SEM	Shannon	SEM	H vs Allergic / H vs Asthmatic
1.96	0.11	1.87	0.19	1.82	0.30	0.351/0.411
HEALTHY IgA-bound (n=40)		ALLERGIC IgA-bound (n=40)		ASTHMA IgA-bound (n=20)		p-value
Shannon	SEM	Shannon	SEM	Shannon	SEM	H vs Allergic / H vs Asthmatic
2.45	0.10	2.38	0.11	2.42	0.18	0.490/0.893
HEALTHY IgA-free (n=40)		ALLERGIC IgA-free (n=40)		ASTHMA IgA-free (n=20)		p-value
Shannon	SEM	Shannon	SEM	Shannon	SEM	H vs Allergic / H vs Asthmatic
2.40	0.09	2.32	0.15	2.18	0.28	0.875/0.859

* Mann-Whitney U test.

Fig. S1. IgA coating levels in the breastmilk of mothers treated, or not, with *L. reuteri*.

(A) Proportions of IgA bound bacteria in breastmilk of mothers treated with *L. reuteri* probiotics and mothers treated with placebo. $p=0.04$; $n_{\text{Placebo}}=27$, $n_{\text{Probiotics}}=24$ (B) Proportions of IgA bound bacteria in breastmilk of mothers, treated with probiotics or not, whose children stayed healthy ($p=0.02$) or developed allergic diseases

during the first 7 years of age. Healthy= n_{Placebo}=11, n_{Probiotics}=13; Allergic: n_{Placebo}=13, n_{Probiotics}=14. Media with interquartile ranges are indicated; * Mann-Whitney U test.

Fig.S2. Bacterial load in breastmilk collected at one month *post partum*. Quantification of bacterial numbers was obtained by using qPCR detection with universal primers targeting the 16S rRNA bacterial gene. n_{Healthy} =19; and n_{Allergic} = 19; n_{Asthmatic} =9. Media with interquartile ranges are indicated.

Fig. S3. Selected bacterial genera found in milk of mothers whose children stay healthy or develop allergic and/or asthmatic manifestations. Mean with SEM are presented. * $p < 0.05$, Mann-Whitney U test, FDR adjusted p-value.

Fig. S4. Breastmilk microbiota composition patterns of sorted IgA fractions, IgA-coated and IgA-free, in allergy development. Constrained correspondence analyses (CCA) based on breastmilk microbiota patterns of the dominant bacterial genera (>0.5% of total) coated with IgA (A) or not coated with IgA (B), from mothers whose children did or did not develop allergic diseases.

REFERENCES

1. Abrahamsson TR, Jakobsson T, Böttcher MF, Fredrikson M, Jenmalm MC, Björkstén B, et al. Probiotics in prevention of IgE-associated eczema: A double-blind, randomized, placebo-controlled trial. *J Allergy Clin Immunol.* 2007 May;119(5):1174–80.
2. Abrahamsson TR, Jakobsson T, Björkstén B, Oldaeus G, Jenmalm MC. No effect of probiotics on respiratory allergies: a seven-year follow-up of a randomized controlled trial in infancy. *Pediatr Allergy Immunol.* 2013 Sep;24(6):556–61.
3. Böttcher MF, Abrahamsson TR, Fredriksson M, Jakobsson T, Björkstén B. Low breast milk TGF- β 2 is induced by *Lactobacillus reuteri* supplementation and associates with reduced risk of sensitization during infancy. *Pediatr Allergy Immunol.* 2008 Aug;19(6):497–504.
4. Simón-Soro Á, D'Auria G, Collado MC, Džunková M, Culshaw S, Mira A. Revealing microbial recognition by specific antibodies. *BMC Microbiol.* 2015 Jul;15(1):132.
5. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* 2013 Jan;41(1):e1.
6. Dzidic M, Collado MC, Abrahamsson T, Artacho A, Stensson M, Jenmalm MC, et al. Oral microbiome development during childhood: an ecological succession influenced by postnatal factors and associated with tooth decay. *ISME J.* 2018;12(9):2292–306.
7. Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets. *Bioinformatics.* 2011 Mar;27(6):863–4.
8. Edgar RC. UNCROSS: Filtering of high-frequency cross-talk in 16S amplicon reads. *bioRxiv [Internet].* 2016;
9. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods.* 2010 Apr;7:335.
10. Oksanen J. *Vegan: community ecology.* R package version 2.4-6. 2018. p. <<http://cran.r-project.org/web/packages/vegan/>>.
11. Dzidic M, Abrahamsson TRTR, Artacho A, Björkstén B, Collado MCMC, Mira A, et al. Aberrant IgA responses to the gut microbiota during infancy precede asthma and allergy development. *J Allergy Clin Immunol.* 2017 Mar;139(3):1017–1025.e14.
12. Bunker JJ, Flynn TM, Koval JC, Shaw DG, Meisel M, McDonald BD, et al. Innate and Adaptive

- 197 Humoral Responses Coat Distinct Commensal Bacteria with Immunoglobulin A. *Immunity*.
198 2015;43(3):541–53.
199 13. Morgan X, Tickle T, Sokol H, Gevers D, Devaney K, Ward D. Dysfunction of the intestinal microbiome
200 in inflammatory bowel disease and treatment. *Genome Biol* [Internet]. 2012;13(R79).
201