Author Query Form

Journal: JEN

Article: 12751

Dear Author,

During the copyediting of your manuscript, the following queries arose.

Please refer to the query reference callout numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools.

Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

AUTHOR: Please note that missing content in references have been updated where we have been able to match the missing elements without ambiguity against a standard citation database, to meet the reference style requirements of the journal. It is your responsibility to check and ensure that all listed references are complete and accurate.

Query reference	Query	Remarks
1	AUTHOR: Please confirm that given names (blue) and surnames/family names (vermillion) have been identified correctly.	
2	AUTHOR: Please verify that the linked ORCID identifiers are correct for each author. The ORCID ID for Suárez seems to be invalid. Please check and supply the correct ORCID ID for this author.	
3	AUTHOR: Please check that authors and their affiliations are correct.	
4	AUTHOR: Please check corresponding author details are correct.	
5	AUTHOR: Kitahiro, 2014 has been changed to Kitahiro et al., 2014 so that this citation matches the Reference List. Please confirm that this is correct.	
6	AUTHOR: Reference González-Martín & Espadaler (2011) has not been included in the Reference List, please supply full publication details.	
7	AUTHOR: The funder details, 'Consejería de Política Territorial, Sostenibilidad y Seguridad. Gobierno de Canarias' and the grant number '151/2016 and 4/2017' supplied during the manuscript submission stage do not match with the funder information found in the Acknowledgements/Funding information section. Please confirm that all funding information is correct as typeset. If you have not provided the required funding information and wish to do so, please check and amend the 'Acknowledgements' section as appropriate.'	

Funding Info Query Form

Please confirm that the funding sponsor list below was correctly extracted from your article: that it includes all funders and that the text has been matched to the correct FundRef Registry organization names. If a name was not found in the FundRef registry, it may not be the canonical name form, it may be a program name rather than an organization name, or it may be an organization not yet included in FundRef Registry. If you know of another name form or a parent organization name for a “not found” item on this list below, please share that information.

FundRef name	FundRef Organization Name
Consejería de Política Territorial, Sostenibilidad y Seguridad. Gobierno de Canarias	

Six new non-native ants (Formicidae) in the Canary Islands and their possible impacts

David Hernández-Teixidor^{1,2} | Antonio José Pérez-Delgado^{1,2} | Daniel Suárez^{1,2} |
1 2 Joaquin Reyes-López³

¹Departamento de Biología Animal, Edafología y Geología, Sección de Biología, Facultad de Ciencias, Universidad de La Laguna, La Laguna, Spain

²Grupo de Investigaciones Entomológicas de Tenerife (GIET), La Laguna, Spain

³Área de Ecología, Universidad de Córdoba, Córdoba, Spain

Correspondence

David Hernández-Teixidor, Departamento de Biología Animal, Edafología y Geología, Sección de Biología, Facultad de Ciencias, Universidad de La Laguna, 38200 La Laguna Tenerife, Spain.
Email: davihdez@ull.es

Present address

David Hernández-Teixidor, Antonio José Pérez-Delgado and Daniel Suárez, Island Ecology and Evolution Research Group (IPNA-CSIC), La Laguna, Spain

Funding information

Consejería de Política Territorial, Sostenibilidad y Seguridad. Gobierno de Canarias, Grant/Award Number: 151/2016 and 4/2017

Abstract

Biological invasions are one of the main causes of biodiversity loss, especially on oceanic islands. Ants are among the most damaging pests in the world. After systematic sampling of more than 1,000 localities in the Canary Islands, six new exotic ant species are reported for the first time: *Pheidole bilimeki* (Myrmicinae), *Pheidole navigans* (Myrmicinae), *Strumigenys membranifera* (Myrmicinae), *Brachymyrmex cordemoyi* (Formicinae), *Tapinoma darioi* (Dolichoderinae) and *Technomyrmex pallipes* (Dolichoderinae). Moreover, another two recently reported species have been genetically confirmed. Morphological and genetic data were analysed to confirm the identity of the new records. For each species, information regarding identification, distribution, global invasive records and possible impacts is given. The arrival of these species may endanger local biodiversity.

KEYWORDS

barcoding, exotic ants, introduced species, Macaronesia, New record, oceanic islands

1 | INTRODUCTION

Exotic species, by definition, are those that are outside their natural distribution, introduced intentionally or unintentionally by human action (IUCN, 2000). The expansion of these species around the world is directly related to transport by humans (Hulme, Pyšek, Nentwig, & Vilà, 2009; Kairo, Cock, & Quinlan, 2003). The greatest number of introductions has occurred in the last few decades, mainly due to the growth of more frequent means of transportation and altered ecological niches (Holway, Lach, Suarez, Ysutsui, & Case, 2002). In fact, this is currently one of the main issues concerning biodiversity at the international level (Madjidian, Morales, & Smith, 2008).

Biological invasions are among the most important causes of the current biodiversity crisis, being involved in over half of the

contemporary extinctions (Bertelsmeier, Luque, Hoffmann, & Courchamp, 2015). They constitute a risk to endemic fauna and flora, since they can displace native species by competition for resources, predation, hybridization and pathogen transmission (Goulson, 2003).

Although all regions of the planet are suffering impacts derived from biological invasions, this problem is especially serious on oceanic islands (IUCN, 2000). Insular ecosystems host unique flora and fauna, which have generally less adaptable species, fewer populations and/or smaller population size than continental species. This has led to a particular vulnerability of insular species to the arrival of exotic species (Reaser et al., 2007).

Invasive ants are among the most damaging pests in the world. In fact, 19 have been listed as highly problematic by the IUCN SSC Invasive Species Specialist Group (GISD, 2019). Ants are easily

JEN	12751	WILEY	Dispatch: 16-3-2020	CE: Jenifar Seles S
	Journal Name	Manuscript No.		No. of pages: 8	PE: Mohanapriya L.

transported by humans because of their small size and the ability to nest in superficial or ephemeral sites such as root masses, leaf litter, logs and plant debris. In fact, exotic ants intercepted at the ports of entry are frequently detected in plant material (Bertelsmeier, Blight, & Courchamp, 2016).

In the Canary Islands, fifty-four ant species have been reported, of which 21 are considered introduced (Gobierno de Canarias, 2018; Schifani, Gentile, Scupola, & Espadaler, 2018; Staab, 2019). However, this number could increase in the coming years due to the continued commerce in agricultural and other carrier materials (including importation of foreign plant species). There is a lack of control measures for entry of exotic species, combined with an appropriate subtropical climate that favours species establishment. In this article, we provide information regarding morphological and genetic identification, distribution, global invasive records and possible impacts of several species that are reported for the first time for the archipelago.

2 | MATERIALS AND METHODS

2.1 | Sampling area

The Canary Islands are a volcanic archipelago comprising seven major islands off the south-west Atlantic coast of Morocco (Figure 1). In order to know the biodiversity of exotic ants in this archipelago, systematic sampling was conducted throughout all islands (including islets) during spring and summer of the years 2016–2018, periods in which ant species are more active at these latitudes (Angulo, Boulay, Ruano, Tinaut, & Cerdá, 2016). Since exotic ant species tend to colonize altered or anthropogenic environments first, especially those with high humidity (Mack et al., 2000), these kinds of habitats were the most sampled. Areas of possible arrival and dispersion such as ports, airports and plant nurseries were also surveyed. Lastly, natural areas were sampled to confirm whether they had been colonized by exotic species.

2.2 | Sampling methods

Two main sampling methods were carried out: pitfall traps and active searching. Pitfall traps are known to be an effective way of obtaining a representative sample of ant communities (quantitative and qualitative data) and are recommended as part of the standard protocol for measuring biodiversity (Alonso & Agosti, 2000; Clarke, Fisher, & LeBuhn, 2008). Traps were sunk into the ground with the top flush with the soil surface. Each trap consisted of a 150-ml plastic container, filled with 25 ml of water and a few drops of detergent to decrease water surface tension. In order to maximize the diversity of ants attracted by the traps, two types of bait were used: (a) tinned tuna, for species that are attracted mainly by protein foods; and (b) sugar water, for the species that prefer sugary foods. In each locality, 10 pitfall traps were placed in line at a distance of 5 m from

each other, aiming at the greatest diversity of possible microhabitats (under trees/shrubs, under stones, near water source, etc.). Traps 1, 5 and 9 contained tuna baits and the others sugar water. Once placed, they stayed functional in the field for 48 hr. These traps are not always useful to detect all species (Alonso & Agosti, 2000), and therefore, active searching has to be carried out as well (Majer, 1997). This consists of a thorough 20-min search of each locality at any ant-susceptible site. In addition, occasional sampling was conducted in buildings or areas where it was not possible to carry out a complete survey.

2.3 | Morphological and genetic identification

Specimens were identified morphologically and by DNA barcoding, as the morphological taxonomy of some genera is unclear or complex, such as *Brachymyrmex* (Ortiz-Sepulveda, Bocxlaer, Meneses, & Fernández, 2019), *Pheidole* (Sarnat, Fischer, Guénard, & Economo, 2015), *Tapinoma* (Seifert, D'Eustacchio, Kaufmann, Centorame, & Modica, 2017) and *Technomyrmex* (Bolton, 2007). Firstly, external morphological characters were reviewed following some identification keys, and then, a genetic identification was carried out to confirm the previous one. DNA was extracted using QIAGEN DNeasy Blood and Tissue Kit, and COI was amplified using LCO1490/HCO2980 primers (Folmer, Black, Hoch, Lutz, & Vrijenhoek, 1994). The reaction product was Sanger sequenced, and the obtained sequence was blasted against the GenBank database for a proper taxonomic identification. Only, unambiguously identified sequences were considered, especially in complex genera.

3 | RESULTS

A total of 1,016 localities were sampled, 103 of them by pitfall traps. This survey revealed six new exotic species for the first time in the Canary Islands, more particularly in urban areas on the islands of La Gomera, Tenerife and Gran Canaria (Table 1, Figure 1 and Table S1). This is a 22% increase in the number of introduced ants present in the Canaries. Moreover, three of the six species belong to genera not previously detected in the archipelago (*Strumigenys*, *Technomyrmex* and *Brachymyrmex*). Except for *Technomyrmex pallipes* and *Strumigenys membranifera*, the rest of the species here reported have never been discovered before in the Macaronesian archipelagos. In addition, two other new exotic ants detected in our survey and recently reported by other authors, namely *Plagiolepis alluaudi* (Staab, 2019) and *Lepisiota frauenfeldi* (Schifani et al., 2018), were genetically confirmed, and their distribution is broadened to other islands.

All identifications were confirmed by morphological and genetic data. Sequences were compared with verified GenBank entries (Table S2). Both approaches gave consistent results. Biological and ecological information is presented hereafter.

FIGURE 1 Ant sampling sites in the Canary Islands where new exotic ant species were detected. *Brachymyrmex cordemoyi* populations indicated with brown squares, *Pheidole bilimeki* green circles, *Pheidole navigans* blue diamond, *Strumigenys membranifera* yellow circles, *Tapinoma darioi* black crosses, and *Technomyrmex pallipes* pink pentagons. Localities with more than one new exotic ant species detected are marked with red triangles. For details about localities, see Tables S1 and S2

TABLE 1 New records of exotic ants in the Canary Islands, with information about the distribution, and negative effects previously recorded in other countries

Species	Subfamily	Native from	Negative effects	Distribution
<i>Pheidole bilimeki</i>	Myrmicinae	Neotropical region	Yes	TF, LG
<i>Pheidole navigans</i>	Myrmicinae	Neotropical region	Yes	TF
<i>Strumigenys membranifera</i>	Myrmicinae	Afrotropical region	No	TF, GC, LG
<i>Brachymyrmex cordemoyi</i>	Formicinae	Neotropical region	Yes	TF, GC
<i>Tapinoma darioi</i>	Dolichonerinae	Mediterranean region	No	TF
<i>Technomyrmex pallipes</i>	Dolichonerinae	Afrotropical region	Yes	GC

Note: Native distribution data taken from Guénard et al. (2017).
Abbreviations: GC, Gran Canaria; LG, La Gomera; TF, Tenerife.

3.1 | New records

3.1.1 | *Pheidole bilimeki* (Myrmicinae)

The main feature differentiating *P. bilimeki* from other *Pheidole* species is the shorter scapes (scape length 0.40–0.60 mm; scape index 95–125) (Longino & Cox, 2009). *Pheidole bilimeki* is usually reddish

brown, rarely yellowish brown, and with the face and mesosoma uniformly foveolate.

This species is native to the Neotropical region, from the north of South America to the south of North America, and is present throughout the Caribbean (Guénard, Weiser, Gomez, Narula, & Economo, 2017). It has been introduced into the Nearctic and Palaearctic regions (Antweb, 2019; Guénard et al., 2017; Sarnat

et al., 2015). In the Canaries, it has only appeared in three localities on Tenerife and ten on La Gomera (Figure 1 and Table S1).

In the regions where it has been introduced, it is occasionally found inside houses, but more frequently in greenhouses (Sarnat et al., 2015). There is little indication that this species causes a significant impact on agricultural systems or native ecosystems. As it has low tolerance for cold climates, it is not considered a highly invasive species (Sarnat et al., 2015).

3.1.2 | *Pheidole navigans* (Myrmicinae)

Pheidole navigans is a small, short-limbed, reddish-brown species belonging to the *P. flavens* complex. Minor workers are impossible to distinguish on the basis of morphological characters, but major can be separated from other species of the *P. flavens* complex. The distinctive characters are the combination of predominantly longitudinal rugae on their posterolateral lobes, the more distinct and narrower antennal scrobe bordered mesially by a strong, unbroken frontal carina and the more continuously glossy scrobe depression (Sarnat et al., 2015).

Pheidole navigans is thought to be native to the Neotropical region (centre and north of South America) and has been introduced into the Palaearctic and Nearctic regions (Sarnat et al., 2015; Wetterer, 2017). In the Canary Islands, it was only collected in a farming area in the north of Tenerife (Figure 1 and Table S1).

This species occurs in both disturbed areas and moist woodlands but rarely enters houses (Deyrup, Davis, & Cover, 2000). It is not considered a major pest; however, in the south-east of the United States, it has been expanding its distribution area since its first record (Sarnat et al., 2015).

3.1.3 | *Strumigenys membranifera* (Myrmicinae)

Strumigenys differs from the rest of the ants present in the Canaries by its characteristic triangular head, six-segmented antennae with a two-segmented smaller antennal club and spongi-form appendages on the propodeum and waist. Within this genus, *S. membranifera* differs from other synanthropic species by having mandibles with 12 teeth and its cephalic dorsum pilosity restricted to a pair of hairs at the highest point of the vertex (Sharaf, Fisher, & Aldawood, 2014).

Strumigenys membranifera has achieved broad distributions in both the Old and New World, spread through human commerce (Wetterer, 2011). Although its geographical origin remains unclear (Wetterer, 2011), it seems to be native to the Afrotropical region (Guénard et al., 2017). In the Canary archipelago, it has been found on the islands of Gran Canaria (two localities), Tenerife (one locality) and La Gomera (one locality) (Figure 1 and Table S1).

This species has been reported in gardens, cultivated areas and forests (Sharaf et al., 2014), although open areas seem to be its preferred habitat worldwide (Kitahiro, Yamamoto, Touyama, & Ito,

2014). There is no information regarding impacts (Wetterer, 2011). Nevertheless, its hypogeal and leaf-litter lifestyle makes it a difficult species to detect and therefore to study its possible invasiveness.

3.1.4 | *Brachymyrmex cordemoyi* (Formicinae)

As with all *Brachymyrmex* species, the presence of only nine antennal segments with a lack of a differentiated antennal club distinguishes workers of *Brachymyrmex* from other Formicinae genera (Bolton, 1994) and from all other species in the Canary Islands. This species resembles *B. obscurior* and *B. patagonicus* but differs from the former by having a denser pubescence on the gaster and from the latter by its larger head, more ommatidia along the maximal diameter of the eye and lighter-coloured pubescence. The latter feature is denser on the dorsum of the entire body and appressed on the gaster instead of decumbent in *B. obscurior* (Ortiz-Sepulveda et al., 2019).

Brachymyrmex cordemoyi is native to the Neotropical region and has been introduced in parts of the Afrotropical, Australasian and Palaearctic regions (Guénard et al., 2017; Husemann & Ortiz-Sepulveda, 2019; Ortiz-Sepulveda et al., 2019). Only two populations have been detected on Tenerife and one on Gran Canaria (Figure 1 and Table S1).

In its introduced range, it has been collected both indoors (Husemann & Ortiz-Sepulveda, 2019) and outdoors, nesting in soil (Sharaf, Salman, Aldhafer, Yousef, & Aldawood, 2016). There is not much information about its effects as an invasive species, but it seems they are mild (Jacquot et al., 2017). However, other species of this genus can act as urban pests, such as *B. obscurior* or *B. patagonicus* (Klotz, Mangold, Vail, Davis, & Patterson, 1995; MacGown, Hill, & Deyrup, 2007).

3.1.5 | *Tapinoma darioi* (Dolichoderinae)

The genus *Tapinoma* is easily identified by its flattened petiole and without a conspicuous, dorsally protruding scale. The dorsal face of propodeum is much shorter than the declivitous (posterior) face. The shape of the clypeal excision is one of the characteristics that differentiate species from this genus (Gómez & Espadaler, 2018). *Tapinoma darioi*, which has a completely dark body, belongs to the *T. nigerrimum* complex that includes four species that have been recently described (Seifert et al., 2017). Furthermore, the depth-to-width ratio of clypeal excision and length-to-width ratio of second funiculus segment are on average larger than in the other species (Seifert et al., 2017).

This ant is originally from the Mediterranean basin, showing preference for coastal areas (Seifert et al., 2017). Outside its native area, it had been introduced only in the Netherlands (Guénard et al., 2017). On Tenerife, it has only appeared in one locality in a greenhouse in Valle de La Orotava (Figure 1 and Table S1). We found individuals of this species together with others of *Pheidole pallidula* in a group of olive trees introduced from the Iberian Peninsula, where

both species are very common in olive groves (Moreno-García, Reyes-López, Repullo-Ruibérriz de Torres, & Carbonell-Bojollo, 2017).

Tapinoma darioi has never been recorded as invasive, except a possible case in the Netherlands (Seifert et al., 2017). Other species within the complex, such as *T. magnum*, have a considerable invasive potential (Seifert et al., 2017).

3.1.6 | *Technomyrmex pallipes* (Dolichoderinae)

The genus *Technomyrmex* belongs to the subfamily Dolichoderinae and is characterized by its 12-segmented antennae without a terminal club, reduced petiole and gaster with five tergites visible in dorsal view (Bolton, 2007). Within this genus, *T. pallipes* can be distinguished from another widespread introduced species by having the dorsum behind the level of the posterior margin of the eye with two pairs of very short, stubbly setae (Fernández & Guerrero, 2008).

This species is of Afrotropical origin and has been introduced to the Palaearctic, Nearctic, Indo-Malayan and Australian regions (Bolton, 2007; Guénard et al., 2017). In the Canary Islands, it has reported for just one single locality of Gran Canaria (Figure 1 and Table S1).

In its introduced range, this species is found in greenhouses and houses (Wetterer, Espadaler, Wetterer, Aguin-Pombo, & Franquinho-Aguiar, 2007), but it is also common in outdoor areas such as parks and gardens in warmer countries (Bolton, 2007). Moreover, it has been reported as an occasional locally serious pest in citrus orchards (Bolton, 2007; Samways, Nel, & Prins, 1982). Although no major negative impacts have been reported to date, it could act as invasive such as other species of this genus like *T. albipes*.

3.2 | Species confirmation

Although Schifani et al. (2018) and Staab (2019) have already reported *L. frauenfeldi* and *P. alluaudi*, respectively, our data allow us to genetically confirm the identity of the species. They show their establishment and spread in the archipelago by detecting their presence in more locations and islands.

3.2.1 | *Plagiolepis alluaudi* (Formicinae)

Plagiolepis alluaudi is a tiny yellow ant (total length 1.5 mm approximately), whose workers have eleven antennal segments. In addition, this ant has a round head with strongly convex posterior margins unlike *P. exigua* (also yellowish), which has an oblong head (Wetterer, 2014).

This ant is primarily a tropical species, native to the Afrotropical region (Guénard et al., 2017). It has been introduced into the Australasian, Indo-Malayan, Neotropical, Oriental and Palaearctic regions (Wetterer, 2014), and has been spread worldwide through

human commerce (Wetterer, 2014). In the Canary Islands, Staab (2019) recently reported it for Tenerife, but in our sampling, it was detected in three localities on Gran Canaria, one on La Palma and one on Tenerife (Figure 1 and Table S1).

Outside its native area, it has been detected in natural areas such as forests, primarily in the Afrotropical region (Wetterer, 2014), as well as indoors in the Palaearctic region (Guénard et al., 2017). Apart from having a short life cycle and a wide tolerance to temperature and humidity, its success in new localities may be due to its ability to co-exist with dominant invasive ants like *Pheidole megacephala* (Smith, 1957; Wetterer, 2014). It has become a minor household and agricultural pest in some areas, especially in the Pacific islands and in European greenhouses (Wetterer, 2014).

3.2.2 | *Lepisiota frauenfeldi* (Formicinae)

Within the Formicinae subfamily, the genus *Lepisiota* is the only one that has its propodeum armed with spines or teeth (Fisher & Bolton, 2016). It can be distinguished from *L. capensis*, which is reported for the Canary Islands, by having a petiole with asymmetrical spine-shaped dorsolateral apices, while *L. capensis* has a sharply bidentate petiole (Espadaler & Fernández, 2014).

Although its origin seems Palaearctic, this species is more densely distributed in the Mediterranean area and the Middle East and has been introduced on some islands of the Afrotropical, Indo-Malayan and Nearctic region as well as in the Australasia region (Guénard et al., 2017). Schifani et al. (2018) recently discovered it on the islands of Tenerife and Fuerteventura. In our sampling, it was also detected on Lanzarote and La Gomera as well as the islands where it was previously reported.

Lepisiota frauenfeldi is considered an invasive species (FAO, 2018), which eats and displaces native ant species, as well as other insects in the infested area (Commonwealth of Australia, 2019). In Australia, it is a serious environmental pest (Commonwealth of Australia, 2019) and currently is under national eradication there (Commonwealth of Australia, 2018).

4 | DISCUSSION

The geographical situation of the Canary Islands and its links with America, Africa and Europe facilitates the arrival of numerous exotic species on these islands due to the increase in sea and air traffic. Human trading activity has been breaking down biogeographical barriers and spreading species around the world (Bertelsmeier, Ollier, Liebhold, & Keller, 2017). In fact, the new exotic ant species here reported originate from different regions (Table 1): three of them have a Neotropical origin (*B. cordemoyi*, *P. bilimeki* and *P. navigans*), two Afrotropical (*S. membranifera* and *T. pallipes*), and one is from the Mediterranean basin (*T. darioi*). Moreover, the colonization of the Canary Islands by these exotic ants may serve as a focus of dispersion to nearby areas such as

Africa and Europe. For instance, *B. cordemoyi* and both *Pheidole* species, which are barely present in the Mediterranean basin, could become a threat to this region after their establishment as an intermediate step in the Canary Islands.

Together with the current disturbed state of the habitats, the prevailing subtropical climate consisting of relatively constant mild temperatures year-round facilitates the establishment of many of these introduced species and their expansion throughout the archipelago, especially those from warm climates. Moreover, species from places with cold winters can be active the whole year in these islands. Conversely, tropical and subtropical ants cannot survive in geographical areas with cold winter conditions and are restricted to indoor locations (Ivanov, 2016). For example, *P. bilimeki* in central Europe is restricted to houses and greenhouses (AntWeb, 2019; Boer & Vierbergen, 2008). However, the climatic conditions in the Canaries allow their populations to become established in outdoor areas in addition to inside buildings. This facilitates their colonization of different habitats and expansion throughout the archipelago. Currently, the distribution of these six new exotic ants recorded for the Canaries is restricted to a few locations, all of them in human-disturbed areas. This finding reinforces the hypothesis that the arrivals of these species in these islands were relatively recent.

The arrival of alien species on oceanic islands threatens their biodiversity and causes socio-economic impacts. Intensive and long-term surveys allow the study of changes in ant fauna on islands and the early detection of exotic species (Hoffmann, Graham, & Smith, 2017; Moreau, Deyrup, & Davis, 2014). Correct identification is essential to assess the severity of these new introductions. As suggested by our data, the combined use of barcoding identification and recent taxonomic revisions is fundamental in order to properly identify Formicidae specimens, especially in complex genera. Subsequently, risk assessment for these new exotic ants is necessary to minimize potential damage. *Brachymyrmex cordemoyi* is known to cause damage to native fauna in other regions (Jacquot et al., 2017). Moreover, species belonging to genera that include highly invasive species such as *Pheidole* should be considered as potential invasive species. In fact, *P. bilimeki* and *P. navigans* have caused some negative impacts in other warm regions (Sarnat et al., 2015). Species associated with honeydew-producing Hemiptera, such as *T. darioi* and *T. pallipes* (Bolton, 2007; Seifert et al., 2017), could be potential crop pests. Additionally, some of these ants stand out for their predator potential, such as *S. membranifera* in the soil, whose impact on other tiny soil arthropods could be substantial (Wetterer, 2011).

In this study, we provide six new records of ants recently found in the Canaries for the first time, which notably increase the number of exotic ants. Given these new records, exotic ants constitute 45% of all ant species present in the archipelago. The presence of a higher proportion of introduced ants compared with native ants seems to be common in oceanic archipelagos worldwide, such as the other Macaronesian archipelagos, Hawaii or Galapagos (AntWeb, 2019; Borges et al., 2008, 2010). In contrast, the percentage of

native species is much higher in mainland areas such as the Iberian Peninsula and Morocco (González-Martín & Espadaler, 2011; Taheri & Reyes-López, 2018).

To effectively manage exotic ants, a comprehensive plan should be developed with actions at all stages of the invasion. It should at least include prevention, detection, response, containment and management actions (Commonwealth of Australia, 2018). Border controls are the principal defence lines to prevent the arrival of exotic species (Hayden & White, 2003); however, for those already established a rapid response is needed to prevent their expansion. Eradication, the main objective, is only possible when ant populations are established in a small area. However, if the populations have spread, the use of control techniques to mitigate their impact is recommended (Commonwealth of Australia, 2018; Krushelnicky, Loope, & Reimer, 2005). The six species here reported are plausible candidates for wider expansion in the future, so regular monitoring should be maintained, as well as implementing eradication or control efforts when required.

ACKNOWLEDGEMENTS

This study was supported by the Canary Government and cofinanced by the European Regional Development Fund. David Hernández-Teixidor is currently funded by the Cabildo de Tenerife, under the TFinnova Programme supported by MEDI and FDCAN funds. We thank Gloria Ortega (Museum of Nature and Archaeology, Santa Cruz, Tenerife) for providing access to the museum collection, and Pedro Oromí (La Laguna University) for making available his laboratory and collections. Brent Emerson (IPNA-CSIC) allowed us to perform genetic identification in his laboratory. Alfredo Reyes (Orotava Acclimatisation Garden) and María del Rosario Fresno (Canary Islands Agricultural Research Institute, ICIA) facilitated our research at their facilities. Javier García, Irene Santos, David Lugo, Pedro Oromí, Heriberto López, Manuel Naranjo, Alejandro Escáñez, Kyrie Hampton and Vanessa Pérez collaborated in fieldwork. Santa Cruz city council and the various Island Councils granted collection permits and support. El Palmetum, La Marquesa, Viera y Clavijo and Agaete Botanical Gardens, and Maspalomas, Costa Tegui, and Lanzarote golf courses, Hesperia Playa Dorada hotel and other private property owners all kindly permitted access for sampling. Guido Jones (IPNA-CSIC) revised the English text of the manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

AUTHOR CONTRIBUTIONS

DH-T, AJP-D and DS surveyed the sampling areas and collected data. JR-L led the morphological identifications. DS and DH-T did the molecular analysis. All authors contributed to the writing. All authors read and approved the manuscript.

DATA AVAILABILITY STATEMENT

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

ORCID

David Hernández-Teixidor <https://orcid.org/0000-0003-3114-7892>

Antonio José Pérez-Delgado <https://orcid.org/0000-0002-3797-4058>

Daniel Suárez <https://orcid.org/0000-0003-3417-1169>

Joaquín Reyes-López <https://orcid.org/0000-0002-0702-4588>

REFERENCES

- Alonso, L. E., & Agosti, D. (2000). Biodiversity studies, monitoring and ants: An overview. In D. Agosti, J. D. Majer, L. E. Alonso, & R. Shultz (Eds.), *Ants. Standard methods for measuring and monitoring biodiversity* (pp. 1–8). Washington, DC: Smithsonian Institution Press.
- Angulo, E., Boulay, R., Ruano, F., Tinaut, A., & Cerdá, X. (2016). Anthropogenic impacts in protected areas: Assessing the efficiency of conservation efforts using Mediterranean ant communities. *PeerJ*, 4, e2773. <https://doi.org/10.7717/peerj.2773>
- AntWeb (2019). Antweb v. 8.8.4. Retrieved from <https://www.antweb.org/>
- Bertelsmeier, C., Blight, O., & Courchamp, F. (2016). Invasions of ants (Hymenoptera: Formicidae) in light of global climate change. *Myrmecological News*, 22, 25–42.
- Bertelsmeier, C., Luque, G. M., Hoffmann, B. D., & Courchamp, F. (2015). Worldwide ant invasions under climate change. *Biodiversity and Conservation*, 24, 117–128. <https://doi.org/10.1007/s10531-014-0794-3>
- Bertelsmeier, C., Ollier, S., Liebhold, A., & Keller, L. (2017). Recent human history governs global ant invasion dynamics. *Nature Ecology & Evolution*, 1(7), 0184. <https://doi.org/10.1038/s41559-017-0184>
- Boer, P., & Vierbergen, B. (2008). Exotic ants in the Netherlands (Hymenoptera: Formicidae). *Entomologische Berichten*, 68(2), 121–129.
- Bolton, B. (1994). *Identification guide to the ant genera of the world*. Cambridge, MA: Harvard University Press.
- Bolton, B. (2007). Taxonomy of the Dolichoderine ant genus *Technomyrmex* Mayr (Hymenoptera: Formicidae) based on the worker caste. *Contributions of the American Entomological Institute*, 35(1), 1–149.
- Borges, P. A. V., Abreu, C., Aguiar, A. M. F., Carvalho, P., Jardim, R., Melo, I., ... Vieira, P. (2008). *A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos*. Funchal, Portugal: Direcção Regional do Ambiente da Madeira and Universidade dos Açores, Funchal and Angra do Heroísmo.
- Borges, P. A. V., Costa, A., Cunha, R., Gabriel, R., Gonçalves, V., Martins, A. F., ... Vieira, V. (2010). *A list of the terrestrial and marine biota from the Azores*. Cascais, Portugal: Principia.
- Clarke, K. M., Fisher, B. L., & LeBuhn, G. (2008). The influence of urban park characteristics on ant (Hymenoptera, Formicidae) communities. *Urban Ecosystems*, 11(3), 317–334. <https://doi.org/10.1007/s11252-008-0065-8>
- Commonwealth of Australia (2018). *National Invasive Ant Biosecurity Plan 2018–2028*. Canberra, ACT: Department of the Environment and Energy, Australian Government.
- Commonwealth of Australia (2019). *National pest & disease outbreaks. Browsing ant*. Retrieved from <https://www.outbreak.gov.au/current-responses-to-outbreaks/browsing-ant>
- Deyrup, M., Davis, L., & Cover, S. (2000). Exotic ants in Florida. *Transactions of the American Entomological Society*, 126, 293–326.
- Espadaler, X., & Fernández, G. (2014). *Lepisiota capensis* (Mayr, 1862), a new exotic ant (Hymenoptera, Formicidae) in La Gomera (Canary Islands). *Iberomyrmex*, 6, 5–8.
- FAO (2018). *Invasive ants as a biosecurity threat*. Retrieved from <http://www.fao.org/publications/card/es/c/18939EN/>
- Fernández, F., & Guerrero, R. (2008). *Technomyrmex* (Formicidae: Dolichoderinae) in the New World: Synopsis and description of a new species. *Revista Colombiana De Entomología*, 34(1), 110–115.
- Fisher, B. L., & Bolton, B. (2016). *Ants of Africa and Madagascar. A guide to the genera*. Berkeley, CA: University of California Press.
- Folmer, O., Black, M., Hoch, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294–299.
- GISD (2019). *Global invasive species database*. Retrieved from <http://www.issg.org/database/welcome>
- Gobierno de Canarias (2018). *Banco de Datos de Biodiversidad de Canarias*. Retrieved from www.biodiversidadcanarias.es
- Gómez, K., & Espadaler, X. (2018). *Hormigas ibéricas*. Retrieved from <http://www.hormigas.org>
- Goulson, D. (2003). Effects of introduced bees on native ecosystems. *Annual Review of Ecology, Evolution and Systematics*, 34, 1–26. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132355>
- Guénard, B., Weiser, M., Gómez, K., Narula, N., & Economo, E. P. (2017). The Global Ant Biodiversity Informatics (GABI) database: A synthesis of ant species geographic distributions. *Myrmecological News*, 24, 83–89.
- Hayden, B. J., & White, C. (2003). Invasive species management in New Zealand. In G. M. Ruiz, & J. T. Carlton (Eds.), *Invasive species: Vectors and management strategies* (pp. 270–291). Washington, DC: Island Press.
- Hoffmann, B. D., Graham, R., & Smith, D. (2017). Ant species accumulation on Lord Howe Island highlights the increasing need for effective biosecurity on islands. *NeoBiota*, 34, 41–51. <https://doi.org/10.3897/neobiota.34.10291>
- Holway, D. A., Lach, L., Suarez, A. V., Ysutsui, N. D., & Case, T. J. (2002). The causes and consequences of ant invasions. *Annual Review of Ecology and Systematics*, 33, 181–233. <https://doi.org/10.1146/annurev.ecolsys.33.010802.150444>
- Hulme, P. E., Pyšek, P., Nentwig, W., & Vilà, M. (2009). Will threat of biological invasions unite the European Union? *Science*, 324(5923), 40–41. <https://doi.org/10.1126/science.1171111>
- Husemann, M., & Ortiz-Sepulveda, C. M. (2019). First documented record of the neotropical ant *Brachymyrmex cordemoyi* Forel, 1895 (Formicidae: Formicinae) in Germany. *BiolInvasions Records*, 8(4), 764–773. <https://doi.org/10.3391/bir.2019.8.4.04>
- IUCN (2000). *Guidelines for the prevention of biodiversity loss caused by Alien Invasive Species*. Gland, Switzerland: International Union for the Conservation of Nature.
- Ivanov, K. (2016). Exotic ants (Hymenoptera, Formicidae) of Ohio. *Journal of Hymenoptera Research*, 51, 203–226. <https://doi.org/10.3897/jhr.51.9135>
- Jacquot, M., Tixier, P., Flores, O., Muru, D., Massol, F., Derepas, B., ... Deguine, J. P. (2017). Contrasting predation services of predator and omnivore diversity mediated by invasive ants in a tropical agroecosystem. *Basic and Applied Ecology*, 18, 31–39. <https://doi.org/10.1016/j.baae.2016.09.005>
- Kairo, M. T. K., Cock, M. J. W., & Quinlan, M. M. (2003). An assessment of the use of the Code of Conduct for the Import and Release of Exotic Biological Control Agents (ISPM No. 3) since its endorsement as an international standard. *Biocontrol News and Information*, 24(1), 15–27.
- Kitahiro, S., Yamamoto, K., Touyama, Y., & Ito, F. (2014). Habitat preferences of *Strumigenys* ants in western Japan (Hymenoptera: Formicidae). *Asian Myrmecology*, 6, 91–94. <https://doi.org/10.20362/am.006007>
- Klotz, J. H., Mangold, J. R., Vail, K. M., Davis, L. R. Jr, & Patterson, R. S. (1995). A survey of the urban pest ants (Hymenoptera: Formicidae) of peninsular Florida. *Florida Entomologist*, 78, 109–118. <https://doi.org/10.2307/3495674>

- Krushelnicky, P. D., Loope, L. L., & Reimer, N. J. (2005). The ecology, policy, and management of ants in Hawaii. *Proceedings of the Hawaiian Entomological Society*, 37, 1–25.
- Longino, J. T., & Cox, D. J. (2009). *Pheidole bilimeki* reconsidered (Hymenoptera: Formicidae). *Zootaxa*, 1985, 34–42. <https://doi.org/10.11646/zootaxa.1985.1.3>
- MacGown, J. A., Hill, J. G., & Deyrup, M. A. (2007). *Brachymyrmex patagonicus* (Hymenoptera: Formicidae), an emerging pest species in the southeastern United States. *Florida Entomologist*, 90(3), 457–464. [https://doi.org/10.1653/0015-4040\(2007\)90\[457:BPHFAE\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[457:BPHFAE]2.0.CO;2)
- Mack, R. N., Simberloff, D., Lonsdale, W. M., Evans, H., Clout, M., & Bazzaz, F. A. (2000). Biotic invasions: Causes, epidemiology, global consequences, and control. *Ecological Applications*, 10(3), 689–710. [https://doi.org/10.1890/1051-0761\(2000\)010\[0689:BICEG C\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0689:BICEG C]2.0.CO;2)
- Madjidian, J. A., Morales, C. L., & Smith, H. G. (2008). Displacement of a native by an alien bumblebee: Lower pollinator efficiency overcome by overwhelmingly higher visitation frequency. *Oecologia*, 156(4), 835–845. <https://doi.org/10.1007/s00442-008-1039-5>
- Majer, J. D. (1997). The use of pitfall traps for sampling ants – A critique. *Memoirs of Museum Victoria*, 56, 323–329. <https://doi.org/10.24199/j.mmv.1997.56.20>
- Moreau, C. S., Deyrup, M. A., & Davis, L. R. (2014). Ants of the Florida Keys: Species accounts, biogeography, and conservation (Hymenoptera: Formicidae). *Journal of Insect Science*, 14(295), 2014. <https://doi.org/10.1093/jisesa/ieu157>
- Moreno-García, M., Reyes-López, J., Repullo-Ruibérriz de Torres, M. A., & Carbonell-Bojollo, R. M. (2017). *Comunidades de hormigas en diferentes manejos sobre olivares sin laboreo*. Jaén, Spain: Comunicaciones Científicas Simposium Expoliva 2017.
- Ortiz-Sepulveda, C. M., Van Bocxlaer, B., Meneses, A. D., & Fernández, F. (2019). Molecular and morphological recognition of species boundaries in the neglected ant genus *Brachymyrmex* (Hymenoptera: Formicidae): Toward a taxonomic revision. *Organisms Diversity & Evolution*, 19(3), 447–542. <https://doi.org/10.1007/s13127-019-00406-2>
- Reaser, J. K., Meyerson, L. A., Cronk, Q., De Poorter, M., Eldrege, L. G., Green, E., ... Vaiutu, L. (2007). Ecological and socioeconomic impacts of invasive alien species in island ecosystems. *Environmental Conservation*, 34(02), 98–111. <https://doi.org/10.1017/S0376892907003815>
- Samways, M. J., Nel, M., & Prins, A. J. (1982). Ants (Hymenoptera: Formicidae) foraging in citrus trees and attending honeydew-producing Homoptera. *Phytophylactica*, 14(4), 155–157.
- Sarnat, E. M., Fischer, G., Guénard, B., & Economo, E. P. (2015). Introduced *Pheidole* of the world: Taxonomy, biology and distribution. *ZooKeys*, 543, 1–109. <https://doi.org/10.3897/zookeys.543.6050>
- Schifani, E., Gentile, V., Scupola, A., & Espadaler, X. (2018). Yet another alien: A second species of *Lepisiota* spreading across the Canary Islands, Spain (Hymenoptera: Formicidae). *Fragmenta Entomologica*, 50, 61–64. <https://doi.org/10.4081/fe.2018.287>
- Seifert, B., D'Eustachio, D., Kaufmann, B., Centorame, M., & Modica, M. (2017). Four species within the supercolonial ants of the *Tapinoma nigerrimum* complex revealed by integrative taxonomy (Hymenoptera: Formicidae). *Myrmecological News*, 24, 123–144.
- Sharaf, M. R., Fisher, B. L., & Aldawood, A. S. (2014). Notes on Ants of the genus *Strumigenys* F. Smith, 1860 (Hymenoptera: Formicidae) in the Arabian Peninsula, with a key to species. *Sociobiology*, 61(3), 293–299. <https://doi.org/10.13102/sociobiology.v61i3.293-299>
- Sharaf, M. R., Salman, S., Aldhafer, H. M., Yousef, A. F. A., & Aldawood, A. S. (2016). First occurrence of the ant genus *Brachymyrmex* Mayr, 1868 (Hymenoptera: Formicidae) from the Kingdom of Saudi Arabia. *Sociobiology*, 63(2), 800–803. <https://doi.org/10.13102/sociobiology.v63i2.981>
- Smith, M. R. (1957). A contribution to the taxonomy, distribution and biology of the vagrant ant, *Plagiolepis alluaudi* Emery (Hymenoptera, Formicidae). *Journal of the New York Entomological Society*, 65(3/4), 195–198.
- Staab, M. (2019). *Plagiolepis alluaudi* Emery, 1894, a globally spreading exotic ant (Hymenoptera, Formicidae) newly recorded from Tenerife (Canary Islands, Spain). *Journal of Hymenoptera Research*, 74, 83–91. <https://doi.org/10.3897/jhr.74.47315>
- Taheri, A., & Reyes-López, J. (2018). Exotic ants (Hymenoptera: Formicidae) in Morocco: Checklist, comments and new faunistic data. *Transactions of the American Entomological Society*, 144(1), 99–108. <https://doi.org/10.3157/061.144.0104>
- Wetterer, J. K. (2011). Worldwide spread of the membraniferous dactine ant, *Strumigenys membranifera* (Hymenoptera: Formicidae). *Myrmecological News*, 14, 129–135.
- Wetterer, J. K. (2014). Worldwide spread of Alluaud's little yellow ant, *Plagiolepis alluaudi* (Hymenoptera: Formicidae). *Myrmecological News*, 19, 53–59.
- Wetterer, J. K. (2017). Invasive ants of Bermuda revisited. *Journal of Hymenoptera Research*, 54, 33–41.
- Wetterer, J. K., Espadaler, X., Wetterer, A. L., Aguin-Pombo, D., & Franquinho-Aguiar, A. M. (2007). Ants (Hymenoptera: Formicidae) of the Madeiran Archipelago. *Sociobiology*, 49(2), 1–33.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Hernández-Teixidor D, Pérez-Delgado AJ, Suárez D, Reyes-López J. Six new non-native ants (Formicidae) in the Canary Islands and their possible impacts. *J Appl Entomol*. 2020;00:1–8. <https://doi.org/10.1111/jen.12751>