Article

Smaller distance between nest contents and cavity entrance increases risk of ectoparasitism in cavity-nesting birds

**Gustavo Tomás, Cristina Ruiz-Castellano, Magdalena Ruiz-Rodríguez
and Juan J. Soler**

*G. Tomás (<https://orcid.org/0000-0001-6701-2055>) ✉ (gtomas@eeza.csic.es),
C. Ruiz-Castellano, M. Ruiz-Rodríguez and J. J. Soler, Department of
Functional and Evolutionary Ecology, Estación Experimental de Zonas
Áridas (EEZA - CSIC), E-04120 Almería, Spain. Current address of MRR:
Sorbonne Universités. UPMC Univ. Paris 06, Observatoire Océanologique,
Avenue du Fontaulé, 66650, Banyuls-Sur-Mer, France.*

Acknowledgements – We appreciate the constructive criticisms provided by five anonymous reviewers on earlier versions of the manuscript.

Funding – Financial support was provided by grants from Spanish Ministerio de Ciencia e Innovación (MCI) and FEDER (CGL2010-19233-C03-01), from Ministerio de Economía, Industria y Competitividad (CGL2017-89063-P, CGL2017-83103-P), by a predoctoral fellowship from MCI to CRC, by Junta para Ampliación de Estudios programme to MRR, and by Juan de la Cierva and Ramón y Cajal programmes to GT.

Author contributions – GT, CRC and JJS designed the study. GT, CRC, MRR and JJS conducted field work. JJS and GT performed the statistical analyses and wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

Conflict of interest – The authors declare that they have no conflict of interest.

Permits – All applicable institutional and/or national guidelines for the care and use of animals were followed. Authorization to conduct the study was granted by the Dirección General de Gestión del Medio Natural, Consejería de Medio Ambiente, Junta de Andalucía.

Abstract

Determining costs associated with nest-building behaviour and nest characteristics is of prime importance to understand the evolution of such extended phenotypic traits. We explored a cost of nest building experimentally to determine whether large nests attracted more ectoparasites. This cost may be especially important to cavity-nesting species where larger nests likely reduce the distance between nest contents and the cavity entrance, making the host nest more vulnerable to ectoparasite detection. We manipulated the amount of nest material and the distance between nest and cavity entrance and estimated the abundance of ectoparasites (*Carnus hemapterus* flies) before and after manipulation in nests of Spotless Starlings (*Sturnus unicolor*) breeding in nestboxes. Nest size was enlarged and elevated in a first group of nests by adding nest material to the base of the nest (enlarged nests); a second group of nests was elevated with a piece of plastic but without adding nest material (elevated nests); a third group of nests was not manipulated (control nests). Number of flies per nest increased significantly in enlarged and elevated nests compared to control nests. Moreover, distance to cavity entrance after manipulation was negatively associated with the increase in the number of flies per nest (i.e., nests closer to the entrance had more parasites). Thus, nest size, through its effect on distance between nest and cavity entrance, affected fly abundance. Cues produced by nest contents located closer to the cavity entrance may be detected more readily by ectoparasitic flies searching for hosts. These results suggest a novel constraint for the evolution of nest size in cavity-nesting birds.

Keywords: Blood-sucking flies, cavity depth, distance to cavity entrance, ectoparasitism risk, host location, nest building, nest depth, nest size

25 Introduction

26 Avian nest building behaviour and nest characteristics have attracted a great deal of
27 attention by zoologists, ecologists and evolutionary biologists. Nests act as extended
28 phenotypes evolving through both natural and sexual selection, and provide a diverse
29 array of benefits to their builders (Hansell 2000). Nest materials may, for instance,
30 provide thermal insulation (Hoi et al. 1994, Hansell 2000), act as a physical barrier to
31 predators or brood parasites (Soler et al. 2014), be a source of chemical compounds
32 with insecticidal (Clark 1991, Tomás et al. 2012) and/or antimicrobial properties
33 (Mennerat et al. 2009, Peralta-Sánchez et al. 2010, Ruiz-Castellano et al. 2016), or even
34 act as ornaments to attract potential mates (Brouwer and Komdeur 2004, Veiga et al.
35 2006) and/or to elicit differential reproductive investment by mates in a post-mating
36 sexual selection scenario (Soler et al. 1998, 2001, De Neve and Soler 2002, Moreno
37 2012, Tomás et al. 2013). Yet there is considerable intraspecific variability in avian nest
38 features, including nest size, which is not fully understood (e.g., Soler et al. 1998,
39 Tomás et al. 2006, Cantarero et al. 2013, Maziarz et al. 2015).

40 Evolution of avian nests is constrained by costs associated with nest-building
41 activity and exaggeration of nest characteristics (Soler et al. 1998). Research on these
42 costs is therefore essential to understand the evolution of nest-building behaviours
43 and nest features. Most studies exploring the costs associated with nest building have
44 focused on birds, and most of them detected energetic costs (Mainwaring and Hartley
45 2013). Building a large nest might also imply costs in terms of predation (Collias and
46 Collias 1984, Biancucci and Martin 2010) and brood parasitism (Soler et al. 1999,
47 2014), simply because associated nest building activity and large nests are more easily
48 detected.

Flying ectoparasites may locate active nests by following cues derived from nest-building activities or nest size. Several correlative studies have addressed a potential relationship between nest size and ectoparasite abundance, based on the hypothesis that larger nests can sustain larger ectoparasite populations due to density-dependent processes (Eeva et al. 1994, Heeb et al. 2000, Kleindorfer and Dudaniec 2009), or because ectoparasites have more substrate in which to hide and escape nest sanitation behaviours performed by birds (Soler et al. 2007). Another rarely explored possibility is that larger nests attract more ectoparasites due to easier detectability (Tomás et al. 2008a), as suggested in the case of predators (Collias and Collias 1984, Biancucci and Martin 2010) and brood parasites (Soler et al. 1999, 2014).

Ectoparasites inflict severe costs to breeding adults and nestlings, either directly through blood consumption (Merino and Potti 1995), or indirectly acting as vectors of harmful internal parasites (Merino et al. 2000). Blood-sucking flying ectoparasites, or biting flies, are assumed to locate hosts mainly by means of visual and olfactory cues (Allan et al. 1987, Bowen 1991, Gibson and Torr 1999, Lehane 2005). Recently, a potential role for acoustic cues has also been suggested in host detection by ectoparasites (Tomás and Soler 2016). A large nest can be more easily detected than a small one, either visually or because of the increased number of volatiles released by nest materials (see Tomás et al. 2008a). Because nest materials may also trap odours released by birds, larger nests may also retain increased amounts of odorous cues. Relying on chemical or acoustic cues for host detection could be especially important for parasites of hole-nesting birds, where visual cues from nests are constrained.

In hole-nesting birds, nest size should be related to olfactory cues not only because of the association between the amount of nest material and odour intensity,

but also because of a relationship between the amount of nest material and distance from nest contents to cavity entrance. Olfactory cues emitted by nests and nestlings would be easier to locate when distance between nest and cavity entrance is reduced because of air flux patterns inside the cavity (Conover 2007). Similarly, acoustic cues emitted by nestlings would be easier to detect when distance between nest and cavity entrance is reduced. Again, natural nests show a remarkable variability regarding the distance between the nest and the cavity entrance (e.g., Tomás et al. 2006, Maziarz et al. 2015). Therefore, because distance from the nest to cavity entrance is strongly determined by the amount of nest material (Tomás et al. 2006), nest size of hole-nesting birds may be positively related to risk of parasitism.

Carnus hemapterus (hereafter “*Carnus*”) has important negative effects on nestlings (Hoi et al. 2018) and is the most abundant ectoparasite in the Spotless Starling (*Sturnus unicolor*) population reported here (Avilés et al. 2009, Azcárate-García et al. 2019). *Carnus* is a 2-mm fly in which the adult is the blood-sucking stage, and is found in nests of an extremely wide diversity of birds (Brake 2011, Calero-Torralbo 2011). After emergence from overwintering pupae inside or around nests, winged adults may disperse, losing their wings once a suitable host is found (Grimaldi 1997, Calero-Torralbo 2011). *Carnus* feeds mainly on nestlings, where it can be found quickly searching for cover in suitable body areas such as the axilla, inguinal region, around the neck, or inside ears. It feeds also on incubating adults (López-Rull et al. 2007), yet flies are not found on birds outside the nest (Grimaldi 1997). This suggests that, although there is still a poor understanding of its colonization patterns (Veiga et al. 2020), the main dispersion mode of *Carnus hemapterus* is by active flight, which can be over relatively long distances.

In this study, we tested experimentally the association between nest size, distance to cavity entrance, and intensity of parasitism in a population of starlings breeding in nestboxes of identical dimensions and characteristics. We manipulated nest size and distance between nest and cavity entrance shortly after hatching and estimated abundance of *Carnus* ectoparasitic flies in nests before and after manipulation. Nest size was enlarged in a first group of nests by adding nest materials under the base of the nest (enlarged nests). A second group of nests was elevated without adding nest materials (elevated nests). In both groups, distance between nest and cavity entrance was reduced. A third group of nests were maintained as controls. We predicted that distance from the nest to cavity entrance was a major determinant of biting fly abundance. Consequently, ectoparasite abundance should be higher in enlarged and elevated nests as compared with control nests. On the contrary, if nest size rather than distance from nest to cavity entrance determines abundance of ectoparasites, enlarged nests should harbour more parasites than elevated or control nests because the amounts of volatiles released would be positively related to the amount of nest materials. Finally, if distance from nest to cavity entrance predicts probability of nest location by ectoparasites, this distance should be related to abundance of ectoparasites.

Material and methods

Study area and species

Fieldwork was conducted during the 2012 breeding season (April-June) in Guadix, southeastern Spain (37°18'N, 3°11'W). Spotless Starlings (hereafter starlings) bred in

artificial nestboxes that were installed in 2005 in agro-pastoral habitat. Nestboxes made of cork (internal dimensions: 350 mm [height] x 180 mm [width] x 210 mm [depth]; distance from bottom to hole: 240 mm) were attached to walls of abandoned buildings or tree trunks at 3–4 m above the ground. Typical clutch sizes are four to five eggs and incubation usually starts with the third or fourth egg and lasts 12–13 days (Soler et al. 2008). Nests are built by both sexes, but females carry most nest material (Veiga and Polo 2011) which, in our study area, is mainly dry straw. Other nest material found in starling nests, such as green plants, fresh flowers and feathers, has been interpreted as ornamentation from sexual displays of males (plants) and females (feathers) (Polo and Veiga 2006, Veiga et al. 2006). These hypothetical ornamental materials may also function in protecting offspring from microorganisms (Mennerat et al. 2009, Peralta-Sánchez et al. 2010, Ruiz-Castellano et al. 2016, 2019, Soler et al. 2017) and/or ectoparasitic arthropods (Clark 1991, Tomás et al. 2012).

Nest manipulations

Old nest material was removed from nestboxes before the breeding season (on the last week of March) and transported to the lab, thus ensuring that most parasites overwintering in nest material were removed. Thus, any flies quantified in this study colonized nestboxes from outside experimental nests. From April 4th, nestboxes were checked on a regular basis to allow determining hatching dates (i.e., every four days until egg laying was detected and daily close to the expected hatching date). Three days after the first egg hatched, nestlings were removed temporarily and placed in a container covered with a white paper napkin. Nests were also removed carefully from nestboxes and alternately assigned to one of three treatments, assigning the first nest

randomly: i) enlarged ($n = 18$), ii) elevated ($n = 17$), and iii) control ($n = 16$). In enlarged nests, approximately 10 cm of nest material was added under the base of the nest; dry straw, a common nest material in our starling population, were used for this purpose. This material was collected from the ground in a nearby area but far from any nest, and was thus free of ectoparasites because *Carnus* pupae overwinters in nests (Roulin 1998). In elevated nests, a piece of plastic of approximately 10-cm height was inserted under the nest; we used an empty 5-L plastic container turned upside down and “covered” with a 2-cm-thick polystyrene sheet to provide horizontal support for the nest. In control nests, original nest material was returned to nestboxes without further manipulation. Distance between the nest rim and the cavity entrance was measured with a ruler to the nearest 0.5 cm before and after the manipulation. The manipulation led to final nest sizes and distances to cavity entrance that were within the natural range recorded in the study area (pers. obs.). Starlings occupy a great diversity of nesting sites and show a remarkable variability in these nest features.

Cavity temperature and humidity

The microclimate of the cavity may cause differences in abundance and survival of certain ectoparasites (Bennett and Whitworth 1991, Chilton et al. 2000, Heeb et al. 2000, Castaño-Vázquez et al. 2018). Therefore, temperature and humidity were recorded in a subsample of nests ($n = 26$; 9 enlarged, 9 elevated and 8 control nests) to test whether nest manipulations affected microclimate of the cavity, which could subsequently affect ectoparasite abundance. For this purpose, a datalogger (Thermocron iButton, Dallas Semiconductor, Dallas, TX), was attached to the inner lateral wall of the nestbox above the nest rim (e.g., Ardia and Clotfelter 2007). The

datalogger recorded temperature (± 0.5 °C) and relative humidity (± 0.01 %) at 12-min intervals over 48 h, beginning at 2000 hr on day 3 of the nestling stage, thus providing 240 measurements per nest. Mean, minimum and maximum temperature and humidity for each nest were obtained.

Ectoparasite Abundance

Although patterns of *Carnus* colonization vary among nests, in starling nests, *Carnus* abundance often peaks shortly after hatching, reaching highest abundance when day 5 of the nestling stage (pers. obs., see also Dawson and Bortolotti 1997, Roulin 1998, Liker et al. 2001). As nestlings age, *Carnus* abundance decreases but estimation is less reliable with increasing feather development in nestlings. Therefore, the abundance of *Carnus* flies was estimated twice: when nestlings were 3 days (immediately before nest manipulation) and 5 days old to look for experimental differences. The window for nest manipulations and ectoparasite quantification was narrow out of necessity, as manipulating nests before day 3 is not recommended due to risk of abandonment, and estimating *Carnus* abundance after day 5, when feathers start growing, becomes less reliable (pers. obs.). Nestling mass was recorded at day 3 with an electronic balance (Pocket-240; Gram Precision, Barcelona, Spain) to the nearest 0.1 g, and within-nest means were used in analyses to avoid pseudoreplication.

Nestlings were removed simultaneously to avoid any *Carnus* falling off nestlings, and placed in a container covered with a clean white paper napkin. No fly was observed in nests after removal of nestlings. One nestling at a time was carefully examined, forcing all *Carnus* to fall off into another container until no flies were present on any nestling. Finally, all *Carnus* were counted in the container and returned

to the nestlings that were then placed back in the nest. This method employed for estimation of *Carnus* abundance has been previously shown to be reliable (Roulin 1998) and highly repeatable (Veiga et al. 2020). Five of the 51 nests were not visited on nestling day 5 because of cold and/or rainy weather (which may increase probability of nest abandonment), and 5 additional nests were depredated or abandoned. Nests that were not measured on day 5 were not biased relative to their assigned experimental treatment (four enlarged, two elevated and four control nests) and thus, 41 nests were used to explore the effect of the experimental treatments.

At nestling day 8, a sticky trap was installed on nestboxes to sample biting flies other than *Carnus*. The trap consisted of a plastic Petri dish placed inside the nestbox close to the roof facing downwards, supported on a wire mesh and covered with oil (Tomás et al. 2008b). This sampling method has been designed to estimate abundances of biting flies such as Simuliid blackflies and Ceratopogonid biting midges that rest on the roof inside nestboxes (Tomás et al. 2008b). The trap was in use until nestling day 14, when Petri dishes were collected and screened for biting flies. Latex gloves were always worn when handling nests and nestlings to avoid releasing odorous cues that might interfere with the experiment.

Statistical Analyses

Mean, maximum and minimum temperature and humidity inside cavities were compared among treatments with ANOVAs including hatching date (1 = 1 April) as a continuous variable. Comparisons of ectoparasite abundance at days 3 and 5 after hatching were performed using repeated measures ANOVAs, with number of flies per nest at the two sampling ages as the repeated measures (dependent variable), and

hatching date, brood size and mean nestling body mass as between-factor variables. The effect of the experimental treatments on ectoparasite abundance (i.e., response variable) was therefore estimated by the interaction between experimental treatment and the repeated measure of number of flies per nest (i.e., difference in total number of flies counted in all nestlings of a nest at nestling days 3 and 5). The interactions between repeated measures and between factors therefore estimated the effects of hatching date, brood size and nestling body mass on the change detected in ectoparasite abundance estimated at the two different nestling ages.

Change in parasitism intensity experienced by enlarged and elevated nests was similar (see Results) and, thus, these two groups were considered together when analysing predictions derived from the hypothesis that distance from the nest to cavity entrance, rather than amount of nest material, affects detectability by parasites. The effect of the experiment on ectoparasite abundance was also explored using initial and final distance to entrance rather than experimental treatments. In this General Linear Model, change in number of flies per nest between days 3 and 5 of nestling age was used as the dependent variable, distance to entrance before and after the manipulation as continuous independent variables, and hatching date, brood size and nestling body mass as covariables.

All statistical tests were performed in Statistica 8.0 (Statsoft Inc. 2007). Residuals of the models were tested for normality (Kolmogorov-Smirnov tests: all $P > 0.10$).

Results

Before the manipulation, distance to entrance did not differ between nests subsequently assigned to enlarged (mean $\pm$ SE: 17.4 ± 0.5 cm), elevated (16.5 ± 0.5 cm)

and control nests (16.9 ± 0.5 cm; ANOVA: $F_{2,47} = 0.88$, $P = 0.42$). After the manipulation, distance to entrance differed significantly among treatments: (enlarged nests: 7.1 ± 0.5 cm; elevated nests: 3.6 ± 0.5 cm; control nests: 16.7 ± 0.6 cm; ANOVA: $F_{2,47} = 143.22$, $P < 0.001$). Distance to entrance of elevated nests was slightly shorter than distance to entrance of enlarged nests (post-hoc Unequal N Honestly Significant Difference test: $P < 0.001$), while distance to entrance of the two experimental treatments was approximately half the distance to entrance of control nests (enlarged vs. control: $P < 0.001$; elevated vs. control: $P < 0.001$).

Experimental treatments did not create significant differences in microclimate of the cavities. Mean, maximum and minimum temperature and relative humidity did not differ among treatments (Table 1). Hatching date was positively related to mean and maximum temperature, and negatively related to mean, maximum and minimum relative humidity (Table 1).

Abundance of ectoparasitic flies other than *Carnus* was very low in starling nests: Simuliid biting flies were only found in sticky traps in one control nest, and thus we will not refer to them further. No other parasites (e.g., Calliphoridae blowflies) were apparently present in starling nests. On the contrary, *Carnus* was highly prevalent in nests, both at nestling day 3 (94.1%, $n = 51$) and 5 (95.1%, $n = 41$) of nestling age. Before the experiment, three days after nestlings hatched, abundance of *Carnus* parasites in nests under different experimental treatments did not differ significantly (enlarged nests: 12.4 ± 3.0 ; elevated nests: 8.8 ± 3.1 ; control nests: 16.0 ± 3.2 ; ANOVA: $F_{2,48} = 1.34$, $P = 0.27$). Similarly, there were no significant differences among experimental treatments in hatching date (enlarged nests: 39.6 ± 1.0 ; elevated nests: 38.0 ± 1.0 ; control nests: 39.1 ± 1.0 ; ANOVA: $F_{2,48} = 0.65$, $P = 0.53$), brood size (enlarged

264 nests: 3.2 ± 0.2 ; elevated nests: 3.5 ± 0.2 ; control nests: 3.4 ± 0.2 ; ANOVA: $F_{2,48} = 0.82$,
265 $P = 0.45$) and mean nestling mass (enlarged nests: 19.6 ± 1.0 g; elevated nests: $19.7 \pm$
266 1.1 g; control nests: 19.8 ± 1.1 ; ANOVA: $F_{2,48} = 0.01$, $P = 0.99$) at three days of age.

267 Number of flies per nest at day 3 was higher in nests with larger brood sizes (β
268 $[SE] = 0.32 [0.15]$) and tended to be higher in nests with heavier nestlings ($\beta [SE] = 0.35$
269 $[0.15]$) (between effects in Table 2). Number of flies per nest tended to decrease with
270 age (day 3: 12.00 ± 1.99 , $n = 51$; day 5: 10.68 ± 1.39 , $n = 41$), mostly in control nests as
271 compared to enlarged and elevated nests (Table 2; Figure 1). Change in number of flies
272 per nest in enlarged (-1.0 ± 2.9 flies) and elevated nests (2.5 ± 2.8 flies) was similar
273 (repeated measures ANOVA, interaction with treatment, $F_{1,27} = 1.10$, $P = 0.30$). Thus,
274 these two groups were considered together when analyzing the prediction that
275 distance from the nest to cavity entrance, rather than amount of nest material, affects
276 detectability by parasites. In this case, treatment explained change in number of flies
277 per nest between days 3 and 5 after hatching (repeated measures ANOVA, interaction
278 between age and treatment, $F_{1,36} = 4.35$, $P = 0.04$). Number of flies per nest was
279 increased in enlarged and elevated nests compared to control nests (Figure 1) and was
280 higher in nests with larger brood sizes ($F_{1,36} = 5.22$, $P = 0.03$).

281 Finally, and in accordance with the hypothesis that distance to cavity entrance
282 affects risk of ectoparasitism, we found a statistically significant negative association
283 between difference in number of flies per nest and distance to entrance after
284 manipulation ($\beta [SE] = -0.45 [0.16]$; $F_{1,34} = 8.33$, $P = 0.007$; Figure 2), once the negative
285 effect of nestling body mass was controlled for, i.e., number of flies per nest decreased
286 more in nests with heavier nestlings ($\beta [SE] = -0.41 [0.16]$; $F_{1,34} = 6.84$, $P = 0.01$).

Discussion

Our main findings are that ectoparasite abundance in experimentally enlarged and elevated nests was larger than in control nests, and that distance from nest to cavity entrance after the experimental manipulation was negatively related to change in ectoparasite abundance. The abundance of *Carnus hemapterus* flies was not different between the two experimental treatments (enlarged and elevated nests), which indicates that the effects detected were not due to adding additional nesting material (i.e., to chemicals originated from nest material; Tomás et al. 2008b), but to olfactory cues derived from nestling physiological activity (Azcárate-García et al. 2019), and/or auditory cues derived from begging behaviour (Tomás and Soler 2016). Nestlings in nests closer to the cavity entrance would be more easily located by ectoparasites. These results cannot be attributed to differences in cavity microclimate derived from the experimental manipulations. Our findings may have broad implications for understanding the ecology and evolution of cavity-nesting birds. Below we discuss alternative explanations, and the importance of considering the selective pressure exerted by parasitism for understanding the evolution of nest-building behaviour in hole-nesting birds, where larger nests would attract more parasites due to a reduction in the distance from nest contents to cavity entrance.

Given that old nest material with any overwintering parasites was removed from experimental nestboxes before nests had been initiated, we are confident that the vast majority of ectoparasites detected during the nestling phase were from nearby areas. Abundance of *Carnus* at a given nestling stage is a balance between increasing abundance by flies that are newly arrived (immigration) and decreasing abundance due to completion of life cycle, death, or elimination by host birds. Thus, we did not

necessarily expect a differential increase among treatments, but rather differences between post- and pre- experimental abundances, due to this dynamic balance regulating ectoparasite abundance in every nest at a given time (Dawson and Bortolotti 1997, Liker et al. 2001, Veiga et al. 2018). Moreover, it has been previously shown that *Carnus* has detrimental effects on their bird hosts (Wiebe 2009, Hoi et al. 2018), including starlings (Avilés et al. 2009, Tomás et al. 2018). Consequently, our experimental results suggest the existence of costs for starlings building large nests, mediated by blood-sucking parasites.

In an attempt to differentiate between the effects of olfactory cues derived from nest material and those derived from nestling activity as attractors of flying ectoparasites, we elevated the nests with either plant nest material or with artificial plastic material, while assuming that the former but not the latter may include detectable olfactory cues for parasites. We found non-significant differences among these experimental treatments, suggesting that the distance between the nest and the cavity entrance, rather than amount of volatiles derived from nest materials, was the cause of the patterns detected. However, we cannot rule out the hypothesis that parasites cue on volatile emissions from nest material, vegetable or other. This alternative explanation could be a function of the nest material used for the experimental manipulation of nest size. We chose dry straw, following the rationale that it is the main nest material used by starlings in this and other populations (Cramp 1998, Veiga and Polo 2011). However, this is apparently a nest material with limited odorous properties (Tomás et al. 2012), which may make it difficult to detect. This hypothesis is however non-exclusive and may operate additively with the effect of distance to the cavity entrance, or with other nest materials when considering other

species. Another hypothetical scenario predicting a positive relationship between nest size and parasite abundance is that ectoparasites in larger nests have more space to hide and escape from nest sanitation behaviours performed by adult birds (Heeb et al. 2000, Soler et al. 2007). We found that number of flies per nest increased in elevated nests with no additional material, which does not support this hypothesis.

Finally, our results may also be of interest in the debate regarding suitability of using artificial nestboxes for studying factors affecting the risk of parasitism or related life-history traits. Lambrechts et al. (2010) urged researchers to report detailed characteristics of nestboxes because those may affect or confound conclusions derived from nestbox studies. Our results indicate that distance to the cavity entrance should be reported as an important parameter affecting ectoparasite load, therefore supporting this claim. Furthermore, cavity-nesting birds may prefer deeper cavities for breeding to reduce risk of ectoparasitism and associated diseases, in line with what has been suggested regarding predation risk (Wesołowski 2002, Wesołowski and Rowinski 2012, Kaliński et al. 2014, Mainwaring et al. 2015, Fokkema et al. 2018). Other characteristics of the nesting cavity such as orientation of the entrance hole (Goodenough et al. 2008), location of the cavity (e.g., tree or wall) (Lambrechts et al. 2010), or proximity to other cavities occupied in the current or the previous season (Veiga et al, 2020), will surely affect nest colonization success by ectoparasites and deserve further research.

To conclude, our results suggest that building a large nest in a cavity carries an important and overlooked cost by reducing the distance between nest contents (eggs, nestlings, or incubating/brooding/roosting adult) and the cavity entrance, facilitating nest detection by ectoparasites. This may be a strong evolutionary force restricting

nest size in cavities, and therefore the benefits of building larger nests derived from its functioning in scenarios of natural and/or sexual selection (Soler et al. 1998, Moreno 2012, Mainwaring et al. 2014).

Data availability statement

Data used in this paper are available from the CSIC Institutional Repository (<http://dx.doi.org/10.20350/digitalCSIC/12512>).

Acknowledgements – This section has been deleted for blind review.

References

- Allan, S. A., Day, J. F. and Edman, J. D. 1987. Visual ecology of biting flies. – *Ann. Rev. Entomol.* 32: 297–316.
- Ardia, D. R. and Clotfelter, E. D. 2007. Individual quality and age affect responses to an energetic constraint in a cavity-nesting bird. – *Behav. Ecol.* 18: 260–266.
- Avilés, J. M., Pérez-Contreras, T., Navarro, C. and Soler, J. J. 2009. Male spotless starlings adjust feeding effort based on egg spots revealing ectoparasite load. – *Anim. Behav.* 78: 993–999.
- Azcárate-García, M., Ruiz-Rodríguez, M., Díaz-Lora, S., Ruiz-Castellano, C. and Soler, J. J. 2019. Experimentally broken faecal sacs affect nest bacterial environment, development and survival of spotless starling nestlings. – *J. Avian Biol.* 50: e02044.
- Bennett, G. F. and Whitworth, T. L. 1991. Studies on the life history of some species of *Protocalliphora* (Diptera: Calliphoridae). – *Can. J. Zool.* 69: 2048–2058.

383 Biancucci, L. and Martin, T. E. 2010. Can selection on nest size from nest predation
 384 explain the latitudinal gradient in clutch size? – J. Anim. Ecol. 79: 1086–1092.
 385 Bowen, M. F. 1991. The sensory physiology of host-seeking behavior in mosquitoes. –
 386 Ann. Rev. Entomol. 36: 139–158.
 387 Brake, I. 2011. World catalog of the family Carnidae (Diptera, Schizophora). – Myia 12:
 388 113–169.
 389 Brouwer, L. and Komdeur, J. 2004. Green nesting material has a function in attraction
 390 in the European starling. – Anim. Behav. 67: 539–548.
 391 Calero-Torralbo, M. A. 2011. Factores ecológicos y mecanismos implicados en la
 392 variabilidad de la interacción entre un ectoparásito generalista (*Carnus*
 393 *hemapterus*) y sus hospedadores. – Ph.D. dissertation, Univ. Granada, Spain.
 394 Cantarero, A., López-Arrabé, J., Rodríguez-García, V., González-Braojos, S., Ruiz-de-
 395 Castañeda, R., Redondo, A. J. and Moreno, J. 2013. Factors affecting the presence
 396 and abundance of generalist ectoparasites in nests of three sympatric hole-
 397 nesting bird species. – Acta Ornithol. 48: 39–54.
 398 Castaño-Vázquez, F., Martínez, J., Merino, S. and Lozano, M. 2018. Experimental
 399 manipulation of temperature reduce ectoparasites in nests of blue tits (*Cyanistes*
 400 *caeruleus*). – J. Avian Biol. 49: e01695.
 401 Chilton, N. B., Andrews, R. H. and Bull, C. M. 2000. Influence of temperature and
 402 relative humidity on the moulting success of *Amblyomma limbatum* and
 403 *Aponomma hydrosauri* (Acari: Ixodidae) larvae and nymphs. – Int. J. Parasitol. 30:
 404 973–979.

405 Clark, L. 1991. The nest protection hypothesis: the adaptive use of plant secondary
 406 compounds by European starlings. – In: Loye, J. E. and Zuk, M. (eds.), Bird-parasite
 407 interactions: ecology, evolution and behaviour. Oxford Univ. Press, pp. 205–221.
 408 Collias, N. E. and Collias, E. C. 1984. Nest building and bird behaviour. – Princeton Univ.
 409 Press.
 410 Conover, M. 2007. Predator-prey dynamics: the role of olfaction. – CRC Press.
 411 Cramp, S. 1998. The complete birds of the Western Palearctic. – Oxford Univ. Press.
 412 Dawson, R. D. and Bortolotti, G. R. 1997. Ecology of parasitism of nestling American
 413 kestrels by *Carnus hemapterus* (Diptera: Carnidae). – Can. J. Zool. 75: 2021–2026.
 414 De Neve, L. and Soler, J. J. 2002. Nest-building activity and laying date influence female
 415 reproductive investment in magpies: an experimental study. – Anim. Behav. 63:
 416 975–980.
 417 Eeva, T., Lehikoinen, E. and Nurmi, J. 1994. Effects of ectoparasites on breeding
 418 success of great tits (*Parus major*) and pied flycatchers (*Ficedula hypoleuca*) in an
 419 air pollution gradient. – Can. J. Zool. 72: 624–635.
 420 Fokkema, R. W., Ubels, R. and Tinbergen, J. M. 2018. Experimentally quantifying the
 421 effect of nest-site depth on the predation risk and breeding success of Blue Tits. –
 422 Auk 135: 919–932.
 423 Gibson, G. and Torr, S. J. 1999. Visual and olfactory responses of haematophagous
 424 diptera to host stimuli. – Med. Vet. Entomol. 13: 2–23.
 425 Goodenough, A. E., Maitland, D. P., Hart, A. G. and Elliot, S. L. 2008. Nestbox
 426 orientation: a species-specific influence on occupation and breeding success in
 427 woodland passerines. – Bird Study 55: 222–232.

428 Grimaldi, D. 1997. The bird flies, genus *Carnus*: species revision, generic relationships,
 429 and a fossil *Meoneura* in amber (Diptera: Carnidae). – Am. Mus. Novit. 3190: 1–
 430 30.
 431 Hansell, M. 2000. Bird nests and construction behaviour. – Cambridge Univ. Press.
 432 Heeb, P., Kolliker, M. and Richner, H. 2000. Bird-ectoparasite interactions, nest
 433 humidity and ectoparasite community structure. – Ecology 81: 958–968.
 434 Hoi, H., Schleicher, B. and Valera, F. 1994. Female mate choice and nest desertion in
 435 penduline tits, *Remiz pendulinus*: the importance of nest quality. – Anim. Behav.
 436 48: 743–746.
 437 Hoi, H., Darolová, A., Krištofík, J. and Hoi, C. 2018. The effect of the ectoparasite
 438 *Carnus hemapterus* on immune defence, condition, and health of nestling
 439 European Bee-eaters. – J. Ornithol. 159: 291–302.
 440 Kaliński, A., Wawrzyniak, J., Bańbura, M., Skwarska, J., Zieliński, P., Gładalski, M. and
 441 Bańbura, J. 2014. Does the threat of European Pine Marten (*Martes martes*)
 442 predation influence the height of nests built by Blue Tits (*Cyanistes caeruleus*) and
 443 Great Tits (*Parus major*)? – Avian Biol. Res. 7: 83–90.
 444 Kleindorfer, S. and Dudaniec, R. Y. 2009. Love thy neighbour? Social nesting pattern,
 445 host mass and nest size affect ectoparasite intensity in Darwin's tree finches. –
 446 Behav. Ecol. Sociobiol. 63: 731–739.
 447 Lambrechts, M. M., Adriaensen, F., Ardia, D. R., Artemyev, A. V., Atiénzar, F., Bańbura,
 448 J., Barba, E., Bouvier, J. C., Camprodon, J., Cooper, C. B., Dawson, R. D. et al. 2010.
 449 The design of artificial nestboxes for the study of secondary hole-nesting birds: a
 450 review of methodological inconsistencies and potential biases. – Acta Ornithol. 45:
 451 1–26.

452 Lehane, M. J. 2005. The evolution of blood-sucking in insects. – Cambridge Univ. Press.
 453 Liker, A., Márkus, M., Vozár, Á., Zemankovics, E. and Rózsa, L. 2001. Distribution of
 454 *Carnus hemapterus* in a starling colony. – Can. J. Zool. 79: 574–580.
 455 López-Rull, I., Gil, M. and Gil, D. 2007. Spots in Starling *Sturnus unicolor* eggs are good
 456 indicators of ectoparasite load by *Carnus hemapterus* (Diptera: Carnidae). –
 457 Ardeola 54: 131–134.
 458 Mainwaring, M. C. and Hartley, I. R. 2013. The energetic costs of nest building in birds.
 459 – Avian Biol. Res. 6: 12–17.
 460 Mainwaring, M. C., Hartley, I. R., Lambrechts, M. M. and Deeming, D. C. 2014. The
 461 design and function of birds' nests. – Ecol. Evol. 20: 3909–3928.
 462 Mainwaring, M. C., Reynolds, S. J. and Weidinger, K. 2015. The influence of predation
 463 on the location and design of nests. – In: Deeming, D. C. and Reynolds, S. J. (eds.),
 464 Nests, eggs, and incubation: new ideas about avian reproduction. Oxford Univ.
 465 Press, pp. 50–64.
 466 Maziarz, M., Wesołowski, T., Hebda, G. and Cholewa, M. 2015. Natural nest-sites of
 467 Great Tits (*Parus major*) in a primeval temperate forest (Białowieża National Park,
 468 Poland). – J. Ornithol. 156: 613–623.
 469 Mennerat, A., Mirleau, P., Blondel, J., Perret, P., Lambrechts, M. M. and Heeb, P. 2009.
 470 Aromatic plants in nests of the blue tit *Cyanistes caeruleus* protect chicks from
 471 bacteria. – Oecologia 161: 849–855.
 472 Merino, S. and Potti, J. 1995. Mites and blowflies decrease growth and survival in
 473 nestling pied flycatchers. – Oikos 73: 95–103.

474 Merino, S., Moreno, J., Sanz, J. J. and Arriero, E. 2000. Are avian blood parasites
 475 pathogenic in the wild? A medication experiment in blue tits (*Parus caeruleus*). –
 476 Proc. R. Soc. Lond. B 267: 2507–2510.

477 Moreno, J. 2012. Avian nests and nest-building as signals. – Avian Biol. Res. 5: 238–251.

478 Peralta-Sánchez, J. M., Møller, A. P., Martín-Platero, A. M. and Soler, J. J. 2010. Number
 479 and colour composition of nest lining feathers predict eggshell bacterial
 480 community in barn swallow nests: an experimental study. – Funct. Ecol. 24: 426–
 481 433.

482 Polo, V. and Veiga, J. P. 2006. Nest ornamentation by female spotless starlings in
 483 response to a male display: an experimental study. – J. Anim. Ecol. 75: 942–947.

484 Roulin, A. 1998. Cycle de reproduction et abondance du diptère *Carnus hemapterus*
 485 dans les nichées de chouettes effraies *Tyto alba*. – Alauda 66: 265–272.

486 Ruiz-Castellano, C., Tomás, G., Ruiz-Rodríguez, M., Martín-Gálvez, D. and Soler J. J.
 487 2016. Nest material shapes eggs bacterial environment. – PLoS ONE 11: e0148894.

488 Ruiz-Castellano, C., Ruiz-Rodríguez, M., Tomás, G. and Soler, J. J. 2019. Antimicrobial
 489 activity of nest-lining feathers is enhanced by breeding activity in avian nests. –
 490 FEMS Microbiol. Ecol. 95: fiz052.

491 Soler, J. J., Møller, A. P. and Soler, M. 1998. Nest building, sexual selection and parental
 492 investment. – Evol. Ecol. 12: 247–441.

493 Soler, J. J., Martínez, J. G., Soler, M. and Møller, A. P. 1999. Host sexual selection and
 494 cuckoo parasitism: an analysis of nest size in sympatric and allopatric magpie *Pica*
 495 *pica* populations parasitized by the great spotted cuckoo *Clamator glandarius*. –
 496 Proc. R. Soc. Lond. B 266: 1765–1771.

497 Soler, J. J., De Neve, L., Martínez, J. G. and Soler, M. 2001. Nest size affects clutch size
 498 and the start of incubation in magpies: an experimental study. – Behav. Ecol. 12:
 499 301–307.

500 Soler, J. J., Martín-Vivaldi, M., Haussy, C. and Møller, A. P. 2007. Intra- and interspecific
 501 relationships between nest size and immunity. – Behav. Ecol. 18: 781–791.

502 Soler, J. J., Navarro, C., Pérez-Contreras, T., Avilés, J. M. and Cuervo, J. J. 2008. Sexually
 503 selected egg coloration in spotless starlings. – Am. Nat. 171: 184–194.

504 Soler, J. J., Avilés, J. M., Martín-Gálvez, D., De Neve, L. and Soler, M. 2014.
 505 Eavesdropping cuckoos: further insights on great spotted cuckoo preference by
 506 magpie nests and egg colour. – Oecologia 175: 105–115.

507 Soler, J. J., Ruiz-Castellano, C., Figuerola, J., Martín-Vivaldi, M., Martínez-de la Puente,
 508 J., Ruiz-Rodríguez, M. and Tomás, G. 2017. Telomere length and dynamics of
 509 spotless starling nestlings depend on nest-building materials used by parents. –
 510 Anim. Behav. 126: 89–100.

511 StatSoft, Inc. 2007. Statistica (data analysis software system), Version 12. – Available
 512 from: <http://www.statsoft.com>.

513 Tomás, G. and Soler, J. J. 2016. Begging and ectoparasite attraction. – Anim. Behav.
 514 113: 93–98.

515 Tomás, G., Merino, S., Moreno, J., Sanz, J. J., Morales, J. and García-Fraile, S. 2006. Nest
 516 weight and female health in the blue tit (*Cyanistes caeruleus*). – Auk 123: 1013–
 517 1021.

518 Tomás, G., Merino, S., Martínez-de la Puente, J., Moreno, J., Morales, J. and Lobato, E.
 519 2008a. Determinants of abundance and effects of blood-sucking flying insects in
 520 the nest of a hole-nesting bird. – Oecologia 156: 305–312.

521 Tomás, G., Merino, S., Martínez-de la Puente, J., Moreno, J., Morales, J., and Lobato, E.
 522 2008b. A simple trapping method to estimate abundances of blood-sucking flying
 523 insects in avian nests. – Anim. Behav. 75: 723–729.

524 Tomás, G., Merino, S., Martínez-de la Puente, J., Moreno, J., Morales, J., Lobato, E.,
 525 Rivero-de Aguilar, J. and Del Cerro, S. 2012. Interacting effects of aromatic plants
 526 and female age on nest-dwelling ectoparasites and blood-sucking flies in avian
 527 nests. – Behav. Proc. 90: 246–253.

528 Tomás, G., Merino, S., Martínez-de la Puente, J., Moreno, J., Morales, J. and Rivero-de
 529 Aguilar, J. 2013. Nest size and aromatic plants in the nest as sexually selected
 530 female traits in blue tits. – Behav. Ecol. 24: 927–934.

531 Tomás, G., Martín-Gálvez, D., Ruiz-Castellano, C., Ruiz-Rodríguez, M., Peralta-Sánchez,
 532 J. M., Martín-Vivaldi, M. and Soler, J. J. 2018. Ectoparasite activity during
 533 incubation increases microbial growth on avian eggs. – Microb. Ecol. 76: 555–564.

534 Veiga, J. P. and Polo, V. 2011. Estornino negro – *Sturnus unicolor*. – In: Salvador, A. and
 535 Morales, M. B. (eds.), Enciclopedia virtual de los vertebrados españoles. Museo
 536 Nacional de Ciencias Naturales, Madrid.

537 Veiga, J. P., Polo, V. and Viñuela, J. 2006. Nest green plants as a male status signal and
 538 courtship display in the spotless starling. – Ethology 112: 196–204.

539 Veiga, J., Moreno, E., Benzal, J. and Valera, F. 2018. Off-host longevity of the winged
 540 dispersal stage of *Carnus hemapterus* (Insecta: Diptera) modulated by gender,
 541 body size and food provisioning. – Parasitology 146: 241–245.

542 Veiga, J., Václav, R. and Valera, F. 2020. The effect of parasite density on host
 543 colonization success by a mobile avian ectoparasite. – Ecol. Entomol. in press. DOI:
 544 10.1111/een.12864.

- 545 Wesołowski, T. 2002. Anti-predator adaptations in nesting Marsh Tits *Parus palustris*:
546 the role of nest-site security. – Ibis 144: 593–601.
- 547 Wesołowski, T. and Rowinski, P. 2012. The breeding performance of Blue Tits *Cyanistes*
548 *caeruleus* in relation to the attributes of natural holes in a primeval forest. – Bird
549 Study 59: 437–448.
- 550 Wiebe, K. L. 2009. Nest excavation does not reduce harmful effects of ectoparasitism:
551 an experiment with a woodpecker, the northern flicker *Colaptes auratus*. – J.
552 Avian Biol. 40: 166–172.

553 Table 1. Comparison of microclimate of the cavities (temperature and relative humidity) after experimental manipulation of spotless starling
554 nests. Partial correlation coefficients (θ) for the relationships of microclimate variables with hatching date are shown.

		Treatment					Hatching date		
		Enlarged	Elevated	Control	$F_{2,22}$	P	$F_{1,22}$	P	$\theta \pm SE$
Temperature ($^{\circ}\text{C}$)	Mean	22.9 $\pm$ 0.4	22.6 $\pm$ 0.4	22.3 $\pm$ 0.4	0.00	1.00	13.66	0.001	0.63 $\pm$ 0.17
	Maximum	33.3 $\pm$ 1.0	32.3 $\pm$ 1.0	32.8 $\pm$ 1.0	0.47	0.63	14.28	0.001	0.65 $\pm$ 0.17
	Minimum	13.6 $\pm$ 0.7	12.8 $\pm$ 0.7	13.2 $\pm$ 0.7	0.33	0.72	0.34	0.57	0.13 $\pm$ 0.22
Relative humidity (%)	Mean	35.6 $\pm$ 1.2	36.1 $\pm$ 1.2	36.6 $\pm$ 1.2	0.00	1.00	4.98	0.04	-0.44 $\pm$ 0.20
	Maximum	63.0 $\pm$ 2.6	63.7 $\pm$ 2.6	64.3 $\pm$ 2.8	0.03	0.97	4.56	0.04	-0.43 $\pm$ 0.20
	Minimum	17.3 $\pm$ 0.8	17.4 $\pm$ 0.8	17.9 $\pm$ 0.9	0.43	0.65	26.11	<0.001	-0.76 $\pm$ 0.15

555

Table 2. Results from repeated measures ANOVA with number of flies per nest estimated in starling nests three and five days after hatching as repeated measures (i.e., age effect), in relation to body mass, brood size, hatching date and experimental treatments. Interactions between age and between factor variables estimate the effects of between factor variables on change in number of flies per nest associated with age. Effect size (partial eta-squared) and observed power (alpha = 0.05) are also shown.

	<i>SS</i>	<i>df</i>	<i>MS</i>	<i>F</i>	<i>P</i>	Effect Size	Power
BETWEEN EFFECTS							
Treatment	238.0	2,35	119.0	0.72	0.50	0.04	0.16
Body mass	536.3	1,35	536.3	3.23	0.08	0.08	0.42
Brood size	841.7	1,35	841.7	5.08	0.03	0.13	0.59
Hatching date	401.2	1,35	401.2	2.42	0.13	0.06	0.33
WITHIN EFFECTS							
Age	56.4	1,35	56.4	1.01	0.32	0.03	0.16
Age*Treatment	313.6	2,35	156.8	2.81	0.07	0.14	0.52
Age*Body mass	255.8	1,35	255.8	4.59	0.04	0.12	0.55
Age*Brood size	37.7	1,35	37.7	0.68	0.42	0.02	0.13
Age*Hatching date	0.25	1,35	0.25	0.00	0.95	0.00	0.05

Figure captions

Figure 1. Number of *Carnus hemapterus* flies per nest before (day 3 post hatching) and after (day 5) manipulation of distance to cavity entrance in control, enlarged and elevated Spotless Starling nests.

Figure 2. Difference in number of *Carnus hemapterus* flies per nest (number of flies at day 5 after hatching – number of flies at day 3) in relation to distance to cavity entrance after manipulation in Spotless Starling nests; Δ : elevated nests, $\square$: enlarged nests, $\circ$: control nests. Regression line is shown.

Figure 1

Figure 2