Synthesis of oxa-aza spirobicycles by intramolecular hydrogen atom transfer promoted by *N*-radicals in carbohydrate systems

Angeles Marti'n*, Ine's Pérez-Marti'n, Ernesto Sua'rez*

Instituto de Productos Naturales y Agrobiología del C.S.I.C., Carretera de La Esperanza 3, 38206 La Laguna, Tenerife, Spain

article info

Article history:

Received 24 March 2009

Received in revised form 8 May 2009

Accepted 18 May 2009

Available online 27 May 2009

Keywords:

Intramolecular hydrogen atom transfer

N-Radical

(Diacetoxyiodo)benzene

N-Phosphoramidate

N-Cyanamide

abstract

The nitrogen-centred radical generated by reaction of an *N*-phosphoramidate or *N*-cyanamide, attached to a tri- or tetramethylene tether extended from the C-1 of a carbohydrate, with (diacetoxyiodo)benzene (DIB) and iodine can undergo a regio- and stereoselective intramolecular hydrogen atom transfer (HAT) reaction to furnish four different oxa-azaspirobicyclic systems: 1-oxa-6-azaspiro[4.4]nonane, 1-oxa-6-azaspiro[4.5]decane, 6-oxa-1-azaspiro[4.5]decane and 1-oxa-7-azaspiro[5.5]undecane. A tandem 1,5- or 1,6-HAT-oxidation-nucleophilic cyclisation mechanism is proposed.

1. Introduction

Hydrogen abstraction, also referred to as hydrogen atom transfer (HAT), is one of the most common and simplest reactions of a neutral radical with an organic substrate. It is particularly useful since the functionalisation of a position considered to be unreactive under classical conditions could be possible.¹ Within their intramolecular version 1,5- and 1,6-HAT reactions promoted by carbon-centred radicals have driven a good deal of research due to their wide range of synthetic applications,² especially combining this process with a subsequent radical cyclisation to the effective construction of five and six-membered carbo- and heterocyclic compounds.³ On the contrary, few studies have been developed for the intramolecular HAT reactions generated by heteroatom-centred radicals and *N*-radicals in particular.^{1c,4} In our group we have devoted some attention to this field since in previous papers we have reported on the synthesis of different pyrrolidines and piperidines in carbohydrate and steroidal systems employing *N*-radicals.⁵ For that study, intramolecular 1,5- and 1,6-HAT reactions were promoted by aminyl radicals derived from the treatment of a suitably protected amine⁶ with a hypervalent iodine reagent in the presence of iodine. These results together with those previously obtained from the *O*-centred radical studies⁷ have provided an excellent

opportunity to demonstrate the synthetic potential of this methodology for the preparation of different heterocyclic compounds.

In a preliminary communication we have reported the synthesis of some oxa-aza spirobicycles in carbohydrate models performed by employing our 1,5- or 1,6-HAT radical protocol (Scheme 1).^{5d}

Scheme 1. Oxa-azaspirobicycles by 1,5- or 1,6-HAT reaction; PG = PO(OR¹)₂, CN; R¹ = alkyl, aryl.

In an extension to this article, the purposes of the present work were: (a) to show that the electrophilic C-1 propylidene or butylidene *N*-radical abstracts a hydrogen atom in a regio- and stereoselective manner at the *pseudoanomeric* position of a *C*-glycoside, (b) to confirm that intramolecular 1,5-HAT, through a six-

* Corresponding authors. Tel.: þ34 922 251004; fax: þ343 922 260135.

E-mail addresses: angelesmartin@ipna.csic.es (A. Marti'n), esuares@ipna.csic.es (E. Sua'rez).

membered transition state (TS) is more favourable and better yielded process than 1,6-HAT, through a more unstable seven-membered TS,^{8(c)} to explore the influence of the bulkiness of the amino-protective group employed on the hydrogen abstraction process and (d) to investigate whether electron-withdrawing group (EWG) substituents may perhaps influence the reaction course. It therefore offers the opportunity to study the consecutive HAT-oxidation–nucleophilic cyclisation sequence to synthesise four especially interesting oxa-aza spirocompound systems (spiroaminals)⁹ (Scheme 1), which are widespread substructures common to a number of natural products such as: manzamine X,¹⁰ having a 1-oxa-6-azaspiro[4.4]nonane structure A, solanum alkaloids¹¹ and azaspiracid,¹² with a 1-oxa-6-azaspiro[4.5]decane B, stemotinine and isostemotinine alkaloids,¹³ with a 6-oxa-1-azaspiro[4.5]decane structure C and sanglifehrin A,¹⁴ with a 1-oxa-7-azaspiro[5.5]undecane D.

2. Synthesis of substrates and results

Preparation of the majority of the required C-1 three methylene tethered *N*-phosphoramidate C-glycoside derivatives was accomplished following a general well-established four-step protocol starting from suitably protected carbohydrates. A Lewis acid-

Scheme 2. Synthesis of C-1 propylenephosphoramidate precursors: (a) (i) BH₃THF 1 M, THF, 0 °C to rt; (ii) NaOH 3 M, H₂O₂ (30%), 0 °C to rt, 1 h; (b) ZnN₆S₂Py, Ph₃P, DIAD, rt; (c) (i) H₂, Pd/C 10%, EtOAc; (ii) (BnO)₂POCl, TEA, CHCl₃, rt; (d) (i) MsCl, Py, 0 °C to rt; (ii) NaN₃, DMF, 80 °C, 2 h; (e) (i) LiAlH₄, Et₂O, rt; (ii) (R²O)₂POCl, TEA, CHCl₃, rt; (f) (i) CeCl₃·7H₂O, NaBH₄, MeOH, 0 °C to rt; (ii) H₂, Pd/C 10%; (iii) NaH, BnBr, DMF, 0 °C to rt, 71% from 16; (iv) ATMS, BF₃·Et₂O, CH₃CN, 0 °C to rt, 1.5 h, 93%; (v) TBSCl, imidazole, DMF, rt, 3 h, 83%; (g) (i) Bu₃SnH, AIBN, PhH, reflux; (ii) (BnO)₂POCl, TEA, CHCl₃, rt, 95%.

mediated C-glycosidation with allyltrimethylsilane afforded the oct-7-enitols and non-8-enitols, in general with high stereoselectivity,¹⁵ which upon subsequent oxidative hydroboration provided the alcohols 2, 6^{7e} and 18 in good yield, with the exception of the known alcohol 13^{7e} derived from the corresponding butenyl derivative by reductive ozonolysis (Scheme 2). Transformation of the primary unprotected alcohols into the corresponding azide derivatives by Mitsunobu azidation¹⁶ or conversion into the corresponding mesyl product and subsequent nucleophilic substitution with azide ion gave products 3, 7, 14 and 19. Azides 3, 7 and 14 were hydrogenated to the consequent amines whereas 19 was submitted to a radical reduction with *n*-tributyltin hydride to avoid the 5-*O*-benzyl deprotection. On the other hand, the known cyanide derivatives 9^{5b} and 11^{5b,17} were reduced to the respective amine products with LiAlH₄ treatment. The resulting crude free amines were all treated with the corresponding diethyl, diphenyl, or dibenzylchlorophosphate reagent in the presence of TEA to give the required phosphoramidates 4, 8, 10, 12, 15 and 20. It is noteworthy that the choice of phosphonates as protective groups was made regarding their certain application in the amino and amino acid chemistry,¹⁸ to avoid oxidation of the amine group during the formation of the iodoamide intermediate and, at the same time, to control the stability of the *N*-radical during the HAT reaction, but it merits mention that probably the use of other functionalities such as carbamoyl or amide derivatives could be more practical regarding further manipulation at the nitrogen centre because of its facile deprotection. However, in preceding works from this laboratory it was observed that in some cases apparently a lower nucleophilicity of the carbamate group was responsible for the competitive intermolecular attack of the acetate anion coming from the reagent preventing the cyclisation step.^{5a}

To further test the scope of the HAT reaction, we also prepared a number of precursors whose phosphoramidyl groups were attached to a four-carbon tether extended from the C-1 of the sugar (Scheme 3). The alcohol 6 was submitted to iodination and cyanation to obtain 22 in 77% yield. Treatment with LiAlH₄ gave the

Scheme 3. Synthesis of C-1 butylenephosphoramidate precursors: (a) I₂, Ph₃P, imidazole, PhH, reflux, 0.5 h, 75%; (b) NaCN, DMF, rt, 14 h, 99%; (c) (i) LiAlH₄, Et₂O, rt; (ii) TEA, (BnO)₂POCl, CHCl₃, rt; (d) (i) LiAlH₄, Et₂O, rt; (ii) NaOCN, EtOH, H₂O, AcOH, reflux, 1.5 h; (iii) MsCl, Py, 0 °C to rt, 41%; (e) ZnN₆S₂Py, Ph₃P, DIAD, rt; (f) (i) H₂, Pd/C 10%, EtOAc, 23 h; (ii) TEA, (BnO)₂POCl, CHCl₃, rt.

corresponding amine, which was protected without further purification as *N*-phosphoramidate 23 and *N*-cyanamide 24, the latter with minor steric hindrance. The phosphoramidates 27 and 30 were prepared accordingly by azidation–reduction–protection starting from the known alcohols 25^{7e} and 28,^{7e} respectively.

The HAT reactions were performed under the oxidative conditions previously developed in our group, by treatment of the corresponding precursors with (diacetoxyiodo)benzene and iodine, and presented in Tables 1 and 2 for the production of different oxazaspirocompound structures.⁵ Firstly we carried out the synthesis of 1-oxa-6-azaspiro[4.4]nonane and 6-oxa-1-azaspiro[4.5]decane models (substructures A and C, Scheme 1) to verify the feasibility of this methodology, the results being summarised in Table 1.

The 1,5-HAT reaction of the phenyl and benzyl phosphoramidate furanose derivatives 4 and 8, derived from D-ribose and D-mannose, respectively, proceeded smoothly to give the expected spirobicyclic products 31 and 32 in good yields (Table 1, entries 1 and 2). Interestingly, the abstraction, in the case of 32, occurred on the sterically crowded b-side of the furanose ring while the subsequent cyclisation step tentatively took place in a selective manner on the less hindered a-side. In both cases, no NOE correlations were observed between any proton of the furanose ring and 3-H₂ of the pyrrolidine ring and thus the stereochemistry of the quaternary centre was cautiously assigned as indicated.

In order to synthesize a derivative with the spirocentre at C-5, phosphoramidate 15, derived from D-glucose, was prepared in good yield (entry 3). The HAT reaction gave the two corresponding spirocompounds 33 and 34 in a global yield of 84% and a proportion of 1.5:1, respectively. Assignment of the C-5 stereochemistry was made by contrasting the NOE correlation observed in both isomers: epimer 33 showed an NOE interaction between 1-H and the benzylic protons of the phosphoramidyl group while in 34 the interaction was observed between 1-H and 6-H.

The HAT reaction of the diethyl phosphoramidate 10 and dibenzyl derivative 20 gave the oxa-azaspirobicycles 35 and 36 in moderate yields (entries 4 and 5). The stereochemistry of the spirocentre was established on the basis of an NOE correlation between 5-H and the protons of the ethoxyl of the phosphoramidate group in 35, and an NOE interaction of 3-H with a benzylic hydrogen of the 5-OBn group in 36. These results indicate that, analogously to the case of furanose 32 (entry 2), in these two pyranose models the abstraction occurs on the b-side of the molecules while the nucleophilic cyclisation step takes place on the less hindered a-side.

The influence of the protective group at C-5, next to the supposed reacting centre C-4, was subsequently studied in entry 6. The substitution of a methyl ether by an EWG such as an acetyl group inhibited completely the radical abstraction at C-4 and under these conditions the starting material was recovered unchanged.^{5b,7a,c,19}

We next tested the 1,6-HAT reaction to assess whether this methodology could prove practical for the synthesis of 1-oxa-6-azaspiro[4.5]decane and 1-oxa-7-azaspiro[5.5]undecane bicycles (substructures B and D, Scheme 1). The results obtained are summarised in Table 2.

We started our study with phosphoramidate 23 but unfortunately it failed to undergo 1,6-HAT on the sterically crowded b-side of the molecule (Table 2, entry 1). A more unstable seven-membered TS in comparison with precursor 8 (Table 1, entry 2) was postulated as responsible for the failure. In this case only monoiodine 37 and diiodine 38, which are both produced by 1,5-HAT reactions, were isolated in low yields. Their structures were confirmed spectroscopically and by reductive dehalogenation to the starting amide 23 with the *n*-Bu₃SnH/AIBN system. In contrast, the bicycle 39 could be formed in low yield using the 6*S*-isomeric phosphoramidate 27 where the 1,6-HAT reaction proceeded on the less hindered side of the substrate (entry 2). Although the hydrogen abstraction occurred

Table 1
Synthesis of 1-oxa-6-azaspiro[4.4]nonane and 6-oxa-1-azaspiro[4.5]decane bicycles by 1,5-HAT^a

Entry	Substrate	Products	Yield (%)
1			64
2			74 ^b
3			84 (1.5:1)
4			52 ^c
5			47
6			NR

^a All reactions were performed in dry CH₂Cl₂ (80 mL) containing (diacetoxyiodo)benzene (2.5 mmol), and iodine (1.4 mmol) per mmol of substrate and irradiating with two 80 W tungsten filament lamps at room temperature. Yields of pure products isolated after silica gel chromatography.

^b Solid NaHCO₃ (25 mmol %) was added.

^c Acetonitrile (80 mL) was used instead of CH₂Cl₂.

with an overall yield of 47%, the formation of the acetyl side product 40 in a substantial amount (29%) decreased the spirocycle 39 yield (Scheme 4). This product 40 might arise from external nucleophilic attack of the acetate anion, coming from the DIB, to the oxycarbenium ion intermediate in competition with the intramolecular cyclisation reaction. The stereochemistry of the quaternary centre in the acetyl derivative could not be determined on the basis of NOE experiments because no clear interactions were observed. The proposed stereochemistry, where the acetate anion adds to the less hindered side of the molecule, was consistent with the deshielding

Table 2
Synthesis of 1-oxa-6-azaspiro[4.5]decane and 1-oxa-7-azaspiro[5.5]undecane bi-cycles by 1,6-HAT^a

Entry	Substrate	Products	Yield % (dr)
1		 	14 ^b 30
2			18 ^{b,c}
3			48 ^b
4			41 ^d (3:1)

^a All reactions were performed in dry CH₂Cl₂ (80 mL) containing (diacetoxyiodo)benzene (2.5 mmol), and iodine (1.4 mmol) per mmol of substrate and irradiating with two 80 W tungsten filament lamps at room temperature. Yields of pure products isolated after silica gel chromatography.

^b Solid NaHCO₃ (25 mmol %) was added.

^c Acetate 40 (29%) was also obtained.

^d Compound 44 (22%) was also obtained.

Scheme 4. Side product 40 from Table 2, entry 2; R¹PO(OBn)₂.

undergone by the furanose ring protons in its ¹H NMR spectrum according to similar products found in the literature.²⁰ The strong hindrance of the b-side of the molecule may also be responsible for the formation of only one isomer of products 39 and 40.

The model described in entry 3 demonstrates that the steric demand of the amine protector proved to be crucial to the 1,6-HAT and the subsequent cyclisation step. The cyanamide 24 is transformed exclusively into the 1-oxa-6-azaspiro[4.5]decane 41 with absolute regio- and stereoselectivity, albeit in moderate yield. With the voluminous phosphoramidate group only products 37 and 38 derived from more stable 1,6-transition states were formed (entry 1).²¹ Also in this model the hydrogen abstraction took place on the more congested b-side of the furanose ring while the cyclisation step occurred on the opposite a-side. The configuration of the spirocentre was tentatively established as indicated because no NOE interactions were observed between 7-H₂ and any of the furanose protons.

Finally, this HAT-cyclisation sequence was further extended to pyranose model 30 derived from D-glucose yielding smoothly the expected bicycles 42 and 43 in moderate yield together with b-iodo ester 44 as a side product. The stereochemistry of the quaternary centre was established on the basis of an NOE correlation between 1-H and 6-H in 43, an interaction that was not observed in 42. The formation of 44 can be explained by a trans electrophilic addition of acetyl hypoiodite²² to the hypothetical Z-olefin 45, resembling Pre'vost reaction (Scheme 5).²³ Plausibly, this olefin could be generated from the oxycarbenium ion intermediate or by acid-catalysed ring opening of the oxa-azaspirobicycles 42 and 43. The stereochemistry of 44 was assigned taking into account the strong deshielding of the axial protons at C-1 and C-3 in comparison with those of the starting amine 30, which suggest an axial acetyl group at C-5.

Scheme 5. Minor product 44 from Table 2, entry 4; R¹PO(OBn)₂.

It is important to stand out that all the structures were confirmed by ¹H and ¹³C NMR spectra (DEPT, COSY, HSQC and HMBC experiments), revealing in all cases a selective abstraction of the corresponding protons. Likewise, the stereochemistry of the quaternary centre was tentatively assigned in base of intramolecular NOE experiments as it is portrayed in Experimental section in each case.

3. Conclusions

With these examples we have now demonstrated the possibility of using N-radicals, generated in situ from the reaction of N-phosphoramidates and N-cyanamides with the DIB/I₂ system, for the synthesis of four different types of oxa-aza spirobicycles, which are not readily accessible by other methods. The virtues of this reagent are the mildness of the conditions and high protective group tolerance. As observed, the process may be conceptually considered to be an intramolecular N-glycosidation via selective oxidation of the pseudoanomeric position of a C-glycoside using a 1,5- or 1,6-HAT reaction as the key step. The polarity of the amine protecting group may be very important since the nitrogen acts with an umpolung reactivity during the reaction, first as an electrophilic N-radical and then as a nucleophile in the cyclisation step.

4. Experimental

4.1. General

Melting points were determined with a hot-stage apparatus and are uncorrected. Optical rotations were measured at the sodium line at ambient temperature in CHCl₃ solutions. IR spectra were recorded in CCl₄ unless otherwise stated. NMR spectra were determined at 400 MHz for ¹H and 100.6 MHz for ¹³C in CDCl₃ unless otherwise indicated, in the presence of TMS as internal standard. Mass spectra were determined at 70 eV. Merck silica gel 60 PF (0.063–0.2 mm) was used for column chromatography. Circular layers of 1 mm of Merck silica gel 60 PF₂₅₄ were used on a Chromatotron for centrifugally assisted chromatography. Commercially available reagents and solvents were of analytical grade or were purified by standard procedures prior to use. All reactions involving air- or moisture-sensitive materials were carried out under a nitrogen atmosphere. The spray reagents for TLC analysis were conducted with 0.5% vanillin in H₂SO₄/EtOH (4:1) and further heating until development of colour.

4.2. Synthesis of precursors for HAT reactions

4.2.1. General procedure for *N*-phosphoramidate precursors

Starting from azide precursors (1 mmol) in dry EtOAc (29 mL) and with Pd/C 10% (187 mg), the mixture was hydrogenated at rt for 24 h. The suspension was filtered over Celite and concentrated under reduced pressure. Starting from cyanide precursors (1 mmol) in dry THF or Et₂O (6 mL), LiAlH₄ (4 mmol) was portionwise added at 0 °C. The mixture was stirred at rt until completion and then a saturated solution of Na₂SO₄ was dropwise added until the grey mixture turned to white. It was filtered and evaporated. The resulting crude amine (1 mmol) in dry CHCl₃ (16 mL) was treated at 0 °C with triethylamine (3.5 mmol) and (RO)₂POCl (2 mmol) and stirred at rt. Then it was concentrated and poured directly into the column to chromatography (hexanes/EtOAc).

4.2.2. 4,7-Anhydro-1,2,3-trideoxy-1-[[diphenoxyphosphoryl]amino]-5,6,8-tri-*O*-methyl-*D*-altro-octitol (4)

Following the general procedure of phosphoramido protection starting from azide **3** (374 mg, 1.44 mmol), diphenylphosphoramidate **4** (342 mg, 0.74 mmol, 51%) was obtained as a colourless oil: [α]_D²⁰ +16.1 (c, 1.30); IR 3271, 2937, 1591, 1487 cm^{−1}; ¹H NMR d_H 1.48–1.67 (4H, m), 1.93 (1H, m), 3.07–3.13 (2H, m), 3.37 (3H, s), 3.41 (3H, s), 3.42 (1H, dd, *J*_H 10.9, 3.8 Hz), 3.45 (3H, s), 3.50 (1H, dd, *J*_H 10.9, 3.3 Hz), 3.69 (1H, dd, *J*_H 4.5, 4.5 Hz), 3.75 (1H, dd, *J*_H 6.2, 4.8 Hz), 3.91 (1H, ddd, *J*_H 8.6, 4.3, 4.3 Hz), 3.99 (1H, ddd, *J*_H 6.4, 3.8, 3.8 Hz), 7.12–7.33 (10H, m); ¹³C NMR (50.4 MHz) d_C 26.5 (CH₂), 28.1 (CH₂, *J*_C 46.1 Hz), 41.8 (CH₂), 58.5 (CH₃), 59.4 (CH₃), 59.7 (CH₃), 73.1 (CH₂), 79.2 (CH), 79.8 (CH), 80.2 (CH), 81.8 (CH), 120.2 (2×CH), 124.8 (4×CH), 129.6 (4×CH), 150.8 (2×C); MS *m/z* (rel intens) 466 (M⁺_b 1, <1), 262 (100); HRMS *m/z* calcd for C₂₃H₃₃NO₇P 466.1994, found 466.2011. Anal. Calcd for C₂₃H₃₃NO₇P: C, 59.35; H, 6.93; N, 3.01. Found: C, 59.38; H, 6.85; N, 2.92.

4.2.3. 3,6-Anhydro-9-[[bis(benzyloxy)phosphoryl]amino]-7,8,9-trideoxy-1,2:4,5-di-*O*-isopropylidene-*D*-glycero-*D*-manno-nonitol (8)

Following the general procedure of phosphoramido protection starting from azide **7** (85 mg, 0.26 mmol), dibenzylphosphoramidate **8** (85.5 mg, 0.15 mmol, 59%) was obtained as a colourless oil: [α]_D²⁰ −3.4 (c, 1.0); IR 3217, 2937, 1549, 1456 cm^{−1}; ¹H NMR d_H 1.32 (3H, s), 1.35 (1H, m), 1.36 (3H, s), 1.43 (3H, s), 1.46 (1H, m), 1.48 (3H, s), 1.58–1.65 (2H, m), 2.59 (1H, br s), 2.88 (2H, m), 3.67 (1H, dd, *J*_H 7.5, 3.7 Hz), 3.96 (1H, ddd, *J*_H 8.7, 5.0, 0 Hz), 3.98 (1H, dd, *J*_H 8.6, 4.4 Hz), 4.06 (1H, dd, *J*_H 8.6, 6.3 Hz), 4.36 (1H, ddd, *J*_H 6.6, 6.6,

4.7 Hz), 4.38 (1H, dd, *J*_H 6.2, 0 Hz), 4.68 (1H, dd, *J*_H 6.0, 3.8 Hz), 5.04 (4H, d, *J*_P 47.3 Hz), 7.31–7.37 (10H, m); ¹³C NMR d_C 24.5 (CH₃), 25.0 (CH₃), 26.0 (CH₃), 26.8 (CH₃), 27.5 (CH₂), 28.0 (CH₂), 40.9 (CH₂), 66.8 (CH₂), 67.9 (2×CH₂), 73.3 (CH), 79.8 (CH), 80.6 (CH), 83.7 (CH), 85.2 (CH), 109.0 (C), 112.5 (C), 127.7 (4×CH), 128.2 (2×CH), 128.4 (4×CH), 136.3 (2×C); MS *m/z* (rel intens) 560 (M⁺_b −1, <1), 546 (5), 91 (100); HRMS *m/z* calcd for C₂₉H₃₉NO₈P 560.2413, found 560.2404. Anal. Calcd for C₂₉H₄₀NO₈P: C, 62.02; H, 7.18; N, 2.49. Found: C, 62.22; H, 6.82; N, 2.74.

4.2.4. 2,6-Anhydro-7,8,9-trideoxy-9-[[diethoxyphosphoryl]amino]-1,3,4,5-tetra-*O*-methyl-*D*-glycero-*D*-manno-nonitol (10)

Following the general procedure starting from cyanide **9** (1.11 g, 4.07 mmol), **10** (926 mg, 2.24 mmol, 55%) was obtained as a colourless oil: [α]_D²⁰ +7.5 (c, 1.20); IR 3563, 1660, 1455 cm^{−1}; ¹H NMR d_H 1.29 (6H, t, *J*_H 7.1 Hz), 1.46–1.58 (2H, m), 1.58–1.67 (2H, m), 2.75 (1H, m), 2.92 (1H, m), 3.31 (1H, dd, *J*_H 4.8, 3.3 Hz), 3.35 (3H, s), 3.38 (3H, s), 3.39 (1H, dd, *J*_H 6.2, 3.3 Hz), 3.44 (3H, s), 3.45 (3H, s), 3.48 (1H, dd, *J*_H 7.4, 2.6 Hz), 3.52–3.58 (2H, m), 3.62 (1H, m), 3.88 (1H, ddd, *J*_H 9.0, 4.3, 4.3 Hz), 3.97–4.06 (4H, m), 1H from the NH group is missing; ¹³C NMR (50.4 MHz) d_C 16.2 (2×CH₃), 26.4 (CH₂), 28.2 (CH₂), 41.0 (CH₂), 57.5 (CH₃), 57.9 (CH₃), 59.1 (CH₃), 59.5 (CH₃), 62.1 (2×CH₂), 71.5 (CH₂), 71.8 (CH), 72.5 (CH), 76.6 (CH), 78.4 (CH), 79.1 (CH); MS *m/z* (rel intens) 414 (M⁺_b 1, 2), 88 (100); HRMS *m/z* calcd for C₁₇H₃₇NO₈P 414.2256, found 414.2281. Anal. Calcd for C₁₇H₃₆NO₈P: C, 49.38; H, 8.78; N, 3.39. Found: C, 49.44; H, 8.83; N, 3.13.

4.2.5. 2,6-Anhydro-1,3,4-tri-*O*-benzyl-7,8,9-trideoxy-9-[[diphenoxyphosphoryl]amino]-*D*-glycero-*L*-gulo-nonitol (12)

Following the general procedure starting from cyanide **11** (370 mg, 0.70 mmol), a deprotected 5-OH product was obtained (319 mg, 0.44 mmol, 63%), which was submitted to acylation with Ac₂O/Py to give **12** (60% overall yield) as a colourless oil: [α]_D²⁰ +23.1 (c, 0.13); IR 3221, 3066, 2934, 2871, 1746, 1454 cm^{−1}; ¹H NMR d_H 1.42 (1H, m), 1.53 (1H, m), 1.60–1.74 (2H, m), 1.96 (3H, s), 3.10–3.16 (2H, m), 3.32 (1H, m), 3.57–3.67 (4H, m), 3.75 (1H, dd, *J*_H 8.1, 8.1 Hz), 4.07 (1H, m), 4.47 (1H, d, *J*_H 11.0 Hz), 4.48 (1H, d, *J*_H 12.4 Hz), 4.57 (1H, d, *J*_H 12.4 Hz), 4.69 (1H, d, *J*_H 11.4 Hz), 4.73 (1H, d, *J*_H 11.0 Hz), 4.74 (1H, d, *J*_H 11.4 Hz), 5.01 (1H, dd, *J*_H 8.6, 5.2 Hz), 7.13–7.33 (25H, m); ¹³C NMR d_C 20.9 (CH₃), 22.9 (CH₂), 27.5 (CH₂), 41.4 (CH₂), 68.8 (CH₂), 71.9 (CH), 72.1 (CH), 72.8 (CH), 73.4 (CH₂), 74.5 (CH₂), 74.7 (CH₂), 77.4 (CH), 79.8 (CH), 120.2–129.7 (25×CH), 137.9 (2×C), 138.3 (C), 150.8 (2×C), 170.0 (C); MS *m/z* (rel intens) 766 (M⁺_b 1, 1), 91 (100); HRMS *m/z* calcd for C₄₄H₄₉NO₉P, 766.3144, found 766.3145. Anal. Calcd for C₄₄H₄₈NO₉P: C, 69.01; H, 6.32; N, 1.83. Found: C, 69.37; H, 6.49; N, 1.76.

4.2.6. Methyl 8-[[bis(benzyloxy)phosphoryl]amino]-6,7,8-trideoxy-2,3,4-tri-*O*-methyl-*D*-gluco-octopyranoside (15)

Following the general procedure of phosphoramido protection starting from azide **14** (101.5 mg, 0.33 mmol), dibenzylphosphoramidate **15** (66.7 mg, 0.12 mmol, 38%) was obtained as a colourless oil: [α]_D²⁰ −5.3 (c, 0.94); IR 3217, 2933, 1456, 1234 cm^{−1}; ¹H NMR d_H 1.40 (1H, dddd, *J*_H 13.8, 9.0, 9.0, 5.0 Hz), 1.50 (1H, m), 1.66 (1H, m), 1.81 (1H, m), 2.77 (1H, dd, *J*_H 9.2, 9.2 Hz), 2.87–2.93 (2H, m), 2.92 (1H, dd, *J*_H 8.5, 8.5 Hz), 3.01 (1H, ddd, *J*_H 9.3, 9.3, 2.5 Hz), 3.09 (1H, dd, *J*_H 8.9, 8.9 Hz), 3.43 (3H, s), 3.48 (3H, s), 3.54 (3H, s), 3.59 (3H, s), 4.04 (1H, d, *J*_H 7.5 Hz), 5.03 (4H, d, *J*_P 47.0 Hz), 7.30–7.35 (10H, m), 1H from the NH group is missing; ¹³C NMR d_C 27.8 (CH₂), 28.4 (CH₂), 41.2 (CH₂), 56.8 (CH₃), 60.4 (CH₃), 60.6 (CH₃), 60.7 (CH₃), 67.9 (2×CH₂), 74.3 (CH), 83.6 (CH), 83.8 (CH), 86.5 (CH), 104.1 (CH), 127.7 (4×CH), 128.2 (2×CH), 128.5 (4×CH), 136.5 (2×C); MS (FAB) *m/z* (rel intens) 546 (M⁺_b Na, 3), 524 (1), 91 (100). Anal. Calcd for C₂₆H₃₈NO₈P: C, 59.64; H, 7.32; N, 2.68. Found: C, 59.52; H, 7.60; N, 2.54.

4.2.7. 2,6-Anhydro-5-*O*-benzyl-9-[[bis(benzyloxy)phosphoryl]amino]-1-*O*-[tert-butyl(dimethyl)silyl]-3,4,7,8,9-pentadeoxy-*D*-arabino-nonitol (20)

To a solution of the azide 19 (39.4 mg, 0.09 mmol) in dry benzene (3 mL), Bu₃SnH (126 mL, 0.468 mmol) and AIBN (2 mg) were added and the mixture was refluxed for 1.5 h. Then it was evaporated and the obtained crude was submitted to the general protection procedure to give dibenzylphosphoramidate 20 (58.3 mg, 0.09 mmol, 95%) as a colourless oil and as a mixture of two conformers in equilibrium A/B (1:3):²⁴ [α]_D 14.2 (c, 0.39); IR 3220, 2929, 2857, 1455, 1252 cm⁻¹; ¹H NMR d_H 0.04 (12H, s), 0.87 (18H, s), 1.26–1.82 (16H, m), 2.66 (2H, br s), 2.88–2.90 (4H, m), 3.08–3.14 (2H, m), 3.45–3.49 (2H, m), 3.54–3.70 (6H, m), 4.34 (1H, d, *J*/*J*_{12.3} Hz, A), 4.46 (1H, d, *J*/*J*_{12.0} Hz, A), 4.56 (1H, d, *J*/*J*_{12.0} Hz, B), 4.63 (1H, d, *J*/*J*_{12.2} Hz, B), 5.00–5.06 (8H, m), 7.24–7.36 (30H, m); ¹³C NMR d_C—5.29 (4×CH₃), 18.2 (2×C), 23.4 (2×CH₂), 23.6 (4×CH₂), 25.7 (6×CH₃), 27.6 (2×CH₂), 27.8 (2×CH₂), 41.2 (2×CH₂), 64.3 (CH₂, B), 66.5 (CH₂, A), 67.8 (2×CH₂), 70.3 (CH₂, B), 70.5 (CH₂, A), 71.0 (CH, B), 72.1 (CH, A), 74.3 (CH, B), 75.4 (CH, A), 78.5 (CH, B), 79.2 (CH, A), 128.5–127.5 (30×CH), 136.5 (4×C), 138.5 (2×C); MS *m/z* (relintens) 654 (M⁺_B 1, <1), 596 (17); HRMS *m/z* calcd for C₃₆H₅₃NO₆PSi 654.3379, found 654.3370. Anal. Calcd for C₃₆H₅₂NO₆PSi: C, 66.13; H, 8.02; N, 2.14. Found: C, 66.13; H, 8.04; N, 2.12.

4.2.8. 3,6-Anhydro-10-[[bis(benzyloxy)phosphoryl]amino]-7,8,9,10-tetradexy-1,2:4,5-di-*O*-isopropylidene-*D*-glycero-*D*-manno-decitol (23)

Following the general procedure starting from cyanide 22 (213 mg, 0.68 mmol), 23 (228.3 mg, 0.40 mmol, 58%) was obtained as a colourless oil: [α]_D —1.75 (c, 0.286); IR 3219, 2937, 1455, 1380, 1232 cm⁻¹; ¹H NMR d_H 1.25–1.29 (2H, m), 1.32 (3H, s), 1.35–1.40 (2H, m), 1.37 (3H, s), 1.40–1.44 (2H, m), 1.42 (3H, s), 1.49 (3H, s), 2.58 (1H, m), 2.84 (2H, ddd, *J*/*J*_{16.6}, 6.9, 6.9 Hz), 3.67 (1H, dd, *J*/*J*_{7.6}, 3.4 Hz), 3.96 (1H, ddd, *J*/*J*_{9.1}, 5.3, 0 Hz), 4.00 (1H, dd, *J*/*J*_{8.5}, 4.7 Hz), 4.07 (1H, dd, *J*/*J*_{8.5}, 6.4 Hz), 4.37 (1H, ddd, *J*/*J*_{7.0}, 7.0, 4.4 Hz), 4.43 (1H, dd, *J*/*J*_{6.0}, 0 Hz), 4.72 (1H, dd, *J*/*J*_{5.9}, 3.8 Hz), 5.04 (4H, d, *J*/*J*_{8.1} Hz), 7.31–7.37 (10H, m); ¹³C NMR d_C 22.6 (CH₂), 24.5 (CH₃), 25.0 (CH₃), 26.0 (CH₃), 26.8 (CH₃), 30.0 (CH₂), 31.1 (CH₂), 41.2 (CH₂), 66.9 (CH₂), 67.8 (2×CH₂), 73.3 (CH), 79.9 (CH), 80.6 (CH), 83.8 (CH), 85.1 (CH), 109.0 (C), 112.5 (C), 127.7 (4×CH), 128.2 (2×CH), 128.4 (4×CH), 136.4 (2×C); MS *m/z* (rel intens) 575 (M⁺_B, <1), 560 (7), 91 (100); HRMS *m/z* calcd for C₃₀H₄₂NO₈P 575.2648, found 575.3074. Anal. Calcd for C₃₀H₄₂NO₈P: C, 62.60; H, 7.35; N, 2.43. Found: C, 62.44; H, 7.33; N, 2.57.

4.2.9. 3,6-Anhydro-10-cyanoamino-7,8,9,10-tetradexy-1,2:4,5-di-*O*-isopropylidene-*D*-glycero-*D*-manno-decitol (24)

The cyanide 22 (334 mg, 1.07 mmol) was submitted to LiAlH₄ reduction as in the general procedure. The resulting crude amine was dissolved in EtOH (48 mL), water (1.4 mL) and acetic acid (0.13 mL), and sodium cyanate was added (104 mg, 1.60 mmol). The mixture was refluxed for 2 h and then it was poured into brine and extracted with ethyl acetate. The organic solution was dried over Na₂SO₄ and evaporated. To a solution of the crude urea in pyridine (4.8 mL), at 0 °C, methanesulphonyl chloride (0.26 mL, 3.39 mmol) was added. After 30 min, the solution was allowed to warm to rt and then it was stirred for 2 h. It was poured into water and extracted with CHCl₃. The organic layer was washed with aqueous NaHCO₃ and water and evaporated. Column chromatography (hexanes/EtOAc, 7:3) gave the cyanamide 24 (150 mg, 0.44 mmol, 41%) as a colourless oil: [α]_D —7.2 (c, 2.50); IR 3251, 2227, 1381 cm⁻¹; ¹H NMR d_H 1.29 (3H, s), 1.32 (3H, s), 1.32–1.49 (4H, m), 1.40 (3H, s), 1.44 (3H, s), 1.52–1.64 (2H, m), 3.01 (2H, ddd, *J*/*J*_{13.5}, 6.9, 0 Hz), 3.69 (1H, dd, *J*/*J*_{7.4}, 3.7 Hz), 3.97 (2H, dd, *J*/*J*_{8.5}, 4.5 Hz), 4.04 (1H, dd, *J*/*J*_{8.5}, 6.1 Hz), 4.26 (1H, br t, *J*/*J*_{5.0} Hz), 4.33 (1H, m), 4.45 (1H, dd, *J*/*J*_{6.1}, 0 Hz), 4.71 (1H, dd, *J*/*J*_{6.1}, 3.7 Hz); ¹³C NMR d_C 22.4 (CH₂), 24.5 (CH₃), 25.0 (CH₃), 26.0 (CH₃), 26.7 (CH₃), 29.1 (CH₂),

29.9 (CH₂), 45.8 (CH₂), 66.7 (CH₂), 73.3 (CH), 79.9 (CH), 80.6 (CH), 83.8 (CH), 85.1 (CH), 109.0 (C), 112.5 (C), 116.3 (C); MS *m/z* (rel intens) 325 (M⁺_B—Me, 39), 101 (100); HRMS *m/z* calcd for C₁₆H₂₅N₂O₅ 325.1763, found 325.1768. Anal. Calcd for C₁₇H₂₈N₂O₅: C, 59.98; H, 8.29; N, 8.23. Found: C, 59.64; H, 8.31; N, 8.59.

4.2.10. 3,6-Anhydro-10-[[bis(benzyloxy)phosphoryl]amino]-7,8,9,10-tetradexy-1,2:4,5-di-*O*-isopropylidene-glycero-*D*-manno-decitol (23 and 27)

Following the general procedure starting from azide 26 (35 mg, 0.10 mmol), 23 and 27 (32.2 mg, 0.06 mmol, 55%) were obtained as a mixture 23/27 (1.4:1) and as a colourless oil: IR 3217, 2988, 2935, 2871, 1456, 1380, 1232 cm⁻¹; ¹H NMR d_H 1.25–1.29 (2H, m), 1.29 (3H, s), 1.32 (3H, s), 1.33–1.43 (6H, m), 1.34 (3H, s), 1.37 (3H, s), 1.40–1.44 (2H, m), 1.42 (6H, s), 1.43 (3H, s), 1.49 (3H, s), 1.61 (2H, m), 1.80 (2H, br s), 2.85 (4H, m), 3.36 (1H, ddd, *J*/*J*_{6.7}, 6.7, 3.8 Hz), 3.40 (1H, dd, *J*/*J*_{7.4}, 3.3 Hz), 3.67 (1H, dd, *J*/*J*_{7.6}, 3.4 Hz), 3.96 (1H, ddd, *J*/*J*_{9.1}, 5.3, 0 Hz), 4.00 (2H, dd, *J*/*J*_{8.5}, 4.7 Hz), 4.04 (1H, dd, *J*/*J*_{8.0}, 6.5 Hz), 4.07 (1H, dd, *J*/*J*_{8.5}, 6.4 Hz), 4.37 (2H, m), 4.43 (1H, dd, *J*/*J*_{6.0}, 0 Hz), 4.52 (1H, dd, *J*/*J*_{6.2}, 3.8 Hz), 4.69 (1H, dd, *J*/*J*_{6.2}, 3.3 Hz), 4.72 (1H, dd, *J*/*J*_{5.9}, 3.8 Hz), 5.04 (8H, d, *J*/*J*_{8.1} Hz), 7.31–7.37 (20H, m); ¹³C NMR d_C 22.6 (CH₂), 22.8 (CH₂), 24.5 (2×CH₃), 25.0 (CH₃), 25.1 (CH₃), 25.6 (CH₃), 26.0 (CH₃), 26.8 (2×CH₃), 27.5 (CH₂), 30.0 (CH₂), 31.1 (CH₂), 31.3 (CH₂), 41.0 (CH₂), 41.2 (CH₂), 66.8 (CH₂), 66.9 (CH₂), 67.8 (4×CH₂), 73.0 (CH), 73.3 (CH), 79.9 (CH), 80.6 (2×CH), 81.1 (CH), 81.4 (CH), 81.8 (CH), 83.8 (CH), 85.1 (CH), 108.9 (C), 109.0 (C), 112.1 (C), 112.5 (C), 127.7 (8×CH), 128.1 (2×CH), 128.2 (2×CH), 128.4 (8×CH), 136.4 (2×C), 137.7 (2×C); MS *m/z* (rel intens) 560 (M⁺_B—CH₃, 2), 484 (2), 91 (100); HRMS *m/z* calcd for C₂₉H₃₉NO₈P 560.2413, found 560.2419. Anal. Calcd for C₃₀H₄₂NO₈P: C, 62.60; H, 7.35; N, 2.43. Found: C, 62.44; H, 7.73; N, 2.57.

4.2.11. Methyl 9-[[bis(benzyloxy)phosphoryl]amino]-6,7,8,9-tetradexy-2,3,4-tri-*O*-methyl-*D*-gluco-nonopyranoside (30)

Following the general procedure starting from azide 29 (39.3 mg, 0.14 mmol), 30 (36.6 mg, 0.07 mmol, 52%) was obtained as a crystalline solid: mp 59.6–60.6 °C (*n*-hexane/EtOAc); [α]_D —2.6 (c, 1.29); IR 3216, 2934, 1456, 1233 cm⁻¹; ¹H NMR d_H 1.30 (1H, m), 1.36–1.54 (4H, m), 1.75 (1H, m), 2.78 (1H, dd, *J*/*J*_{9.2}, 9.2 Hz), 2.84–2.91 (2H, m), 2.93 (1H, dd, *J*/*J*_{8.4}, 8.4 Hz), 3.01 (1H, ddd, *J*/*J*_{9.3}, 9.3, 2.4 Hz), 3.10 (1H, dd, *J*/*J*_{8.9}, 8.9 Hz), 3.46 (3H, s), 3.50 (3H, s), 3.54 (3H, s), 3.60 (3H, s), 4.05 (1H, d, *J*/*J*_{8.0} Hz), 5.03 (4H, d, *J*/*J*_{7.5} Hz), 7.29–7.36 (10H, m), 1H from the NH group is missing; ¹³C NMR d_C 22.6 (CH₂), 31.1 (CH₂), 31.5 (CH₂), 41.3 (CH₂), 56.7 (CH₃), 60.3 (CH₃), 60.6 (CH₃), 60.7 (CH₃), 68.0 (2×CH₂), 74.4 (CH), 83.7 (CH), 83.9 (CH), 86.5 (CH), 104.0 (CH), 127.7 (4×CH), 128.2 (2×CH), 128.5 (4×CH), 136.4 (2×C); MS *m/z* (rel intens) 537 (M⁺_B, <1), 88 (100); HRMS *m/z* calcd for C₂₇H₄₀NO₈P 537.2491, found 537.2520. Anal. Calcd for C₂₇H₄₀NO₈P: C, 60.32; H, 7.50; N, 2.61. Found: C, 60.42; H, 7.49; N, 2.79.

4.3. General procedure for HAT reactions

To a solution of the *N*-phosphoramidate or *N*-cyanamide precursor (1 mmol) in dry CH₂Cl₂ (1.5 mL), DIB (2.5 mmol) and I₂ (1.5 mmol) were added and the mixture was irradiated with two 80 W tungsten lamps at rt. When the starting material was completely consumed, the mixture was poured into a solution of Na₂S₂O₃ and extracted with CH₂Cl₂. The organic extracts were dried over Na₂SO₄ and evaporated. Column chromatography of the residue (hexanes/EtOAc) gave the title compound.

4.3.1. (4*S*)-1,4:4,7-Dianhydro-1,2,3-trideoxy-1-[[diphenoxyphosphoryl]amino]-5,6,8-tri-*O*-methyl-*D*-ribo-oct-4-ulose (31)

Following the general procedure, diphenylphosphoramidate 4 (69.6 mg, 0.15 mmol) gave 31 (44.4 mg, 0.10 mmol, 64%) as a colourless oil: [α]_D —37.9 (c, 0.14); IR 2933, 1490, 1279 cm⁻¹; ¹H NMR

d_H 1.82 (1H, m), 1.90–1.99 (2H, m), 2.20 (1H, m), 3.22 (1H, m), 3.29 (1H, m), 3.29 (3H, s), 3.30 (3H, s), 3.36 (3H, s), 3.38 (1H, dd, J /4.9.8, 7.4 Hz), 3.62 (1H, m), 3.65 (1H, dd, J /4.5.7, 1.4 Hz), 3.90 (1H, ddd, J /4.7.6, 7.6, 1.4 Hz), 4.73 (1H, d, J /4.5.7 Hz), 7.13–7.32 (10H, m); NOE correlation between 3-H and 5-H was observed; ^{13}C NMR (50.4 MHz) d_C 23.0 (CH₂), 33.9 (CH₂), 48.4 (CH₂), 57.6 (CH₃), 58.4 (CH₃), 59.1 (CH₃), 73.1 (CH₂), 79.1 (CH), 79.2 (CH), 79.5 (CH), 102.9 (C), 120.3 (CH), 120.6 (CH), 124.7 (2×CH), 124.9 (2×CH), 129.3 (2×CH), 129.6 (2×CH), 150.7 (C), 150.9 (C); MS m/z (rel intens) 463 (M⁺, 1), 101 (100); HRMS m/z calcd for C₂₃H₃₀NO₇P: C, 59.61; H, 6.52; N, 3.02. Found: C, 59.69; H, 6.80; N, 3.03.

4.3.2. (4S)-1,4:4,7-Dianhydro-1-[[bis(benzyloxy)phosphoryl]amino]-1,2,3-trideoxy-5,6:8,9-di-O-isopropylidene-D-manno-non-4-ulose (32)

Following the general procedure but adding also NaHCO₃ (25 mmol %), dibenzylphosphoramidate 8 (10.7 mg, 0.02 mmol) gave 32 (5.1 mg, 0.01 mmol, 74%) as a colourless oil: $[\alpha]_D^{25} +22.0$ (c, 1.03); IR 2986, 1380, 1265, 1070 cm⁻¹; ^1H NMR d_H 1.34 (3H, s), 1.36 (3H, s), 1.44 (3H, s), 1.46 (3H, s), 1.73 (1H, m), 1.88 (1H, m), 2.00 (1H, m), 2.09 (1H, m), 3.08 (1H, ddd, J /4.8.8, 8.8, 8.8 Hz), 3.20 (1H, ddd, J /4.8.8, 8.8, 0 Hz), 3.97 (1H, dd, J /4.8.5, 4.6 Hz), 4.04 (1H, dd, J /4.8.5, 6.2 Hz), 4.33 (1H, ddd, J /4.6.4, 6.4, 5.0 Hz), 4.52 (1H, dd, J /4.7.1, 3.8 Hz), 4.96 (1H, d, J /4.1.9 Hz), 4.98 (1H, d, J /4.1.4 Hz), 5.01 (1H, m), 5.02 (2H, d, J /4.2.0 Hz), 5.14 (1H, dd, J /4.5.9, 0 Hz), 7.30–7.38 (10H, m); there was no NOE correlation between any of the hydrogen atoms on the furanose system and 3-H; ^{13}C NMR d_C 22.6 (CH₂, J /4.9.1 Hz), 24.4 (CH₃), 25.4 (CH₃), 25.9 (CH₃), 26.9 (CH₃), 36.6 (CH₂, J /4.9.1 Hz), 48.4 (CH₂), 66.6 (CH₂), 67.3 (CH₂), 67.7 (CH₂), 73.8 (CH), 79.9 (CH), 81.7 (CH), 86.5 (CH), 104.8 (C), 108.8 (C), 112.0 (C), 127.7 (2×CH), 128.1 (2×CH), 128.2 (2×CH), 128.3 (2×CH), 128.4 (2×CH), 136.2 (2×C); MS m/z (rel intens) 559 (M⁺, <1), 544 (4), 91 (100); HRMS m/z calcd for C₂₉H₃₈NO₈P: C, 62.24; H, 6.84; N, 2.50. Found: C, 62.32; H, 6.45; N, 2.68.

4.3.3. Methyl (5R)-5,8-anhydro-8-[[bis(benzyloxy)-phosphoryl]amino]-6,7,8-trideoxy-2,3,4-tri-O-methyl-b-D-xylo-octo-5-olopyranoside (33) and methyl (5S)-5,8-anhydro-8-[[bis(benzyloxy)phosphoryl]amino]-6,7,8-trideoxy-2,3,4-tri-O-methyl-b-D-xylo-octo-5-olopyranoside (34)

Following the general procedure, dibenzylphosphoramidate 15 (72.8 mg, 0.14 mmol) gave 33 (37 mg, 0.07 mmol, 51%) and 34 (16.4 mg, 0.03 mmol, 33%) as colourless oils. Compound 33: $[\alpha]_D^{25} -26.9$ (c, 0.87); IR 2952, 1273, 1090 cm⁻¹; ^1H NMR d_H 1.75 (1H, m), 1.99–2.04 (2H, m), 2.11 (1H, ddd, J /4.2.0, 7.3, 0 Hz), 3.01 (1H, dd, J /4.8.2, 8.2 Hz), 3.26 (1H, m), 3.37 (1H, d, J /4.9.3 Hz), 3.39 (3H, s), 3.43 (1H, ddd, J /4.9.6, 8.7, 0 Hz), 3.56 (3H, s), 3.59 (3H, s), 3.66 (3H, s), 4.79 (1H, dd, J /4.8.7, 8.7 Hz), 4.83 (1H, d, J /4.8.0 Hz), 5.00 (1H, dd, J /4.1.7 Hz, J /4.6.7 Hz), 5.03 (1H, dd, J /4.1.8 Hz, J /4.6.1 Hz), 5.12 (2H, d, J /4.7.0 Hz), 7.29–7.40 (10H, m); an NOE correlation between 1-H and benzylic hydrogen atoms was observed; ^{13}C NMR d_C 21.5 (CH₂, d, J /4.6.1 Hz), 42.0 (CH₂, d, J /4.9.2 Hz), 50.5 (CH₂), 56.4 (CH₃), 59.6 (CH₃), 60.2 (CH₃), 61.5 (CH₃), 67.3 (CH₂), 67.9 (CH₂), 83.3 (CH), 83.6 (CH), 84.4 (CH), 95.4 (C, d, J /4.6.1 Hz), 99.5 (CH), 127.6 (2×CH), 127.7 (2×CH), 127.9 (CH), 128.1 (CH), 128.3 (2×CH), 128.4 (2×CH), 136.6 (C, d, J /4.6.1 Hz), 137.2 (C, d, J /4.6.1 Hz); MS (FAB) m/z (rel intens) 521 (M⁺, 2), 490 (100); HRMS m/z calcd for C₂₅H₃₃NO₇P: C, 59.87; H, 6.96; N, 2.69. Found: C, 59.55; H, 7.21; N, 2.63. Compound 34: $[\alpha]_D^{25} -8.4$ (c, 2.4); IR 2935, 1274, 1094 cm⁻¹; ^1H NMR d_H 1.81–1.90 (3H, m), 2.14 (1H, ddd, J /4.2.2, 8.7, 7.3 Hz), 3.07 (1H, dd, J /4.9.4, 9.4 Hz), 3.14 (1H, dd, J /4.8.5, 8.5 Hz), 3.16 (1H, m), 3.38 (1H, m), 3.41 (3H, s), 3.57 (3H, s), 3.59 (3H, s), 3.63 (3H, s), 4.20 (1H, d, J /4.8.0 Hz), 4.36 (1H, d, J /4.9.8 Hz), 5.03 (1H, dd, J /4.1.9 Hz, J /4.6.9 Hz), 5.07 (1H, dd,

J /4.1.9 Hz, J /4.6.6 Hz), 5.09 (1H, dd, J /4.2.2 Hz, J /4.6.2 Hz), 5.12 (1H, dd, J /4.2.2 Hz, J /4.7.2 Hz), 7.27–7.38 (10H, m); an NOE correlation between 1-H and 6-H was observed; ^{13}C NMR d_C 23.3 (CH₂, d, J /4.9.2 Hz), 30.5 (CH₂, d, J /4.9.2 Hz), 47.9 (CH₂), 56.3 (CH₃), 60.3 (2×CH₃), 60.7 (CH₃), 67.3 (CH₂), 67.4 (CH₂), 80.4 (CH), 83.0 (2×CH), 84.5 (CH), 95.8 (C), 100.7 (CH), 127.5 (2×CH), 127.7 (2×CH), 127.9 (2×CH), 128.1 (CH), 128.3 (2×CH), 136.5 (C, d, J /4.9.2 Hz), 137.1 (C, d, J /4.9.2 Hz); MS m/z (rel intens) 490 (M⁺—OCH₃, 1), 88 (100); HRMS m/z calcd for C₂₅H₃₃NO₇P: C, 59.87; H, 6.96; N, 2.69. Found: C, 59.98; H, 6.67; N, 2.77.

4.3.4. (4S)-1,4:4,8-Dianhydro-1,2,3-trideoxy-1-[[diethoxyphosphoryl]amino]-5,6,7,9-tetra-O-methyl-D-manno-non-4-ulose (35)

Following the general procedure but using acetonitrile as solvent, phosphoramidate 10 (30 mg, 0.07 mmol) gave 35 (15.5 mg, 0.04 mmol, 52%) as a colourless oil: $[\alpha]_D^{25} +44.5$ (c, 1.10); IR 2933, 1715, 1455, 1252 cm⁻¹; ^1H NMR d_H 1.29 (6H, t, J /6.9 Hz), 1.72 (1H, m), 1.85 (1H, m), 1.98 (1H, m), 2.19 (1H, ddd, J /4.1.3.2, 9.0, 9.0 Hz), 3.14–3.18 (2H, m), 3.23 (1H, dd, J /4.9.0, 3.5 Hz), 3.33 (3H, s), 3.41 (3H, s), 3.44–3.46 (2H, m), 3.50 (3H, s), 3.51 (3H, s), 3.83 (1H, dd, J /4.3.5, 3.5 Hz), 3.97–4.10 (5H, m), 4.57 (1H, d, J /4.3.5 Hz); NOE correlations were observed between 5-H and PO(OCH₂CH₃)₂ and between 3-H and the C-5-OMe; ^{13}C NMR (50.4 MHz) d_C 16.1 (2×CH₃), 22.4 (CH₂, J /4.6.1 Hz), 37.1 (CH₂, J /4.9.1 Hz), 47.5 (CH₂), 58.0 (2×CH₃), 59.0 (CH₃), 59.2 (CH₃), 61.9 (2×CH₂), 71.3 (CH), 73.0 (CH₂), 77.8 (CH), 78.6 (CH), 79.4 (CH), 96.9 (C, J /4.9.1 Hz); MS m/z (rel intens) 411 (M⁺, <1), 366 (3), 88 (100); HRMS m/z calcd for C₁₇H₃₄NO₈P: C, 49.63; H, 8.33; N, 3.41. Found: C, 49.33; H, 8.70; N, 3.46.

4.3.5. (4S)-1,4:4,8-Dianhydro-1-[[bis(benzyloxy)phosphoryl]amino]-9-O-[tert-butyl(dimethyl)silyl]-1,2,3,6,7-pentadeoxy-D-threo-non-4-ulose (36)

Following the general procedure, phosphoramidate 20 (16.2 mg, 0.02 mmol) gave 36 (7.6 mg, 0.01 mmol, 47%) as a colourless oil: $[\alpha]_D^{25} +13.5$ (c, 1.0); IR 2929, 2857, 1254, 1101 cm⁻¹; ^1H NMR d_H 0.01 (6H, s), 0.87 (9H, s), 1.65–1.86 (4H, m), 1.86–1.95 (2H, m), 2.21–2.33 (2H, m), 3.23 (1H, m), 3.35 (1H, m), 3.54 (1H, dd, J /4.1.0.5, 5.9 Hz), 3.64 (1H, dd, J /4.1.0.4, 4.8 Hz), 4.23 (1H, m), 4.65 (1H, d, J /4.1.1 Hz), 4.67 (1H, dd, J /4.9.4, 5.2 Hz), 4.73 (1H, d, J /4.1.8 Hz), 4.98 (1H, dd, J /4.1.8 Hz, J /4.6.9 Hz), 5.03 (1H, dd, J /4.1.8 Hz, J /4.6.2 Hz), 5.13 (2H, d, J /4.6.9 Hz), 7.26–7.45 (15H, m); no NOE correlation was observed between 3-H and 5-H; ^{13}C NMR d_C —5.3 (2×CH₃), 18.2 (C), 23.1 (CH₂), 23.3 (CH₂), 23.6 (CH₂), 25.8 (3×CH₃), 35.0 (CH₂), 48.2 (CH₂), 65.9 (CH₂), 67.2 (2×CH₂), 71.2 (CH), 71.7 (CH₂), 75.1 (CH), 98.0 (C), 127.2–128.3 (15×CH), 136.5 (C), 136.9 (C), 139.0 (C); MS m/z (rel intens) 594 (M⁺—^tBu, 6), 91 (100); HRMS m/z calcd for C₃₂H₄₁NO₆PSi: C, 66.33; H, 7.73; N, 2.15. Found: C, 66.00; H, 8.12; N, 2.17.

4.3.6. 5,8-Anhydro-1-[[bis(benzyloxy)phosphoryl]amino]-1,2,3,4-tetradeoxy-4-iodo-6,7:9,10-di-O-isopropylidene-D-erythro-L-altro-decitol (37) and 3,6-anhydro-10-[[bis(benzyloxy)phosphoryl]amino]-7,8,9,10-tetradeoxy-7,7-diiodo-1,2:4,5-di-O-isopropylidene-D-glycero-D-manno-decitol (38)

Following the general procedure but adding also NaHCO₃ (25 mmol %), dibenzylphosphoramidate 23 (204 mg, 0.35 mmol) gave the diastereomeric mixture of monoiodides 37 (34.6 mg, 0.05 mmol, 14%), and diiodide 38 (80.4 mg, 0.10 mmol, 30%) as colourless oils. Compound 37: IR 3216, 2936, 1745, 1252 cm⁻¹; complex ^1H and ^{13}C NMR spectra; MS m/z (rel intens) 686 (M⁺—CH₃, 1), 91 (100); HRMS m/z calcd for C₂₉H₃₈INO₈P: C, 51.36; H, 5.89; N, 2.00. Found: C, 51.25; H, 5.88; N, 1.89. Compound 38: $[\alpha]_D^{25} -2.6$ (c,

0.154); IR 3209, 2989, 1381, 1257 cm^{-1} ; ^1H NMR d_H 1.36 (3H, s), 1.41 (3H, s), 1.43 (3H, s), 1.53 (3H, s), 1.81 (2H, dq, $J/47.3, 7.3, 7.3, 7.3$ Hz), 2.07 (1H, m), 2.36 (1H, m), 2.79 (1H, m), 2.96 (2H, dq, $J/410.2, 6.8, 6.8, 6.8$ Hz), 3.82 (1H, d, $J/42.1$ Hz), 3.99 (1H, dd, $J/48.7, 4.9$ Hz), 4.05 (1H, dd, $J/48.7, 6.1$ Hz), 4.35 (1H, ddd, $J/46.3, 6.3, 5.1$ Hz), 4.77 (1H, dd, $J/47.6, 4.2$ Hz), 4.87 (1H, dd, $J/45.9, 2.5$ Hz), 4.94 (1H, dd, $J/45.9, 3.8$ Hz), 5.05 (4H, d, $J_p/47.6$ Hz), 7.31–7.38 (10H, m); ^{13}C NMR d_C 23.2 (C), 24.9 (CH₃), 25.2 (CH₃), 26.5 (CH₃), 26.8 (CH₃), 33.8 (CH₂), 40.0 (CH₂), 48.1 (CH₂), 66.6 (CH₂), 68.0 (2 $\times$ CH₂), 73.7 (CH), 81.0 (CH), 84.1 (CH), 88.7 (CH), 95.0 (CH), 109.2 (C), 113.4 (C), 127.7 (4 $\times$ CH), 128.3 (2 $\times$ CH), 128.5 (4 $\times$ CH), 136.3 (2 $\times$ C); MS m/z (rel intens) 812 ($\text{M}^{\text{b}}-\text{CH}_3$, 1), 573 (1), 91 (100); HRMS m/z calcd for $\text{C}_{29}\text{H}_{37}\text{I}_2\text{NO}_8\text{P}$, 812.0346, found 812.0307. Anal. Calcd for $\text{C}_{30}\text{H}_{40}\text{I}_2\text{NO}_8\text{P}$: C, 43.55; H, 4.87; N, 1.69. Found: C, 43.59; H, 4.65; N, 1.81.

4.3.7. (5S)-1,5:5,8-Dianhydro-1-[[bis(benzyloxy)phosphoryl]-amino]-1,2,3,4-tetra-deoxy-6,7:9,10-di-O-isopropylidene-D-manno-dec-5-ulose (39) and 5-O-acetyl-1-[[bis(benzyloxy)phosphoryl]-amino]-1,2,3,4-tetra-deoxy-6,7:9,10-di-O-isopropylidene-D-manno-dec-5-ulofuranose (40)

Starting from a mixture of phosphoramidates (23/27% 1.4:1) (98 mg, 0.17 mmol) and following the general procedure adding solid NaHCO_3 (25 mmol%) also, the mixture of diastereomers monoiido 37 (13.6 mg, 0.02 mmol, 25%), diiodo 38 (5.4 mg, 0.01 mmol, 7%), the spirocompound 39 (7.5 mg, 0.01 mmol, 18%) and the acetate 40 (13.2 mg, 0.02 mmol, 29%) were obtained as colourless oils. Compound 39: $[\alpha]_\text{D} -5.2$ (c, 0.65); IR 2926, 2856, 1730, 1456, 1380, 1260 cm^{-1} ; ^1H NMR d_H 1.21–1.75 (4H, m), 1.24 (3H, s), 1.35 (6H, s), 1.44 (3H, s), 1.98–2.07 (2H, m), 3.14 (1H, m), 3.49 (1H, m), 3.86 (1H, dd, $J/47.9, 3.6$ Hz), 3.97 (1H, dd, $J/48.6, 4.3$ Hz), 4.06 (1H, dd, $J/48.6, 6.2$ Hz), 4.33 (1H, ddd, $J/46.9, 6.9, 5.0$ Hz), 4.57 (1H, dd, $J/45.7, 3.8$ Hz), 4.94–5.08 (4H, m), 5.32 (1H, d, $J/45.7$ Hz), 7.32–7.38 (10H, m); no NOE correlation was observed between 4-H and any of the furanose protons; ^{13}C NMR d_C 20.8 (CH₂), 24.4 (CH₃), 25.3 (CH₃), 25.8 (CH₃), 26.9 (CH₃), 29.7 (CH₂), 30.8 (CH₂), 43.1 (CH₂), 67.0 (CH₂), 68.2 (CH₂), 68.3 (CH₂), 73.2 (CH), 78.9 (CH), 80.5 (CH), 83.9 (CH), 95.9 (C), 109.0 (C), 111.9 (C), 127.9 (4 $\times$ CH), 128.3 (2 $\times$ CH), 128.6 (4 $\times$ CH), 136.0 (2 $\times$ C); MS m/z (rel intens) 573 (M^{b} , <1), 558 (2), 91 (100); HRMS m/z calcd for $\text{C}_{30}\text{H}_{40}\text{NO}_8\text{P}$, 573.2491, found 573.2601. Anal. Calcd for $\text{C}_{30}\text{H}_{40}\text{NO}_8\text{P}$: C, 62.81; H, 7.03; N, 2.44. Found: C, 62.66; H, 7.42; N, 2.14. Compound 40: $[\alpha]_\text{D} +13.6$ (c, 0.81); IR 3220, 2937, 1738, 1380, 1252 cm^{-1} ; ^1H NMR d_H 1.31–1.48 (4H, m), 1.32 (3H, s), 1.37 (3H, s), 1.44 (3H, s), 1.46 (3H, s), 2.00 (3H, s), 2.07 (2H, ddd, $J/49.6, 6.8, 6.8$ Hz), 2.70 (1H, m), 2.87–2.91 (2H, m), 3.99 (1H, dd, $J/48.7, 4.5$ Hz), 4.01 (1H, dd, $J/47.0, 4.0$ Hz), 4.07 (1H, dd, $J/48.9, 6.6$ Hz), 4.36 (1H, ddd, $J/47.3, 6.6, 4.7$ Hz), 4.81 (1H, d, $J/46.1$ Hz), 4.88 (1H, dd, $J/45.6, 3.7$ Hz), 5.02 (2H, d, $J_p/47.4$ Hz), 5.04 (2H, d, $J_p/47.4$ Hz), 7.31–7.36 (10H, m); no NOE correlations were observed between any of the furanose protons and 4-H nor -OAc hydrogen atoms; ^{13}C NMR d_C 20.1 (CH₂), 21.8 (CH₃), 24.4 (CH₃), 25.1 (CH₃), 25.8 (CH₃), 26.9 (CH₃), 31.0 (CH₂), 31.2 (CH₂, d, $J_p/46.1$ Hz), 40.9 (CH₂), 66.8 (CH₂), 67.9 (2 $\times$ CH₂), 73.0 (CH), 79.5 (CH), 81.3 (CH), 84.7 (CH), 109.2 (C), 113.0 (C), 113.6 (C), 127.7 (4 $\times$ CH), 128.2 (2 $\times$ CH), 128.5 (4 $\times$ CH), 136.4 (2 $\times$ C), 169.5 (C); MS (FAB) m/z (rel intens) 656 ($\text{M}^{\text{b}}\text{Na}$, 16), 574 (100); HRMS m/z calcd for $\text{C}_{32}\text{H}_{44}\text{NNaO}_{10}\text{P}$, 656.2600, found 656.2568. Anal. Calcd for $\text{C}_{32}\text{H}_{44}\text{NO}_{10}\text{P}$: C, 60.65; H, 7.00; N, 2.21. Found: C, 60.47; H, 7.43; N, 2.11.

4.3.8. (5S)-1,5:5,8-Dianhydro-1-cyanoamino-1,2,3,4-tetra-deoxy-6,7:9,10-di-O-isopropylidene-D-manno-dec-5-ulose (41)

Following the general procedure starting from *N*-cyanamide 24 (69.2 mg, 0.20 mmol) but adding solid NaHCO_3 (25 mmol %) also, oxa-azaspirocompound 41 (33 mg, 0.10 mmol, 48%) was obtained as a colourless oil: $[\alpha]_\text{D} -15.2$ (c, 3.1); IR 2990, 2210, 1372 cm^{-1} ; ^1H NMR d_H 1.34 (3H, s), 1.38 (3H, s), 1.46 (3H, s), 1.48 (3H, s), 1.54–

1.63 (2H, m), 1.64–1.78 (4H, m), 3.19 (1H, m), 3.37 (1H, m), 4.01 (1H, dd, $J/48.8, 4.5$ Hz), 4.04 (1H, dd, $J/47.7, 3.7$ Hz), 4.10 (1H, dd, $J/48.8, 6.4$ Hz), 4.37 (1H, m), 4.87 (1H, d, $J/45.8$ Hz), 4.91 (1H, dd, $J/45.8, 3.4$ Hz); no NOE correlation was observed between 4-H and any of the protons of the sugar ring; ^{13}C NMR d_C 19.7 (CH₂), 23.7 (CH₂), 24.4 (CH₃), 25.3 (CH₃), 25.7 (CH₃), 26.8 (CH₃), 29.5 (CH₂), 47.2 (CH₂), 66.9 (CH₂), 72.9 (CH), 80.1 (CH), 80.2 (CH), 84.0 (CH), 95.7 (C), 109.4 (C), 113.2 (C), 116.0 (C); MS m/z (rel intens) 323 ($\text{M}^{\text{b}}-\text{CH}_3$, 10), 101 (100); HRMS m/z calcd for $\text{C}_{16}\text{H}_{23}\text{N}_2\text{O}_5$, 323.1607, found 323.1601. Anal. Calcd for $\text{C}_{17}\text{H}_{26}\text{N}_2\text{O}_5$: C, 60.34; H, 7.74; N, 8.28. Found: C, 60.04; H, 7.76; N, 8.59.

4.3.9. Methyl (5R)-5,9-anhydro-9-[[bis(benzyloxy)phosphoryl]-amino]-6,7,8,9-tetra-deoxy-2,3,4-tri-O-methyl-D-xylo-non-5-ulopyranoside (42), methyl (5S)-5,9-anhydro-9-[[bis(benzyloxy)phosphoryl]amino]-6,7,8,9-tetra-deoxy-2,3,4-tri-O-methyl-D-xylo-non-5-ulopyranoside (43) and methyl (5S)-5-O-acetyl-9-[[bis(benzyloxy)phosphoryl]amino]-6,7,8,9-tetra-deoxy-2,3,4-tri-O-methyl-6-iodo-D-xylo-non-5-ulopyranoside (44)

Following the general procedure starting from 30 (135 mg, 0.25 mmol), a mixture of spirocompound 42 (41.3 mg, 0.08 mmol, 31%) and its epimer 43 (14 mg, 0.03 mmol, 10%) was obtained as colorless oils, jointly with the iodo-acetate 44 (38.2 mg, 0.05 mmol, 22%) as a yellow oil. Compound 42: $[\alpha]_\text{D} -3.5$ (c, 1.9); IR 2935, 1747, 1456, 1261 cm^{-1} ; ^1H NMR d_H 1.41 (1H, m), 1.49–1.63 (2H, m), 1.92 (1H, m), 2.12–2.18 (2H, m), 3.01 (1H, dd, $J/48.2, 8.2$ Hz), 3.14–3.18 (1H, m), 3.15 (1H, d, $J/48.9$ Hz), 3.43 (1H, m), 3.44 (3H, s), 3.54 (3H, s), 3.56 (3H, s), 3.61 (3H, s), 4.41 (1H, d, $J/48.5$ Hz), 4.46 (1H, dd, $J/48.7, 8.7$ Hz), 4.98–5.10 (4H, m), 7.29–7.38 (10H, m); no NOE correlation was observed between 1-H and 6-H; ^{13}C NMR d_C 20.2 (CH₂), 24.1 (CH₂), 35.5 (CH₂), 43.6 (CH₂), 56.9 (CH₃), 59.9 (CH₃), 60.0 (CH₃), 61.7 (CH₃), 68.0 (CH₂, d, $J_p/46.1$ Hz), 68.1 (CH₂, d, $J_p/46.1$ Hz), 81.5 (CH), 84.5 (CH), 87.3 (CH), 88.4 (C), 99.0 (CH), 127.4 (2 $\times$ CH), 127.8 (CH), 128.0 (2 $\times$ CH), 128.1 (CH), 128.3 (2 $\times$ CH), 128.5 (2 $\times$ CH), 136.6 (C, d, $J_p/46.1$ Hz), 137.1 (C, d, $J_p/46.1$ Hz); MS m/z (rel intens) 504 ($\text{M}^{\text{b}}-\text{OCH}_3$, 1), 91 (100); HRMS m/z calcd for $\text{C}_{26}\text{H}_{35}\text{NO}_7\text{P}$, 504.2151, found 504.2055. Anal. Calcd for $\text{C}_{27}\text{H}_{38}\text{NO}_8\text{P}$: C, 60.55; H, 7.15; N, 2.62. Found: C, 60.19; H, 7.53; N, 2.55. Compound 43: $[\alpha]_\text{D} -3.7$ (c, 1.6); IR 2927, 1747, 1455, 1260, 1087 cm^{-1} ; ^1H NMR d_H 1.44 (1H, m), 1.52–1.68 (3H, m), 1.76 (1H, m), 1.99 (1H, m), 3.12 (1H, dd, $J/49.4, 8.0$ Hz), 3.17–3.29 (2H, m), 3.26 (1H, dd, $J/49.4, 6.4$ Hz), 3.49 (3H, s), 3.57 (3H, s), 3.58 (3H, s), 3.67 (3H, s), 4.24 (1H, d, $J/48.0$ Hz), 4.71 (1H, d, $J/49.4$ Hz), 4.99–5.14 (4H, m), 7.29–7.39 (10H, m); NOE correlation between 1-H and 6-H was observed; ^{13}C NMR d_C 16.2 (CH₂), 23.1 (CH₂), 24.5 (CH₂, d, $J_p/46.1$ Hz), 42.4 (CH₂), 56.4 (CH₃), 60.3 (CH₃), 60.4 (2 $\times$ CH₃), 67.4 (CH₂), 67.5 (CH₂), 81.3 (CH), 83.7 (CH), 84.6 (CH), 88.8 (C), 99.8 (CH), 127.5 (CH), 127.9 (4 $\times$ CH), 128.1 (CH), 128.4 (2 $\times$ CH), 128.5 (2 $\times$ CH), 136.6 (2 $\times$ C); MS m/z (rel intens) 504 ($\text{M}^{\text{b}}-\text{OCH}_3$, 1), 88 (100); HRMS m/z calcd for $\text{C}_{26}\text{H}_{35}\text{NO}_7\text{P}$, 504.2151, found 504.2067. Anal. Calcd for $\text{C}_{27}\text{H}_{38}\text{NO}_8\text{P}$: C, 60.55; H, 7.15; N, 2.62. Found: C, 60.19; H, 7.52; N, 2.55. Compound 44: $[\alpha]_\text{D} +41.8$ (c, 0.3); IR 3331, 2935, 1748, 1456, 1238 cm^{-1} ; ^1H NMR d_H 1.47 (1H, m), 1.60 (1H, m), 1.68 (1H, m), 1.94 (1H, m), 2.10 (3H, s), 2.83–2.99 (2H, m), 3.14 (3H, s), 3.28 (1H, dd, $J/47.0, 1.9$ Hz), 3.43 (3H, s), 3.44 (3H, s), 3.53 (3H, s), 3.67 (1H, dd, $J/44.7, 1.9$ Hz), 4.73 (1H, d, $J/44.7$ Hz), 4.99 (1H, dd, $J/410.3, 3.8$ Hz), 5.03 (2H, d, $J_p/47.5$ Hz), 5.04 (2H, d, $J_p/47.5$ Hz), 5.86 (1H, d, $J/47.0$ Hz), 7.30–7.37 (10H, m), the hydrogen from the NH group is missing; a NOE correlation between 4-H and -OAc hydrogen atoms was observed; ^{13}C NMR d_C 21.1 (CH₃), 29.7 (CH₂), 30.8 (CH₂), 31.8 (CH), 40.4 (CH₂), 57.2 (CH₃), 58.1 (CH₃), 59.5 (CH₃), 60.1 (CH₃), 67.9 (2 $\times$ CH₂), 77.2 (CH), 78.6 (CH), 82.8 (CH), 97.7 (CH), 104.3 (C), 127.8 (4 $\times$ CH), 128.2 (2 $\times$ CH), 128.4 (4 $\times$ CH), 136.4 (2 $\times$ C), 170.7 (C); MS (FAB) m/z (rel intens) 744 ($\text{M}^{\text{b}}\text{Na}$, 10), 722 (3), 91 (100); HRMS m/z calcd for $\text{C}_{29}\text{H}_{41}\text{INO}_{10}\text{P}$, 721.1513, found 721.1481. Anal. Calcd for

C₂₉H₄₁INO₁₀P: C, 48.27; H, 5.73; N, 1.94. Found: C, 48.28; H, 5.90; N, 2.09.

Acknowledgements

This work was supported by the research programmes BQU2000-0650 and BQU2001-1665 of the Direcció'n General de Investigació'n Científica y Te'cnica, Spain. I.P.-M. thanks the I3P-CSIC programme for a fellowship.

Supplementary data

Detailed experimental procedures and spectral and analytical data for all compounds are provided. Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tet.2009.05.049.

References and notes

- Recent reviews: (a) Tanko, J. M. *Annu. Rep. Prog. Chem., Sect. B* 2006, **102**, 247–268; (b) Robertson, J.; Pillai, J.; Lush, R. K. *Chem. Soc. Rev.* 2001, **30**, 94–103; (c) Feray, L.; Kuznersov, N.; Renaud, P. In *Radicals in Organic Synthesis*; Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, 2001; Vol. 2, pp 246–278; (d) Majetich, G. *Tetrahedron* 1995, **51**, 7095–7129.
- For reviews, see: (a) *Radicals in Organic Synthesis*; Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, 2001; (b) Curran, D. P.; Porter, N. A.; Giese, B. *Stereochemistry of Radical Reactions*; VCH: Weinheim, 1996; (c) Snider, B. B. *Chem. Rev.* 1996, **96**, 339–363; (d) Jasperse, C. P.; Curran, D. P.; Fevig, T. L. *Chem. Rev.* 1991, **91**, 1237–1286; (e) Curran, D. P. *Synthesis* 1988, 489–513; (f) Giese, B. *Radicals in Organic Synthesis: Formation of Carbon–Carbon Bonds*; Pergamon: Oxford, 1986.
- For some 1,5-HAT from sp³ to sp² carbons followed by radical cyclisation see: (a) De'ne's, F.; Beaufils, F.; Renaud, P. *Org. Lett.* 2007, **9**, 4375–4378 and references cited therein; (b) Sha, C.-K.; Hsu, C.-W.; Chen, Y.-T.; Cheng, S.-Y. *Tetrahedron Lett.* 2000, **41**, 9865–9869; (c) Robertson, J.; Peplow, M. A.; Pillai, J. *Tetrahedron Lett.* 1996, **37**, 5825–5828; (d) Bosch, E.; Bachi, M. D. *J. Org. Chem.* 1993, **58**, 5581–5582; (e) Brown, C. D. S.; Simpkins, N. S. *Tetrahedron Lett.* 1993, **34**, 131–132; (f) Borthwick, A. D.; Caddick, S.; Parsons, P. J. *Tetrahedron* 1992, **48**, 10655–10666; (g) Curran, D. P.; Abraham, A. C.; Liu, H. *J. Org. Chem.* 1991, **56**, 4335–4337; (h) Snieckus, V.; Cuevas, J.-C.; Sloan, C. P.; Liu, H.; Curran, D. P. *J. Am. Chem. Soc.* 1990, **112**, 896–898; (i) Borthwick, A. D.; Caddick, S.; Parsons, P. J. *Tetrahedron Lett.* 1990, **31**, 6911–6914; (j) Lathbury, D. C.; Parsons, P. J.; Pinto, I. *J. Chem. Soc., Chem. Commun.* 1988, 81–82; (k) Curran, D. P.; Kim, D.; Liu, H. T.; Shen, W. *J. Am. Chem. Soc.* 1988, **110**, 5900–5902.
- For recent reviews on the chemistry of nitrogen-centred radicals see: (a) Stella, L. In *Radicals in Organic Synthesis*; Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, 2001; Vol. 2, pp 407–426; (b) Fallis, A. G.; Brinza, I. M. *Tetrahedron* 1997, **53**, 17543–17594; (c) Zard, S. Z. *Synlett* 1996, 1148–1155; (d) Esker, J. L.; Newcomb, M. In *Advances in Heterocyclic Chemistry*; Katritzky, A. R., Ed.; Academic: New York, NY, 1993; Vol. 58, pp 1–45; (e) Stella, L. *Angew. Chem., Int. Ed. Engl.* 1983, **22**, 337–350; For previous reviews, see: (f) Mackiewicz, P.; Furstoss, R. *Tetrahedron* 1978, **34**, 3241–3260; (g) Neale, R. S. *Synthesis* 1971, 1–15; For a recent review on the synthesis of heterocycles by radical cyclisation see: (h) Bowman, W. R.; Bridge, C. F.; Brookes, P. *J. Chem. Soc., Perkin Trans. 1* 2000, 1–14; For the pioneering works on intramolecular HAT reactions promoted by N-radicals see: (i) Hoffman, A. W. *Ber.* 1883, **16**, 558–560; (j) Loeffler, K.; Freytag, C. *Ber.* 1909, **42**, 3427–3431; (k) Barton, D. H. R.; Beckwith, A. L. J.; Goosen, A. *J. Chem. Soc.* 1965, 181–190; (l) Baldwin, J. E.; Barton, D. H. R.; Dainis, I.; Pereira, J. L. C. *J. Chem. Soc.* 1968, 2283–2289; (m) Carruthers, W. *Some Modern Methods of Organic Synthesis*, 2nd ed.; Cambridge University Press: Cambridge, 1978, pp 255–257; For 1,5-hydrogen atom transfer from sp³ carbons to oxygen radicals followed by radical addition reactions, see: (n) Petrovic, G.; Cekovic, Z. *Tetrahedron* 1999, **55**, 1377–1390; (o) Tsunoi, S.; Ryu, I.; Ohuda, T.; Tanaka, M.; Komatsu, M.; Sonoda, N. *J. Am. Chem. Soc.* 1998, **120**, 8692–8701 and references cited therein.
- (a) Francisco, C. G.; Herrera, A. J.; Pe'rez-Marti'n, I.; Sua'rez, E. *Tetrahedron Lett.* 2007, **48**, 6384–6388; (b) Marti'n, A.; Pe'rez-Marti'n, I.; Sua'rez, E. *Org. Lett.* 2005, **7**, 2027–2030; (c) Francisco, C. G.; Herrera, A. J.; Sua'rez, E. *J. Org. Chem.* 2003, **68**, 1012–1017; (d) Freire, R.; Marti'n, A.; Pe'rez-Marti'n, I.; Sua'rez, E. *Tetrahedron Lett.* 2002, **43**, 5113–5116; (e) Dorta, R. L.; Francisco, C. G.; Sua'rez, E. *J. Chem. Soc., Chem. Commun.* 1989, 1168–1169; (f) Armas, P.; Francisco, C. G.; Herna'ndez, R.; Salazar, J. A.; Sua'rez, E. *J. Chem. Soc., Perkin Trans. 1* 1988, 3255–3265; (g) Carrau, R.; Herna'ndez, R.; Sua'rez, E.; Betancor, C. *J. Chem. Soc., Perkin Trans. 1* 1987, 937–943.
- N-Nitroamines, N-cyanamides, N-phosphoramidates and N-tert-butoxycarbonylamides ethers that were one, two, three, or four carbons long were used.
- (a) Francisco, C. G.; Freire, R.; Herrera, A. J.; Pe'rez-Marti'n, I.; Sua'rez, E. *Tetrahedron* 2007, **63**, 8910–8920; (b) Francisco, C. G.; Herrera, A. J.; Sua'rez, E. *J. Org. Chem.* 2002, **67**, 7439–7445; (c) Francisco, C. G.; Freire, R.; Herrera, A. J.; Pe'rez-Marti'n, I.; Sua'rez, E. *Org. Lett.* 2002, **4**, 1959–1961; (d) Dorta, R. L.; Marti'n, A.; Salazar, J. A.; Sua'rez, E. *J. Org. Chem.* 1998, **63**, 2251–2261; (e) Marti'n, A.; Salazar, J. A.; Sua'rez, E. *J. Org. Chem.* 1996, **61**, 3999–4006; For other HAT reactions in carbohydrate chemistry, see: (f) Robins, M. J.; Guo, Z.; Samano, M. C.; Wnuk, S. F. *J. Am. Chem. Soc.* 1999, **121**, 1425–1433; (g) Chatgililoglu, C.; Giminis, T.; Spada, G. P. *Chem. dEur. J.* 1999, **5**, 2866–2876; (h) Descotes, G. *J. Carbohydr. Chem.* 1988, **7**, 1–20.
- For competition between 1,5-, 1,6- and 1,7-intramolecular hydrogen transfer see: Denenmark, D.; Winkler, T.; Waldner, A.; De Mesmaeker, A. *Tetrahedron Lett.* 1992, **33**, 3613–3616.
- For a recent review on the synthesis of spiroaminals, see: Sinibaldi, M.-E.; Canet, I. *Eur. J. Org. Chem.* 2008, 4391–4399.
- Kobayashi, M.; Yin-Ju, C.; Aoki, S.; In, Y.; Ishida, T.; Kitagawa, I. *Tetrahedron* 1995, **51**, 3727–3736.
- (a) Fujiwara, Y.; Takaki, A.; Uehara, Y.; Ikeda, T.; Okawa, M.; Yamauchi, K.; Ono, M.; Yoshimitsu, H.; Hohara, T. *Tetrahedron* 2004, **60**, 4915–4920; (b) Ripberger, H.; Schreiber, K. Solanum Steroid Alkaloids. In *The Alkaloids*; Rodrigo, R. G. A., Ed.; Academic: New York, NY, 1981; Vol. XIX, pp 81–192.
- For some recent publications on the synthesis of azaspiracid, see: (a) Li, Y. F.; Zhou, F.; Forsyth, C. J. *Angew. Chem., Int. Ed.* 2007, **46**, 279–282; (b) Yadav, J. S.; Venugopal, C. *Synlett* 2007, 2262–2266; (c) Evans, D. A.; Dunn, T. B.; Kvoerno, L.; Beauchemin, A.; Raymer, B.; Ohliva, E. J.; Mulder, J. A.; Juhl, M.; Kagechika, K.; Favor, D. A. *Angew. Chem., Int. Ed.* 2007, **46**, 4698–4703; (d) Zhou, X. T.; Lu, L.; Furkert, D. S.; Wells, C. E.; Carter, R. G. *Angew. Chem., Int. Ed.* 2006, **45**, 7622–7626; (e) Nicolaou, K. C.; Frederick, M. O.; Petrovic, G.; Cole, K. P.; Loizidou, E. Z. *Angew. Chem., Int. Ed.* 2006, **45**, 2609–2615; (f) Oikawa, M.; Uehara, T.; Iwayama, T.; Sasaki, M. *Org. Lett.* 2006, **8**, 3943–3946; (g) Nicolaou, K. C.; Koftis, T. V.; Vyskocil, S.; Petrovic, G.; Tang, W. J.; Frederick, M. O.; Chen, D. Y. K.; Li, Y. W.; Ling, T. T.; Yamada, Y. M. A. *J. Am. Chem. Soc.* 2006, **128**, 2859–2872; (h) Nicolaou, K. C.; Pihko, P. M.; Bernal, F.; Frederick, M. O.; Qian, W. Y.; Uesaka, N.; Diedrichs, N.; Hinrichs, J.; Koftis, T. V.; Loizidou, E.; Petrovic, G.; Rodriguez, M.; Sarlah, D.; Zou, N. *J. Am. Chem. Soc.* 2006, **128**, 2244–2257; (i) Nicolaou, K. C.; Vyskocil, S.; Koftis, T. V.; Yamada, Y. M. A.; Ling, T. T.; Chen, D. Y. K.; Tang, W. J.; Petrovic, G.; Frederick, M. O.; Li, Y. W.; Satake, M. *Angew. Chem., Int. Ed.* 2004, **43**, 4312–4318; (j) Nicolaou, K. C.; Li, Y. W.; Uesaka, N.; Koftis, T. V.; Vyskocil, S.; Ling, T. T.; Govindasamy, M.; Qian, W.; Bernal, F.; Chen, D. Y. K. *Angew. Chem., Int. Ed.* 2003, **42**, 3643–3648; (k) Nicolaou, K. C.; Chen, D. Y. K.; Li, Y. W.; Ling, T. T.; Vyskocil, S.; Koftis, T. V.; Govindasamy, M.; Uesaka, N. *Angew. Chem., Int. Ed.* 2003, **42**, 3649–3653; (l) Forsyth, C. J.; Hao, J. L.; Aiguade, J. *Angew. Chem., Int. Ed.* 2001, **40**, 3663–3667; (m) Nicolaou, K. C.; Pihko, P. M.; Diedrichs, N.; Zou, N.; Bernal, F. *Angew. Chem., Int. Ed.* 2001, **40**, 1262–1265 and references cited in all of them; For the first isolation and characterization of azaspiracid, see: (n) Satake, M.; Ofuji, K.; Naoki, H.; James, K. J.; Furey, A.; McMahon, T.; Silke, J.; Yasumoto, T. *J. Am. Chem. Soc.* 1998, **120**, 9967–9968.
- Xu, R.-S.; Lu, Y.-J.; Chu, J.-H.; Iwashita, T.; Naoki, H.; Naya, Y.; Nakanishi, K. *Tetrahedron* 1982, **38**, 2667–2670.
- Total synthesis: (a) Paquette, L. A.; Duan, M.; Konetzki, I.; Kempmann, C. *J. Am. Chem. Soc.* 2002, **124**, 4257–4270; (b) Duan, M.; Paquette, L. A. *Angew. Chem., Int. Ed.* 2001, **40**, 3632–3636; (c) Nicolaou, K. C.; Murphy, F.; Bar-luenga, S.; Ohshima, T.; Wei, H.; Xu, J.; Gray, D. L. F.; Baudoin, O. *J. Am. Chem. Soc.* 2000, **122**, 3830–3838; (d) Nicolaou, K. C.; Xu, J.; Murphy, F.; Barluenga, S.; Baudoin, O.; Wei, H.; Gray, D. L. F.; Ohshima, T. *Angew. Chem., Int. Ed.* 1999, **38**, 2447–2451.
- (a) Richter, P. K.; Tomaszewski, M. J.; Miller, R. A.; Patron, A. P.; Nicolaou, K. C. *J. Chem. Soc., Chem. Commun.* 1994, 1151–1152; (b) Luengo, J. I.; Gleason, J. G. *Tetrahedron Lett.* 1992, **33**, 6911–6914; (c) Giannis, A.; Sandhoff, K. *Tetrahedron Lett.* 1985, **26**, 1479–1482.
- Viand, M. C.; Rollin, P. *Synthesis* 1990, 130–132.
- Parrish, J. D.; Little, R. D. *Org. Lett.* 2002, **4**, 1439–1442.
- Gupta, A. K.; Acharya, J.; Dubey, D. K.; Kaushik, M. P. *Synth. Commun.* 2007, **37**, 3403–3407.
- (a) Gonza'lez, C. C.; Kennedy, A. R.; Leo'n, E. I.; Riesco-Fagundo, C.; Sua'rez, E. *Angew. Chem., Int. Ed.* 2001, **40**, 2326–2328; (b) Francisco, C. G.; Gonza'lez, C. C.; Sua'rez, E. *J. Org. Chem.* 1998, **63**, 8092–8093; For an example of the influence of the substituents on the 1,5-HAT reaction compared with competing b-fragmentation of alkoxy radicals see: Allen, P. R.; Brimble, M. A.; Fare's, F. A. *J. Chem. Soc., Perkin Trans. 1* 1998, 2403–2411.
- Dondoni, A.; Marra, A.; Rojo, I.; Sherrmann, M.-C. *Tetrahedron* 1996, **52**, 3057–3074.
- The structures of 37 and 38 were confirmed by n-Bu₃SnH reductive deiodination to give the starting material 23 (see Supplementary data section).
- Acetyl hypoiodite has never been isolated but its presence has been demonstrated in solutions of DIB and iodine, see: (a) Ogata, Y.; Aoki, K. *J. Am. Chem. Soc.* 1968, **90**, 6186–6191; (b) Merkushev, E. B.; Simakhina, N. D.; Kovedhnikova, G. M. *Synthesis* 1980, 486–487; (c) Courtneidge, J. L.; Luszyk, J.; Page, D. *Tetrahedron Lett.* 1994, **35**, 1003–1006.
- Although we have only found a brief mention on the reactivity of DIB/I₂ with olefins (Aoki, K.; Ogata, Y. *Bull. Chem. Soc. Jpn.* 1968, **41**, 1476–1477), in our laboratory we have observed that this system reacts with tri-O-acetyl-β-glucal to give a diastereomeric mixture from which the major product 1,3,4,6-tetra-O-acetyl-2-deoxy-2-iodo-α-D-manno-pyranose was isolated in 65% yield. For other related studies see: (a) Kirschning, A.; Plumeier, C.; Rose, L. *Chem. Commun.* 1998, 33–34; (b) Kirschning, A.; Jesberger, M.; Monenschein, H. *Tetrahedron Lett.* 1999, **40**, 8999–9002.
- As observed by ¹H NMR this is presumably a mixture of conformational isomers. Variable temperature ¹H NMR experiments were performed (70 °C) but at this temperature the equilibrium between conformers was not significantly affected.