



Toxoplasma gondii infection and toxoplasmosis in farm animals: Risk factors and economic impact

S. Stelzer^a, W. Basso^b, J. Benavides Silván^c, L.M. Ortega-Mora^d, P. Maksimov^a, J. Gethmann^a, F.J. Conraths^a, G. Schares^{a,*}

^a Friedrich-Loeffler-Institut, Federal Research Institute for Animal Health, Institute of Epidemiology, Südufer 10, 17493 Greifswald - Insel Riems, Germany

^b Institute of Parasitology, University of Bern, Länggassstrasse 122, 3012 Bern, Switzerland

^c Instituto de Ganadería de Montaña (CSIC-Universidad de León) Grulleros, 24346 León, Spain

^d SALUVET, Animal Health Department, Faculty of Veterinary Sciences, Complutense University of Madrid, Ciudad Universitaria s/n, 28040 Madrid, Spain

ARTICLE INFO

Article history:

Received 17 December 2018

Received in revised form 25 January 2019

Accepted 30 January 2019

Keywords:

Zoonosis

Livestock

Prevalence

Natural infection

Experimental infection

Costs

ABSTRACT

The protozoan parasite *Toxoplasma gondii* is a zoonotic parasite that can be transmitted from animals to humans. Felids, including domestic cats, are definitive hosts that can shed oocysts with their feces. In addition to infections that occur by accidental oral uptake of food or water contaminated with oocysts, it is assumed that a large proportion of affected humans may have become infected by consuming meat or other animal products that contained infective parasitic stages of *T. gondii*. Since farm animals represent a direct source of infection for humans, but also a possible reservoir for the parasite, it is important to control *T. gondii* infections in livestock. Moreover, *T. gondii* may also be pathogenic to livestock where it could be responsible for considerable economic losses in some regions and particular farming systems, e.g. in areas where the small ruminant industry is relevant.

This review aims to summarize actual knowledge on the prevalence and effects of infections with *T. gondii* in the most important livestock species and on the effects of toxoplasmosis on livestock. It also provides an overview on potential risk factors favoring infections of livestock with *T. gondii*. Knowledge on potential risk factors is prerequisite to implement effective biosecurity measures on farms to prevent *T. gondii* infections. Risk factors identified by many studies are cat-related, but also those associated with a potential contamination of fodder or water, and with access to a potentially contaminated environment. Published information on the costs *T. gondii* infections cause in livestock production, is scarce. The most recent peer reviewed reports from Great Britain and Uruguay suggest annual cost of about 5–15 million US \$ per country. Since these estimates are outdated, future studies are needed to estimate the present costs due to toxoplasmosis in livestock. Further, the fact that *T. gondii* infections in livestock may affect human health needs to be considered and the respective costs should also be estimated, but this is beyond the scope of this article.

© 2019 The Authors. Published by Elsevier Inc. on behalf of International Association of Food and Waterborne Parasitology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction	0
2. <i>Toxoplasma gondii</i> infections in livestock animals – importance for livestock production and prevalence	0
2.1. Pigs	0
2.1.1. Prevalence in pigs	0

* Corresponding author.

E-mail address: gereon.schares@fli.de. (G. Schares).

2.1.2.	Possible routes of infection in pigs	0
2.1.3.	Disease caused by <i>T. gondii</i> infections in pigs	0
2.1.4.	Effects of experimental infections in pigs	0
2.2.	Sheep and goats	0
2.2.1.	Prevalence in sheep and goats	0
2.2.2.	Possible routes of infection in sheep and goats	0
2.2.3.	Disease caused by <i>T. gondii</i> in naturally infected sheep and goats.	0
2.2.4.	Effects of experimental infections in sheep and goats	0
2.3.	Chickens and other poultry	0
2.3.1.	Prevalence in chickens and other poultry	0
2.3.2.	Possible routes of infection in chickens and other poultry	0
2.3.3.	Disease caused by <i>T. gondii</i> in naturally infected poultry.	0
2.3.4.	Effects of experimental infections in livestock poultry.	0
2.4.	Cattle.	0
2.4.1.	Prevalence in cattle	0
2.4.2.	Possible routes of infection in cattle.	0
2.4.3.	Disease caused by <i>T. gondii</i> infections in cattle	0
2.4.4.	Effects of experimental infections in cattle.	0
2.5.	Horses and other equids	0
2.5.1.	Prevalence in horses and other equids.	0
2.5.2.	Possible routes of infection in horses and other equids	0
2.5.3.	Disease caused by <i>T. gondii</i> infections in horses and other equids.	0
2.5.4.	Effects of experimental infections in horses	0
3.	Potential risk factors for infection in livestock	0
3.1.	General factors	0
3.1.1.	Age	0
3.1.2.	Gender.	0
3.1.3.	Geographic and regional characteristics	0
3.1.4.	Farm management	0
3.2.	Factors related to the <i>T. gondii</i> life cycle	0
3.2.1.	Definitive hosts (cats)	0
3.2.2.	Other on-farm intermediate hosts, such as rodents and their control	0
3.2.3.	Feed-related parameters	0
3.2.4.	Water-related parameters	0
3.2.5.	Soil contact, outside access and pasturing	0
4.	Economic impact of toxoplasmosis in livestock	0
5.	Conclusion and future prospects	0
	Conflict of interest statement.	0
	Acknowledgements	0
	References	0

1. Introduction

Toxoplasma gondii is a zoonotic apicomplexan parasite that is able to infect probably all warm-blooded animals, including livestock (Dubey, 2010b; Schlüter et al., 2014). Domestic cats and other felids are definitive hosts of *T. gondii*. This implies that the parasite is only able to complete its sexual life cycle in these species, i.e. environmentally resistant oocysts can only be shed with the feces of infected felids (Dubey, 2010b). Oocysts are central in the life cycle of *T. gondii*. After one up to a few days of maturation (sporulation) in the environment, they become infective to a large variety of warm-blooded intermediate hosts (livestock, synanthropic and wild living animals such as rodents or birds, poultry or humans) if ingested (Dubey, 2010b). In addition to oocysts, there are two further stages of *T. gondii*, which are infective, i.e. tachyzoites and bradyzoites, the latter being present in tissue cysts. After infection, tachyzoites invade host cells, in which they multiply. This replication is strictly intracytoplasmatic in parasitophorous vacuoles formed by the parasite (Schlüter et al., 2014). In parallel, after several rounds of multiplication, the parasite establishes intracellular tissue cysts, which contain slowly or no longer replicating bradyzoites (Jerome et al., 1998; Radke et al., 2003; Watts et al., 2015). Tissue cyst formation preferentially occurs in brain tissue, the skeletal and heart muscle or also in the retina of infected intermediate hosts (Schlüter et al., 2014).

While the tachyzoite stages are repressed after the onset of the immune response of the host, the dormant tissue cysts ensure that *T. gondii* infections persist inside the host cells, where they are protected from the immune system, possibly for the rest of the life of the intermediate host (Rougier et al., 2017). Tissue cysts may contain hundreds of bradyzoites (Dubey, 2010b). If carnivorous or omnivorous animals feed on material that contains tissue cysts, encysted bradyzoites can survive the gastric passage and initiate parasite multiplication in the intestine of the infected intermediate or definitive host (Dubey, 2010b). In some regions of

the world, in particular in Europe, risk factor studies suggested that most humans become infected by ingesting bradyzoites present in undercooked or not sufficiently inactivated meat (Cook et al., 2000; Kapperud et al., 1996).

Toxoplasma gondii is transmitted vertically in several intermediate host species including humans (Dubey, 2010b). The transplacental vertical transmission is facilitated by tachyzoites usually after the primary and during the acute phase of infection (Dubey, 2010b). Tachyzoites circulating after a reactivated chronic infection also seem to be able to facilitate vertical transmission in some livestock species, although experimental findings suggest that this might be a rare event in *T. gondii* (Dubey, 1982).

As farm animals represent a source of infection for humans and reservoirs of *T. gondii* for wildlife it has been proposed to reduce *T. gondii* infections in livestock as much as possible, particularly in pigs (Anonymous, 2011a,b). Potential risk factors for *T. gondii* infections in livestock have been studied and were reviewed in recent years (Guo et al., 2015; Opsteegh et al., 2016), but since the knowledge on the epidemiology of toxoplasmosis in animals and humans is rapidly evolving, these reviews deserved an update. Therefore, the main objectives of this study were (i) to briefly summarize the recent gain in knowledge on the prevalence of *T. gondii* in the most important livestock species and on the importance of infection with this parasite in livestock production, (ii) to compile existing knowledge on the effect of natural and experimental *T. gondii* infections in the dominant livestock species, and finally (iii) to provide an up-to-date summary on potential risk factors and risk factor studies for *T. gondii* infection in livestock.

2. *Toxoplasma gondii* infections in livestock animals – importance for livestock production and prevalence

2.1. Pigs

2.1.1. Prevalence in pigs

Seroepidemiological surveys have provided evidence for a worldwide distribution of *T. gondii* in pigs, with prevalences varying according to age, pig categories, geography and management system. In general, a low prevalence of *T. gondii* infections (<1%) is observed in pigs reared in confinement with controlled management conditions, preventing access of rodents and cats, whereas higher prevalence values (>60%) can be found in farms with free-ranging pigs, farms without controlled conditions allowing outdoor access and in backyard holdings (De Berardinis et al., 2017).

Worldwide information on *T. gondii* infection in pigs up to 2009 was reviewed several times in the past (Dubey, 1986b, 2009a; Dubey and Beattie, 1988). The situation in the USA was also reviewed more recently (Hill and Dubey, 2013). A systematic review, based on reports on a direct identification of *T. gondii* in pigs and pork identified a pooled *T. gondii* prevalence of 12.3% with a 95% prediction interval ranging from 0 to 55% (Belluco et al., 2017). In Europe, the seroprevalence for *T. gondii*-specific antibodies was reported to range from 0 to 64% in fattening pigs and from 3 to 31% in breeding sows (De Berardinis et al., 2017). In Africa, a systematic review and meta-data analysis of seroepidemiological studies reported seroprevalences from 9.3 to 40.6%, with an overall estimated *T. gondii* seroprevalence in pigs of 26% (Tonouhewa et al., 2017). In China, pigs are assumed to represent one of the farm animal species most frequently infected with *T. gondii*. It was estimated that 30 to 50% of the raised pigs are seropositive, reaching a prevalence of 70% in some areas and farms (Pan et al., 2017). There is quite a large number of further and more recent local studies on prevalence in pigs, but there is little information in addition to that what has been reported and summarized before.

2.1.2. Possible routes of infection in pigs

Most *T. gondii* infections in pigs are acquired postnatally, either by ingestion of sporulated oocysts in contaminated soil, feed and water, or by ingestion of cysts in the tissues of infected intermediate hosts (e.g. rodents, birds, meat and cannibalism). Pigs can also become infected prenatally by transplacental transmission of the parasite (Dubey, 2009a). The occurrence of galactogenic transmission of *T. gondii* from the sow to the piglets has been reported, but might be a rare event (Basso et al., 2017). It is presumed that infection through oocysts accounts for most infections in conventional pig breeding systems and especially in outbreaks of clinical toxoplasmosis involving several animals (Kim et al., 2009; Li et al., 2010; Okamoto et al., 1989; Weissenböck and Dubey, 1993).

2.1.3. Disease caused by *T. gondii* infections in pigs

Toxoplasma gondii infections in pigs are commonly subclinical; nevertheless, several cases of clinical disease after natural infection have been recorded worldwide (Dubey, 1986b, 2009a; Dubey and Beattie, 1988). Clinical manifestations seem to occur more frequently in neonatal and weaned pigs, but also cases of clinical toxoplasmosis affecting sows have been described. Common signs observed in clinically infected pigs include anorexia, apathy, fever, ocular and nasal discharge, dyspnoea, cyanosis, diarrhoea, limb weakness, neurological signs and sometimes death (Dubey, 1986b, 2009a; Dubey and Beattie, 1988). None of these signs is pathognomonic for toxoplasmosis. Besides, *T. gondii* infections may be associated with reproductive failure in sows characterized by abortion, fetal mummification, stillbirth and neonatal mortality (Basso et al., 2015; Dubey, 1986b, 2009a; Dubey and Beattie, 1988). Clinical disease is believed to occur only during the acute phase of infection as result of necrotic and inflammatory processes during tachyzoite multiplication in several tissues. Chronically infected animals do not have clinical signs, but they represent an important source of infection for humans, in particular, if undercooked pork or insufficiently treated meat products containing tissue cysts are consumed (Dubey, 2009a). Different factors are believed to influence the clinical presentation of *T. gondii* infection in pigs such as age and immune status of the host, co-infection with other agents, the parasitic stage of *T. gondii* (i.e. oocyst, tissue cyst), infection dose, and the strain or the genetic background of *T. gondii*. In some cases, viral

infections such as Porcine Circovirus type 2 (Klein et al., 2010) and Porcine Parvovirus (Basso et al., 2015) were also associated with clinical manifestations of toxoplasmosis in pigs.

Reports prior to 2009 were summarized previously (Dubey, 1986b, 2009a; Dubey and Beattie, 1988). Since then, further confirmed cases of clinical toxoplasmosis involving (i) suckling piglets in Brazil (Olinda et al., 2016), (ii) fattening pigs in China (Li et al., 2010), Germany (Klein et al., 2010) and Brazil (Olinda et al., 2016) and (iii) *T. gondii* infections associated with reproduction failure in sows in Switzerland (Basso et al., 2015) were published.

A severe outbreak of toxoplasmosis in fattening pigs was reported from Gansu Province, China, with morbidity affecting 549/960 (57%) fattening pigs (Li et al., 2010). The pigs had fever (40–42.2 °C), anorexia and depression, and 19 of the affected pigs died. Serological analysis of 154 clinically ill animals had *T. gondii* IgG or IgM positive ELISA results in 142 (92.2%) and 147 (95.4%) of the animals, respectively. Moreover, *T. gondii* could be isolated in mice by intraperitoneal inoculation of pooled heart, liver, spleen and brain tissues from two pigs which showed clinical signs. The source of infection was assumed to be feed contaminated with cat feces. A controlled feeding experiment administering randomly collected feed to five seronegative piglets lead to development of fever, depression and seroconversion of three of the animals. In another study (Klein et al., 2010), systemic toxoplasmosis was diagnosed in a 3.5-month-old fattening pig suffering from post-weaning multisystemic wasting syndrome associated with PCV-2 infection in Germany. The pig had severe respiratory signs and died suddenly. Immunohistochemically, *T. gondii* was detected associated with interstitial and necrotizing pneumonia, lymphadenitis and adrenal necrosis. It was assumed that immunosuppression caused by primary PCV-2 infection may have triggered secondary systemic toxoplasmosis (Klein et al., 2010).

Interestingly, many of these cases of clinical toxoplasmosis in pigs were registered in Asia (i.e. Japan, Taiwan, China, Korea, Thailand), although there are reports of clinical toxoplasmosis in pigs from several countries around the world (Dubey, 2009a; Dubey and Beattie, 1988). It is not known if specific *T. gondii* genotypes circulating in Asia may be more prone to cause clinical infections in pigs (Basso et al., 2017).

Recently, *T. gondii* parasites belonging to the Chinese 1 genotype (synonymous to ToxoDB#9), a frequent genotype in Asia and especially in China (Hou et al., 2018; Wang et al., 2017), were detected in apparently independent fatal cases of toxoplasmosis in two pigs in Brazil (Olinda et al., 2016). The animals (aged one and four months) showed apathy, dyspnoea, poor general condition and died after a few days. The main lesions in both pigs consisted of severe diffuse necrotizing bronchointerstitial pneumonia associated with numerous *T. gondii* tachyzoites present in the lesions. Interestingly, the cases occurred 3 months apart from each other and both animals derived from two different farms, showing that *T. gondii* resembling a ToxoDB#9-like genotype is circulating in Brazil (Olinda et al., 2016). This genotype was also identified in 16 out of 17 samples from infected pigs with high fever, dyspnoea, subcutaneous haemorrhage, abortion, enlargement and necrosis of liver and spleen suspected of having clinical toxoplasmosis in China (Jiang et al., 2013). In China, *T. gondii* infections in pigs are very common and outbreaks of clinical toxoplasmosis with death of numerous pigs have been reported on several occasions (summarized by Pan et al. (2017) and Li et al. (2010)). Moreover, there are reports of repeated outbreaks over 5 years in an individual pig farm in the Shandong Province (Li et al., 2010). These outbreaks of fatal toxoplasmosis were thought to be related to consumption of feed contaminated with oocysts from cat feces (Li et al., 2010). Unfortunately, no molecular characterization of the isolates involved in these outbreaks was performed. Studies in Jiangsu province, Eastern China, revealed high positive rates of *T. gondii* infection in sick pigs (showing poor mental state, fever, and/or dyspnoea) with 46.8% (66/141) PCR positive animals in various tissues (Hou et al., 2018). In 58 pigs, coinfection with other pathogens was observed but in seven animals *T. gondii* was the only agent detected, suggesting that it could be involved in the aetiology of sickness or death of pigs in that region. Molecular analysis of *T. gondii* from 17 sick pigs showed that *T. gondii* Chinese 1 (ToxoDB#9) was the most frequently (11/17) detected genotype (Hou et al., 2018). In China, parasites with this genotype were also isolated from one case of human toxoplasmosis, but in several American countries also from subclinically infected livestock animals (summarized in Olinda et al. (2016)).

Contrary to the vast knowledge about the importance of vertical transmission of *T. gondii* in small ruminants and humans, the role of *T. gondii* as cause of reproductive disorders in sows and the epidemiological significance of intrauterine and galactogenic infections in piglets, showing no clinical signs are less understood (Basso et al., 2017). Reports of reproductive failure due to toxoplasmosis and congenital infection in piglets are well documented, but the experimental reproduction of vertical transmission in pregnant sows is often not successful (Basso et al., 2017). In general, sows that abort or deliver infected offspring usually do not show further clinical signs, but fever, anorexia, neurological signs and even death were observed on some occasions in sows that aborted and transmitted the infection to the fetuses (Kim et al., 2009). In China, abortions caused by *T. gondii* in sows are considered common and assumed to cause economic losses (Pan et al., 2017). In Europe, reports of reproductive problems due to *T. gondii* infection in pigs are scarce. A large epidemiological study in 94 pig breeding farms in Germany suggested an association of *T. gondii* with reproductive failure in sows. The within-farm seroprevalence to *T. gondii* was significantly higher in farms experiencing reproductive disorders (repeat-breeders, abortion, neonatal mortality), than in farms without such problems, but the role of *T. gondii* in causing these reproductive problems was not further assessed (Damriyasa et al., 2004). Recently, *T. gondii* was detected in the placenta or in fetuses of 34 out of 113 sows that had aborted or delivered a high number of stillborn or weak piglets in Switzerland (Basso et al., 2015). By real time PCR, *T. gondii* DNA was detected in three placentas from one seropositive sow (abortion at 71 days of gestation [dg]), and in brain tissues from one fetus (abortion at 76 dg), one stillborn (116 dg) and one mummy (112 dg) originating from three further seropositive sows, but in no sample derived from the seronegative dams (Basso et al., 2015). By contrast, the examination of fetal tissues and fluids from 32 sow abortions in Romania by PCR did not yield any *T. gondii* positive samples (Iovu et al., 2010).

2.1.4. Effects of experimental infections in pigs

Pigs can be experimentally infected with any *T. gondii* stage (i.e. oocysts, tissue cysts, tachyzoites). Most experimentally inoculated pigs, including animals inoculated with very low infection doses (as few as 1 or 10 oocysts), seroconverted after 2–4 weeks and the parasite could be successfully recovered from different tissues. However, experimental reproduction of clinical toxoplasmosis, vertical transmission and congenital toxoplasmosis in pigs is considered difficult (Boch et al., 1965b; Dubey, 2009a; Dubey and Beattie, 1988; Dubey and Urban, 1990; Moller et al., 1970; Sanger and Cole, 1955; Work et al., 1970). Various parasite related factors (i.e. *T. gondii* stage, dose, infection route, virulence and the genetic background of the strain) and host related factors (i.e. breed, age, immune status and stage of gestation) may influence the outcome of an experimental infection (Dubey, 1986b, 2009a). Weaned pigs fed oocysts or tissue cysts often developed transient clinical signs such as weight loss, anorexia and/or fever, independent of the *T. gondii* isolate in the inoculum and generally recovered by three weeks post inoculation (Basso et al., 2013; Dubey, 2009a).

Experimental infections with *T. gondii* in pigs were performed within the framework of numerous studies aiming to reveal different aspects of the biology of the parasite (pathogenesis, immune response, persistence of the infection in the tissues, reproduction of congenital toxoplasmosis, development and evaluation of diagnostic methods) or aiming to establish vaccines (Dubey, 2009a; Dubey and Beattie, 1988).

It seems that clinical toxoplasmosis in any pig category and vertical transmission of *T. gondii* in pregnant sows can be more frequently reproduced by intravenous inoculation of high doses of tachyzoites than by feeding tissue cysts or oocysts. Furthermore, the potential occurrence of vertical transmission may be influenced by the *T. gondii* isolate used in the inoculations (Basso et al., 2017; Dubey and Urban, 1990; Jungersen et al., 2001; Work et al., 1970). Oral inoculations with 10^3 oocysts of the GT-1 strain (Type I strain; ToxoDB#10) led to a transplacental infection in five out of 11 inoculated pregnant sows and to transient lethargy, anorexia and respiratory distress between 5 and 15 days post infection (dpi) (Dubey and Urban, 1990), while inoculations with 10^4 to 10^5 oocysts of the CZ isolate (a European Type II isolate, Toxo DB#3) were not able to reproduce vertical transmission or other clinical signs in any of the 8 pregnant and infected sows (Basso et al., 2017). Likewise, feeding of 5×10^3 oocysts of the CZ isolate to six 4.5 week-old piglets caused infection in all animals but only transiently fever (in all animals); apathy, anorexia and soft feces (in only one piglet) were observed, suggesting a low virulence of this isolate for pigs (Basso et al., 2013). Nevertheless, some authors considered low pathogenic *T. gondii* strains as good candidates to reproduce vertical transmissions in sows as these parasites might produce a subclinical infection in the dam, having a better chance of establishing placental infections and congenital toxoplasmosis in the piglets before development of a limiting immune response in the sow (Jungersen et al., 2001).

Experimental infections of pigs have recently been performed to evaluate viability of *T. gondii* in meat after processing techniques. Twelve pigs were inoculated with 10^3 *T. gondii* oocysts of a type II field isolate from cat feces and slaughtered 4 months after inoculation. Clinical signs were not reported, but the pigs seroconverted post inoculation and PCR positive results were obtained from most thighs, both at slaughter and post curing (Genchi et al., 2017). In two further experimental studies conducted to test vaccination or to assess swine as an experimental model for human ocular toxoplasmosis, no clinical signs and also no ocular toxoplasmosis were reported after experimental infection with either 10^3 oocysts per animal or 10^3 tissue cysts per animal of the M4 strain (a *T. gondii* Type II strain) of pigs (Burrells et al., 2015; Garcia et al., 2017).

2.2. Sheep and goats

Sheep and goats are highly susceptible for infections with *T. gondii* and this protozoan parasite is considered a major cause of reproductive losses in small ruminants worldwide. While most descriptions and investigations have been carried out in sheep (Dubey, 2009b), toxoplasmosis has a similar or even greater importance as an abortive disease in goats (Dubey, 2010b). In addition, toxoplasmosis is a relevant zoonosis and infection in small ruminants may play a major role in its transmission to humans (Belluco et al., 2016; Opsteegh et al., 2016).

2.2.1. Prevalence in sheep and goats

Antibodies to *T. gondii* have been found in sheep and goats worldwide. More than 200 articles reported seroprevalence studies in these domestic ruminant species before 2010, as reviewed by Dubey (2009b, 2010b). At that time, areas of the world with a large number of seroprevalence reports were Brazil, Europe, North America, and the Middle East. From 2010 to 2018, further epidemiological studies in small ruminants have been published, including reports from areas where information was scarce and regions, where sheep and goats are relevant livestock species. These studies are from different parts of Asia (i.e. China, Pakistan, South East Asia), Sub-Saharan Africa and countries from the Mediterranean (Ahmed et al., 2016; Dong et al., 2018; Garcia-Bocanegra et al., 2013; Gazzonis et al., 2015; Kantzoura et al., 2013; Khan et al., 2017; Tilahun et al., 2018; Tzanidakis et al., 2012). Although differences in study design, purpose of the study, serological methodology and cut off points applied make it difficult or even impossible to compare data, these as well as the previous studies clearly show that *T. gondii* infections are highly prevalent in small ruminants (Dubey, 2010b). In the following, representative examples of recent studies conducted on different continents are summarized.

In Africa, in a recent meta-analysis, summarizing data from 1969 to 2016, the overall estimated prevalence was 26.1% (17.0–37.0%) for sheep and 22.9% (12.3–36.0%) for goats (Tonouhewa et al., 2017). In Egypt, antibody prevalence was higher in goats (62%) than in sheep (between 4.1 and 26%) (Al-Kappany et al., 2018). In Tunisia, antibodies to *T. gondii* were found in 40.2% sheep and 34.5% goats (Lahmar et al., 2015). In Ethiopia, the seroprevalence of *T. gondii* infection in sheep and goats was

33.7% and 27.6%, respectively (Tilahun et al., 2018). A further study from this country reported high flock (59.7%) and individual animal (31.8%) *T. gondii* seroprevalences associated with abortion in some districts (Gebremedhin et al., 2014). A lower seroprevalence was reported from South Africa with 8% in sheep (Hammond-Aryee et al., 2015).

In America, a systematic meta-analysis provided estimates on *T. gondii* infection in food animals produced in the United States, including small ruminants. *T. gondii* infection seroprevalence in goats (30.7%) was higher than in sheep or lambs (22.0%) (Guo et al., 2016). Further studies report *T. gondii* seroprevalences in sheep and goats from the Caribbean Islands Dominica (67%, 58%), Grenada (48%, 57%), Montserrat (89%, 80%) and St. Kitts and Nevis (57%, 42%) (Hamilton et al., 2014). In another study, antibodies to *T. gondii* (Modified Agglutination Test (MAT) titre 1:≥25) were found in 44.1% of sheep and 42.8% goats in Grenada and Carriacou (Chikweto et al., 2011). In Brazil, serum samples of 930 sheep were tested in two regions of Rio Grande do Norte (Northeastern Brazil), with different climatic conditions and the overall estimated prevalence was 22.1% (Andrade et al., 2013).

Regarding Asia, the situation of *T. gondii* infections has recently been reviewed for China. Seroprevalence for *T. gondii* in sheep has been estimated to be 11.8% (2305/19,565) and the overall estimated seroprevalence for *T. gondii* in goats was 17.6% (3260/18,556) (Dong et al., 2018). In Myanmar, an 11.4% seroprevalence has been reported in goats (Bawm et al., 2016). In other South Asian countries, reported prevalence in sheep and goats was 21.1% and 25.4%, respectively (Khan et al., 2017). In Pakistan, the results also showed higher seroprevalence of *T. gondii* in goats (42.8%) as compared to sheep (26.2%) (Ahmed et al., 2016). In Saudi Arabia, antibodies to *T. gondii* were found in 36.4% (325/891) of sheep and 35.3% (196/555) of goats (Alanazi, 2013).

In Europe, high prevalence values have been observed in both, sheep and goats in Mediterranean countries. In Greece, in one study, sheep had a higher seroprevalence (48.6% [729/1501]) for *T. gondii* than goats (30.7% [166/541]) (Tzanidakis et al., 2012). In Thessaly, a total of 540 sheep and goat serum samples were examined and the seroprevalence was 24.5% (Kantzoura et al., 2013). In another study, specific IgG against *T. gondii* were detected in 53.71% and 61.3% of the sheep and goats from mixed flocks (Diakoua et al., 2013). In Northern Italy, antibodies were found in 96.6% of goat farms and in 87.5% of sheep farms; 41.7% goats and 59.3% sheep had a positive result. The seroprevalence was significantly higher in sheep than in goats (Gazzonis et al., 2015). In Portugal, 33.6% of sheep and 18.5% of goats were seropositive by a modified agglutination test (MAT) (A.P. Lopes et al., 2013). In Southern Spain, 248 (49.3%) of 503 sheep, and 124 (25.1%) of 494 goats were seropositive. The herd seroprevalence was 84.7% (61/72), and 72.2% (52/72) for sheep and goats, respectively (Garcia-Bocanegra et al., 2013). In another study in the same region, the seroprevalence was 41.2% in sheep and 5.6% in goats (Almeria et al., 2018). In the northwestern part of Spain, individual (48%) and flock-level (74%) *T. gondii* seroprevalence values in goats were high; the within-flock prevalence was 53% (Diaz et al., 2016). In Eastern Europe as Poland, seroprevalences of 21% in goats and 47% in sheep have been reported (Moskwa et al., 2018). In Romania, the seroprevalence in sheep varied with the region, age and the serological methods from 6.9 to 71% (Dubey et al., 2014a). In the UK, of the 3539 sera collected from 227 sheep flocks, 2619 (74%) were found to be positive for *T. gondii* specific antibodies (Hutchinson et al., 2011). In France, applying a low cut off titre of 1:≥6 in MAT the overall estimate of the *T. gondii* seroprevalence was 17.7% (11.6–31.5%) for lambs and 89% (73.5–100%) for adult sheep (Halos et al., 2010). In a Scandinavian country (Norway), 55 of 73 flocks (75%) had one or more serologically positive animals, while 377 of 2188 (17%) of the individual samples tested positive for *T. gondii* antibodies (Stormoen et al., 2012).

In Oceania, 1917 out of 2254 (85%) sheep sera tested in New Zealand were positive, using a titre of 1:≥16, and 1384/2254 (61%) with a titre of 1:≥64 using a latex agglutination test. All 198 ewe flocks tested were seropositive for antibodies to *T. gondii*, at a titre of 1:≥16, and all but three were at a titre of 1:≥64 (Dempster et al., 2011).

Isolation of viable parasites from tissues of small ruminants corroborate serological findings and confirm that these species are important intermediate hosts. In sheep, viable *T. gondii* has been detected in brain, heart, diaphragm and different muscles (Dubey, 2010b; Opsteegh et al., 2016). Due to the fact that *T. gondii* readily disseminates into the edible tissues of sheep, this parasite represents a risk for consumers (Belluco et al., 2016; Opsteegh et al., 2016). In goats, brain and heart also rank high on the list of predilection organs and muscle tissues had high within study scores, and ranked first when combined in the meat/muscle category (Opsteegh et al., 2016). These results are corroborated by studies in different areas of the world. For instance, the proportion sheep carcasses in France carrying live parasites according to bioassay results was estimated at 5.4% (3–7.5%) (Halos et al., 2010). In the US, 53 isolates of *T. gondii* were obtained from 68 seropositive lambs sampled at the slaughterhouse (Dubey et al., 2008). In another study in this country, hearts of goats obtained from a local grocery store were examined for *T. gondii* infection and the parasite was isolated from 29 out of 112 animals (Dubey et al., 2011).

2.2.2. Possible routes of infection in sheep and goats

Horizontal transmission of *T. gondii* to small ruminants by the oral uptake of environmentally resistant oocysts through contaminated fodder or water is considered the most important route of transmission (Buxton and Losson, 2007; Dubey, 2010b). It is generally assumed that <2% of sheep become infected congenitally and <4% of the persistently infected sheep transmit the infection to their offspring (reviewed in Dubey (2010b) and Innes et al. (2009)). Recrudescence of a chronic infection and the endogenous trans-placental transmission of the parasite to offspring was described in goats (Dubey, 1982). In addition, it was proposed some years ago that a repeated transplacental transmission in sheep was more common than previously thought (Williams et al., 2005) and recent descriptions from Brazil seem to corroborate this hypothesis (Dos Santos et al., 2016; Klauck et al., 2016). Further studies are needed to assess the possibility that certain breeds are more susceptible to endogenous vertical transmission in chronically infected ewes or that vertical transmission is a trait of particular *T. gondii* strains or genotypes.

Possible alternative routes are venereal or galactogenic transmission. Several studies have identified *T. gondii* DNA in semen samples from rams and male goats, either from natural cases of infection (Bezerra et al., 2014) or from animals experimentally

inoculated (W.D. Lopes et al., 2013; Santana et al., 2010). Furthermore, the transmission of the infection to sheep and goats through semen has also been proven, both under mating with experimentally infected rams (W.D. Lopes et al., 2013) or through artificial insemination with semen spiked with *T. gondii* tachyzoites (Consalter et al., 2017; Wanderley et al., 2013). On the other hand, the epidemiological significance of this route might be limited (Buxton, 1998). Similarly, milk may also pose a risk of infection to lambs or goat kids, as *T. gondii* DNA has been identified in milk samples from naturally infected ewes and goats (de Santana Rocha et al., 2015; Saad et al., 2018), and bioassay results in raw milk suggest its infective potential (Chiari and Neves, 1984; Dehkordi et al., 2013). However, it needs to be mentioned that the latter findings have been challenged and their epidemiological significance has been questioned (Dubey and Jones, 2014). Even if these alternative routes of transmission are possible in small ruminants, it still needs to be established, to which extent they contribute to infection.

2.2.3. Disease caused by *T. gondii* in naturally infected sheep and goats

It has been estimated that toxoplasmosis is responsible of 10 to 23% of ovine abortions in Europe or USA (Dubey, 2009b). Recent reports have shown that also in other regions of the world, as in the Middle East and South America, *T. gondii* infections are associated with 3 to 54% of ovine abortion (Table 1).

The only evident clinical sign associated with acquired toxoplasmosis (horizontal transmission), is a brief episode of fever and lack of appetite from about 3–6 days after infection and lasting usually for about 4–5 days, but sometimes also for up to 10 days (Buxton et al., 1982; Buxton et al., 1988; Castano et al., 2016; Dubey, 1981; Esteban-Redondo et al., 1999; McColgan et al., 1988). In contrast, congenital transmission has severe consequences for the fetus. Whether the trans-placental transmission causes the death of the fetus partly depends on the time of gestation when infection occurs. If the dam was infected in early gestation, at a time when the immune system of the fetus is still immature, vertical transmission commonly results in fetal death and resorption. However, when infection occurs at mid gestation, abortion or the delivery of a stillborn lamb are the most common outcomes, while dams infected late in gestation may deliver a stillborn or congenitally infected weak or clinically normal offspring (Buxton and Losson, 2007). Macroscopic lesions in cases of abortion are restricted to the cotyledons, where multifocal areas of necrosis, macroscopically visible as white foci of variable size are suggestive for toxoplasmosis (Buxton and Finlayson, 1986; Buxton et al., 1982). Microscopically, multifocal necrosis, commonly associated with the infiltration of non-purulent lymphoid cells, could be found in placentomes or fetal organs, mainly brain, liver or lung (Buxton and Finlayson, 1986).

2.2.4. Effects of experimental infections in sheep and goats

The precise mechanisms responsible for *T. gondii* abortion in small ruminants are not yet fully understood. The most recent studies, employing the oral route for administering oocysts as infective parasitic stage are summarized in Table 2. The outcome of experimental infections might be affected by the viability of oocysts which needs to be confirmed prior to use (Dubey,

Table 1
Reports of *Toxoplasma gondii* induced abortions in small ruminants since 2010.

Country	No. of placentas, fetuses and stillborn lambs examined (sheep/goats)	No. farms tested (sheep/goats)	No. of submissions (sheep/goats)	% positive, total or ovine/caprine	Diagnostic methods				Observations	Reference
					Other causes investigated	IHC	PCR	Fetal serology		
Brazil	35	n.a.	n.a.	14.3	No	No	Yes	No	Ovine abortions	de Moraes et al. (2011)
Great Britain	n.a. ^a	n.a.	n.a.	23.7	Yes ^b	n.a. ^a	Yes ^b	Yes ^b	Ovine abortions	Carson and Group (2017)
Iran	325	n.a.	n.a.	5	No	No	No	Yes	Ovine abortions	Razmi et al. (2010)
	18	n.a.	n.a.	66	No	No	Yes	No	Ovine abortions	Habibi et al. (2012)
	200	n.a.	n.a.	13.5	No	No	Yes	No	Ovine abortions	Rassouli et al. (2011)
	37	n.a.	n.a.	54	No	No	Yes	Yes	Ovine abortions	Danehchin et al. (2016)
Ireland		66	n.a.	17	Yes	Yes	Yes	Yes	Ovine abortions	Gutierrez et al. (2012)
Jordan	106 (66/40)	n.a.	n.a.	31	No	No	Yes	No	Ovine/caprine abortions	Abu-Dalbouh et al. (2012)
Netherlands	n.a.	n.a.	452 (282/170)	10.6/5.9	Yes	Yes	No	No	Ovine/caprine abortions	van den Brom et al. (2012)
	n.a.	n.a.	81 (57/24)	16.7/14.0	Yes	Yes	No	No	Ovine/caprine abortions	van Engelen et al. (2014)
Spain	100 (74/26)	n.a.	n.a.	5.4/3	Yes	No	Yes	No	Ovine/caprine abortions	Moreno et al. (2012)
Switzerland	30	n.a.	n.a.	10	Yes	No	Yes	No	Ovine/caprine abortions	Schnydrig et al. (2017)

^a n.a. = not applicable.

^b Based on <https://veterinaryrecord.bmj.com/content/vetrec/183/17/528.full.pdf> (last accessed 2019-01-22).

Table 2Experimental studies in sheep orally inoculated with *Toxoplasma gondii* oocysts or tissue cysts. Studies published since 2010.

Designation of <i>Toxoplasma gondii</i> isolate	Dose	Numbers and age category	Remarks	Reference
M4	3000	28 sheep	Inoculated with sporulated oocysts at day 90 of gestation	Gutierrez et al. (2010)
	500,000; 5000	16 lambs	Inoculated with sporulated oocysts.	Benavides et al. (2011)
	3000	9 sheep	Inoculated with sporulated oocysts at day 90 of gestation	O'Donovan et al. (2012)
	3000	15 sheep	Inoculated with sporulated oocysts at day 90 of gestation	Marques et al. (2012)
	2000; 500	24; 24 sheep	Inoculated with sporulated oocysts at day 90 of gestation (n = 24) and at day 120 (n = 24)	Castano et al. (2014)
	500 50	33 lambs 27 sheep	Inoculated with sporulated oocysts Inoculated with sporulated oocysts at three terms of gestation	Katzer et al. (2014) Castano et al. (2016)
PRU	400; 400; 100	36; 54; 33 sheep	Inoculated with sporulated oocysts at mid-gestation	Mevelec et al. (2010)
	3000 3000	13 lambs 4 sheep	Inoculated with tissue cysts Inoculated with tissue cysts	Verhelst et al. (2014) Verhelst et al. (2015)
P	200,000	4 rams	Inoculated with sporulated oocysts.	Lopes et al. (2011)
ME49; VEG	2500; 2500	20 sheep	Inoculated with sporulated oocysts at three terms of gestation of chronically infected ewes.	Dos Santos et al. (2016)
	500; 50; 10	5; 5; 5 sheep	Inoculated with sporulated oocysts at day 90 of gestation	Sánchez-Sánchez et al. (2019)
TgShSp1	500; 50; 10	6; 6; 6 sheep	Inoculated with sporulated oocysts at day 90 of gestation	Sánchez-Sánchez et al. (2019)

2010b). In addition, the *T. gondii* strain characteristics including the virulence, which seems to change after repeated passages (Saraf et al., 2017), need to be taken into account. Results of experimental infections have clearly shown that the gestational age, in particular the stage of maturation of the fetal immune system has an important effect on the pathogenesis (Buxton and Finlayson, 1986). In addition, the cellular immune response of the dam, mainly mediated by IFN- γ , is of importance in controlling the parasite multiplication (Innes et al., 1995). The experimental inoculation of sheep and goats has also helped to demonstrate that toxoplasmosis in small ruminants could also cause early abortion shortly after infection. In these early abortions, invasion and multiplication of the parasite in the placenta or fetus could not be demonstrated (Owen et al., 1998; Rosa et al., 2001). Although the cause of these early abortions was thought to be high fever or hormonal dysregulation (Dubey, 2010b), recent studies have shown that they are related to vascular lesions in the placenta and leukomalacia in the fetal brain (Castano et al., 2014). All together, these results suggest that the pathogenesis of early abortion is different from the classically described, which is based on the multiplication of the parasite and subsequent lesions in the placenta and target fetal organs. The mechanisms underlying the early abortions in this disease are still unknown. Bearing these observations in mind, there are still several gaps in the knowledge of small ruminant toxoplasmosis that warrant further characterization of the experimental models for ovine and caprine toxoplasmosis and investigation on how different variables, e.g. *T. gondii* strain or isolate virulence, previous immunization or individual susceptibility could affect the pathogenesis of this disease.

2.3. Chickens and other poultry

2.3.1. Prevalence in chickens and other poultry

Toxoplasma gondii infection in free-ranging poultry is an indicator of environmental contamination with *T. gondii* oocysts (Dubey, 2010b). Strains isolated from local poultry probably mirror the *T. gondii* genotypes prevailing in a region (Dubey, 2010a,b). The prevalence of *T. gondii* infections in poultry depends on a number of factors. The type of husbandry seems to be very important. In poultry originating from free-range or small backyard farms, the *T. gondii* prevalence is usually higher than in poultry kept indoors (Guo et al., 2015; Maksimov et al., 2011; Schares et al., 2017a; Yang et al., 2012).

In chickens, there is a number of recent articles summarizing the seroprevalence of antibodies to *T. gondii* (Bangoura et al., 2011; Deng et al., 2018; Dubey, 2010a; Guo et al., 2015; Shokri et al., 2017). Prevalence estimates are often not comparable among studies because different serological tests have been applied and sampled farms may differ for example in farm type and size, feed source, presence or absence of cats, rodent or bird control and water quality (Bangoura et al., 2011; Dubey, 2010a; Schares et al., 2017a). In some studies, a low specificity of the serological tests may have overestimated the seroprevalence or a low sensitivity may have led to an underestimation. Overall, the *T. gondii* seroprevalences in chickens ranged between 0 and 100% (Ayinmode and Olaosebikan, 2014; Bangoura et al., 2011; Deng et al., 2018; Dubey, 2010a; Matsuo et al., 2014).

Only few studies on *T. gondii* prevalence in turkeys have been published. Apparent seroprevalence varies largely between studies and ranges from 0% (Czech Republic), 11% (Brazil), 20% (Germany) to 59% (Egypt) (Bangoura et al., 2011; El-Massry et al., 2000; Harfoush and Tahoon Ael, 2010; Koethe et al., 2011; Sa et al., 2016).

The *T. gondii* seroprevalence in ducks and geese, as summarized for Poland, Czech Republic, Germany and Norway, varied between 1.7 and 21% in ducks and between 5.9 and 43% in geese (reviewed by Bangoura et al. (2011)). Only 3.5% of geese in Kazakhstan were seropositive (Bangoura et al., 2011). *T. gondii* seroprevalences reported for China in ducks and geese were in

the range of 6.3–13.9% (Cong et al., 2012; Yang et al., 2012). The highest seroprevalences in ducks were reported from Egypt (50–55%) (El-Massry et al., 2000; Harfoush and Tahooon Ael, 2010) and Malaysia (30%) (Puvanuesuaran et al., 2013).

2.3.2. Possible routes of infection in chickens and other poultry

Due to the ground feeding behavior of poultry, the oral ingestion of material or water contaminated with *T. gondii* oocysts is most likely the main route of infection (Dubey, 2010b). Water may be contaminated with *T. gondii* oocysts (Isaac-Renton et al., 1998). Thus, oocyst contaminations of water can be of particular importance as a source of infection for waterfowl. Infected rodents and other intermediate hosts on farms may serve as a reservoir (Schaes et al., 2017a). Poultry such as chickens, turkeys, ducks and geese are omnivorous, i.e. they also may feed on earthworms, cockroaches and other insects, which may harbor, or could be contaminated with oocysts (Bettiol et al., 2000; Ruiz and Frenkel, 1980; Wallace, 1971, 1972). In addition, poultry may predate on small rodents as putative carriers of *T. gondii* tissue cysts. In an experimental setting, turkeys became infected after inoculation with brains of chronically infected mice (Koethe et al., 2015) and also chickens fed tissue cysts became infected (summarized by Dubey (2010a)). There is, however, lack of information, to which extent such different routes of infection (i.e. infections via tissue cysts) are relevant under natural conditions. Vertical transmission of *T. gondii* in poultry has been discussed in the past, but extensive experiments in chickens indicated that this route of infection can be left outside the focus (Biancifiori et al., 1986; Boch et al., 1966).

2.3.3. Disease caused by *T. gondii* in naturally infected poultry

In general, chickens, turkeys, ducks and geese rarely develop clinical signs after infection with *T. gondii* (Dubey, 2010a). Worldwide, there are only few reports on clinical toxoplasmosis in naturally infected poultry (Dubey, 2010a,b). It has to be kept in mind, however, that some of the clinical cases reported as caused by *T. gondii* may have been triggered by other infections (e.g. viral) or complicated by other diseases (Dubey, 2010a).

No *T. gondii* genotype-dependent virulence in adult chickens has been recorded and even South-American strains, highly virulent to mice, seem to be avirulent in chickens (Hamilton et al., 2017; Vaudaux et al., 2010). However, it has to be mentioned here that young chickens (1-month-old), infected by oocysts of *T. gondii* Type I (GT1 strain) developed clinical toxoplasmosis, whereas those infected by oocysts of *T. gondii* Type II (ME49) did not develop any clinical signs (Dubey et al., 1993b). Chicken-line dependent differences in mortality after experimental inoculation of chicks with recombinant *T. gondii* clones suggested that, in addition to the parasites genotype, also genetic factors of the host may play an important role in the development of clinical toxoplasmosis in chickens (Schaes et al., 2017b). Age of the chicken seems to be a very because also in another study, 10-day old chickens showed mortality in a dose-dependent fashion (S. Wang et al., 2015).

Reports of toxoplasmosis in magpie (*Anseranas semipalmata*) and Hawaiian geese (*Branta sandvicensis*) (Dubey et al., 2001; Work et al., 2015) suggest that there might be differences in susceptibility for *T. gondii* infection and toxoplasmosis between different species of Anseriformes. By contrast, we could not find reports on clinical toxoplasmosis in domestic geese (*Anser anser*).

2.3.4. Effects of experimental infections in livestock poultry

The susceptibility of poultry to experimentally induced toxoplasmosis depends on the infectious dose, the parasite strain, stage, the route of infection and, as mentioned above, the age of the animal (Dubey et al., 1993b; Schaes et al., 2017b). In chickens, parenteral infection with *T. gondii* tachyzoites or oral infection with oocysts rarely cause clinical signs (Dubey, 2010a). However, in the case of intracranial infections using encysted *T. gondii*, the animals developed severe cerebral toxoplasmosis (Bickford and Saunders, 1966; Dubey, 2010a).

No clinical toxoplasmosis was reported in turkeys either after experimental oral infection with different doses of *T. gondii* oocysts (Bangoura et al., 2013; Dubey, 2010b; Dubey et al., 1993a), or intravenous inoculation with varying doses of tachyzoites (with strains representative for *T. gondii* clonal Types I, II and III) (Maksimov et al., 2018; Zöller et al., 2013).

The results of experimental infections in ducks and geese with oocysts (Bartova et al., 2004; Maksimov et al., 2011) and intravenous infections with tachyzoites (Maksimov et al., 2011) showed that also these animal species were resistant against clinical toxoplasmosis regardless of the parasite stage used for infection.

2.4. Cattle

2.4.1. Prevalence in cattle

Seroprevalence estimates in cattle, if obtained by highly specific tests, can be of value to monitor the exposure of cattle to *T. gondii*. However, these serological data must be interpreted with care as studies conducted with bioassay experiments suggest that in the vast majority of seropositive cattle there was no evidence for the presence of viable *T. gondii* infections. This has been also shown by analyses of naturally exposed animals, in some studies with very large populations of cattle (Boch et al., 1965a; Dubey et al., 2005; Dubey and Streitl, 1976; Fortier et al., 1990; Freyre et al., 1990; Jacobs and Moyle, 1963; Passos et al., 1984). There are only a few reports on naturally exposed cattle, in which positive *T. gondii* bioassays indicated viable infections (Arias et al., 1994; Catar et al., 1969; de Macedo et al., 2012; Dubey, 1992; Jacobs et al., 1960). This is in a sharp contrast to the findings in small ruminants as discussed above.

With the advent of new methodologies, i.e. genome detection by PCR, a number of studies utilizing these techniques yielded very high proportions (up to 10 or 20%) of *T. gondii* genome positive samples in cattle tissues (Amdouni et al., 2017; Azizi et al., 2014; Berger-Schoch et al., 2011; Campo-Portacio et al., 2014; Ergin et al., 2009; Hosein et al., 2016; Mahami-Oskouei et al., 2017;

Opsteegh et al., 2011; Rahdar et al., 2012; Wyss et al., 2000). Keeping in mind the failure of many large-scale studies to find viable *T. gondii* in bovine tissues, the validity of these reports on the detection of *T. gondii* genome fragments has to be questioned. Detection of genome fragments of *T. gondii* in absence of positive bioassays should not be regarded as conclusive.

A recent meta-analysis revealed the possibility of geographic differences in the proportion of *T. gondii*-positive cattle. A significantly higher proportion of positive cattle was found in Central America as compared to North America (Belluco et al., 2016). This may indicate that the susceptibility of cattle to *T. gondii* is influenced by the genotype of the parasite, which largely varies in different regions of the world (Bertranpetit et al., 2017; Chaichan et al., 2017; Shwab et al., 2014). However, these considerations are hypothetical and need to be addressed in future studies. In addition, differences in husbandry conditions, hygienic situations, in climate and in other factors may affect the extent, to which cattle from different regions are exposed to *T. gondii*. Therefore, there is a need to confirm the detection of *T. gondii* genome positive samples in cattle by additional experiments, thus assessing the presence of viable parasites.

Over the past decades, numerous articles have been published on the seroprevalence of *T. gondii*-specific antibodies in taurine cattle. Many of these publications have been reviewed before, with a global scope (Dubey, 1986a, 2010b; Tenter et al., 2000) or, more recently, by focusing on the situation in particular regions of the world like China (Deng et al., 2018), South Asia (Khan et al., 2017) and Africa (Tonouhewa et al., 2017). Overall, these summaries show a large variation in the reported proportions of positive findings and the summarizing estimates were 9.5% for cattle in China (Deng et al., 2018), 27.9% in South Asia (Khan et al., 2017) or 12% in Africa (Tonouhewa et al., 2017).

2.4.2. Possible routes of infection in cattle

It is generally accepted that most cattle become infected orally, through ingestion of feed or water contaminated with *T. gondii* oocysts. Many experimental infections in cattle, especially earlier ones, used oocysts as the inoculum, thus demonstrating that cattle are susceptible to this infective stage (Burrells et al., 2018; Costa et al., 2011; de Oliveira et al., 2001; Dubey and Thulliez, 1993; Esteban-Redondo et al., 1999; Stalheim et al., 1980). However, usually large numbers of oocysts were administered, but we are not aware of any experiments that aimed at establishing the minimum infective dose for cattle.

There are also reports on bovine infections with viable *T. gondii* caused by oral inoculation with *T. gondii* tissue cysts (10^3) (Costa et al., 2011) or brains of chronically infected mice (Rommel et al., 1966). Although cattle are herbivores, infections through accidental ingestion of tissue cysts may occur, i.e. if cattle feed is contaminated with fresh tissue of an infected intermediate host.

In infection with *Neospora caninum*, an apicomplexan parasite closely related to *T. gondii*, vertical transmission after acute or chronic infection is of utmost importance in cattle (Dubey et al., 2017). However, for *T. gondii* the situation seems to be different. Although there are reports on the detection of *T. gondii* genome fragments in aborted bovine fetuses (Ellis, 1998; Gottstein et al., 1998), the isolation of viable *T. gondii* parasites from bovine fetuses was achieved only occasionally (Canada et al., 2002; Costa et al., 1977) or not at all (Conrad et al., 1993). Experimental inoculations with tachyzoites resulted in abortion or vertical transmission (Stalheim et al., 1980; Wiengcharoen et al., 2011), but the epidemiological significance of these findings is not clear, because the presence of *T. gondii* in the aborted fetuses was not confirmed. Overall, if vertical transmission of *T. gondii* naturally occurs in cattle, it seems to be a rare event. However, the large genetic variability between *T. gondii* populations worldwide should be kept in mind, which may result in a variety of biological traits that may also include differences in virulence in cattle. In the light of this variation, findings in North America and Europe with isolates prevailing in these regions should not be generalized without confirmation.

2.4.3. Disease caused by *T. gondii* infections in cattle

Reports on clinical toxoplasmosis in naturally infected cattle are rare. This suggests that cattle are resistant to infection and to clinical toxoplasmosis. Although clinical signs and histological alterations were recorded after experimental infection, natural cases of clinical toxoplasmosis in cattle comprised only of abortions in association with the isolation of *T. gondii* from the aborted fetuses (Canada et al., 2002). It is not clear, however, whether the infection with *T. gondii* had caused the abortions.

2.4.4. Effects of experimental infections in cattle

After experimental infection, febrile reactions starting 2 days post inoculation at the earliest and lasting up to 15 days p.i. have been regularly reported (Burrells et al., 2018; Costa et al., 1977; Esteban-Redondo et al., 1999; Munday, 1978; Rommel et al., 1966; Stalheim et al., 1980; Wiengcharoen et al., 2011). Bovine infection with *T. gondii* regularly leads to a parasitemia, which seems to be responsible for the elevated temperatures observed in inoculated animals shortly after infection (de Oliveira et al., 2001; Stalheim et al., 1980). In one study, the parasitemia was even recorded up to 62 days p.i. (Costa et al., 1977). In addition, respiratory distress, nasal discharge, and hyperemia of the conjunctivae were reported in the latter study (Costa et al., 1977).

Reports on mortality in inoculated animals are rare. It occurred in cases of calves inoculated with oocysts or intravenously with tachyzoites, but only in the latter the infection was confirmed (Stalheim et al., 1980). In another experiment, two out of four dams inoculated with *T. gondii* tachyzoites became recumbent and were euthanized (Wiengcharoen et al., 2011). In this case, adult cattle were affected a long time after inoculation (2 to 3 month p.i.) and this finding represented a surprising exception among a series of experiments, where inoculated cattle developed no or only mild clinical signs (Beverley et al., 1977; Burrells et al., 2018; Costa et al., 1977; Dubey, 1983; Esteban-Redondo et al., 1999; Munday, 1978; Rommel et al., 1966).

2.5. Horses and other equids

2.5.1. Prevalence in horses and other equids

A relatively large number of studies report on the seroprevalence of antibodies against *T. gondii* in horses, mules and donkeys world-wide. Most but not all of the publications have been reviewed previously (Dubey, 2010b; Dubey et al., 2014b; Tassi, 2007; Tenter et al., 2000). The study results are difficult to compare because different, not always validated serological methods and various cut-offs have been applied. In addition, the equids selected for the studies differed largely in number, age, origin and the purpose for keeping the animals. Currently, there is no reference standard available to validate serological tests in horses properly. A recent attempt to correlate serological test results (i.e. results by MAT and a commercially available ELISA) with those of *T. gondii* PCR on horse meat samples largely failed (Aroussi et al., 2015). There was almost no correlation between the serological data and *T. gondii* genome detection using a highly sensitive magnetic capture PCR (Aroussi et al., 2015). Nevertheless, there is no doubt that horses can harbor viable *T. gondii*, which could be isolated from tissues of both, naturally (Evers et al., 2013; Gennari et al., 2015; Klun et al., 2017; Shaapan and Ghazy, 2007) or experimentally infected animals (Al-Khalidi et al., 1980; Dubey, 1985). The results indicated that viable *T. gondii* can persist in edible tissues up to 476 days after infection (Dubey, 1985). In addition, imported meat from infected horses was suspected as cause of toxoplasmosis in France (Elbez-Rubinstein et al., 2009; Pomares et al., 2011). A recent example on isolation of viable *T. gondii* from horse shows that truly infected horse may remain seronegative or develop only a low specific antibody titre in particular serological tests such as the MAT (Klun et al., 2017). Currently, serological responses in equids do not seem to be reliable indicators for viable infections; this is similar to the situation in cattle. Nevertheless, positive antibody responses indicate the exposure of equids to *T. gondii* and could thus be used to identify risk factors for their exposure to *T. gondii*. Reported seroprevalence for equids range in South America from 3 to 90% (Cazarotto et al., 2016; Costa et al., 2012; Dangoudoubiyam et al., 2011; de Oliveira et al., 2013; Dubey, 2010b; Evers et al., 2013; Finger et al., 2013; Gennari et al., 2015; Portella et al., 2017; Ribeiro et al., 2016; Tassi, 2007; Venturi et al., 2017), in North America from 0 to 73% (Alvarado-Esquivel et al., 2015; Alvarado-Esquivel et al., 2012b; Dubey, 2010b; Dubey et al., 2014b; James et al., 2017; Schale et al., 2018; Tassi, 2007), in Europe from 0 to 55% (Bartova et al., 2015; Dubey, 2010b; Garcia-Bocanegra et al., 2012; Karatepe et al., 2010; Klun et al., 2017; Kouam et al., 2010; Machacova et al., 2014; Mancianti et al., 2014; Papini et al., 2015; Pastiu et al., 2015; Tassi, 2007; Zhou et al., 2017), in Asia from 0 to 71% (Aharonson-Raz et al., 2015; Alanazi and Alyousif, 2011; Dubey, 2010b; Hajjalilo et al., 2010; Lee et al., 2014; Mancianti et al., 2014; Masatani et al., 2016; Matsuo et al., 2014; Miao et al., 2013; Raeghi et al., 2011; Razmi et al., 2016; Saqib et al., 2015; Tassi, 2007; Tavalla et al., 2015; J.L. Wang et al., 2015; Yang et al., 2013), in Africa from 14 to 45% (Ayinmode et al., 2014; Boughattas et al., 2011; Haridy et al., 2010) and 2% in Australia (Tassi, 2007).

2.5.2. Possible routes of infection in horses and other equids

In the case of equids, oral infection by oocysts is the most probable route as it has been confirmed by a number experimental infections using different doses of oocysts (Al-Khalidi et al., 1980; Altan et al., 1977; Dubey, 1985; Marques et al., 1995), i.e. doses of 10^5 (Al-Khalidi et al., 1980), 10^4 (Dubey, 1985), 10^6 (Altan et al., 1977), or up to 1.5×10^5 (Marques et al., 1995).

Rodents are intermediate hosts of *T. gondii* and regarded as a source of infection especially in omnivorous animals like pigs. Since it has been demonstrated that a large proportion of horses would eat meat and may become infected by *Trichinella spiralis* via this route (Murrell et al., 2004), it is tempting to speculate that the oral ingestion of carcasses of *T. gondii* infected rodents or other small intermediate hosts may represent another potential source of infection for equids.

Reports on transplacental *T. gondii* transmission in experimentally infected mares (Marques et al., 1995) have to be interpreted carefully and need further investigation because infections with other related parasites like *Sarcocystis neurona* or *N. caninum* need to be ruled out.

2.5.3. Disease caused by *T. gondii* infections in horses and other equids

The general view is that toxoplasmosis, i.e. disease caused by *T. gondii* infection, is rather rare in equids, after both natural and experimental infection (Dubey, 2010b; Tassi, 2007). *T. gondii* DNA has been detected in the eyes of an aborted foal (Turner and Savva, 1992) and in the placenta of a mare that foaled normally (Turner and Savva, 1990). Together with the transplacental transmission reported in an experimental study, these results may indicate that *T. gondii* could be occasionally involved in equine abortion (Marques et al., 1995), but further studies are necessary to clarify this issue. It was recently discussed if *T. gondii* is involved in equine protozoal myeloencephalitis (EPM) (James et al., 2017; Schale et al., 2018). A case control study conducted in California found an association between high levels of *T. gondii* IFAT titres ($1:\geq 160$ or $1:\geq 320$) and the presence of neurologic signs compatible with EPM (James et al., 2017). Another study, not designed as a case-control study but thoroughly assessing EPM cases, revealed only low proportions of *T. gondii* (and also *Neospora* sp.) positive animals in this group of patients, contradicting an involvement of *T. gondii* in EPM. In this study, *S. neurona* was identified as the most probable cause of EPM.

An earlier study conducted in the UK reported on the presence of *T. gondii* DNA in the eye of a pony (Turner and Savva, 1991). The significance of this finding and the involvement of *T. gondii* as a possible cause of blindness in horse are unknown. However, after experimental infection of a pony with *T. gondii*, the infection was also observed in the eye (Dubey, 1985).

2.5.4. Effects of experimental infections in horses

In equids experimentally inoculated with large numbers of oocysts (10^4), mild fever was observed in few animals, while the others remained clinically normal (Dubey, 1985; Dubey and Desmonts, 1987). In ponies orally inoculated with a high number of

oocysts and in addition immunosuppressed by corticosteroid, 8 out of 9 ponies developed fever between days 2 and 15 p.i. (Al-Khalidi et al., 1980). In an earlier study, ponies, not immunosuppressed but orally inoculated by 10^6 oocysts did not develop clinical signs (Altan et al., 1977). Horses inoculated intravenously with tachyzoites (i.e. 3.28×10^7 or 2.19×10^7 , strain RH) developed fever between 4 to 8 days after inoculation and ocular discharge from 10 to 26 days post inoculation (Sposito Filha et al., 1992).

3. Potential risk factors for infection in livestock

Raw and undercooked meat are regarded as one of the main sources of *T. gondii* infections for humans. Knowledge on risk factors for the infection with *T. gondii* in livestock and an assessment of the importance of these risk factors is essential to ensure safe food and intervene effectively. This section is partially based on previous reviews (Guo et al., 2015; Opsteegh et al., 2016) and was extended including the most recent reports in the field. We present a brief overview on the existing literature, but no deeper meta-analysis. We chose to categorize the literature data on individual risk factors only into “statistically significant” and “not statistically significant”, not discriminating between the statistical testing methods used. Mainly reports were included, in which risk analysis was based on the seropositivity of livestock animals. In addition, the review was restricted to reports on the main livestock species, i.e. pigs, small ruminants (sheep and goats), cattle, equids (horses, donkey and mules) and poultry (chickens, ducks and geese).

To identify specific risk factors for infection, it is crucial to know the most important routes, by which livestock can acquire the infection. These routes include oocyst contaminations of feed, water or the environment, and the ingestion of tissues of infected intermediate hosts like rodents (Fig. 1).

The risk of infection and particularly of infection routes on individual farms (Fig. 1) are influenced by several indirectly acting factors. They include factors related to particular rearing systems of specific livestock groups. There are no general rules to define or to assess such risk factors, which makes a comparison of the results between studies difficult or even impossible. Although the spectrum of risk factors reported in the literature is very heterogenic, we tried to categorize them as much as possible to come to some more general conclusions. In our analysis, we looked for the basic, overarching factors and stratified the records by livestock species. If both, univariable and multivariable statistical analyses had been reported for a specific factor, we included only the results of multivariable statistical analyses, i.e. the results of analyses, in which the specific factor had been examined together with at least one other factor. Because almost all of the studies included were retrospective studies it has to be kept in mind that associations or statistical effects of factors identified by this type of studies are not entirely conclusive and only allow the generation of hypotheses.

3.1. General factors

3.1.1. Age

The association of age with the risk of a *T. gondii* infection has been examined in a number of studies (Table 3). Generally, the wide spread of *T. gondii* and the extremely broad host spectrum of the parasite leads to a time of exposure to the infective stages of the parasite that is proportional to the age of the animal. The age of the animals often depends on the production type, i.e. meat-producing animals are usually slaughtered at younger age, whereas animals reared for dairy production, reproduction or recreational purposes often live much longer. Overall, literature data confirm the expected association with age, i.e. in most studies age appeared as a risk factor for infection with *T. gondii* (Table 3). Due to its general importance, the factor “age” should be




Fig. 1. *Toxoplasma gondii* life cycle highlighting conditions of horizontal transmission concerning livestock infection.

Table 3The effect of age on the seropositivity for *T. gondii* in livestock species.

Factor	Species	Statistically significant	Not statistically significant
Higher age	Pigs	17, 20, 25, 34, 33, 39, 53, 57 (p), 75, 80, 86, 94	6, 28, 73, 96
	Sheep	13, 15, 23, 35, 36, 41, 42, 45, 46, 50, 57, 70, 85, 88, 90	7, 68, 72, 92, 95, 98
	Goats	3, 8, 19 (p), 24, 30, 35, 47, 57, 68, 88	92, 95
	Cattle	–	79, 95, 99
	Equids	102, 104	5, 11, 31, 40, 54, 100, 101, 103
Finishing period	Chicken	4, 93	–
	Pigs	75	–
Age below 12 months	Sheep	16 (p)	–
Younger age	Cattle	38, 79	53
Age <24 months	Cattle	48 (p)	–
Age >24 months–96 months + >120 months (dairy and mixed dairy)	Cattle	48	–
Age >48 months–72 months (beef and mixed beef)	Cattle	48	–

(p) = protective factor; coding of references: 3 = Alvarado-Esquivel et al. (2011), 4 = Alvarado-Esquivel et al. (2012a), 5 = Alvarado-Esquivel et al. (2012b), 6 = Alvarado-Esquivel et al. (2014), 7 = Alvarado-Esquivel et al. (2013a), 8 = Alvarado-Esquivel et al. (2013b), 11 = Cazarotto et al. (2016), 13 = Andrade et al. (2013), 15 = Cosendey-KezenLeite et al. (2014), 16 = D'Alencar Mendonca et al. (2013), 17 = Damriyasa et al. (2004), 19 = de Moura et al. (2016), 20 = de Sousa et al. (2014), 23 = Deksne et al. (2017), 24 = Deng et al. (2016), 25 = Djokic et al. (2016), 28 = Esteves et al. (2014), 30 = Garcia et al. (2012), 31 = Garcia-Bocanegra et al. (2012), 33 = Garcia-Bocanegra et al. (2010a), 34 = Garcia-Bocanegra et al. (2010b), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 38 = Gilot-Fromont et al. (2009), 39 = Goerlich (2011), 40 = Guerra et al. (2018), 41 = Guimaraes et al. (2013), 42 = Hammond-Aryee et al. (2015), 45 = Holec-Gasior et al. (2015), 46 = Hutchinson et al. (2011), 47 = Iovu et al. (2012), 48 = Jokelainen et al. (2017), 50 = Katzer et al. (2011), 53 = Klun et al. (2006), 54 = Kouam et al. (2010), 57 = A.P. Lopes et al. (2013), 68 = Rego et al. (2016), 70 = Romanelli et al. (2007), 72 = Sakata et al. (2012), 73 = Santoro et al. (2017), 75 = Schulzig and Fehlhaber (2005), 79 = Tan et al. (2015), 80 = Tao et al. (2011), 85 = Vesco et al. (2007), 86 = Villari et al. (2009), 88 = Xu et al. (2015), 90 = Zhang et al. (2016), 92 = Zou et al. (2015), 93 = Schares et al. (2017a), 94 = Samico-Fernandes et al. (2017), 95 = Tilahun et al. (2018), 96 = Bawm et al. (2016), 98 = Yin et al. (2015), 99 = Schoonman et al. (2010), 101 = Bartova et al. (2017), 102 = Machacova et al. (2014), 103 = Miao et al. (2013), 104 = Ribeiro et al. (2016).

always included as one of the explanatory variables, if risk factor analyses on postnatal infection among animals with varying age are performed. Age is prone to act as a confounder or effect modifier in statistical analyses (Schaes et al., 2017a).

3.1.2. Gender

The gender of livestock animals as a putative risk factor has been studied only occasionally (Table 4). Experimental studies in mice and guinea pigs showed a higher susceptibility of females to infection with *T. gondii* (Kittas and Henry, 1979, 1980; Roberts et al., 1995; Roberts et al., 2001). In most of the published studies on livestock recorded, a significant effect for female animals to be serologically positive for *T. gondii* was not detected. Nevertheless, with the exception of two studies, in which males showed an increased risk, females were more frequently seropositive in a few studies conducted with pigs, sheep and goats or equids (Table 4). Whether these apparent associations were in fact related to gender or to other underlying factors, e.g. the way animals of different gender are reared, needs to be questioned. It must be noted that gender frequently shows up as a confounder in epidemiological studies because “gender” may mask these underlying factors (Thrusfield, 2007).

3.1.3. Geographic and regional characteristics

For all species taken into consideration in this review, there were studies reporting on significant differences in seroprevalence with respect to regions or geographic characteristics of the farm locations (Table 5). Many region- and geography-related variables that could possibly affect the survival and presence of *T. gondii* or the exposure of animals to the parasite must therefore

Table 4The effect of female gender on the seropositivity for *T. gondii* in livestock species.

Factor	Species	Statistically significant	Not statistically significant
Female gender	Pigs	73	6, 20, 28, 43, 53, 86, 94
	Sheep	15 (p), 36, 45 (p), 68, 95	7, 23, 35, 41, 59, 72, 81, 88, 90, 92, 98
	Goats	19, 30, 68	8, 35, 81, 88, 92, 96
	Cattle	–	21, 53
	Equids	102, 103	5, 11, 31, 40, 54, 100, 101

(p) = protective factor; coding of references: 5 = Alvarado-Esquivel et al. (2012b), 6 = Alvarado-Esquivel et al. (2014), 7 = Alvarado-Esquivel et al. (2013a), 8 = Alvarado-Esquivel et al. (2013b), 11 = Cazarotto et al. (2016), 15 = Cosendey-KezenLeite et al. (2014), 19 = de Moura et al. (2016), 20 = de Sousa et al. (2014), 21 = de Souza et al. (2016), 23 = Deksne et al. (2017), 28 = Esteves et al. (2014), 30 = Garcia et al. (2012), 31 = Garcia-Bocanegra et al. (2012), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 40 = Guerra et al. (2018), 41 = Guimaraes et al. (2013), 43 = Herrero et al. (2016), 45 = Holec-Gasior et al. (2015), 53 = Klun et al. (2006), 54 = Kouam et al. (2010), 59 = Magalhaes et al. (2016), 68 = Rego et al. (2016), 72 = Sakata et al. (2012), 73 = Santoro et al. (2017); 81 = Tzanidakis et al. (2012), 86 = Villari et al. (2009), 88 = Xu et al. (2015), 90 = Zhang et al. (2016), 92 = Zou et al. (2015), 94 = Samico-Fernandes et al. (2017), 95 = Tilahun et al. (2018), 96 = Bawm et al. (2016), 98 = Yin et al. (2015), 100 = Almeida et al. (2017), 101 = Bartova et al. (2017), 102 = Machacova et al. (2014), 103 = Miao et al. (2013).

Table 5
The effect of geographic parameters on the seropositivity for *T. gondii* in livestock species.

Factor	Species	Statistically significant	Not statistically significant
Region, province, municipality, prefecture or district	Pigs	17, 20, 22, 53, 67	6
	Sheep	15, 32, 35, 45, 53, 50, 59, 69, 83, 90, 92, 98, 106	23, 81, 88, 95
	Goats	8, 19, 26, 32, 35, 47, 92	81, 88, 95, 107
	Cattle	32, 48, 53, 95	78, 79
	Equids	40, 54, 101	31
	Chicken	4	89
Altitude	Pigs	6, 86	–
	Sheep	2, 7, 35, 36, 49, 77	14, 81
	Goats	8, 49	35, 81
Mean monthly temperatures	Pigs	34	6
	Sheep	7	–
Mean annual rainfall	Pigs	34	6
	Sheep	7	–
Climate	Pigs	6	–
	Goats	8	–
Relative humidity	Pigs	34	–
	Hills relative to plains	Pigs	–
Generalized land cover	Sheep	49	–
	Goats	49	–
Distance to next village	Sheep	–	81
	Goats	–	81
Rural environment relative to urban environment	Sheep	–	95
	Goats	–	95
	Cattle	–	95
	Equids	5	–
Terrain waterlogged (versus rough and flat)	Sheep	69	–
	Goats	3	–

Coding of references: 2 = Alvarado-Esquivel et al. (2013a), 3 = Alvarado-Esquivel et al. (2011), 4 = Alvarado-Esquivel et al. (2012a), 5 = Alvarado-Esquivel et al. (2012b), 6 = Alvarado-Esquivel et al. (2014), 7 = Alvarado-Esquivel et al. (2013b), 8 = Alvarado-Esquivel et al. (2013c), 14 = Condoleo et al. (2016), 15 = Cosendey-KezenLeite et al. (2014), 17 = Damriyasa et al. (2004), 19 = de Moura et al. (2016), 20 = de Sousa et al. (2014), 22 = Deksne and Kirjusina (2013), 23 = Deksne et al. (2017), 26 = Djokic et al. (2014), 31 = Garcia-Bocanegra et al. (2012), 32 = Garcia-Bocanegra et al. (2013), 34 = Garcia-Bocanegra et al. (2010b), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 40 = Guerra et al. (2018), 45 = Holec-Gasior et al. (2015), 47 = Iovu et al. (2012), 48 = Jokelainen et al. (2017), 49 = Kantzoura et al. (2013), 50 = Katzer et al. (2011), 53 = Klun et al. (2006), 54 = Kouam et al. (2010), 59 = Magalhaes et al. (2016), 67 = Pastiu et al. (2013), 69 = Rizzo et al. (2017), 77 = Skjerve et al. (1998), 78 = Sun et al. (2015), 79 = Sun et al. (2015), 80 = Tao et al. (2011), 81 = Tzanidakis et al. (2012), 83 = Verhelst et al. (2014), 86 = Villari et al. (2009), 88 = Xu et al. (2015), 90 = Zhang et al. (2016), 92 = Zou et al. (2015), 95 = Tilahun et al. (2018), 98 = Yin et al. (2015), 101 = Bartova et al. (2017), 106 = Jokelainen et al. (2010), 107 = Shuralev et al. (2018).

be taken into account here. Few studies not only evaluated the differences between certain regions, but also looked into more details concerning the most likely underlying variables, such as mean temperatures, mean rainfall, humidity, altitude or terrain characteristics (Table 5). Most of them found a statistically significant influence of these parameters on the proportion of

Table 6
Production system as a putative risk factor for *T. gondii* seropositivity in livestock.

Factor	Species	Statistically significant	Not statistically significant
Intensive	Pigs	9 (p), 20 (p), 28, 37 (p), 51 (p), 75 (p), 82 (p)	94
	Sheep	15 (p), 35 (p), 61, 69 (p), 81, 88 (p), 90 (p)	18, 57
	Goats	35 (p), 61, 68 (p), 81	–
Semi-intensive	Sheep	2, 35 (p), 68, 69, 81	36
	Goats	30, 35, 81	–
Extensive/animal friendly/organic/transhumance	Pigs	22, 51, 105	94
	Cattle	59, 78	21, 97
	Sheep	35, 58, 68, 88	14, 56
	Goats	35 (p), 68, 88	–
Backyard	Pigs	67	–
	Goats	47	–
	Chicken	4, 89, 91	–

(p) = protective factor; coding of references: 2 = Alvarado-Esquivel et al. (2013a), 4 = Alvarado-Esquivel et al. (2012a), 9 = Assadi-Rad et al. (1995), 14 = Condoleo et al. (2016), 15 = Cosendey-KezenLeite et al. (2014), 18 = de Barros Correia et al. (2015), 20 = de Sousa et al. (2014), 21 = de Souza et al. (2016), 22 = Deksne and Kirjusina (2013), 28 = Esteves et al. (2014), 30 = Garcia et al. (2012), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 37 = Gebreyes et al. (2008), 47 = Iovu et al. (2012), 51 = Kijlstra et al. (2004), 56 = Liu et al. (2015), 57 = A.P. Lopes et al. (2013) and W.D. Lopes et al. (2013), 58 = Lopes et al. (2010), 59 = Magalhaes et al. (2016), 61 = Mainar et al. (1996), 67 = Pastiu et al. (2013), 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 75 = Schulzig and Fehlhaber (2005), 78 = Sun et al. (2015), 81 = Tzanidakis et al. (2012), 82 = van der Giessen et al. (2007), 86 = Villari et al. (2009), 88 = Xu et al. (2015), 89 = Xu et al. (2012), 90 = Zhang et al. (2016), 91 = Zhu et al. (2008), 94 = Samico-Fernandes et al. (2017), 97 = Fajardo et al. (2013), 105 = Wallander et al. (2016).

seropositivity in livestock. Consequently, it is important to take regional differences into consideration but the underlying true effect factors such as climatic factors or variables related to differences in animal husbandry need to be included.

3.1.4. Farm management

3.1.4.1. Production system. The production systems, in which livestock is kept, may have a major impact on the risk for *T. gondii* infection (Table 6). However, the association of seropositivity in livestock within a particular production system provides no clear picture on the routes by which the animals became infected. Production systems are often related to specific conditions under which the animals are reared, fed or provided with water. These conditions may influence the likelihood of a contamination of feed, water or farmland with oocysts of *T. gondii* and the possibility of contact with the matrices mentioned above or with other infected intermediate hosts, e.g. rodents. In an intensive production system for example, the level of confinement is very high, at least for the respective livestock species, and thus, exposure of the animals to infective stages of the parasite is presumably lower as compared to extensive or other production system, where the animals have access to outdoor facilities. Intensive farming usually requires storing supplements. If these materials are stored open or in places where they may attract rodents or cats, additional routes of transmission may become relevant. Contaminated supplements may represent one explanation why intensive farming was not associated with a protective statistical effect in some studies (Table 6).

3.1.4.2. Specific farming conditions. It was observed in several studies that differences in farming conditions, which are often livestock species-specific, were statistically significantly associated with differences in the risk of infection with *T. gondii* for various livestock species (Table 7). However, many of these associations are difficult to explain. Like production systems, livestock species-specific farming conditions may influence the probability of contamination of feed, water or farmland with *T. gondii* oocysts. This holds also true for other potential sources of exposure to *T. gondii*, e.g. the presence of infected intermediate hosts such as rodents. Specific conditions related to more extensive farming may also represent risk factors for seropositivity in livestock (Table 7).

3.1.4.3. Herd and flock size. Herd or flock size is also related to the management and production system. Larger herds are more likely to be intensively managed. It can be assumed that farms with smaller herds often have a lower level of specialization and might be less confined. As farms applying animal welfare-oriented production techniques need more space to rear animals, this may restrict the size of the herd or flock. Again, herd or flock size may have an often un-explained link to conditions that can influence the probability of exposure of livestock to *T. gondii* due to contamination of feed, water or farmland with oocysts and contact to other infected intermediate hosts, e.g. rodents. There is a clear tendency showing that the smaller the herd or flock, the higher is the chance of seropositivity (Table 8).

3.1.4.4. Presence of other animal species and contact to other herds. The presence of several animal species on a farm or other animals kept close to the livestock species may serve as an indicator of low farming intensity. Even contact to or mixing with other herds

Table 7
Livestock species-specific parameters as putative risk factors for *T. gondii* seropositivity.

Species	Factor	Statistically significant	Not statistically significant
Pigs	Complete production cycle performed on farm (farrow-to-finish)	17 (p), 34, 55 (p), 86 (p)	–
	Only part of production cycle performed on farm	17, 53, 55	–
	Origin of replacement sows, Source of pigs (own farm versus outside)	–	17, 55
Cattle	Mixed farming	–	78
	Beef farm (relative to dairy and mixed)	48 (p)	32, 78
Sheep	Feeder/stocker/backgrounder (versus feeder/stocker)	21	–
	Purpose subsistence (versus breeding/rebreeding/fattening)	69	–
	Purpose (meat, milk, mixed)	–	68
Goats	Mixed (milk, meat)	14	–
	Additional uses to dairy	26	–
	Purpose meat (versus genetic enhancement)	68	–
Chicken	Purpose (milk/meat/mix)	–	19, 68
	Breeders	89	–
	Layers	89	–
Equids	Broilers	89 (p)	–
	Racing	–	54
	Recreation	–	54
	Farming	54	–
	Use (breeding versus ceremonial, research or sports) ^a	101	–

(p) = protective factor; coding of references: 14 = Condoleo et al. (2016), 17 = Damriyasa et al. (2004), 19 = de Moura et al. (2016), 21 = de Souza et al. (2016), 26 = Djokic et al. (2014), 32 = Garcia-Bocanegra et al. (2013), 34 = Garcia-Bocanegra et al. (2010a,b), 48 = Jokelainen et al. (2017), 53 = Klun et al. (2006), 54 = Kouam et al. (2010), 55 = Limon et al. (2017), 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 78 = Sun et al. (2015), 80 = Tao et al. (2011), 86 = Villari et al. (2009), 89 = Xu et al. (2012), 101 = Bartova et al. (2017).

^a Statistics not conclusive.

Table 8The effect of herd size on the seropositivity for *T. gondii* in livestock species.

Factor	Species	Statistically significant	Not statistically significant
Herd/flock size (numbers of animals as continuous variable)	Pigs	–	73
	Sheep	35, 46, 81	14, 32, 45, 56
	Goats	35 (p), 81 (p)	32, 56
	Cattle	–	78
	Equids	5	–
Small herd/flock size	Pigs	9, 25, 43, 55, 65, 86	–
	Sheep	12, 36, 76, 88	95
	Goats	49, 88	95
	Cattle	38, 53, 95, 99 (p)	–
	Chicken	93	–
	Equids	104	102
Large flock size	Sheep	1, 18, 23	–
	Goats	1	–
Higher number of sows	Pigs	25	–
Increased breeding density	Pigs	80	–
Dairy herds with a size ≥40–105 and >384 animals versus herds with a size <40 animals	Cattle	48	–
Mixed beef herd with a size of <50–200 animals versus herds with a size of >400 animals	Cattle	48	–
Flock size 150–300 versus >300 animals	Sheep	49 (p)	–
Low animal density in herd	Cattle	32	–

(p) = protective factor; coding of references: 1 = Abu-Dalbouh et al. (2012), 5 = Alvarado-Esquivel et al. (2012b), 9 = Assadi-Rad et al. (1995), 12 = Cenci-Goga et al. (2013), 14 = Condoleo et al. (2016), 18 = de Barros Correia et al. (2015), 23 = Deksné et al. (2017), 25 = Djokic et al. (2016), 32 = Garcia-Bocanegra et al. (2013), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 38 = Gilot-Fromont et al. (2009), 43 = Herrero et al. (2016), 45 = Holec-Gasior et al. (2015), 46 = Hutchinson et al. (2011), 48 = Jokelainen et al. (2017), 49 = Kantzoura et al. (2013), 53 = Klun et al. (2006), 55 = Limon et al. (2017), 56 = Liu et al. (2015), 65 = Ortega-Pacheco et al. (2013), 73 = Santoro et al. (2017), 76 = Sechi et al. (2013), 78 = Sun et al. (2015), 80 = Tao et al. (2011), 81 = Tzanidakis et al. (2012), 86 = Villari et al. (2009), 88 = Xu et al. (2015), 93 = Schares et al. (2017a), 95 = Tilahun et al. (2018), 99 = Schoonman et al. (2010), 102 = Machacova et al. (2014), 104 = Ribeiro et al. (2016).

(of the same species, or from other farms) was established as a risk factor. As already discussed, low intensity was often related to a higher risk of *T. gondii* seropositivity (Table 6). The overall tendency that presence of or contact to other animal species may pose a risk for livestock to be *T. gondii*-seropositive (Table 9), could therefore be explained by the specific conditions in low-intensive farming. These conditions may be associated with an increased risk of exposure of livestock to *T. gondii* (e.g. through contamination of feed, water or farmland with oocysts and contact to rodents). On the other hand, the presence of other animal species could also have a direct effect on the risk of infection for livestock animals, as they may represent reservoirs for *T. gondii* and can thus be involved in the establishment of an on-farm multiplication of *T. gondii* (Table 9).

3.1.4.5. Biosecurity, farm buildings, staff hygiene, animal restocking. Biosecurity, the structural condition of farm buildings, staff hygiene and a restrictive policy of restocking seem to be associated with a lower risk of seropositivity in livestock (Table 10). In most cases, staff hygiene and biosecurity measures may have no direct effect on the exposure of animals to *T. gondii*, except for e.g. cleaning methods or the floor type (Table 10). However, the implementation of biosecurity and hygiene measures could represent an indirect indicator for the level of confinement under which livestock are reared. A restricted restocking policy by avoiding the purchase of animals from other farms may prevent the accidental incorporation of infected animals into the herd or flock.

3.1.4.6. Hygiene, cleaning and disinfection measures. Measures of hygiene and regimes of cleaning and disinfection applied at the farms may play an important role in the infection of livestock with *T. gondii* because cleaning reduces the probability of contamination of the facilities with oocysts and may also reduce exposure to infected intermediate hosts, e.g. rodents. There is a clear tendency that a high hygienic status and the implementation of cleaning and disinfection measures has a protective effect (Table 11).

3.1.4.7. Disease and treatment parameters and other factors characterizing herd health and veterinary care. Disease and treatment parameters and other factors characterizing herd health and veterinary care are often unrelated to *T. gondii* seropositivity in livestock (Table 12). Even when parameters characterizing reproductive problems in small ruminants were assessed, statistically significant associations were hardly observed. However, a few studies that had specifically looked at abortion in general, serial abortions or neonatal mortality, revealed associations to *T. gondii* seropositivity (Table 12).

3.2. Factors related to the *T. gondii* life cycle

3.2.1. Definitive hosts (cats)

Cats as definitive hosts of *T. gondii* may shed oocysts with their feces. Excreted oocysts are environmentally resistant and become infective after a short period of sporulation. Thus, a large number of on-farm studies assessed cat-related factors in relation to seropositivity in livestock animals (Table 13). Surprisingly, about half of the studies which considered the presence of cats on the farms failed to find an association with seropositivity regarding *T. gondii*, especially in small ruminants, poultry and equids

Table 9Parameters related to the presence of other animals (livestock, non-livestock) close to livestock species as putative risk factors for seropositivity to *T. gondii*.

Species	Presence of other species, contact with other herds/flocks	Statistically significant	Not statistically significant	
Pigs	Presence of dairy cattle	–	6, 25	
	Presence of poultry	86	–	
	Exposure to wild animals	–	25	
	Other livestock	–	6, 25, 55	
	Presence of dogs	34, 43, 86	6	
Cattle	Presence of sheep/goats/pigs/poultry/canids; number of additional species on farm	97	78	
Sheep	Contact with other flocks	50	14	
	Grazing with other herds	49	18, 81	
	Presence of different species	–	14	
	Presence of animals (other than sheep) from other farms	35	14, 18	
	Presence of wild animals	–	14	
	Presence of cattle	22, 46	14	
	Presence of goats	–	14, 35	
	Presence of dogs	–	14, 68, 81	
	Feeding of dogs with pet food (versus feeding food waste)	68	–	
	Presence of wild dogs	–	68	
	Presence of poultry	23 (p)	81	
	Presence of pigs	–	14, 81	
	Goats	Common grazing with animals from other herds	49	81
		Presence of sheep	35	–
		Presence of dogs	–	81
Number of dogs ≤ 2		68	–	
Feeding of the dogs with pet food/leftovers		–	68	
Dogs have access to water		68 (p)	–	
Dogs have access to facilities		–	68	
Presence of poultry		–	81	
Presence of pigs		–	81	
Presence of wild dogs		–	68	
Presence of other species		35, 96	–	
Solely goats on the farm		26	24	
Chicken		Presence of sheep	71	–
	Birth of sheep on property	71	–	
	Reproductive disorders in sheep	71	–	
Equids	Presence of domestic ruminants	31	102	

(p) = protective factor; coding of references: 6 = Alvarado-Esquivel et al. (2014), 14 = Condoleo et al. (2016), 18 = de Barros Correia et al. (2015), 23 = Deksne et al. (2017), 24 = Deng et al. (2016), 25 = Djokic et al. (2016), 26 = Djokic et al. (2014), 31 = Garcia-Bocanegra et al. (2012), 34 = Garcia-Bocanegra et al. (2010a,b), 35 = Gazzonis et al. (2015), 43 = Herrero et al. (2016), 46 = Hutchinson et al. (2011), 49 = Kantzoura et al. (2013), 50 = Katzer et al. (2011), 55 = Limon et al. (2017), 68 = Rego et al. (2016), 71 = Sa et al. (2017), 78 = Sun et al. (2015), 81 = Tzanidakis et al. (2012), 86 = Villari et al. (2009), 96 = Bawm et al. (2016), 97 = Fajardo et al. (2013), 102 = Machacova et al. (2014).

(Table 13). In addition, cat-related factors had a protective statistical effect in some investigations (Table 13). This shows that not just the presence of cats, but more specifically the realistic chance of cats to contaminate farmland, feed or water provided to livestock needs to be examined. A study in 12 pig farms in China indicated for example that the seroprevalence in pigs was higher in farms with a high cat density and with high soil contamination with *T. gondii* oocysts (as determined by PCR and loop-mediated isothermal amplification (LAMP)) than in those with low cat density (Du et al., 2012). If only a small number of pigs in the herd were infected, ingestion of tissue cysts, present in accidentally eaten intermediate hosts (e.g. birds or rodents), need to be taken into account as a potential source of infection. Although some research on serological methods to detect the source of infection (oocysts vs. cysts) in animals was reported (Hill et al., 2011; Munoz-Zanzi et al., 2012), such tools are hardly available or validated for epidemiological studies, unfortunately.

Studies that looked in more detail at the routes, by which cats could expose livestock to *T. gondii*, detected more frequently statistically significant associations. In addition, the chance to find a significant association was increased, when the number of cats present on a farm was taken into account (Table 13).

3.2.2. Other on-farm intermediate hosts, such as rodents and their control

Rodents like mice and rats are intermediate hosts of *T. gondii* and may serve as a reservoir for the parasite on-farms. Cats, if allowed to ingest rodents that carry tissue cysts of *T. gondii*, may become infected and eventually shed oocysts. Omnivorous animals like pigs are at risk of getting infected through rodents. Overall, the recorded studies mainly showed that the presence of rodents and the absence of rodent control pose a risk for livestock to be *T. gondii*-seropositive (Table 14).

3.2.3. Feed-related parameters

The uptake of infective stages of *T. gondii* with animal feed represents an important route, by which animals can get infected. Feed-related parameters are also influenced by the production system, which is in place on the farm. The evaluated studies suggest that open or less confined feed storage or feeding area represent an increased risk for exposure of livestock to the parasite (Table 15). In addition, feeding particular materials like goat whey may pose an infection risk as shown for pigs (Table 15). This

Table 10Parameters related to biosecurity, structural condition of farm buildings, staff hygiene and restricted animal restocking as putative risk factors for *T. gondii* seropositivity in livestock.

Species	Management, biosecurity and staff	Statistically significant	Not statistically significant
Pigs	Biosecurity	–	55
	Staff restriction	–	55
	Low level of staff hygiene	39	–
	Specialized boots, clothes	–	25, 43
	Proper maintenance of farm facilities	43 (p)	–
	Control of mosquitos and flies	34, 80	84
	Bird proof nets	34 (p)	–
	Removal of dead animals	84 (p)	–
	Floor type	–	17, 25, 84
	Farm holdings (one or more sites)	–	55
Cattle	Danish entry	25	–
	Presence of birds in stables	78	–
	Farm neighborhood (isolation)	38	–
	Work clothes available	–	78
	Slaughter on property	–	21
	Cattle introduced from other farms	78	–
Sheep	Dirty floor versus concrete floor	99	–
	State-owned farms (versus private)	53	–
	Commercial (versus family)	41	–
	Large size	70, 85	–
	Agriculture is main occupation	70	–
	Lambing in paddocks or parks	–	50
	Slatted floor	16, 68	–
	Presence of pen	–	41, 95
	Cement	68 (p)	–
	Dirt floor versus concrete floor	68	–
	Multiple boundaries	50	–
	Technified rearing	–	18
	Educational level of farmer	49	–
	Farm recently created	61	–
	Replacements during preceding year	61	18
	Use of exchanged or borrowed breeding males	16	–
	Leaving aborted fetuses on ground	1	81
	Predominantly external replacement	–	81
	Stocking rate (<1 versus ≥1)	41 (p)	–
	Use of quarantine	69 (p)	81
Frequency of domestic slaughtering	–	14	
Goats	Availability of a special place for parturition	–	81
	Pen flooring dirt (to suspended slat, mix, cement)	68	–
	Leaving aborted fetuses on ground	1	81
	Predominantly external replacement	–	81
	Purchase of spare breeding animals	–	35
	No replacements during preceding year	61	24
	Use of quarantine	–	81
	Availability of a special place for parturition	–	81
	Animals born on farm	26	–
	Educational level of farmer	49	–
Chicken	Farm recently created	61	–
	Intake of fetal adnexa, fluids and placentas	71	–
Equids	Slaughter of animals on property	71	–
	Animals of replenishments from other districts or states	29	–
	Use of studs from other stables	–	29
	Acquisition of female breeders in the last 5 years	–	29
	Introduced breeders in the last 5 years	–	29
	Treatment, cleaning and care area	104	–

(p) = protective factor; coding of references: 1 = Abu-Dalbouh et al. (2012), 14 = Condoleo et al. (2016), 16 = D'Alencar Mendonca et al. (2013), 17 = Damriyasa et al. (2004), 18 = de Barros Correia et al. (2015), 21 = de Souza et al. (2016); 24 = Deng et al. (2016), 25 = Djokic et al. (2016), 26 = Djokic et al. (2014), 29 = Fonseca de Araujo Valenca et al. (2015), 34 = Garcia-Bocanegra et al. (2010a,b), 35 = Gazzonis et al. (2015), 38 = Gilot-Fromont et al. (2009); 39 = Goerlich (2011), 41 = Guimaraes et al. (2013), 43 = Herrero et al. (2016), 49 = Kantzoura et al. (2013), 50 = Katzer et al. (2011), 53 = Klun et al. (2006), 55 = Limon et al. (2017), 61 = Mainar et al. (1996), 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 70 = Romanelli et al. (2007), 71 = Sa et al. (2017), 78 = Sun et al. (2015); 80 = Tao et al. (2011), 81 = Tzanidakis et al. (2012), 84 = Veronesi et al. (2011), 85 = Vesco et al. (2007), 95 = Tilahun et al. (2018), 99 = Schoonman et al. (2010), 104 = Ribeiro et al. (2016).

suggests that goats excrete viable *T. gondii* stages in their milk, that may remain infective even after the whey has been produced (Table 15).

Table 11Parameters related to hygiene, cleaning and disinfection measures as putative risk or protective factors for *T. gondii* seropositivity in livestock.

Species	Hygiene, disinfection and cleaning measures	Statistically significant	Not statistically significant
Pigs	Manual cleaning	86	–
	Frequency of disinfection	–	80
	Empty period length (short versus long)	39	25
	All-in/all-out	34 (p), 39 (p), 84 (p)	43, 55
	No cleaning and disinfection	34	–
	Cleaning method (only mechanical)	–	17
Cattle	Only removing manure	39	–
	Cleaning method	–	78
Sheep	Poor hygiene conditions	15, 56	–
	Hygiene level	–	81
	Disinfection of installations	69 (p)	–
	Use of dunghill	69 (p)	–
Goats	Feces management	69 (p)	–
	Feces management	–	68
	Poor hygiene level	56	81
Equids	No dunghill	29	–
Chicken	Service period prior to restocking	93	–

(p) = protective factor; coding of references: 15 = Cosendey-KezenLeite et al. (2014), 17 = Damriyasa et al. (2004), 25 = Djokic et al. (2016), 29 = Fonseca de Araujo Valenca et al. (2015), 34 = Garcia-Bocanegra et al. (2010a,b), 39 = Goerlich (2011), 43 = Herrero et al. (2016), 55 = Limon et al. (2017), 56 = Liu et al. (2015), 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 78 = Sun et al. (2015), 80 = Tao et al. (2011), 81 = Tzanidakis et al. (2012), 84 = Veronesi et al. (2011), 86 = Villari et al. (2009), 93 = Schares et al. (2017a).

3.2.4. Water-related parameters

Since oocysts of *T. gondii* can remain infective in water for a long time (i.e., under optimal conditions several months or even years (Dubey, 2010b)), it is hypothesized that the water supply for livestock may represent a risk factor for infection. Water can be supplied

Table 12Disease and treatment parameters and other factors characterizing herd health and their association with *T. gondii* seropositivity in livestock.

Species	Factors	Statistically significant	Not statistically significant
Pigs	Deworming	–	6, 55
	Cannibalism	65	43
Cattle	Number of pregnancies (0 or 1 versus 3 or more pregnancies)	79	–
	Reproductive disorder	–	32, 21
	Brucellosis status	–	32
Sheep	Veterinary service	–	78
	No veterinary care	15	–
	Vaccination status (multivariable testing including age)	46	–
	Anthelmintic treatment	49 (p)	68
	Treatment with albendazoles (versus salicylanilides and imidazothiazoles)	49 (p)	–
	Previous history of serial abortions	61	–
	Unusual episodes of neonatal mortality	61	–
	Proportion of abortions (high versus low)	18	81
	Occurrence of stillbirths	–	18
	Occurrence of death at weaning	–	18
	Neurological problems observed	–	81
	Number of abortion waves per year	–	81
	Laboratory investigation of causes of abortion	–	81
	Reproductive disorder	15	32, 69, 81
	Brucellosis status	–	32
	Goats	Proportion of abortions	–
Neurological problems observed		–	81
Number of abortion waves per year		–	81
Laboratory investigation of causes of abortion		–	81
Reproductive disorder		–	32, 81
Brucellosis status		–	32
Anthelmintic treatment		49	–
Deworming		68 (p)	–
Previous history of serial abortions		61	–
Unusual episodes of neonatal mortality		61	–
Equids	Vaccination (tetanus, influenza)	104 (p)	–
	Use of embryo transfer	104 (p)	–

(p) = protective factor; coding of references: 6 = Alvarado-Esquivel et al. (2014), 15 = Cosendey-KezenLeite et al. (2014), 18 = de Barros Correia et al. (2015), 21 = de Souza et al. (2016); 32 = Garcia-Bocanegra et al. (2013), 43 = Herrero et al. (2016); 46 = Hutchinson et al. (2011), 49 = Kantzoura et al. (2013), 55 = Limon et al. (2017), 61 = Mainar et al. (1996), 65 = Ortega-Pacheco et al. (2013); 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 78 = Sun et al. (2015); 79 = Tan et al. (2015); 81 = Tzanidakis et al. (2012), 104 = Ribeiro et al. (2016).

Table 13
Cat-related parameters as putative risk factors for *T. gondii* seropositivity in livestock.

Factor	Species	Statistically significant	Not statistically significant
Presence of cats on farm	Pigs	9, 20, 27, 33, 34, 39, 43, 65, 87	6, 17, 55, 84, 86, 94
	Cattle	38, 59, 78	21, 32, 38, 99
	Sheep	1, 13, 16, 56, 58, 61, 77, 85, 90	14, 15, 32, 35, 36, 41, 72, 81
	Goats	1, 61, 24, 56, 96	32, 35, 81
	Chicken	60, 93	71
	Equids	102	11, 31, 40, 29
Number of cats on farm (as continuous variable or >2-3 cats/>2/>3)	Pigs	27, 39, 63, 65	-
	Cattle	59, 97	-
	Sheep	41, 68	-
	Goats	24	68
	Chicken	60	-
	Equids	63, 80	55
Contact/exposure to cats or cat feces (plus frequency of exposure)	Sheep	59	95
	Goats	19, 30, 68 (p)	24, 95
	Cattle	-	21, 95
	Pigs	55	-
	Equids	102	29
Contact of cats with feed	Pigs	55	-
	Cattle	59	-
	Sheep	69, 70	68
	Goats	30	-
	Equids	102	29
	Pigs	-	55
Contact of cats with water	Sheep	12, 76, 68	41, 69
	Goats	68 (p)	-
	Equids	-	102
	Pigs	62 (p)	-
Vaccination of cats on farm	Pigs	27, 87	-
Detection of oocysts in soil, cat feces, water	Sheep	-	81
Cat-proof storage of feed supplements	Goats	-	81
	Sheep	-	81
Cats seen in hay	Goats	-	81
	Goats	-	81
Cats are used for rodent control	Cattle	97	-
	Chicken	93	-
Population control of cats	Sheep	68 (p)	-
	Goats	-	68
	Sheep	68	-
Feeding of cats with pet food (versus food waste)	Goats	-	68
	Sheep	-	68
Cats feed on placenta remains	Goats	68	-
	Sheep	68	-
Presence of wild cats	Goats	68 (p)	-
	Goats	68	-
Stray cats or wild felids on farm or farm land	Sheep	41	-
Occurrence of birth of cats on property	Chicken	-	71

(p) = protective factor; coding of references: 1 = Abu-Dalbouh et al. (2012), 6 = Alvarado-Esquivel et al. (2014), 9 = Assadi-Rad et al. (1995), 11 = Cazarotto et al. (2016), 12 = Cenci-Goga et al. (2013), 13 = Andrade et al. (2013), 14 = Condoleo et al. (2016), 15 = Cosendey-KezenLeite et al. (2014), 16 = D'Alencar Mendonca et al. (2013), 17 = Damriyasa et al. (2004), 19 = de Moura et al. (2016), 20 = de Sousa et al. (2014), 21 = de Souza et al. (2016), 24 = Deng et al. (2016), 27 = Du et al. (2012), 29 = Fonseca de Araujo Valenca et al. (2015), 30 = Garcia et al. (2012), 31 = Garcia-Bocanegra et al. (2012), 32 = Garcia-Bocanegra et al. (2013), 33 = Garcia-Bocanegra et al. (2010a), 34 = Garcia-Bocanegra et al. (2010b), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 38 = Gilot-Fromont et al. (2009), 39 = Goerlich (2011), 40 = Guerra et al. (2018), 41 = Guimaraes et al. (2013), 43 = Herrero et al. (2016), 55 = Limon et al. (2017), 56 = Liu et al. (2015), 58 = Lopes et al. (2010), 59 = Magalhaes et al. (2016), 60 = Magalhães et al. (2016), 61 = Mainar et al. (1996), 62 = Mateus-Pinilla et al. (1999), 63 = Meerburg et al. (2006), 65 = Ortega-Pacheco et al. (2013), 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 71 = Sa et al. (2017), 72 = Sakata et al. (2012), 76 = Sechi et al. (2013), 77 = Skjerve et al. (1998), 78 = Sun et al. (2015), 80 = Tao et al. (2011), 81 = Tzanidakis et al. (2012), 84 = Veronesi et al. (2011), 85 = Vesco et al. (2007), 86 = Villari et al. (2009), 87 = Weigel et al. (1995), 90 = Zhang et al. (2016), 93 = Schares et al. (2017a), Samico-Fernandes et al. (2017), 95 = Tilahun et al. (2018), 96 = Bawm et al. (2016), 97 = Fajardo et al. (2013), 99 = Schoonman et al. (2010), 102 = Machacova et al. (2014).

to the animals from a variety of sources such as tap water, wells or surface water and on different ways, which may depend on various factors like the production system or specific regional parameters. It is important to establish whether cats have access to the water at any stage before it reaches the livestock animals. Overall, from the recorded studies it is hard to quantify the risk for infection of the animals through contaminated water (Table 16). Often, the outcomes of the studies are contradicting. For example, well water was associated with an increased risk in some studies, while it seemed to have a protective statistical effect in others (Table 16).

3.2.5. Soil contact, outside access and pasturing

Exposure to contaminated soil or pastures seems to be another important factor (Table 17). Especially in pigs, extensive management or outdoor access seems to increase the chance for animals to get in contact with infective *T. gondii* stages (Table 17). Both, oocyst contaminations on farmland and, in the case of omnivorous livestock species, tissues of infected intermediate hosts represent likely sources of infection for livestock.

Table 14Parameters related to intermediate hosts (rodents) and their control on farms as putative risk factors for *T. gondii* seropositivity in livestock.

Factor	Species	Statistically significant	Not statistically significant
Presence of rodents (mice, rats)	Pigs	–	87
	Cattle	78	–
	Sheep	15	–
Access of rodents to feed	Cattle	59	–
	Sheep	70	–
No rodent control measures	Pigs	9, 34, 43, 52, 86	51
	Sheep	77	–
Cats and dogs as a measure of rodent control/cats used for rodent control	Pigs	39, 44	–
	Chickens	93	–
Problems with mice and rats	Goats	–	24

Coding of references: 9 = Assadi-Rad et al. (1995), 15 = Cosendey-KezenLeite et al. (2014), 24 = Deng et al. (2016), 34 = Garcia-Bocanegra et al. (2010a,b), 39 = Goerlich (2011), 43 = Herrero et al. (2016), 44 = Hill et al. (2010), 51 = Kijlstra et al. (2004), 52 = Kijlstra et al. (2008), 59 = Magalhaes et al. (2016), 70 = Romanelli et al. (2007), 75 = Schulzig and Fehlhaber (2005), 77 = Skjerve et al. (1998), 78 = Sun et al. (2015), 86 = Villari et al. (2009), 87 = Weigel et al. (1995), 93 = Schares et al. (2017a).

4. Economic impact of toxoplasmosis in livestock

In this review, we focused on articles that assessed the costs of *T. gondii* in livestock animals. It is important to stress, however, that *T. gondii* infections in animals used for food production may also affect human health and cause costs in this respect. These aspects are very difficult to assess and beyond the scope of the article, but have been addressed by others (Buzby and Roberts, 1997; Todd, 1989).

As summarized in Section 2.2, *T. gondii* is considered a major cause of reproductive losses in the small ruminant industry worldwide and infections in small ruminants may play a major role in the transmission of the parasite to humans (Belluco et al., 2016; Opsteegh et al., 2016). Especially in China, abortions caused by *T. gondii* in sows seem to be common and may lead to huge losses (Pan et al., 2017). There is one report on severe clinical signs in a fattening pig farm in China (Li et al., 2010) indicating that toxoplasmosis is of economic importance on these farms.

Table 15Feeding-related parameters as putative risk factors for *T. gondii* seropositivity in livestock.

Species	Feeding characteristics	Statistically significant	Not statistically significant
Pigs	Food storage open	39	55, 65
	Food storage in owner's home	6	–
	Roughage not covered	63	–
	Manual feeder type (versus automatic feeder)	39	25, 55, 65
	Fluid feed (versus dry feed)	39 (p)	–
	Trough	39	–
	Feeding human food	6, 20	–
	Ration, mixed versus human food waste	–	94
	Feeding goat whey	63	–
Cattle	Use of silage	38	53
	Raw milk consumption	–	21
Sheep	Food storage uncovered	69	–
	Food through covered trough	69 (p)	–
	Feeding concentrate	81	–
	No mineral supplementation	58, 70	18
	Food source common	–	41
Goats	Atypical grazing	77	–
	Feeding concentrate	35, 81	19
	Use of mixer feeder	–	24
Chicken	Use of silo	–	24
	Feeding from ground	4	–
Equids	Poultry feed indoors	4 (p)	–
	Feed (with or without supplements)	–	40
	Mix of collective and individual troughs	29 (p)	–
	Ration consumption	–	29
	Ration storage location open	29	–
	Hay consumption	29	–
	Hay storage location open	29	–

(p) = protective factor; coding of references: 4 = Alvarado-Esquivel et al. (2012a), 6 = Alvarado-Esquivel et al. (2014), 18 = de Barros Correia et al. (2015), 19 = de Moura et al. (2016), 20 = de Sousa et al. (2014), 21 = de Souza et al. (2016), 24 = Deng et al. (2016), 25 = Djokic et al. (2016), 29 = Fonseca de Araujo Valenca et al. (2015), 35 = Gazzonis et al. (2015), 38 = Gilot-Fromont et al. (2009), 39 = Goerlich (2011), 40 = Guerra et al. (2018), 41 = Guimaraes et al. (2013), 53 = Klun et al. (2006), 55 = Limon et al. (2017), 58 = Lopes et al. (2010), 63 = Meerburg et al. (2006), 65 = Ortega-Pacheco et al. (2013), 69 = Rizzo et al. (2017), 70 = Romanelli et al. (2007), 72 = Sakata et al. (2012), 77 = Skjerve et al. (1998), 81 = Tzanidakis et al. (2012), 94 = Samico-Fernandes et al. (2017).

Table 16Water-related parameters as putative risk factors for *T. gondii* seropositivity in livestock.

Species	Water supply characteristics	Statistically significant	Not statistically significant
Pigs	Water supply (various sources assessed)	–	6, 55, 80, 84, 87
	Water from wells (versus municipal water)	86	–
Cattle	Water supply from pond or well	78	–
	Water point on pasture	38	–
	Water from reservoir	59	–
	Access to surface water	99	–
	Stagnant/pond water versus mixed water sources (river, stream, well)	95	–
	Tap water versus mixed water sources (river, stream, well)	95	–
Sheep	Use of surface water	85	–
	Water from the public supply	81	41
	River water	36	41
	Tap water	36	–
	Water directly from the source (well)	15 (p), 16 (p)	69
	Still water (versus running water)	13 (p), 76	41
	Water from deep well	15, 16 (p)	–
	Water from sluice	15	–
	Stagnant/pond water versus mixed water sources (river, stream, well)	95	–
	Tap water versus mixed water sources (river, stream, well)	95	–
	Location of drinking trough	68	–
	Dogs and wild dogs have access to water	68	–
	Water sources in main grazing area	–	14
	Location of drinking trough	68	–
Goats	Water source (various sources assessed)	–	19, 24
	Water from river	35	–
	Water from the public supply	81	–
	Stagnant/pond water versus mixed water sources (river, stream, well)	95	–
Chicken	Tap water versus mixed water sources (river, stream, well)	95	–
	Water source (dam)	60	–
Equids	Drinking from a mix of individual and collective troughs	29 (p)	–
	Water well versus public system	–	100
	Tank or river/stream versus public system	100	–

(p) = protective factor; coding of references: 6 = Alvarado-Esquivel et al. (2014), 13 = Andrade et al. (2013), 14 = Condoleo et al. (2016), 15 = Cosendey-KezenLeite et al. (2014), 16 = D'Alencar Mendonca et al. (2013), 19 = de Moura et al. (2016), 24 = Deng et al. (2016), 29 = Fonseca de Araujo Valenca et al. (2015), 35 = Gazzonis et al. (2015), 36 = Gebremedhin et al. (2013), 38 = Gilot-Fromont et al. (2009), 41 = Guimaraes et al. (2013), 55 = Limon et al. (2017), 59 = Magalhaes et al. (2016), 60 = Magalhães et al. (2016), 68 = Rego et al. (2016), 69 = Rizzo et al. (2017), 76 = Sechi et al. (2013), 78 = Sun et al. (2015), 80 = Tao et al. (2011), 81 = Tzanidakis et al. (2012), 84 = Veronesi et al. (2011), 85 = Vesco et al. (2007), 86 = Villari et al. (2009), 87 = Weigel et al. (1995), 95 = Tilahun et al. (2018), 99 = Schoonman et al. (2010), 100 = Almeida et al. (2017).

The number of studies assessing the economic impact of *T. gondii* infection in livestock is scarce. To estimate the economic consequences of an infection exclusively for the affected livestock species, i.e. leaving out any potential effect on human health, the clinical consequences, their severity and impact on the performance of the animals have to be analyzed. Small ruminants may suffer from a *T. gondii* infection and the infection can therefore cause economic losses to farmers. Experimentally infected

Table 17Soil-contact, outside access, pasturing and related parameters as putative risk factors for *T. gondii* seropositivity in livestock.

Species	Soil-contact, outside access, pasturing	Statistically significant	Not statistically significant
Pigs	Outdoor facilities	6, 9, 22, 33, 53, 55	43
	Detection of oocysts in soil, cat feces, water	27, 87	–
	Scavenging	80	–
	Pasture length month	105	–
Sheep	Size of the grazing area	–	14
	Frequency of grazing	–	14
	Pasture	58	18, 41
Goats	Outdoor access	–	24
	Grazing	35	–
Cattle	Access to pasture (relative to stable only), grazing	53, 99	–
Equids	Shelter (in- or outdoor)	–	31, 40
	Pasture versus stable	101	–
Chicken	Size of chicken run per animal (≥ 10 sqm versus < 10 sqm), multivariable analysis including age	93	–

Coding of references: 6 = Alvarado-Esquivel et al. (2014), 9 = Assadi-Rad et al. (1995), 14 = Condoleo et al. (2016), 18 = de Barros Correia et al. (2015), 22 = Deksne and Kirjusina (2013), 24 = Deng et al. (2016), 27 = Du et al. (2012), 31 = Garcia-Bocanegra et al. (2012), 33 = Garcia-Bocanegra et al. (2010a, b), 35 = Gazzonis et al. (2015), 40 = Guerra et al. (2018), 41 = Guimaraes et al. (2013), 43 = Herrero et al. (2016), 53 = Klun et al. (2006), 55 = Limon et al. (2017), 58 = Lopes et al. (2010), 80 = Tao et al. (2011); 87 = Weigel et al. (1995), 93 = Schares et al. (2017a), 99 = Schoonman et al. (2010), 101 = Bartova et al. (2017), 105 = Wallander et al. (2016).

sheep showed a number of clinical signs, which included fever, diarrhea, abortion, stillbirth, and fetal mummification (Dubey, 2009b). There is also one report indicating that *T. gondii* infected rams may become infertile (Savvulidi et al., 2018). It has also been suggested that *T. gondii* could be transmitted via semen (de Moraes et al., 2010; W.D. Lopes et al., 2013).

In general, the economic impact of diseases has different aspects that need to be taken into account. Direct costs of a disease (C) include not only losses (L), but also costs for the treatment of animals (T) and costs for disease prevention (P). The first aspect is 'the value of the loss in expected output and/or of resource wastage due to the disease (L)' (Bennett et al., 1999). In sheep, for example, lambs, wool, milk and meat represent the main output of a flock. As an example, in the case of a primary infection of sheep with *T. gondii*, there is a high probability of abortion (Dubey, 2009b) and the loss is therefore at least the value of the newborn lamb. Moreover, in dairy flocks, fever after acute infection, but mainly the occurrence of abortion, is related to the complete or partial loss of milk production for that lactation, i.e. the main source of income for these farms.

'The costs for treatment of affected animals (T)' represent the second aspect in an economic analysis (Bennett, 2003; Dijkhuizen and Morris, 1997). In the case of toxoplasmosis, the treatment costs include costs for anti-inflammatory substances to reduce fever or other veterinary services (e.g. removing mummified lambs, treatment of fertility problems after abortion, costs of laboratory diagnosis etc.).

'The costs associated with specific disease prevention (P)' form the third aspect in the economic analysis (Bennett et al., 1999). In the case of toxoplasmosis, vaccination of sheep on a regular base may represent such a preventive measure to reduce *T. gondii*-associated abortion.

There are only two formally published studies on the economic impact of toxoplasmosis in sheep and they refer only to two countries, Great Britain and Uruguay. Both studies focused on the losses that were due to abortion. Freyre et al. (1997), who analyzed the situation in Uruguay, estimated that about 10 million lambs were born in 1993 and that 14,000–400,000 sheep fetuses were aborted due to toxoplasmosis. They assumed a loss of 10 to 12 US \$ per fetus, resulting in a loss of 1.4–4.68 million US \$. They took neither the retardation of the genetic progress, nor the costs for replacement animals nor for husbandry into consideration.

Bennett and colleagues published several studies on the direct costs associated with endemic diseases of livestock in Great Britain, including toxoplasmosis in sheep (Bennett, 2003; Bennett et al., 1997, 1999; Bennett and Ijpelaar, 2005). In the first study, referring to the year 1996, annual costs of 12–18 million £ for output losses (L) and <1 million £ for treatment (T), but no costs for prevention (P) were estimated. An incidence of ovine toxoplasmosis in ewes of 1.2 and 2.2% was assumed. In an update of this study, the authors estimated that about 334,000 (range: 136,000–526,000) sheep were affected per year, with disease effects of 9.1 (range: 3.7–14.1) million £ due to abortion or stillbirth and 3.2 (range: 2.7–5.6) million £ for disease control. It was assumed that toxoplasmosis caused 50,000 (range 8000–116,000) severely affected sheep in Great Britain per year. According to a recent ABC News report (<https://www.abc.net.au/news/rural/2017-02-07/toxoplasmosis-costs-south-australian-sheep-producers/8245676>, assessed on 11/12/2018, 12:32), a study carried out by Ryan O'Handley in South Australia in 2017 estimated toxoplasmosis costs in sheep at 70 million Australian \$.

In summary, there are only a few formally published studies that assessed the costs of toxoplasmosis in livestock. They focused on sheep and peer-reviewed reports on this topic are >20 years old.

To improve the assessment of the economic impact of *T. gondii* on animals, it is necessary to carry out further studies including all aspects of the infection (Perry and Randolph, 1999) and all cost categories that can lead to losses. Information at the country

Table 18

Factors possibly causing costs in livestock due to infection with *T. gondii*.

Type of costs	Costs	Comments
Direct costs (production losses caused by disease)	Reduced milk yield	Only in dairy sheep and dairy goats
	Weight loss in infected animals	Caused by fever and inappetence
	Reduced fertility	Cause of an increased replacement rate and retardation of the breeding progress
	Abortion/stillbirth	Cause of an increased replacement rate, loss of sells (e.g. lamb sells) and in retardation of the breeding progress
	Increased mortality	Reduced profit due to loss of animals and increased replacement rate
	Prolonged fattening periods in infected animals	Additional costs for feeding and reduced profit
Indirect costs (reaction to disease)	Weak or malformed progeny	Loss of progeny that causes additional costs, e.g. if a caesarian section has to be carried out or animals need veterinary service
	Optimization of herd management	Improvements in biosecurity, hygiene and farm buildings
	Treatment of diseased animals	Cost for veterinarian, drugs, etc.
	Control measures	For example vaccination
	Monitoring and diagnosis	For example costs for sampling and laboratory testing (also to achieve a differential diagnosis); testing of animals before housing
	Slaughter of infected animals	For example in animals with reduced fertility, increase in replacement costs
	Impact on trade (both national or international)	For example if meat of infected animals would be excluded from slaughter for specific meat products, or if international trade becomes restricted to avoid introduction of virulent types of <i>T. gondii</i>

level is scarce. Estimates are often based on studies conducted on (heavily) affected farms. This may lead to an overestimation of the total costs. It also remains unclear, if all recorded abortions have been caused by infection with *T. gondii*, even if part of the aborted fetuses has tested positive for toxoplasmosis. The lack of information on the effect of potential co-infections, e.g. border disease or bacterial infections that may also cause abortion, represents another factor of uncertainty.

In many publications with statements on costs the parameters included in the cost calculations remain unclear. It is therefore difficult to compare these analyses. Furthermore, they may include only costs directly related to the animals, but often ignore potential consequences on trade, a possible decrease of the value of infected flocks or herds and any potential impact on human health, if products contaminated with infectious stages of *T. gondii* are placed on the market. Consequently, we suggest that such studies list all parameters in a given livestock production system that might be affected by infection with *T. gondii* and indicate how they were taken into account in the cost calculations. As example, we provide such information categorized as direct and indirect costs in Table 18. It has to be noted, however, that some factors (e.g. herd management, hygiene) may have a general influence on the herd health status. This will make it hard to allocate any potential costs related to these factors to specific infections, for example with *T. gondii*.

5. Conclusion and future prospects

Although many studies on *T. gondii* infections and toxoplasmosis in livestock are available in literature, there are still several important gaps in our current knowledge. The routes of infection seem to be clear in herbivores, as oocysts can be presumed to be the main source. Nevertheless, there is uncertainty regarding many aspects, e.g. about the role of contamination of water and pastures and concerning the intensity of farming. In pigs, it seems clear that outdoor access is an important risk factor for *T. gondii* infection. It is not clear, however, to which extent oocysts contamination or the presence or uptake of infected intermediate hosts such as rodents play a role. Overall, further epidemiological studies are necessary and the standards in these studies need to be raised to obtain a better knowledge and more confidence regarding the epidemiologically relevant routes of *T. gondii* infection in livestock. In addition, more prospective studies including the assessment of interventions are needed to confirm previous findings and to determine the feasibility and the efficiency of control measures. Furthermore, the economic impact of toxoplasmosis in livestock, e.g. in small ruminants, has never been assessed in most of the regions worldwide, although especially small ruminants are economically important species in many countries. There is a clear need for further assessments of economic consequences of *T. gondii* infections and toxoplasmosis in livestock.

Conflict of interest statement

None of the authors of this paper has a financial or personal relationship with other people or organizations that could inappropriately influence or bias the content of the paper.

Acknowledgements

We wish to thank the FLI librarians (Viola Damrau, Mandy Nass) who supported our work tremendously. In addition, we thank Rob van Spronsen, Rijksinstituut voor Volksgezondheid en Milieu, Bilthoven, The Netherlands, for supporting our search for relevant literature. Luis-Miguel Ortega-Mora and Julio Benavides are supported by grant AGL2016-75935 from the Spanish Ministry of Science. Sandra Stelzer was funded by JIP1 ORION within the One Health EJP. One Health EJP has received funding from the European Union Horizon 2020 research and innovation program under grant agreement No 773830.

References

- Abu-Dalbouh, M.A., Ababneh, M.M., Giadinis, N.D., Lafi, S.Q., 2012. Ovine and caprine toxoplasmosis (*Toxoplasma gondii*) in aborted animals in Jordanian goat and sheep flocks. *Trop. Anim. Health Prod.* 44, 49–54.
- Aharonson-Raz, K., Baneth, G., Lopes, A.P., Brancal, H., Schallig, H., Cardoso, L., Steinman, A., 2015. Low seroprevalence of *Leishmania infantum* and *Toxoplasma gondii* in the horse population in Israel. *Vector Borne Zoonotic Dis.* 15, 726–731.
- Ahmed, H., Malik, A., Arshad, M., Mustafa, I., Khan, M.R., Afzal, M.S., Ali, S., Mobeen, M., Simsek, S., 2016. Seroprevalence and spatial distribution of toxoplasmosis in sheep and goats in north-eastern region of Pakistan. *Korean J. Parasitol.* 54, 439–446.
- Alanazi, A.D., 2013. Determination of seropositivity for *Toxoplasma gondii* in sheep, goats and camels slaughtered for food and human consumptions in Riyadh municipal abattoirs, Saudi Arabia. *J. Egypt. Soc. Parasitol.* 43, 569–576.
- Alanazi, A.D., Alyousif, M.S., 2011. Prevalence of antibodies to *Toxoplasma gondii* in horses in Riyadh Province, Saudi Arabia. *J. Parasitol.* 97, 943–945.
- Al-Kappany, Y.M., Abbas, I.E., Devleeschauwer, B., Dorny, P., Jennes, M., Cox, E., 2018. Seroprevalence of anti-*Toxoplasma gondii* antibodies in Egyptian sheep and goats. *BMC Vet. Res.* 14, 120.
- Al-Khalidi, N.W., Weisbrode, S.E., Dubey, J.P., 1980. Pathogenicity of *Toxoplasma gondii* oocysts to ponies. *Am. J. Vet. Res.* 41, 1549–1551.
- Almeida, J.C., Vidotto, O., Ferreira, E.P., Ribeiro, L.P., Mongruel, A.C., Vieira, T.S., Freire, R.L., Mota, R.A., Vieira, R.F., 2017. Serosurvey of anti-*Toxoplasma gondii* antibodies in sport horses from Paraíba state, Northeastern Brazil. *Acta Parasitol.* 62, 225–227.
- Almeria, S., Cabezon, O., Paniagua, J., Cano-Terriza, D., Jimenez-Ruiz, S., Arenas-Montes, A., Dubey, J.P., Garcia-Bocanegra, I., 2018. *Toxoplasma gondii* in sympatric domestic and wild ungulates in the Mediterranean ecosystem. *Parasitol. Res.* 117, 665–671.
- Altan, Y., Heydorn, A.O., Janitschke, K., 1977. Zur Infektiosität von *Toxoplasma*-Oozysten für das Pferd [Infectivity of *Toxoplasma* oocysts for horses]. *Berl. Muench. Tierärztl. Wschr.* 90, 433–435.
- Alvarado-Esquivel, C., Garcia-Machado, C., Vitela-Corrales, J., Villena, I., Dubey, J.P., 2011. Seroprevalence of *Toxoplasma gondii* infection in domestic goats in Durango State, Mexico. *Vet. Parasitol.* 183, 43–46.
- Alvarado-Esquivel, C., Gonzalez-Salazar, A.M., Alvarado-Esquivel, D., Ontiveros-Vazquez, F., Vitela-Corrales, J., Villena, I., Dubey, J.P., 2012a. Seroprevalence of *Toxoplasma gondii* infection in chickens in Durango State, Mexico. *J. Parasitol.* 98, 431–432.

- Alvarado-Esquivel, C., Rodriguez-Pena, S., Villena, I., Dubey, J.P., 2012b. Seroprevalence of *Toxoplasma gondii* infection in domestic horses in Durango State, Mexico. *J. Parasitol.* 98, 944–945.
- Alvarado-Esquivel, C., Estrada-Malacón, M.A., Reyes-Hernández, S.O., Perez-Ramirez, J.A., Trujillo-Lopez, J.I., Villena, I., Dubey, J.P., 2013a. Seroprevalence of *Toxoplasma gondii* in domestic sheep in Oaxaca State, Mexico. *J. Parasitol.* 99, 151–152.
- Alvarado-Esquivel, C., Silva-Aguilar, D., Villena, I., Dubey, J.P., 2013b. Seroprevalence and correlates of *Toxoplasma gondii* infection in domestic sheep in Michoacán State, Mexico. *Prev. Vet. Med.* 112, 433–437.
- Alvarado-Esquivel, C., Silva-Aguilar, D., Villena, I., Dubey, J.P., 2013c. Seroprevalence of *Toxoplasma gondii* infection in dairy goats in Michoacan State, Mexico. *J. Parasitol.* 99, 540–542.
- Alvarado-Esquivel, C., Romero-Salas, D., Garcia-Vazquez, Z., Crivelli-Diaz, M., Barrientos-Morales, M., Lopez-de-Buen, L., Dubey, J.P., 2014. Seroprevalence and correlates of *Toxoplasma gondii* infection in domestic pigs in Veracruz State, Mexico. *Trop. Anim. Health Prod.* 46, 705–709.
- Alvarado-Esquivel, C., Alvarado-Esquivel, D., Dubey, J.P., 2015. Prevalence of *Toxoplasma gondii* antibodies in domestic donkeys (*Equus asinus*) in Durango, Mexico slaughtered for human consumption. *BMC Vet. Res.* 11, 6.
- Amoudni, Y., Rjeibi, M.R., Rouatbi, M., Amairia, S., Awadi, S., Garbi, M., 2017. Molecular detection of *Toxoplasma gondii* infection in slaughtered ruminants (sheep, goats and cattle) in Northwest Tunisia. *Meat Sci.* 133, 180–184.
- Andrade, M.M.C., Carneiro, M., Medeiros, A.D., Neto, V.A., Vitor, R.W.A., 2013. Seroprevalence and risk factors associated with ovine toxoplasmosis in Northeast Brazil. *Parasite* 20, 20.
- Anonymous, 2011a. Scientific opinion on the public health hazards to be covered by inspection of meat (swine). EFSA J. 9 (10), 2351.
- Anonymous, 2011b. Technical specifications on harmonised epidemiological indicators for public health hazards to be covered by meat inspection of swine. EFSA J. 9 (10), 2351.
- Arias, M.L., Chinchilla, M., Reyes, L., Sabah, J., Guerrero, O.M., 1994. Determination of *Toxoplasma gondii* in several organs of cattle by carbon immunoassay (CIA) testing. *Vet. Parasitol.* 55, 133–136.
- Aroussi, A., Vignoles, P., Dalmay, F., Wimmel, L., Darde, M.L., Mercier, A., Ajzenberg, D., 2015. Detection of *Toxoplasma gondii* DNA in horse meat from supermarkets in France and performance evaluation of two serological tests. *Parasite* 22, 14.
- Assadi-Rad, A.M., New, J.C., Patton, S., 1995. Risk factor associated with transmission of *Toxoplasma gondii* to sows kept in different management systems in Tennessee. *Vet. Parasitol.* 57, 289–297.
- Ayinmode, A.B., Olaosebikan, R.I., 2014. Seroprevalence of *Toxoplasma gondii* infection in free ranged chicken from rural and urban settlements in Oyo State, Nigeria. *Afr. J. Med. Med. Sci.* 43 (Suppl), 51–57.
- Ayinmode, A.B., Oluwayelu, D.O., Sule, W.F., Obebe, O.O., 2014. Seroprevalence of *Toxoplasma gondii* in recreational horses in two metropolitan cities of Southwestern Nigeria. *Afr. J. Med. Med. Sci.* 43 (Suppl), 47–50.
- Azizi, H., Shiran, B., Boroujeni, A.B., Jafari, M., 2014. Molecular survey of *Toxoplasma gondii* in sheep, cattle and meat products in Chaharmahal va Bakhtiari Province, southwest of Iran. *Iran. J. Parasitol.* 9, 429–434.
- Bangoura, B., Zoller, B., Dausgchies, A., 2011. Prevalence and relevance of avian *Toxoplasma gondii* infections in Europe. *Berl. Munch. Tierarztl. Wochenschr.* 124, 485–496.
- Bangoura, B., Zoller, B., Koethe, M., Ludewig, M., Pott, S., Fehlhaber, K., Straubinger, R.K., Dausgchies, A., 2013. Experimental *Toxoplasma gondii* oocyst infections in turkeys (*Meleagris gallopavo*). *Vet. Parasitol.* 196, 272–277.
- Bartova, E., Dvorakova, H., Barta, J., Sedlak, K., Literak, I., 2004. Susceptibility of the domestic duck (*Anas platyrhynchos*) to experimental infection with *Toxoplasma gondii* oocysts. *Avian Pathol.* 33, 153–157.
- Bartova, E., Machacova, T., Sedlak, K., Budikova, M., Mariani, U., Veneziano, V., 2015. Seroprevalence of antibodies of *Neospora* spp. and *Toxoplasma gondii* in horses from southern Italy. *Folia Parasitol.* 62.
- Bartova, E., Sedlak, K., Kobedova, K., Budikova, M., Joel Atuman, Y., Kamani, J., 2017. Seroprevalence and risk factors of *Neospora* spp. and *Toxoplasma gondii* infections among horses and donkeys in Nigeria, West Africa. *Acta Parasitol.* 62, 606–609.
- Basso, W., Hartnack, S., Pardini, L., Maksimov, P., Koudela, B., Venturini, M.C., Schares, G., Sidler, X., Lewis, F.I., Deplazes, P., 2013. Assessment of diagnostic accuracy of a commercial ELISA for the detection of *Toxoplasma gondii* infection in pigs compared with IFAT, TgSAG1-ELISA and Western blot, using a Bayesian latent class approach. *Int. J. Parasitol.* 43, 565–570.
- Basso, W., Handke, M., Sydler, T., Borel, N., Grimm, F., Sidler, X., Deplazes, P., 2015. Involvement of *Toxoplasma gondii* in reproductive disorders in Swiss pig farms. *Parasitol. Int.* 64, 157–160.
- Basso, W., Grimm, F., Ruettten, M., Djokic, V., Blaga, R., Sidler, X., Deplazes, P., 2017. Experimental *Toxoplasma gondii* infections in pigs: humoral immune response, estimation of specific IgG avidity and the challenges of reproducing vertical transmission in sows. *Vet. Parasitol.* 236, 76–85.
- Bawm, S., Maung, W.Y., Win, M.Y., Thu, M.J., Chel, H.M., Khaing, T.A., Wai, S.S., Htun, L.L., Myaing, T.T., Tiwananthagorn, S., Igarashi, M., Katakura, K., 2016. Serological survey and factors associated with *Toxoplasma gondii* infection in domestic goats in Myanmar. *Scientifica (Cairo)* 2016, 4794318.
- Belluco, S., Mancin, M., Conficoni, D., Simonato, G., Pietrobelli, M., Ricci, A., 2016. Investigating the determinants of *Toxoplasma gondii* prevalence in meat: a systematic review and meta-regression. *PLoS ONE* 11, e0153856.
- Belluco, S., Simonato, G., Mancin, M., Pietrobelli, M., Ricci, A., 2017. *Toxoplasma gondii* infection and food consumption: a systematic review and meta-analysis of case-controlled studies. *Crit. Rev. Food Sci. Nutr.* 1–12.
- Benavides, J., Maley, S., Pang, Y., Palarea, J., Eaton, S., Katzer, F., Innes, E.A., Buxton, D., Chianini, F., 2011. Development of lesions and tissue distribution of parasite in lambs orally infected with sporulated oocysts of *Toxoplasma gondii*. *Vet. Parasitol.* 179, 209–215.
- Bennett, R., 2003. The 'direct costs' of livestock disease: the development of a system of models for the analysis of 30 endemic livestock diseases in Great Britain. *J. Agric. Econ.* 54, 55–71.
- Bennett, R., Ijpelaar, J., 2005. Updated estimates of the costs associated with thirty four endemic livestock diseases in Great Britain: a note. *J. Agric. Econ.* 56, 135–144.
- Bennett, R., Christiansen, K., Clifton-Hadley, R., 1997. An economic study of the importance of non-notifiable diseases of farm animals in Great Britain. Proceedings of the 8th Symposium of the International Society for Veterinary Epidemiology and Economics (ISVEE), Paris, France.
- Bennett, R., Christiansen, K., Clifton-Hadley, R., 1999. Preliminary estimates of the direct costs associated with endemic diseases of livestock in Great Britain. *Prev. Vet. Med.* 39, 155–171.
- Berger-Schoch, A.E., Herrmann, D.C., Schares, G., Muller, N., Bernet, D., Gottstein, B., Frey, C.F., 2011. Prevalence and genotypes of *Toxoplasma gondii* in feline faeces (oocysts) and meat from sheep, cattle and pigs in Switzerland. *Vet. Parasitol.* 177, 290–297.
- Bertranpetit, E., Jombart, T., Paradis, E., Pena, H., Dubey, J., Su, C., Mercier, A., Devillard, S., Ajzenberg, D., 2017. Phylogeography of *Toxoplasma gondii* points to a South American origin. *Infect. Genet. Evol.* 48, 150–155.
- Bettiol, S.S., Obendorf, D.L., Nowarkowski, M., Milstein, T., Goldsmid, J.M., 2000. Earthworms as paratenic hosts of toxoplasmosis in eastern barred bandicoots in Tasmania. *J. Wildl. Dis.* 36, 145–148.
- Beverly, J.K., Henry, L., Hunter, D., Brown, M.E., 1977. Experimental toxoplasmosis in calves. *Res. Vet. Sci.* 23, 33–37.
- Bezerra, M.J., Cruz, J.A., Kung, E.S., Albuquerque, P.P., Kim, P.C., Moraes, E.P., Pinheiro Junior, J.W., Mota, R.A., 2014. Detection of *Toxoplasma gondii* DNA in fresh and frozen semen from rams in Brazil. *Reprod. Domest. Anim.* 49, 753–755.
- Biancifiore, F., Rondini, C., Grelloni, V., Frescura, T., 1986. Avian toxoplasmosis: experimental infection of chicken and pigeon. *Comp. Immunol. Microbiol. Infect. Dis.* 9, 337–346.
- Bickford, A.A., Saunders, J.R., 1966. Experimental toxoplasmosis in chickens. *Am. J. Vet. Res.* 27, 308–318.
- Boch, J., Janitschke, K., Rommel, M., Sommer, R., 1965a. Untersuchungen über das Vorkommen von *Toxoplasma*-Infektionen bei Schlachtrindern. *Wien. Tierarztl. Monatsschr.* 12, 1477–1480.
- Boch, J., Rommel, M., Janitschke, K., 1965b. Contributions on pig toxoplasmosis. 3. Studies on the possibility of connatal infection. *Berl. Munch. Tierarztl. Wochenschr.* 78, 115–120.

- Boch, J., Rommel, M., Weiland, G., Janitschke, K., Sommer, R., 1966. Experimentelle *Toxoplasma*-Infektion bei Legehennen. Berl. Münch. Tierärztl. Wochenschr. 79, 352–356.
- Boughattas, S., Bergaoui, R., Essid, R., Aoun, K., Bouratbine, A., 2011. Seroprevalence of *Toxoplasma gondii* infection among horses in Tunisia. Parasit. Vectors 4, 218.
- Burrells, A., Benavides, J., Canton, G., Garcia, J.L., Bartley, P.M., Nath, M., Thomson, J., Chianini, F., Innes, E.A., Katzer, F., 2015. Vaccination of pigs with the S48 strain of *Toxoplasma gondii*—safer meat for human consumption. Vet. Res. 46, 47.
- Burrells, A., Taroda, A., Opsteegh, M., Schares, G., Benavides, J., Dam-Deisz, C., Bartley, P.M., Chianini, F., Villena, I., van der Giessen, J., Innes, E.A., Katzer, F., 2018. Detection and dissemination of *Toxoplasma gondii* in experimentally infected calves, a single test does not tell the whole story. Parasit. Vectors 11, 45.
- Buxton, D., 1998. Protozoan infections (*Toxoplasma gondii*, *Neospora caninum* and *Sarcocystis* spp.) in sheep and goats: recent advances. Vet. Res. 29, 289–310.
- Buxton, D., Finlayson, J., 1986. Experimental infection of pregnant sheep with *Toxoplasma gondii*: pathological and immunological observations on the placenta and foetus. J. Comp. Pathol. 96, 319–333.
- Buxton, D., Losson, B., 2007. Toxoplasmosis: general considerations. In: Ortega-Mora, L.M., Gottstein, B., Conraths, F.J., Buxton, D. (Eds.), Protozoal Abortifacients in Farm Ruminants: Guidelines for Diagnosis and Control. CAB International, Wallingford, UK, pp. 122–131.
- Buxton, D., Gilmour, J.S., Angus, K.W., Blewett, D.A., Miller, J.K., 1982. Perinatal changes in lambs infected with *Toxoplasma gondii*. Res. Vet. Sci. 32, 170–176.
- Buxton, D., Blewett, D.A., Trees, A.J., McColgan, C., Finlayson, J., 1988. Further studies in the use of monensin in the control of experimental ovine toxoplasmosis. J. Comp. Pathol. 98, 225–236.
- Buzby, J.C., Roberts, T., 1997. Economic costs and trade impacts of microbial foodborne illness. World Health Stat. Q. 50, 57–66.
- Campo-Portacio, D.M., Discuciviche-Rebolledo, M.A., Blanco-Tuirán, P.J., Montero-Pérez, Y.M., Orozco-Méndez, K.E., Assia-Mercado, Y.M., 2014. *Toxoplasma gondii* detection by gene B1 amplification in meat human consumption. Asociación Colombiana de Infectología 18, 93–99.
- Canada, N., Meireles, C.S., Rocha, A., da Costa, J.M., Erickson, M.W., Dubey, J.P., 2002. Isolation of viable *Toxoplasma gondii* from naturally infected aborted bovine fetuses. J. Parasitol. 88, 1247–1248.
- Carson, A., Group, A.S.R.E., 2017. Focus on ovine abortions. Vet. Rec. 180, 246–247.
- Castano, P., Fuertes, M., Ferre, I., Fernandez, M., Ferreras Mdel, C., Moreno-Gonzalo, J., Gonzalez-Lanza, C., Katzer, F., Regidor-Cerrillo, J., Ortega-Mora, L.M., Perez, V., Benavides, J., 2014. Placental thrombosis in acute phase abortions during experimental *Toxoplasma gondii* infection in sheep. Vet. Res. 45, 9.
- Castano, P., Fuertes, M., Regidor-Cerrillo, J., Ferre, I., Fernandez, M., Ferreras, M.C., Moreno-Gonzalo, J., Gonzalez-Lanza, C., Pereira-Bueno, J., Katzer, F., Ortega-Mora, L.M., Perez, V., Benavides, J., 2016. Experimental ovine toxoplasmosis: influence of the gestational stage on the clinical course, lesion development and parasite distribution. Vet. Res. 47, 43.
- Catar, G., Bergendi, L., Holkova, R., 1969. Isolation of *Toxoplasma gondii* from swine and cattle. J. Parasitol. 55, 952–955.
- Cazarotto, C.J., Balzan, A., Grosskopf, R.K., Boito, J.P., Portella, L.P., Vogel, F.F., Favero, J.F., de C Cucco, D., Biazus, A.H., Machado, G., Da Silva, A.S., 2016. Horses seropositive for *Toxoplasma gondii*, *Sarcocystis* spp. and *Neospora* spp.: possible risk factors for infection in Brazil. Microb. Pathog. 99, 30–35.
- Cenci-Goga, B.T., Ciampelli, A., Sechi, P., Veronesi, F., Moretta, I., Cambiotti, V., Thompson, P.N., 2013. Seroprevalence and risk factors for *Toxoplasma gondii* in sheep in Grosseto district, Tuscany, Italy. BMC Vet. Res. 9, 25.
- Chaichan, P., Mercier, A., Galal, L., Mahittikorn, A., Arie, F., Morand, S., Boumediene, F., Udonsom, R., Hamidovic, A., Murat, J.B., Sukthana, Y., Darde, M.L., 2017. Geographical distribution of *Toxoplasma gondii* genotypes in Asia: a link with neighboring continents. Infect. Genet. Evol. 53, 227–238.
- Chiari, C. de A., Neves, D.P., 1984. Human toxoplasmosis acquired by ingestion of goat's milk. Mem. Inst. Oswaldo Cruz 79, 337–340.
- Chikweto, A., Kuntthekar, S., Tiwari, K., Nyack, B., Deakar, M.S., Stratton, G., Macpherson, C.N., Sharma, R.N., Dubey, J.P., 2011. Seroprevalence of *Toxoplasma gondii* in pigs, sheep, goats, and cattle from Grenada and Carriacou, West Indies. J. Parasitol. 97, 950–951.
- Condoleo, R., Musella, V., Maurelli, M.P., Bosco, A., Cringoli, G., Rinaldi, L., 2016. Mapping, cluster detection and evaluation of risk factors of ovine toxoplasmosis in Southern Italy. Geospat. Health 11, 432.
- Cong, W., Huang, S.Y., Zhou, D.H., Xu, M.J., Wu, S.M., Yan, C., Zhao, Q., Song, H.Q., Zhu, X.Q., 2012. First report of *Toxoplasma gondii* infection in market-sold adult chickens, ducks and pigeons in northwest China. Parasit. Vectors 5, 110.
- Conrad, P.A., Barr, B.C., Sverlow, K.W., Anderson, M., Draft, B., Kinde, H., Dubey, J.P., Munson, L., Ardans, A., 1993. In vitro isolation and characterization of a *Neospora* sp. from aborted bovine foetuses. Parasitology 106, 239–249.
- Consalter, A., Silva, A.F., Frazao-Teixeira, E., Matos, L.F., de Oliveira, F.C.R., Leite, J.S., Silva, F.B.F., Ferreira, A.M.R., 2017. *Toxoplasma gondii* transmission by artificial insemination in sheep with experimentally contaminated frozen semen. Theriogenology 90, 169–174.
- Cook, A.J.C., Gilbert, R.E., Buffolano, W., Zufferey, J., Petersen, E., Jenum, P.A., Foulon, W., Sempirini, A.E., Dunn, D.T., 2000. Sources of toxoplasma infection in pregnant women: European multicentre case-control study. Br. Med. J. 321, 142–147.
- Cosendey-KezenLeite, R.I., de Oliveira, F.C., Frazao-Teixeira, E., Dubey, J.P., de Souza, G.N., Ferreira, A.M., Lilienbaum, W., 2014. Occurrence and risk factors associated to *Toxoplasma gondii* infection in sheep from Rio de Janeiro, Brazil. Trop. Anim. Health Prod. 46, 1463–1466.
- Costa, A.J., Araujo, F.G., Costa, J.O., Lima, J.D., Nascimento, E., 1977. Experimental infection of bovines with oocysts of *Toxoplasma gondii*. J. Parasitol. 63, 212–218.
- Costa, G.H., da Costa, A.J., Lopes, W.D., Bresciani, K.D., dos Santos, T.R., Esper, C.R., Santana, A.E., 2011. *Toxoplasma gondii*: infection natural congenital in cattle and an experimental inoculation of gestating cows with oocysts. Exp. Parasitol. 127, 277–281.
- Costa, D.G., Marvulo, M.F., Silva, J.S., Santana, S.C., Magalhaes, F.J., Filho, C.D., Ribeiro, V.O., Alves, L.C., Mota, R.A., Dubey, J.P., Silva, J.C., 2012. Seroprevalence of *Toxoplasma gondii* in domestic and wild animals from the Fernando de Noronha, Brazil. J. Parasitol. 98, 679–680.
- D'Alencar Mendonca, C.E., Bomfim Barros, S.L., Accioly Guimarães, V.A., Ferraudo, A.S., Munhoz, A.D., 2013. Prevalence and risk factors associated to ovine toxoplasmosis in northeastern Brazil. Rev. Bras. Parasitol. Vet. 22, 230–234.
- Damriyasa, I.M., Bauer, C., Edelhofer, R., Failing, K., Lind, P., Petersen, E., Schares, G., Tenter, A.M., Volmer, R., Zahner, H., 2004. Cross-sectional survey in pig breeding farms in Hesse, Germany: seroprevalence and risk factors of infections with *Toxoplasma gondii*, *Sarcocystis* spp. and *Neospora caninum* in sows. Vet. Parasitol. 126, 271–286.
- Danehchin, L., Razmi, G., Naghibi, A., 2016. Isolation and genotyping of *Toxoplasma gondii* strains in ovine aborted fetuses in Khorasan Razavi Province, Iran. Korean J. Parasitol. 54, 15–20.
- Dangoudoubiyam, S., Oliveira, J.B., Viquez, C., Gomez-Garcia, A., Gonzalez, O., Romero, J.J., Kwok, O.C., Dubey, J.P., Howe, D.K., 2011. Detection of antibodies against *Sarcocystis neurona*, *Neospora* spp., and *Toxoplasma gondii* in horses from Costa Rica. J. Parasitol. 97, 522–524.
- de Barros Correia, E.L., Feitosa, T.F., dos Santos, F.A., de Azevedo, S.S., de Jesus Pena, H.F., Gennari, S.M., Mota, R.A., Alves, C.J., 2015. Prevalence and risk factors for *Toxoplasma gondii* in sheep in the State of Paraíba, Northeastern Brazil. Rev. Bras. Parasitol. Vet. 24, 383–386.
- De Berardinis, A., Paludi, D., Pennisi, L., Vergara, A., 2017. *Toxoplasma gondii*, a foodborne pathogen in the swine production chain from a European perspective. Foodborne Pathog. Dis. 14, 637–648.
- de Macedo, M.F., de Macedo, C.A., Ewald, M.P., Martins, G.F., Zulpo, D.L., da Cunha, I.A., Taroda, A., Cardim, S.T., Su, C., Garcia, J.L., 2012. Isolation and genotyping of *Toxoplasma gondii* from pregnant dairy cows (*Bos taurus*) slaughtered. Rev. Bras. Parasitol. Vet. 21, 74–77.
- de Moraes, E.P., Batista, A.M., Faria, E.B., Freire, R.L., Freitas, A.C., Silva, M.A., Braga, V.A., Mota, R.A., 2010. Experimental infection by *Toxoplasma gondii* using contaminated semen containing different doses of tachyzoites in sheep. Vet. Parasitol. 170, 318–322.
- de Moraes, E.P., da Costa, M.M., Dantas, A.F., da Silva, J.C., Mota, R.A., 2011. *Toxoplasma gondii* diagnosis in ovine aborted fetuses and stillborns in the State of Pernambuco, Brazil. Vet. Parasitol. 183, 152–155.
- de Moura, A.B., Ribeiro, A., de Souza, A.P., da Silva, M.O., Machado, G., Klauk, V., Pazinato, R., Da Silva, A.S., 2016. Seroprevalence and risk factors for *Toxoplasma gondii* infection in goats in Southern Brazil. Acta Sci. Vet. 44.
- de Oliveira, F.C.R., da Coosta, A.J., Bechara, G.H., Sabatini, G.A., 2001. Distribuicao e viabilidade de cistos de *Toxoplasma gondii* (Apicomplexa: Toxoplasmatinae) em tecidos de *Bos indicus*, *Bos taurus* e *Bos bulbalis* infectados com oocistos. Rev. Bras. Med. Vet. 23, 28–34.
- de Oliveira, E., de Albuquerque, P.P., de Souza Neto, O.L., Faria, E.B., Junior, J.W., Mota, R.A., 2013. Occurrence of antibodies to *Toxoplasma gondii* in mules and donkeys in the northeast of Brazil. J. Parasitol. 99, 343–345.

- de Santana Rocha, D., de Sousa Moura, R.L., Maciel, B.M., Guimaraes, L.A., O'Dwyer, H.N., Munhoz, A.D., Albuquerque, G.R., 2015. Detection of *Toxoplasma gondii* DNA in naturally infected sheep's milk. *Genet. Mol. Res.* 14, 8658–8662.
- de Sousa, R.A., Lemos Jda, F., Farias, L.A., Lopes, C.D., dos Santos, K.R., 2014. Seroprevalence and risk factors for *Toxoplasma gondii* infection in pigs in southern Piauí. *Rev. Bras. Parasitol.* Vet. 23, 98–100.
- de Souza, J.B., Soares, V.E., Maia, M.O., Pereira, C.M., Ferraudo, A.S., Cruz, B.C., Pires Teixeira, W.F., Felippelli, G., Maciel, W.G., Goncalves, W.A.J., da Costa, A.J., Zanetti Lopes, W.D., 2016. Spatial distribution and risk factors for *Toxoplasma gondii* seropositivity in cattle slaughtered for human consumption in Rondonia, north region, Brazil. *Vet. Parasitol.* 226, 145–149.
- Dehkordi, F.S., Borujeni, M.R., Rahimi, E., Abdzadeh, R., 2013. Detection of *Toxoplasma gondii* in raw caprine, ovine, buffalo, bovine, and camel milk using cell cultivation, cat bioassay, capture ELISA, and PCR methods in Iran. *Foodborne Pathog. Dis.* 10, 120–125.
- Deksne, G., Kirjusina, M., 2013. Seroprevalence of *Toxoplasma gondii* in domestic pigs (*Sus scrofa domestica*) and wild boars (*Sus scrofa*) in Latvia. *J. Parasitol.* 99, 44–47.
- Deksne, G., Ligere, B., Sneidera, A., Jokelainen, P., 2017. Seroprevalence and factors associated with *Toxoplasma gondii* infections in sheep in Latvia: Latvian dark headed sheep breed associated with higher seroprevalence. *Vector Borne Zoonotic Dis.* 17, 478–482.
- Dempster, R.P., Wilkins, M., Green, R.S., de Lisle, G.W., 2011. Serological survey of *Toxoplasma gondii* and *Campylobacter fetus fetus* in sheep from New Zealand. *N. Z. Vet. J.* 59, 155–159.
- Deng, H., Dam-Deisz, C., Luttiikholt, S., Maas, M., Nielen, M., Swart, A., Vellema, P., van der Giessen, J., Opsteegh, M., 2016. Risk factors related to *Toxoplasma gondii* seroprevalence in indoor-housed Dutch dairy goats. *Prev. Vet. Med.* 124, 45–51.
- Deng, H., Devleeschauwer, B., Liu, M., Li, J., Wu, Y., van der Giessen, J.W.B., Opsteegh, M., 2018. Seroprevalence of *Toxoplasma gondii* in pregnant women and livestock in the mainland of China: a systematic review and hierarchical meta-analysis. *Sci. Rep.* 8, 6218.
- Diakoua, A., Papadopoulos, E., Panousis, N., Karatzias, C., Giadinis, N., 2013. *Toxoplasma gondii* and *Neospora caninum* seroprevalence in dairy sheep and goats mixed stock farming. *Vet. Parasitol.* 198, 387–390.
- Diaz, P., Cabanelas, E., Diaz-Cao, J.M., Vina, M., Bejar, J.P., Perez-Creo, A., Prieto, A., Lopez, C.M., Panadero, R., Fernandez, G., Diez-Banos, P., Morrondo, P., 2016. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in goats from north-western Spain. *Ann. Agric. Environ. Med.* 23, 587–590.
- Dijkhuizen, A.A., Morris, R.S., 1997. *Animal Health Economics: Principles and Applications*. Wageningen Pers., Sydney.
- Djokic, V., Klun, I., Musella, V., Rinaldi, L., Cringoli, G., Sotiraki, S., Djurkovic-Djakovic, O., 2014. Spatial epidemiology of *Toxoplasma gondii* infection in goats in Serbia. *Geospat. Health* 8, 479–488.
- Djokic, V., Fablet, C., Blaga, R., Rose, N., Perret, C., Djurkovic-Djakovic, O., Boireau, P., Durand, B., 2016. Factors associated with *Toxoplasma gondii* infection in confined farrow-to-finish pig herds in western France: an exploratory study in 60 herds. *Parasit. Vectors* 9, 466.
- Dong, H., Su, R., Lu, Y., Wang, M., Liu, J., Jian, F., Yang, Y., 2018. Prevalence, risk factors, and genotypes of *Toxoplasma gondii* in food animals and humans (2000–2017) from China. *Front. Microbiol.* 9, 2108.
- Dos Santos, T.R., Faria, G.D., Guerreiro, B.M., Dal Pietro, N.H., Lopes, W.D., da Silva, H.M., Garcia, J.L., Luvizotto, M.C., Bresciani, K.D., da Costa, A.J., 2016. Congenital toxoplasmosis in chronically infected and subsequently challenged and persistently infected ewes. *PLoS ONE* 11, e0165124.
- Du, F., Zhang, Q., Yu, Q., Hu, M., Zhou, Y., Zhao, J., 2012. Soil contamination of *Toxoplasma gondii* oocysts in pig farms in central China. *Vet. Parasitol.* 187, 53–56.
- Dubey, J.P., 1981. Protective immunity against clinical toxoplasmosis in dairy goats vaccinated with *Hammondia hammondi* and *Hammondia heydorni*. *Am. J. Vet. Res.* 42, 2068–2070.
- Dubey, J.P., 1982. Repeat transplacental transfer of *Toxoplasma gondii* in dairy goats. *J. Am. Vet. Med. Assoc.* 180, 1220–1221.
- Dubey, J.P., 1983. Distribution of cysts and tachyzoites in calves and pregnant cows inoculated with *Toxoplasma gondii* oocysts. *Vet. Parasitol.* 13, 199–211.
- Dubey, J.P., 1985. Persistence of encysted *Toxoplasma gondii* in tissues of equids fed oocysts. *Am. J. Vet. Res.* 46, 1753–1754.
- Dubey, J.P., 1986a. A review of toxoplasmosis in cattle. *Vet. Parasitol.* 22, 177–202.
- Dubey, J.P., 1986b. A review of toxoplasmosis in pigs. *Vet. Parasitol.* 19, 181–223.
- Dubey, J.P., 1992. Isolation of *Toxoplasma gondii* from a naturally infected beef cow. *J. Parasitol.* 78, 151–153.
- Dubey, J.P., 2009a. Toxoplasmosis in pigs—the last 20 years. *Vet. Parasitol.* 164, 89–103.
- Dubey, J.P., 2009b. Toxoplasmosis in sheep—the last 20 years. *Vet. Parasitol.* 163, 1–14.
- Dubey, J.P., 2010a. *Toxoplasma gondii* infections in chickens (*Gallus domesticus*): prevalence, clinical disease, diagnosis and public health significance. *Zoonoses Public Health* 57, 60–73.
- Dubey, J.P., 2010b. *Toxoplasmosis of Animals and Humans*. 2nd edition. CRC Press, Boca Raton.
- Dubey, J.P., Beattie, C.P., 1988. *Toxoplasmosis of Animals and Man*. CRC Press, Boca Raton, Florida, USA, pp. 1–220.
- Dubey, J.P., Desmonts, G., 1987. Serological responses of equids fed *Toxoplasma gondii* oocysts. *Equine Vet. J.* 19, 337–339.
- Dubey, J.P., Jones, J.L., 2014. Comments on “detection of *Toxoplasma gondii* in raw caprine, ovine, buffalo, bovine, and camel milk using cell cultivation, cat bioassay, capture ELISA, and PCR methods in Iran”. *Foodborne Pathog. Dis.* 11, 500–501.
- Dubey, J.P., Streitl, R.H., 1976. Prevalence of *Toxoplasma* infection in cattle slaughtered at an Ohio abattoir. *J. Am. Vet. Med. Assoc.* 169, 1197–1199.
- Dubey, J.P., Thulliez, P., 1993. Persistence of tissue cysts in edible tissues of cattle fed *Toxoplasma gondii* oocysts. *Am. J. Vet. Res.* 54, 270–273.
- Dubey, J.P., Urban Jr., J.F., 1990. Diagnosis of transplacentally induced toxoplasmosis in pigs. *Am. J. Vet. Res.* 51, 1295–1299.
- Dubey, J.P., Camargo, M.E., Ruff, M.D., Wilkins, G.C., Shen, S.K., Kwok, O.C., Thulliez, P., 1993a. Experimental toxoplasmosis in turkeys. *J. Parasitol.* 79, 949–952.
- Dubey, J.P., Ruff, M.D., Camargo, M.E., Shen, S.K., Wilkins, G.L., Kwok, O.C., Thulliez, P., 1993b. Serologic and parasitologic responses of domestic chickens after oral inoculation with *Toxoplasma gondii* oocysts. *Am. J. Vet. Res.* 54, 1668–1672.
- Dubey, J.P., Garner, M.W., Willette, M.M., Batey, K.L., Gardiner, C.H., 2001. Disseminated toxoplasmosis in magpie geese (*Anseranas semipalmata*) with large numbers of tissue cysts in livers. *J. Parasitol.* 87, 219–223.
- Dubey, J.P., Hill, D.E., Jones, J.L., Hightower, A.W., Kirkland, E., Roberts, J.M., Marcet, P.L., Lehmann, T., Vianna, M.C., Miska, K., Sreekumar, C., Kwok, O.C., Shen, S.K., Gamble, H.R., 2005. Prevalence of viable *Toxoplasma gondii* in beef, chicken, and pork from retail meat stores in the United States: risk assessment to consumers. *J. Parasitol.* 91, 1082–1093.
- Dubey, J.P., Sundar, N., Hill, D., Velmurugan, G.V., Bandini, L.A., Kwok, O.C., Majumdar, D., Su, C., 2008. High prevalence and abundant atypical genotypes of *Toxoplasma gondii* isolated from lambs destined for human consumption in the USA. *Int. J. Parasitol.* 38, 999–1006.
- Dubey, J.P., Rajendran, C., Ferreira, L.R., Martins, J., Kwok, O.C., Hill, D.E., Villena, I., Zhou, H., Su, C., Jones, J.L., 2011. High prevalence and genotypes of *Toxoplasma gondii* isolated from goats from a retail meat store, destined for human consumption in the USA. *Int. J. Parasitol.* 41, 827–833.
- Dubey, J.P., Hotea, I., Olariu, T.R., Jones, J.L., Darabus, G., 2014a. Epidemiological review of toxoplasmosis in humans and animals in Romania. *Parasitology* 141, 311–325.
- Dubey, J.P., Ness, S.L., Kwok, O.C., Choudhary, S., Mittel, L.D., Divers, T.J., 2014b. Seropositivity of *Toxoplasma gondii* in domestic donkeys (*Equus asinus*) and isolation of *T. gondii* from farm cats. *Vet. Parasitol.* 199, 18–23.
- Dubey, J.P., Hemphill, A., Calero-Bernal, R., Schares, G., 2017. *Neosporosis in Animals*. CRC Press, Boca Rotan.
- Elbez-Rubinstein, A., Aizenberg, D., Darde, M.L., Cohen, R., Dumetre, A., Yera, H., Gondon, E., Janaud, J.C., Thulliez, P., 2009. Congenital toxoplasmosis and reinfection during pregnancy: case report, strain characterization, experimental model of reinfection, and review. *J. Infect. Dis.* 199, 280–285.
- Ellis, J.T., 1998. Polymerase chain reaction approaches for the detection of *Neospora caninum* and *Toxoplasma gondii*. *Int. J. Parasitol.* 28, 1053–1060.
- El-Massry, A., Mahdy, O.A., El-Ghaysh, A., Dubey, J.P., 2000. Prevalence of *Toxoplasma gondii* antibodies in sera of turkeys, chickens, and ducks from Egypt. *J. Parasitol.* 86, 627–628.
- Ergin, S., Ciftcioglu, G., Midilli, K., Issa, G., Gargili, A., 2009. Detection of *Toxoplasma gondii* from meat and meat products by the nested-PCR method and its relationship with seroprevalence in slaughtered animals. *Bull. Vet. Inst. Pulawy* 53, 657–661.
- Esteban-Redondo, I., Maley, S.W., Thomson, K., Nicoll, S., Wright, S., Buxton, D., Innes, E.A., 1999. Detection of *T. gondii* in tissues of sheep and cattle following oral infection. *Vet. Parasitol.* 86, 155–171.
- Esteves, F., Aguiar, D., Rosado, J., Costa, M.L., de Sousa, B., Antunes, F., Matos, O., 2014. *Toxoplasma gondii* prevalence in cats from Lisbon and in pigs from centre and south of Portugal. *Vet. Parasitol.* 200, 8–12.

- Evers, F., Garcia, J.L., Navarro, I.T., Zulplo, D.L., Nino Bde, S., Ewald, M.P., Pagliari, S., Almeida, J.C., Freire, R.L., 2013. Diagnosis and isolation of *Toxoplasma gondii* in horses from Brazilian slaughterhouses. *Rev. Bras. Parasitol. Vet.* 22, 58–63.
- Fajardo, H.V., D'Avila, S., Bastos, R.R., Cyrino, C.D., de Lima Detoni, M., Garcia, J.L., das Neves, L.B., Nicolau, J.L., Amendoeira, M.R., 2013. Seroprevalence and risk factors of toxoplasmosis in cattle from extensive and semi-intensive rearing systems at Zona da Mata, Minas Gerais state, Southern Brazil. *Parasit. Vectors* 6, 191.
- Finger, M.A., Villalobos, E.M.C., Lara, M.D.C.S.H., Cunha, E.M.S., De Barros, I.R., Dornbusch, P.T., Ullmann, L.S., Biondo, A.W., 2013. Detection of anti-*Toxoplasma gondii* antibodies in carthorses in the metropolitan region of Curitiba, Parana, Brazil. *Rev. Bras. Parasitol. Vet.* 22, 179–181.
- Fonseca de Araujo Valenca, S.R., Barreto Valenca, R.M., Pinheiro Junior, J.W., Feitosa de Albuquerque, P.P., Souza Neto, O.L., Mota, R.A., 2015. Risk factors of occurrence of *Toxoplasma gondii* among horses in the state of Alagoas, Brazil. *Acta Parasitol.* 60, 707–711.
- Fortier, B., Dealmeida, E., Pinto, I., Ajana, F., Camus, D., 1990. Prevalence de la toxoplasmose porcine et bovine a Porto. *Med. Mal. Infect.* 20, 551–554.
- Freyre, A., Falcon, J., Cedda, C., 1990. El papel de la carne vacuna en la adquisicion de la infeccion toxoplasmica. *Rev. Iber. Parasitol.* 50, 15–24.
- Freyre, A., Bonino, J., Falcon, J., Castells, D., Correa, O., Casaretto, A., 1997. The incidence and economic significance of ovine toxoplasmosis in Uruguay. *Vet. Parasitol.* 73, 13–15.
- Garcia, G., Sotomaior, C., do Nascimento, A.J., Navarro, I.T., Soccol, V.T., 2012. *Toxoplasma gondii* in goats from Curitiba, Parana, Brazil: risks factors and epidemiology. *Rev. Bras. Parasitol. Vet.* 21, 42–47.
- Garcia, J.L., Burrells, A., Bartley, P.M., Bartley, K., Innes, E.A., Katzer, F., 2017. The use of ELISA, nPCR and qPCR for diagnosis of ocular toxoplasmosis in experimentally infected pigs. *Res. Vet. Sci.* 115, 490–495.
- Garcia-Bocanegra, I., Dubey, J.P., Simon-Grife, M., Cabezon, O., Casal, J., Allepuz, A., Napp, S., Almeria, S., 2010a. Seroprevalence and risk factors associated with *Toxoplasma gondii* infection in pig farms from Catalonia, north-eastern Spain. *Res. Vet. Sci.* 89, 85–87.
- Garcia-Bocanegra, I., Simon-Grife, M., Dubey, J.P., Casal, J., Martin, G.E., Cabezon, O., Perea, A., Almeria, S., 2010b. Seroprevalence and risk factors associated with *Toxoplasma gondii* in domestic pigs from Spain. *Parasitol. Int.* 59, 421–426.
- Garcia-Bocanegra, I., Cabezon, O., Arenas-Montes, A., Carbonero, A., Dubey, J.P., Perea, A., Almeria, S., 2012. Seroprevalence of *Toxoplasma gondii* in equids from Southern Spain. *Parasitol. Int.* 61, 421–424.
- Garcia-Bocanegra, I., Cabezon, O., Hernandez, E., Martinez-Cruz, M.S., Martinez-Moreno, A., Martinez-Moreno, J., 2013. *Toxoplasma gondii* in ruminant species (cattle, sheep, and goats) from southern Spain. *J. Parasitol.* 99, 438–440.
- Gazzonis, A.L., Veronesi, F., Di Cerbo, A.R., Zanzani, S.A., Molineri, G., Moretta, I., Moretti, A., Piergili Fioretti, D., Invernizzi, A., Manfredi, M.T., 2015. *Toxoplasma gondii* in small ruminants in Northern Italy - prevalence and risk factors. *Ann. Agric. Environ. Med.* 22, 62–68.
- Gebremedhin, E.Z., Agonafir, A., Tessema, T.S., Tilahun, G., Medhin, G., Vitale, M., Di Marco, V., Cox, E., Vercurysse, J., Dorny, P., 2013. Seroprevalence of ovine toxoplasmosis in East and West Sheva Zones of Oromia Regional State, Central Ethiopia. *BMC Vet. Res.* 9, 117.
- Gebremedhin, E.Z., Abdurahman, M., Tessema, T.S., Tilahun, G., Cox, E., Goddeeris, B., Dorny, P., De Craeye, S., Darde, M.L., Ajzenberg, D., 2014. Isolation and genotyping of viable *Toxoplasma gondii* from sheep and goats in Ethiopia destined for human consumption. *Parasit. Vectors* 7, 425.
- Gebreyes, W.A., Bahnsen, P.B., Funk, J.A., McKean, J., Patchanee, P., 2008. Seroprevalence of Trichinella, Toxoplasma, and Salmonella in antimicrobial-free and conventional swine production systems. *Foodborne Pathog. Dis.* 5, 199–203.
- Genchi, M., Vismarra, A., Mangia, C., Faccini, S., Vicari, N., Rigamonti, S., Prati, P., Marino, A.M., Kramer, L., Fabbri, M., 2017. Lack of viable parasites in cured 'Parma Ham' (PDO), following experimental *Toxoplasma gondii* infection of pigs. *Food Microbiol.* 66, 157–164.
- Gennari, S.M., Esmerini, P.D., Lopes, M.G., Soares, H.S., Vitaliano, S.N., Cabral, A.D., Pena, H.F.J., Horta, M.C., Cavalcante, P.H., Fortes, K.P., Villalobos, E.M.C., 2015. Occurrence of antibodies against *Toxoplasma gondii* and its isolation and genotyping in donkeys, mules, and horses in Brazil. *Vet. Parasitol.* 209, 129–132.
- Gilot-Fromont, E., Aubert, D., Belkilani, S., Hermitte, P., Gibout, O., Geers, R., Villena, I., 2009. Landscape, herd management and within-herd seroprevalence of *Toxoplasma gondii* in beef cattle herds from Champagne-Ardenne, France. *Vet. Parasitol.* 161, 36–40.
- Goerlich, K., 2011. Toxoplasma-Infektionen bei Schweinen: Semi-automatisiertes Testsystem für Surveillance und Monitoring. Querschnittsstudie in Niedersachsen. Tierärztliche Hochschule Hannover, Hannover.
- Gottstein, B., Hentrich, B., Wyss, R., Thur, B., Busato, A., Stark, K.D.C., Muller, N., 1998. Molecular and immunodiagnostic investigations on bovine neosporosis in Switzerland. *Int. J. Parasitol.* 28, 679–691.
- Guerra, N.R., Almeida, J.C., Silva, E.L., Silva, E.M., Santos, J.A., Lepold, R., Mota, R.A., Alves, L.C., 2018. Soroprevalência de *Toxoplasma gondii* em equídeos do Nordeste do Brasil. *Pesqui. Vet. Bras.* 38, 400–406.
- Guimaraes, L.A., Bezerra, R.A., Rocha Dde, S., Albuquerque, G.R., 2013. Prevalence and risk factors associated with anti-*Toxoplasma gondii* antibodies in sheep from Bahia state, Brazil. *Rev. Bras. Parasitol. Vet.* 22, 220–224.
- Guo, M., Dubey, J.P., Hill, D., Buchanan, R.L., Gamble, H.R., Jones, J.L., Pradhan, A.K., 2015. Prevalence and risk factors for *Toxoplasma gondii* infection in meat animals and meat products destined for human consumption. *J. Food Prot.* 78, 457–476.
- Guo, M., Mishra, A., Buchanan, R.L., Dubey, J.P., Hill, D.E., Gamble, H.R., Jones, J.L., Pradhan, A.K., 2016. A systematic meta-analysis of *Toxoplasma gondii* prevalence in food animals in the United States. *Foodborne Pathog. Dis.* 13, 109–118.
- Gutierrez, J., O'Donovan, J., Williams, E., Proctor, A., Brady, C., Marques, P.X., Worrall, S., Nally, J.E., McElroy, M., Bassett, H., Sammin, D., Buxton, D., Maley, S., Markey, B.K., 2010. Detection and quantification of *Toxoplasma gondii* in ovine maternal and foetal tissues from experimentally infected pregnant ewes using real-time PCR. *Vet. Parasitol.* 172, 8–15.
- Gutierrez, J., O'Donovan, J., Proctor, A., Brady, C., Marques, P.X., Worrall, S., Nally, J.E., McElroy, M., Bassett, H., Fagan, J., Maley, S., Buxton, D., Sammin, D., Markey, B.K., 2012. Application of quantitative real-time polymerase chain reaction for the diagnosis of toxoplasmosis and enzootic abortion of ewes. *J. Vet. Diagn. Investig.* 24, 846–854.
- Habibi, G., Imani, A., Gholami, M., Hablolvarid, M., Behroozkikhah, A., Lotfi, M., Kamalzade, M., Najjar, E., Esmaeil-Nia, K., Bozorgi, S., 2012. Detection and identification of *Toxoplasma gondii* type one infection in sheep aborted fetuses in Qazvin province of Iran. *Iran. J. Parasitol.* 7, 64–72.
- Hajjalilo, E., Ziaali, N., Harandi, M.F., Saraei, M., Hajjalilo, M., 2010. Prevalence of anti-*Toxoplasma gondii* antibodies in sport horses from Qazvin, Iran. *Trop. Anim. Health Prod.* 42, 1321–1322.
- Halos, L., Thebault, A., Aubert, D., Thomas, M., Perret, C., Geers, R., Alliot, A., Escotte-Binet, S., Ajzenberg, D., Darde, M.L., Durand, B., Boireau, P., Villena, I., 2010. An innovative survey underlining the significant level of contamination by *Toxoplasma gondii* of ovine meat consumed in France. *Int. J. Parasitol.* 40, 193–200.
- Hamilton, C.M., Katzer, F., Innes, E.A., Kelly, P.J., 2014. Seroprevalence of *Toxoplasma gondii* in small ruminants from four Caribbean islands. *Parasit. Vectors* 7, 449.
- Hamilton, C.M., Kelly, P.J., Boey, K., Corey, T.M., Huynh, H., Metzler, D., Villena, I., Su, C., Innes, E.A., Katzer, F., 2017. Predominance of atypical genotypes of *Toxoplasma gondii* in free-roaming chickens in St. Kitts, West Indies. *Parasit. Vectors* 10, 104.
- Hammond-Aryee, K., Van Helden, L.S., Van Helden, P.D., 2015. The prevalence of antibodies to *Toxoplasma gondii* in sheep in the Western Cape, South Africa. *Onderstepoort J. Vet. Res.* 82, e1–e5.
- Harfoush, M., Tahoou Ael, N., 2010. Seroprevalence of *Toxoplasma gondii* antibodies in domestic ducks, free-range chickens, turkeys and rabbits in Kafr El-Sheikh Governorate Egypt. *J. Egypt. Soc. Parasitol.* 40, 295–302.
- Haridy, F.M., Saleh, N.M., Khalil, H.H., Morsy, T.A., 2010. Anti-*Toxoplasma gondii* antibodies in working donkeys and donkey's milk in greater Cairo, Egypt. *J. Egypt. Soc. Parasitol.* 40, 459–464.
- Herrero, L., Gracia, M.J., Perez-Arquillue, C., Lazaro, R., Herrera, M., Herrera, A., Bayarri, S., 2016. *Toxoplasma gondii*: pig seroprevalence, associated risk factors and viability in fresh pork meat. *Vet. Parasitol.* 224, 52–59.
- Hill, D.E., Dubey, J.P., 2013. *Toxoplasma gondii* prevalence in farm animals in the United States. *Int. J. Parasitol.* 43, 107–113.
- Hill, D.E., Haley, C., Wagner, B., Gamble, H.R., Dubey, J.P., 2010. Seroprevalence of and risk factors for *Toxoplasma gondii* in the US swine herd using sera collected during the National Animal Health Monitoring Survey (swine 2006). *Zoonoses Public Health* 57, 53–59.
- Hill, D., Coss, C., Dubey, J.P., Wroblewski, K., Sautter, M., Hosten, T., Munoz-Zanzi, C., Mui, E., Withers, S., Boyer, K., Hermes, G., Coyne, J., Jagdis, F., Burnett, A., McLeod, P., Morton, H., Robinson, D., McLeod, R., 2011. Identification of a sporozoite-specific antigen from *Toxoplasma gondii*. *J. Parasitol.* 97, 328–337.

- Holec-Gasior, L., Dominiak-Gorski, B., Kur, J., 2015. First report of seroprevalence of *Toxoplasma gondii* infection in sheep in Pomerania, northern Poland. *Ann. Agric. Environ. Med.* 22, 604–607.
- Hosein, S., Limon, G., Dadios, N., Guitian, J., Blake, D.P., 2016. *Toxoplasma gondii* detection in cattle: a slaughterhouse survey. *Vet. Parasitol.* 228, 126–129.
- Hou, Z.F., Su, S.J., Liu, D.D., Wang, L.L., Jia, C.L., Zhao, Z.X., Ma, Y.F., Li, Q.Q., Xu, J.J., Tao, J.P., 2018. Prevalence, risk factors and genetic characterization of *Toxoplasma gondii* in sick pigs and stray cats in Jiangsu Province, eastern China. *Infect. Genet. Evol.* 60, 17–25.
- Hutchinson, J.P., Wear, A.R., Lambton, S.L., Smith, R.P., Pritchard, G.C., 2011. Survey to determine the seroprevalence of *Toxoplasma gondii* infection in British sheep flocks. *Vet. Rec.* 169, 582.
- Innes, E.A., Panton, W.R., Thomson, K.M., Maley, S., Buxton, D., 1995. Kinetics of interferon gamma production in vivo during infection with the S48 vaccine strain of *Toxoplasma gondii*. *J. Comp. Pathol.* 113, 89–94.
- Innes, E.A., Bartley, P.M., Buxton, D., Katzer, F., 2009. Ovine toxoplasmosis. *Parasitology* 136, 1887–1894.
- Iovu, A., Titilincu, A., Mircean, V., Junie, M., Cozma, V., 2010. Identification of *Toxoplasma gondii* and *Hammondia hammondi* in sow abortions. *Int. J. Mol. Med.* 26, S71.
- Iovu, A., Gyorke, A., Mircean, V., Gavrea, R., Cozma, V., 2012. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in dairy goats from Romania. *Vet. Parasitol.* 186, 470–474.
- Isaac-Renton, J., Bowie, W.R., King, A., Irwin, G.S., Ong, C.S., Fung, C.P., Shokeir, M.O., Dubey, J.P., 1998. Detection of *Toxoplasma gondii* oocysts in drinking water. *Appl. Environ. Microbiol.* 64, 2278–2280.
- Jacobs, L., Moyle, G.G., 1963. The prevalence of toxoplasmosis in New Zealand sheep and cattle. *Am. J. Vet. Res.* 24, 673–675.
- Jacobs, L., Remington, J.S., Melton, M.L., 1960. A survey of meat samples from swine, cattle, and sheep for the presence of encysted *Toxoplasma*. *J. Parasitol.* 46, 23–28.
- James, K.E., Smith, W.A., Packham, A.E., Conrad, P.A., Pusterla, N., 2017. *Toxoplasma gondii* seroprevalence and association with equine protozoal myeloencephalitis: a case-control study of Icelandic horses. *Vet. J.* 224, 38–43.
- Jerome, M.E., Radke, J.R., Bohne, W., Roos, D.S., White, M.W., 1998. *Toxoplasma gondii* bradyzoites form spontaneously during sporozoite-initiated development? *Infect. Immun.* 66, 4838–4844.
- Jiang, H.H., Huang, S.Y., Zhou, D.H., Zhang, X.X., Su, C., Deng, S.Z., Zhu, X.Q., 2013. Genetic characterization of *Toxoplasma gondii* from pigs from different localities in China by PCR-RFLP. *Parasit. Vectors* 6, 227.
- Jokelainen, P., Nareaho, A., Knaapi, S., Oksanen, A., Rikula, U., Sukura, A., 2010. *Toxoplasma gondii* in wild cervids and sheep in Finland: north-south gradient in seroprevalence. *Vet. Parasitol.* 171, 331–336.
- Jokelainen, P., Tagel, M., Motus, K., Viltrop, A., Lassen, B., 2017. *Toxoplasma gondii* seroprevalence in dairy and beef cattle: large-scale epidemiological study in Estonia. *Vet. Parasitol.* 236, 137–143.
- Jungersen, G., Bille-Hansen, V., Jensen, L., Lind, P., 2001. Transplacental transmission of *Toxoplasma gondii* in minipigs infected with strains of different virulence. *J. Parasitol.* 87, 108–113.
- Kantzoura, V., Diakou, A., Kouam, M.K., Feidas, H., Theodoropoulou, H., Theodoropoulos, G., 2013. Seroprevalence and risk factors associated with zoonotic parasitic infections in small ruminants in the Greek temperate environment. *Parasitol. Int.* 62, 554–560.
- Kapperud, G., Jenum, P.A., Stray-Pedersen, B., Melby, K.K., Eskild, A., Eng, J., 1996. Risk factors for *Toxoplasma gondii* infection in pregnancy: results of a prospective case-control study in Norway. *Am. J. Epidemiol.* 144, 405–412.
- Karatepe, B., Babur, C., Karatepe, M., Kilic, S., 2010. Seroprevalence of toxoplasmosis in horses in Nigde Province of Turkey. *Trop. Anim. Health Prod.* 42, 385–389.
- Katzer, F., Brulisaue, F., Collantes-Fernandez, E., Bartley, P.M., Burrells, A., Gunn, G., Maley, S.W., Cousens, C., Innes, E.A., 2011. Increased *Toxoplasma gondii* positivity relative to age in 125 Scottish sheep flocks; evidence of frequent acquired infection. *Vet. Res.* 42, 121.
- Katzer, F., Canton, G., Burrells, A., Palarea-Albaladejo, J., Horton, B., Bartley, P.M., Pang, Y., Chianini, F., Innes, E.A., Benavides, J., 2014. Immunization of lambs with the S48 strain of *Toxoplasma gondii* reduces tissue cyst burden following oral challenge with a complete strain of the parasite. *Vet. Parasitol.* 205, 46–56.
- Khan, M.U., Rashid, I., Akbar, H., Islam, S., Riaz, F., Nabi, H., Ashraf, K., Singla, L.D., 2017. Seroprevalence of *Toxoplasma gondii* in South Asian countries. *Rev. Sci. Tech.* 36, 981–996.
- Kijlstra, A., Eissen, O.A., Cornelissen, J., Munniksma, K., Eijck, I., Kortbeek, T., 2004. *Toxoplasma gondii* infection in animal-friendly pig production systems. *Invest. Ophthalmol. Vis. Sci.* 45, 3165–3169.
- Kijlstra, A., Meerburg, B., Cornelissen, J., De Craey, S., Vereijken, P., Jongert, E., 2008. The role of rodents and shrews in the transmission of *Toxoplasma gondii* to pigs. *Vet. Parasitol.* 156, 183–190.
- Kim, J.H., Kang, K.I., Kang, W.C., Sohn, H.J., Jean, Y.H., Park, B.K., Kim, Y., Kim, D.Y., 2009. Porcine abortion outbreak associated with *Toxoplasma gondii* in Jeju Island, Korea. *J. Vet. Sci.* 10, 147–151.
- Kittas, C., Henry, L., 1979. Effect of sex hormones on the immune system of guinea-pigs and on the development of toxoplasmic lesions in non-lymphoid organs. *Clin. Exp. Immunol.* 36, 16–23.
- Kittas, C., Henry, L., 1980. Effect of sex hormones on the response of mice to infection with *Toxoplasma gondii*. *Br. J. Exp. Pathol.* 61, 590–600.
- Klauck, V., Pazinato, R., Radavelli, W.M., Custodio, E., Bianchi, A.E., Camillo, G., Cezar, A.S., Vogel, F.F., Tonin, A.A., Ferreira, R., Stefani, L.M., Da Silva, A.S., 2016. *Toxoplasma gondii* infection in dairy ewes: vertical transmission and influence on milk production and reproductive performance. *Microb. Pathog.* 99, 101–105.
- Klein, S., Wendt, M., Baumgartner, W., Wohlsein, P., 2010. Systemic toxoplasmosis and concurrent porcine circovirus-2 infection in a pig. *J. Comp. Pathol.* 142, 228–234.
- Klun, I., Djurkovic-Djakovic, O., Katic-Radivojevic, S., Nikolic, A., 2006. Cross-sectional survey on *Toxoplasma gondii* infection in cattle, sheep and pigs in Serbia: seroprevalence and risk factors. *Vet. Parasitol.* 135, 121–131.
- Klun, I., Uzelac, A., Villena, I., Mercier, A., Bobic, B., Nikolic, A., Rajnpreht, I., Opsteegh, M., Aubert, D., Blaga, R., van der Giessen, J., Djurkovic-Djakovic, O., 2017. The first isolation and molecular characterization of *Toxoplasma gondii* from horses in Serbia. *Parasit. Vectors* 10, 167.
- Koethe, M., Pott, S., Ludewig, M., Bangoura, B., Zoller, B., Dausgies, A., Tenter, A.M., Spekker, K., Bittame, A., Mercier, C., Fehlhauer, K., Straubinger, R.K., 2011. Prevalence of specific IgG-antibodies against *Toxoplasma gondii* in domestic turkeys determined by kinetic ELISA based on recombinant GRA7 and GRA8. *Vet. Parasitol.* 180, 179–190.
- Koethe, M., Straubinger, R.K., Pott, S., Bangoura, B., Geuthner, A.C., Dausgies, A., Ludewig, M., 2015. Quantitative detection of *Toxoplasma gondii* in tissues of experimentally infected turkeys and in retail turkey products by magnetic-capture PCR. *Food Microbiol.* 52, 11–17.
- Kouam, M.K., Diakou, A., Kanzoura, V., Papadopoulos, E., Gajadhar, A.A., Theodoropoulos, G., 2010. A seroepidemiological study of exposure to *Toxoplasma*, *Leishmania*, *Echinococcus* and *Trichinella* in equids in Greece and analysis of risk factors. *Vet. Parasitol.* 170, 170–175.
- Lahmar, I., Lachkhem, A., Slama, D., Sakly, W., Haouas, N., Gorcii, M., Pfaff, A.W., Candolfi, E., Babba, H., 2015. Prevalence of toxoplasmosis in sheep, goats and cattle in Southern Tunisia. *J. Bacteriol. Parasitol.* 6.
- Lee, S.H., Lee, S.E., Seo, M.G., Goo, Y.K., Cho, K.H., Cho, G.J., Kwon, O.D., Kwak, D., Lee, W.J., 2014. Evidence of *Toxoplasma gondii* exposure among horses in Korea. *J. Vet. Med. Sci.* 76, 1663–1665.
- Li, X., Wang, Y., Yu, F., Li, T., Zhang, D., 2010. An outbreak of lethal toxoplasmosis in pigs in the Gansu province of China. *J. Vet. Diagn. Investig.* 22, 442–444.
- Limon, G., Beauvais, W., Dadios, N., Villena, I., Cockle, C., Blaga, R., Guitian, J., 2017. Cross-sectional study of *Toxoplasma gondii* infection in pig farms in England. *Foodborne Pathog. Dis.* 14, 269–281.
- Liu, Z.K., Li, J.Y., Pan, H., 2015. Seroprevalence and risk factors of *Toxoplasma gondii* and *Neospora caninum* infections in small ruminants in China. *Prev. Vet. Med.* 118, 488–492.
- Lopes, W.D., Santos, T.R., da Silva Rdos, S., Rossanese, W.M., de Souza, F.A., de Faria Rodrigues, J.D., de Mendonca, R.P., Soares, V.E., Costa, A.J., 2010. Seroprevalence of and risk factors for *Toxoplasma gondii* in sheep raised in the Jaboticabal microregion, Sao Paulo State, Brazil. *Res. Vet. Sci.* 88, 104–106.
- Lopes, W.D., Santos, T.R., Luvizotto, M.C., Sakamoto, C.A., Oliveira, G.P., Costa, A.J., 2011. Histopathology of the reproductive system of male sheep experimentally infected with *Toxoplasma gondii*. *Parasitol. Res.* 109, 405–409.
- Lopes, A.P., Dubey, J.P., Neto, F., Rodrigues, A., Martins, T., Rodrigues, M., Cardoso, L., 2013. Seroprevalence of *Toxoplasma gondii* infection in cattle, sheep, goats and pigs from the north of Portugal for human consumption. *Vet. Parasitol.* 193, 266–269.

- Lopes, W.D., Rodriguez, J.D., Souza, F.A., dos Santos, T.R., dos Santos, R.S., Rosanese, W.M., Lopes, W.R., Sakamoto, C.A., da Costa, A.J., 2013. Sexual transmission of *Toxoplasma gondii* in sheep. *Vet. Parasitol.* 195, 47–56.
- Machacova, T., Bartova, E., Di Loria, A., Sedlak, K., Mariani, U., Fusco, G., Fulgione, D., Veneziano, V., Dubey, J.P., 2014. Seroprevalence of *Toxoplasma gondii* in donkeys (*Equus asinus*) in Italy. *J. Vet. Med. Sci.* 76, 265–267.
- Magalhaes, F.J., Ribeiro-Andrade, M., Alcantara, A.M., Pinheiro, J.W.J., Sena, M.J., Porto, W.J., Vieira, R.F., Mota, R.A., 2016. Risk factors for *Toxoplasma gondii* infection in sheep and cattle from Fernando de Noronha Island, Brazil. *Rev. Bras. Parasitol. Vet.* 25, 511–515.
- Magalhães, F.J.R., da Silva, J.G., Ribeiro-Andrade, M., Pinheiro, J.W., Aparecido Mota, R., 2016. High prevalence of toxoplasmosis in free-range chicken of the Fernando de Noronha Archipelago, Brazil. *Acta Trop.* 159, 58–61.
- Mahami-Oskouei, M., Moradi, M., Fallah, E., Hamidi, F., Asl Rahnamaye Akbari, N., 2017. Molecular detection and genotyping of *Toxoplasma gondii* in chicken, beef, and lamb meat consumed in Northwestern Iran. *Iran. J. Parasitol.* 12, 38–45.
- Mainar, R.C., Delacruz, C., Asensio, A., Dominguez, L., Vazquezboland, J.A., 1996. Prevalence of agglutinating antibodies to *Toxoplasma gondii* in small ruminants of the Madrid region, Spain, and identification of factors influencing seropositivity by multivariate analysis. *Vet. Res. Commun.* 20, 153–159.
- Maksimov, P., Buschtöns, S., Herrmann, D.C., Conraths, F.J., Görlich, K., Tenter, A.M., Dubey, J.P., Nagel-Kohl, U., Thoms, B., Bötcher, L., Kühne, M., Schares, G., 2011. Serological survey and risk factors for *Toxoplasma gondii* in domestic ducks and geese in Lower Saxony, Germany. *Vet. Parasitol.* 182, 140–149.
- Maksimov, P., Basso, W., Zerweck, J., Schutkowski, M., Reimer, U., Maksimov, A., Conraths, F.J., Schares, G., 2018. Analysis of *Toxoplasma gondii* clonal type-specific antibody reactions in experimentally infected turkeys and chickens. *Int. J. Parasitol.* 48, 845–856.
- Mancianti, F., Nardoni, S., Papini, R., Mugnaini, L., Martini, M., Altomonte, I., Salari, F., D'Ascenzi, C., Dubey, J.P., 2014. Detection and genotyping of *Toxoplasma gondii* DNA in the blood and milk of naturally infected donkeys (*Equus asinus*). *Parasit. Vectors* 7, 165.
- Marques, L.C., Costa, A.J., Lopes, C.W.G., de Moraes, F.R., de Moraes, J.R.E., 1995. Experimental toxoplasmosis in pregnant mares: a study of fetuses and placentas. *Braz. J. Vet. Res. Anim. Sci.* 32, 246–250.
- Marques, P.X., O'Donovan, J., Williams, E.J., Gutierrez, J., Worrall, S., McElroy, M., Proctor, A., Brady, C., Sammin, D., Bassett, H., Buxton, D., Maley, S., Markey, B.K., Nally, J.E., 2012. Detection of *Toxoplasma gondii* antigens reactive with antibodies from serum, amniotic, and allantoic fluids from experimentally infected pregnant ewes. *Vet. Parasitol.* 185, 91–100.
- Masatani, T., Takashima, Y., Takasu, M., Matsuu, A., Amaya, T., 2016. Prevalence of anti-*Toxoplasma gondii* antibody in domestic horses in Japan. *Parasitol. Int.* 65, 146–150.
- Mateus-Pinilla, N.E., Dubey, J.P., Choromanski, L., Weigel, R.M., 1999. A field trial of the effectiveness of a feline *Toxoplasma gondii* vaccine in reducing *T. gondii* exposure for swine. *J. Parasitol.* 85, 855–860.
- Matsuo, K., Kamai, R., Uetsu, H., Goto, H., Takashima, Y., Nagamune, K., 2014. Seroprevalence of *Toxoplasma gondii* infection in cattle, horses, pigs and chickens in Japan. *Parasitol. Int.* 63, 638–639.
- McColgan, C., Buxton, D., Blewett, D.A., 1988. Titration of *Toxoplasma gondii* oocysts in non-pregnant sheep and the effects of subsequent challenge during pregnancy. *Vet. Rec.* 123, 467–470.
- Meerburg, B.G., van Riel, J.W., Cornelissen, J.B., Kijlstra, A., Mul, M.F., 2006. Cats and goat whey associated with *Toxoplasma gondii* infection in pigs. *Vector Borne Zoonotic Dis.* 6, 266–274.
- Mevelec, M.N., Ducourmau, C., Bassuny Ismael, A., Olivier, M., Seche, E., Lebrun, M., Bout, D., Dimier-Poisson, I., 2010. Mic1-3 knockout *Toxoplasma gondii* is a good candidate for a vaccine against *T. gondii*-induced abortion in sheep. *Vet. Res.* 41, 49.
- Miao, Q., Wang, X., She, L.N., Fan, Y.T., Yuan, F.Z., Yang, J.F., Zhu, X.Q., Zou, F.C., 2013. Seroprevalence of *Toxoplasma gondii* in horses and donkeys in Yunnan Province, Southwestern China. *Parasit. Vectors* 6, 168.
- Moller, T., Fennestad, K.L., Eriksen, L., Work, K., Siim, J.C., 1970. Experimental toxoplasmosis in pregnant sows. *Acta Pathol. Microbiol. Scand.* A 78, 241–255.
- Moreno, B., Collantes-Fernandez, E., Villa, A., Navarro, A., Regidor-Cerrillo, J., Ortega-Mora, L.M., 2012. Occurrence of *Neospora caninum* and *Toxoplasma gondii* infections in ovine and caprine abortions. *Vet. Parasitol.* 187, 312–318.
- Moskwa, B., Kornacka, A., Cybulska, A., Cabaj, W., Reiterova, K., Bogdaszewski, M., Steiner-Bogdaszewska, Z., Bien, J., 2018. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* infection in sheep, goats, and fallow deer farmed on the same area. *J. Anim. Sci.* 96, 2468–2473.
- Munday, B.L., 1978. Bovine toxoplasmosis: experimental infections. *Int. J. Parasitol.* 8, 285–288.
- Munoz-Zanzi, C., Tamayo, R., Balboa, J., Hill, D., 2012. Detection of oocyst-associated toxoplasmosis in swine from southern Chile. *Zoonoses Public Health* 59, 389–392.
- Murrell, K.D., Djordjevic, M., Cuperlovic, K., Sofronic, L., Savic, M., Djordjevic, M., Damjanovic, S., 2004. Epidemiology of *Trichinella* infection in the horse: the risk from animal product feeding practices. *Vet. Parasitol.* 123, 223–233.
- O'Donovan, J., Proctor, A., Gutierrez, J., Worrell, S., Nally, J., Marques, P., Brady, C., McElroy, M., Sammin, D., Buxton, D., Maley, S., Bassett, H., Markey, B., 2012. Distribution of lesions in fetal brains following experimental infection of pregnant sheep with *Toxoplasma gondii*. *Vet. Pathol.* 49, 462–469.
- Okamoto, I., Suzuki, Y., Itho, H., Fukuura, H., Yokoyama, I., 1989. A collective outbreak of porcine toxoplasmosis due to supplied feed supplements contaminated with toxoplasma oocysts. *J. Jpn. Vet. Med. Assoc.* 42, 729–732.
- Olinda, R.G., Pena, H.F., Frade, M.T., Ferreira, J.S., Maia, L.A., Gennari, S.M., Oliveira, S., Dantas, A.F., Riet-Correa, F., 2016. Acute toxoplasmosis in pigs in Brazil caused by *Toxoplasma gondii* genotype Chinese 1. *Parasitol. Res.* 115, 2561–2566.
- Opsteegh, M., Teunis, P., Zuchner, L., Koets, A., Langelaar, M., van der Giessen, J., 2011. Low predictive value of seroprevalence of *Toxoplasma gondii* in cattle for detection of parasite DNA. *Int. J. Parasitol.* 41, 343–354.
- Opsteegh, M., Maas, M., Schares, G., van der Giessen, J., 2016. Relationship between seroprevalence in the main livestock species and presence of *Toxoplasma gondii* in meat (GP/EFSa/BIOHAZ/2013/01) an extensive literature review. Final report. EFSa Supporting Publications 13 (996E-n/a).
- Ortega-Pacheco, A., Acosta Viana, K.Y., Guzman-Marin, E., Segura-Correa, J.C., Alvarez-Fleites, M., Jimenez-Coello, M., 2013. Prevalence and risk factors of *Toxoplasma gondii* in fattening pigs farm from Yucatan, Mexico. *Biomed. Res. Int.* 2013, 231497.
- Owen, M.R., Clarkson, M.J., Trees, A.J., 1998. Acute phase toxoplasma abortions in sheep. *Vet. Rec.* 142, 480–482.
- Pan, M., Lyu, C., Zhao, J., Shen, B., 2017. Sixty years (1957–2017) of research on toxoplasmosis in China—an overview. *Front. Microbiol.* 8, 1825.
- Papini, R.A., Buzzone, G., Nardoni, S., Rocchigiani, G., Mancianti, F., 2015. Seroprevalence and genotyping of *Toxoplasma gondii* in horses slaughtered for human consumption in Italy. *J. Equine Vet. Sci.* 35, 657–661.
- Passos, L., Lima, J., Figueiredo, B., 1984. Determinacao da infeccao por *Toxoplasma gondii* em bovinos abatidos em Belo Horizonte (MG) através da frequencia de anticorpos e tentativa de isolamento a partir de musculatura diafragmática. *Arq. Bras. Med. Vet. Zoot.* 36, 581–589.
- Pastiu, A.I., Gyorke, A., Blaga, R., Mircean, V., Rosenthal, B.M., Cozma, V., 2013. In Romania, exposure to *Toxoplasma gondii* occurs twice as often in swine raised for familial consumption as in hunted wild boar, but occurs rarely, if ever, among fattening pigs raised in confinement. *Parasitol. Res.* 112, 2403–2407.
- Pastiu, A.I., Gyorke, A., Kalmar, Z., Bolf, P., Rosenthal, B.M., Oltean, M., Villena, I., Spinu, M., Cozma, V., 2015. *Toxoplasma gondii* in horse meat intended for human consumption in Romania. *Vet. Parasitol.* 212, 393–395.
- Perry, B.D., Randolph, T.F., 1999. Improving the assessment of the economic impact of parasitic diseases and of their control in production animals. *Vet. Parasitol.* 84, 145–168.
- Pomares, C., Ajzenberg, D., Bornard, L., Bernardin, G., Haseine, L., Darde, M.L., Marty, P., 2011. Toxoplasmosis and horse meat, France. *Emerg. Infect. Dis.* 17, 1327–1328.
- Portella, L.P., Cadore, G.C., Sangioni, L.A., Pellegrini, L.F., Figuera, R., Ramos, F., Vogel, F.S., 2017. Antibodies against apicomplexa protozoa and absence sarcocysts in heart tissues from horses in southern Brazil. *Rev. Bras. Parasitol. Vet.* 26, 100–103.
- Puvanesuaran, V.R., Noordin, R., Balakrishnan, V., 2013. Genotyping of *Toxoplasma gondii* isolates from wild boars in Peninsular Malaysia. *PLoS ONE* 8, e61730.
- Radke, J.R., Guerini, M.N., Jerome, M., White, M.W., 2003. A change in the premitotic period of the cell cycle is associated with bradyzoite differentiation in *Toxoplasma gondii*. *Mol. Biochem. Parasitol.* 131, 119–127.
- Raeghi, S., Akaberi, A., Sedeghi, S., 2011. Seroprevalence of *Toxoplasma gondii* in sheep, cattle and horses in Urmia North-West of Iran. *Iran. J. Parasitol.* 6, 90–94.
- Rahdar, M., Samarbaf-Zadeh, A.R., Arab, L., 2012. Evaluating the prevalence of *Toxoplasma gondii* in meat and meat products in Ahvaz by PCR method. *Jundishapur J. Microb.* 5, 570–573.

- Rassouli, M., Razmi, G.R., Bassami, M.R., Movassaghi, A.R., Azizzadeh, M., 2011. Study on ovine abortion associated with *Toxoplasma gondii* in affected herds of Khorasan Razavi Province, Iran based on PCR detection of fetal brains and maternal serology. *Parasitology* 138, 691–697.
- Razmi, G.R., Ghezi, K., Mahooti, A., Naseri, Z., 2010. A serological study and subsequent isolation of *Toxoplasma gondii* from aborted ovine fetuses in Mashhad area, Iran. *J. Parasitol.* 96, 812–814.
- Razmi, G.R., Abedi, V., Yaghfoori, S., 2016. Serological study of *Toxoplasma gondii* infection in Turkoman horses in the North Khorasan Province, Iran. *J. Parasit. Dis.* 40, 515–519.
- Rego, W.M.F., Paula, N.R.O., Vitor, R.W.A., Silva, R.A.B., Diniz, B.L.M., Sousa, M.M., Coelho, W.A.C., Porfirio, K.P., Pinheiro, R.R., Alves, F.S.F., Cavalcante, A.C.R., Cardoso, J.F.S., 2016. Risk factors for *Toxoplasma gondii* infection in goats and sheep raised in the State of Piauí in northeast Brazil. *Small Rumin. Res.* 141, 17–23.
- Ribeiro, M.J., Rosa, M.H., Bruhn, F.R., Garcia Ade, M., Rocha, C.M., Guimaraes, A.M., 2016. Seroepidemiology of *Sarcocystis neurona*, *Toxoplasma gondii* and *Neospora* spp. among horses in the south of the state of Minas Gerais, Brazil. *Rev. Bras. Parasitol. Vet.* 25, 142–150.
- Rizzo, H., Gaeta, N.C., Hora, J.H.C., Carvalho, J.S., Pinheiro Júnior, J.W., Gennari, S.M., Pena, H.F.J., Villalobos, E.M.C., Gregory, L., 2017. Risk factors for *Toxoplasma gondii* infection in sheep in the northeastern region of Brazil. *Braz. J. Vet. Res. Anim. Sci.* 54, 139–146.
- Roberts, C.W., Cruickshank, S.M., Alexander, J., 1995. Sex-determined resistance to *Toxoplasma gondii* is associated with temporal differences in cytokine production. *Infect. Immun.* 63, 2549–2555.
- Roberts, C.W., Walker, W., Alexander, J., 2001. Sex-associated hormones and immunity to protozoan parasites. *Clin. Microbiol. Rev.* 14, 476–488.
- Romanelli, P.R., Freire, R.L., Vidotto, O., Marana, E.R., Ogawa, L., De Paula, V.S., Garcia, J.L., Navarro, I.T., 2007. Prevalence of *Neospora caninum* and *Toxoplasma gondii* in sheep and dogs from Guarapuava farms, Parana State, Brazil. *Res. Vet. Sci.* 82, 202–207.
- Rommel, M., Sommer, R., Janitschke, K., Müller, I., 1966. Experimentelle Toxoplasma-Infektion bei Kälbern. *Berl. Muench. Tierarztl. Wschr.* 79, 41–45.
- Rosa, C., Kasai, N., Souza, S.L.P., Guerra, J.L., Rego, A.A., Gennari, S.M., 2001. Comparação das técnicas de imuno-histoquímica e bioensaio em camundongos para pesquisa de *Toxoplasma gondii* em tecidos de caprinos, experimentalmente inoculados. *Arq. Inst. Biol. (São Paulo)* 68, 13–17.
- Rougier, S., Montoya, J.G., Peyron, F., 2017. Lifelong persistence of toxoplasma cysts: a questionable dogma? *Trends Parasitol.* 33, 93–101.
- Ruiz, A., Frenkel, J.K., 1980. Intermediate and transport hosts of *Toxoplasma gondii* in Costa Rica. *Am. J. Trop. Med. Hyg.* 29, 1161–1166.
- Sa, S.G., Lima, D.C., Silva, L.T., Pinheiro Junior, J.W., Dubey, J.P., Silva, J.C., Mota, R.A., 2016. Seroprevalence of *Toxoplasma gondii* among turkeys on family farms in the state of Northeastern Brazil. *Acta Parasitol.* 61, 401–405.
- Sa, S.G., Ribeiro-Andrade, M., Silva, L.T.R., Souza, O.L.N., Lima, D.C.V., Pedrosa, C.M., Bezerra, M.J.G., Mota, R.A., 2017. Risk factors associated with *Toxoplasma gondii* infection in free-range chickens in the semi-arid region of Brazil. *Rev. Bras. Parasitol. Vet.* 26, 221–225.
- Saad, N.M., Hussein, A.A.A., Ewida, R.M., 2018. Occurrence of *Toxoplasma gondii* in raw goat, sheep, and camel milk in Upper Egypt. *Vet World* 11, 1262–1265.
- Sakata, F.B., Bellato, V., Sartor, A.A., de Moura, A.B., de Souza, A.P., Farias, J.A., 2012. *Toxoplasma gondii* antibodies sheep in Lages, Santa Catarina, Brazil, and comparison using IFA and ELISA. *Rev. Bras. Parasitol. Vet.* 21, 196–200.
- Samico-Fernandes, E.F.T., Samico-Fernandes, M.F.T., de Albuquerque, P.P.F., de Almeida, J.C., de Souza Santos, A., da Rocha Mota, A., de Souza Neto, O.L., Mota, R.A., 2017. *Toxoplasma gondii* in backyard pigs: seroepidemiology and mouse bioassay. *Acta Parasitol.* 62, 466–470.
- Sánchez-Sánchez, R., Ferré, I., Regidor-Cerrillo, J., Gutiérrez-Expósito, D., Ferrer, L.M., Artech-Villasol, N., Moreno-Gonzalo, J., Müller, J., Aguado-Martínez, A., Pérez, V., Hemphill, A., Ortega-Mora, L.M., Benavides, J., 2019. Virulence in mice of a *Toxoplasma gondii* type II isolate does not correlate with the outcome of experimental infection in pregnant sheep. *Front. Cell. Infect. Microbiol.* 8, 436.
- Sanger, V.L., Cole, C.R., 1955. Toxoplasmosis. VI. Isolation of toxoplasma from milk, placentas, and newborn pigs of asymptomatic carrier sows. *Am. J. Vet. Res.* 16, 536–539.
- Santana, L.F., da Costa, A.J., Pieroni, J., Lopes, W.D., Santos, R.S., de Oliveira, G.P., de Mendonca, R.P., Sakamoto, C.A., 2010. Detection of *Toxoplasma gondii* in the reproductive system of male goats. *Rev. Bras. Parasitol. Vet.* 19, 179–182.
- Santoro, A., Tagel, M., Must, K., Laine, M., Lassen, B., Jokelainen, P., 2017. *Toxoplasma gondii* seroprevalence in breeding pigs in Estonia. *Acta Vet. Scand.* 59, 82.
- Saqib, M., Hussain, M.H., Sajid, M.S., Mansoor, M.K., Asi, M.N., Fadya, A.A., Zohaib, A., Sial, A.U., Muhammad, G., Ullah, I., 2015. Sero-epidemiology of equine toxoplasmosis using a latex agglutination test in the three metropolises of Punjab, Pakistan. *Trop. Biomed.* 32, 276–285.
- Saraf, P., Shwab, E.K., Dubey, J.P., Su, C., 2017. On the determination of *Toxoplasma gondii* virulence in mice. *Exp. Parasitol.* 174, 25–30.
- Savvulidi, F., Ptáček, M., Stádník, L., 2018. Pathogens in processed rm semen and approaches for their elimination. *Acta Univ. Agric. Silv. Mendel Brun* 66, 1065–1072.
- Schale, S., Howe, D., Yeargan, M., Morrow, J.K., Graves, A., Johnson, A.L., 2018. Protozoal coinfection in horses with equine protozoal myeloencephalitis in the eastern United States. *J. Vet. Intern. Med.* 32, 1210–1214.
- Schares, G., Bangoura, B., Randau, F., Goroll, T., Ludewig, M., Maksimov, P., Matzkeit, B., Sens, M., Bärwald, A., Conraths, F.J., Opsteegh, M., Van der Giessen, J., 2017a. High seroprevalence of *Toxoplasma gondii* and probability of detecting tissue cysts in backyard laying hens compared with hens from large free-range farms. *Int. J. Parasitol.* 47, 765–777.
- Schares, G., Herrmann, D.C., Maksimov, P., Matzkeit, B., Conraths, F.J., Moré, G., Preisinger, R., Weigend, S., 2017b. Chicken line-dependent mortality after experimental infection with three type IIxIII recombinant *Toxoplasma gondii* clones. *Exp. Parasitol.* 180, 101–111.
- Schlüter, D., Däubener, W., Schares, G., Gross, U., Pleyer, U., Lüder, C., 2014. Animals are key to human toxoplasmosis. *Int. J. Med. Microbiol.* 304, 917–929.
- Schnydrig, P., Vidal, S., Brodard, I., Frey, C., Posthaus, H., Perreten, V., Rodriguez-Campos, S., 2017. Bacterial, fungal, parasitological and pathological analyses of abortions in small ruminants from 2012–2016. *Schweiz. Arch. Tierheilkd.* 159, 647–656.
- Schoonman, L.B., Wilsmore, T., Swai, E.S., 2010. Sero-epidemiological investigation of bovine toxoplasmosis in traditional and smallholder cattle production systems of Tanga Region, Tanzania. *Trop. Anim. Health Prod.* 42, 579–587.
- Schulzig, H.S., Fehlhaber, K., 2005. Longitudinal study on the seroprevalence of *Toxoplasma gondii* infection in four German pig breeding and raising farms. *Berl. Munch. Tierarztl. Wochenschr.* 118, 399–403.
- Sechi, P., Ciampelli, A., Cambiotti, V., Veronesi, F., Cenci-Goga, B.T., 2013. Seroepidemiological study of toxoplasmosis in sheep in rural areas of the Grosseto district, Tuscany, Italy. *Ital. J. Anim. Sci.* 12.
- Shaapan, R.M., Ghazy, A.A., 2007. Isolation of *Toxoplasma gondii* from horse meat in Egypt. *Pak. J. Biol. Sci.* 10, 174–177.
- Shokri, A., Sharif, M., Teshnizi, S.H., Sarvi, S., Rahimi, M.T., Mizani, A., Ahmadvpour, E., Montazeri, M., Daryani, A., 2017. Birds and poultries toxoplasmosis in Iran: a systematic review and meta-analysis. *Asian Pac J Trop Med* 10, 635–642.
- Shuralev, E.A., Shamaev, N.D., Mukminov, M.N., Nagamune, K., Taniguchi, Y., Saito, T., Kitoh, K., Arleevskaia, M.I., Fedotova, A.Y., Abdulmanova, D.R., Aleksandrova, N.M., Efimova, M.A., Yarulinn, A.I., Valeeva, A.R., Khaertynov, K.S., Takashima, Y., 2018. *Toxoplasma gondii* seroprevalence in goats, cats and humans in Russia. *Parasitol. Int.* 67, 112–114.
- Shwab, E.K., Zhu, X.Q., Majumdar, D., Pena, H.F., Gennari, S.M., Dubey, J.P., Su, C., 2014. Geographical patterns of *Toxoplasma gondii* genetic diversity revealed by multilocus PCR-RFLP genotyping. *Parasitology* 141, 453–461.
- Skjerve, E., Waldeland, H., Nesbakken, T., Kapperud, G., 1998. Risk factors for the presence of antibodies to *Toxoplasma gondii* in Norwegian slaughter lambs. *Prev. Vet. Med.* 35, 219–227.
- Sposito Filha, E., Do Amaral, V., Macruz, R., Reboucas, M.M., Santos, S.M., Borgo, F., 1992. Infecção experimental de equinos com taquizoítos de *Toxoplasma gondii*. *Rev. Bras. Parasitol. Vet.* 1, 51–54.
- Stalheim, O.H., Hubbert, W.T., Boothe, A.D., Zimmermann, W.J., Hughes, D.E., Barnett, D., Riley, J.L., Foley, J., 1980. Experimental toxoplasmosis in calves and pregnant cows. *Am. J. Vet. Res.* 41, 10–13.
- Stormoen, M., Tharaldsen, J., Hopp, P., 2012. Seroprevalence of *Toxoplasma gondii* infection in Norwegian dairy goats. *Acta Vet. Scand.* 54, 75.
- Sun, W.W., Meng, Q.F., Cong, W., Shan, X.F., Wang, C.F., Qian, A.D., 2015. Herd-level prevalence and associated risk factors for *Toxoplasma gondii*, *Neospora caninum*, *Chlamydia abortus* and bovine viral diarrhoea virus in commercial dairy and beef cattle in eastern, northern and northeastern China. *Parasitol. Res.* 114, 4211–4218.
- Tan, Q.D., Yang, X.Y., Yin, M.Y., Hu, L.Y., Qin, S.Y., Wang, J.L., Zhou, D.H., Zhu, X.Q., 2015. Seroprevalence and correlates of *Toxoplasma gondii* infection in dairy cattle in northwest China. *Acta Parasitol.* 60, 618–621.

- Tao, Q., Wang, Z., Feng, H., Fang, R., Nie, H., Hu, M., Zhou, Y., Zhao, J., 2011. Seroprevalence and risk factors for *Toxoplasma gondii* infection on pig farms in central China. *J. Parasitol.* 97, 262–264.
- Tassi, P., 2007. *Toxoplasma gondii* infection in horses. A review. *Parassitologia* 49, 7–15.
- Tavalla, M., Sabaghan, M., Abdizadeh, R., Khademvatan, S., Rafiei, A., Razavi Piranshahi, A., 2015. Seroprevalence of *Toxoplasma gondii* and *Neospora* spp. infections in Arab horses, Southwest of Iran. *Jundishapur. J. Microbiol.* 8, e14939.
- Tenter, A.M., Heckeroth, A.R., Weiss, L.M., 2000. *Toxoplasma gondii*: from animals to humans. *Int. J. Parasitol.* 30, 1217–1258.
- Thrusfield, M.V., 2007. *Veterinary Epidemiology*. 3rd edition. Edition. Blackwell Science Ltd, Oxford.
- Tilahun, B., Tolossa, Y.H., Tilahun, G., Ashenafi, H., Shimelis, S., 2018. Seroprevalence and risk factors of *Toxoplasma gondii* infection among domestic ruminants in East Hararghe zone of Oromia region, Ethiopia. *Vet. Med. Int.* 2018, 4263470.
- Todd, E.C.D., 1989. Preliminary estimates of costs of foodborne disease in the United-States. *J. Food Prot.* 52, 595–601.
- Tonouhewa, A.B., Akpo, Y., Sessou, P., Adoligbe, C., Yessinou, E., Hounmanou, Y.G., Assogba, M.N., Youssao, I., Farougou, S., 2017. *Toxoplasma gondii* infection in meat animals from Africa: systematic review and meta-analysis of sero-epidemiological studies. *Vet. World* 10, 194–208.
- Turner, C.B., Savva, D., 1990. Evidence of *Toxoplasma gondii* in an equine placenta. *Vet. Rec.* 127, 96.
- Turner, C.B., Savva, D., 1991. Detection of *Toxoplasma gondii* in equine eyes. *Vet. Rec.* 129, 128.
- Turner, C.B., Savva, D., 1992. Transplacental infection of a foal with *Toxoplasma gondii*. *Vet. Rec.* 131, 179–180.
- Tzanidakis, N., Maksimov, P., Conraths, F.J., Kiossis, E., Brozos, C., Sotiraki, S., Schares, G., 2012. *Toxoplasma gondii* in sheep and goats: seroprevalence and potential risk factors under dairy husbandry practices. *Vet. Parasitol.* 190, 340–348.
- van den Brom, R., Lievaart-Peterson, K., Lutikholt, S., Peperkamp, K., Wouda, W., Vellema, P., 2012. Abortion in small ruminants in the Netherlands between 2006 and 2011. *Tijdschr. Diergeneesk.* 137, 450–457.
- van der Giessen, J., Fonville, M., Bouwknegt, M., Langelaar, M., Vollema, A., 2007. Seroprevalence of *Trichinella spiralis* and *Toxoplasma gondii* in pigs from different housing systems in The Netherlands. *Vet. Parasitol.* 148, 371–374.
- van Engelen, E., Lutikholt, S., Peperkamp, K., Vellema, P., Van den Brom, R., 2014. Small ruminant abortions in The Netherlands during lambing season 2012–2013. *Vet. Rec.* 174, 506.
- Vaudaux, J.D., Muccioli, C., James, E.R., Silveira, C., Magargal, S.L., Jung, C., Dubey, J.P., Jones, J.L., Doymaz, M.Z., Bruckner, D.A., Belfort Jr., R., Holland, G.N., Grigg, M.E., 2010. Identification of an atypical strain of *Toxoplasma gondii* as the cause of a waterborne outbreak of toxoplasmosis in Santa Isabel do Ivaí, Brazil. *J. Infect. Dis.* 202, 1226–1233.
- Venturi, S.S., da Silva, A.F., Frazao-Teixeira, E., de Oliveira, F.C.R., Consalter, A., Padilha, F.G.F., Fonseca, A.B.M., Ferreira, A.M.R., 2017. Characterization of the zoonotic potential of *Toxoplasma gondii* in horses from Rio de Janeiro State. *Acta Trop.* 171, 159–162.
- Verhelst, D., De Craeye, S., Vanrobaeys, M., Czapllicki, G., Dorny, P., Cox, E., 2014. Seroprevalence of *Toxoplasma gondii* in domestic sheep in Belgium. *Vet. Parasitol.* 205, 57–61.
- Verhelst, D., De Craeye, S., Jennes, M., Dorny, P., Goddeeris, B., Cox, E., 2015. Interferon-gamma expression and infectivity of toxoplasma infected tissues in experimentally infected sheep in comparison with pigs. *Vet. Parasitol.* 207, 7–16.
- Veronesi, F., Ranucci, D., Branciarri, R., Miraglia, D., Mammoli, R., Fioretti, D.P., 2011. Seroprevalence and risk factors for *Toxoplasma gondii* infection on finishing swine reared in the Umbria region, central Italy. *Zoonoses Public Health* 58, 178–184.
- Vesco, G., Buffolano, W., La Chiusa, S., Mancuso, G., Caracappa, S., Chianca, A., Villari, S., Curro, V., Liga, F., Petersen, E., 2007. *Toxoplasma gondii* infections in sheep in Sicily, southern Italy. *Vet. Parasitol.* 146, 3–8.
- Villari, S., Vesco, G., Petersen, E., Crispo, A., Buffolano, W., 2009. Risk factors for toxoplasmosis in pigs bred in Sicily, Southern Italy. *Vet. Parasitol.* 161, 1–8.
- Wallace, G.D., 1971. Experimental transmission of *Toxoplasma gondii* by filth-flies. *Am. J. Trop. Med. Hyg.* 20, 411–413.
- Wallace, G.D., 1972. Experimental transmission of *Toxoplasma gondii* by cockroaches. *J. Infect. Dis.* 126, 545–547.
- Wallander, C., Frossling, J., Dorea, F.C., Uggla, A., Vagsholm, I., Lunden, A., 2016. Pasture is a risk factor for *Toxoplasma gondii* infection in fattening pigs. *Vet. Parasitol.* 224, 27–32.
- Wanderley, F.S., Porto, W.J., Camara, D.R., da Cruz, N.L., Feitosa, B.C., Freire, R.L., de Moraes, E.P., Mota, R.A., 2013. Experimental vaginal infection of goats with semen contaminated with the “CPG” strain of *Toxoplasma gondii*. *J. Parasitol.* 99, 610–613.
- Wang, J.L., Zhou, D.H., Chen, J., Liu, G.X., Pu, W.B., Liu, T.Y., Qin, S.Y., Yin, M.Y., Zhu, X.Q., 2015. The prevalence of antibodies to *Toxoplasma gondii* in horses in Changji Hui Autonomous Prefecture, Xinjiang, northwestern China. *Rev. Bras. Parasitol. Vet.* 24, 298–302.
- Wang, S., Zhao, G.W., Wang, W., Zhang, Z.C., Shen, B., Hassan, I.A., Xie, Q., Yan, R.F., Song, X.K., Xu, L.X., Li, X.R., 2015. Pathogenicity of five strains of *Toxoplasma gondii* from different animals to chickens. *Korean J. Parasitol.* 53, 155–162.
- Wang, H., Zhang, L., Ren, Q., Yu, F., Yang, Y., 2017. Diagnosis of swine toxoplasmosis by PCR and genotyping of *Toxoplasma gondii* from pigs in Henan, Central China. *BMC Vet. Res.* 13, 152.
- Watts, E., Zhao, Y., Dhara, A., Eller, B., Patwardhan, A., Sinai, A.P., 2015. Novel approaches reveal that *Toxoplasma gondii* bradyzoites within tissue cysts are dynamic and replicating entities in vivo. *MBio* 6 (e01155-01115).
- Weigel, R.M., Dubey, J.P., Siegel, A.M., Kitron, U.D., Mannelli, A., Mitchell, M.A., Mateus-Pinilla, N.E., Thulliez, P., Shen, S.K., Kwok, O.C.H., Todd, K.S., 1995. Risk factors for transmission of *Toxoplasma gondii* on swine farms in Illinois. *J. Parasitol.* 81, 736–741.
- Weissenböck, H., Dubey, J.P., 1993. Toxoplasmosis epizootic in a fattening swine herd. *Dtsch. Tierarztl. Wochenschr.* 100, 370–374.
- Wiengcharoen, J., Thompson, R.C., Nakhthong, C., Rattanakorn, P., Sukthana, Y., 2011. Transplacental transmission in cattle: is *Toxoplasma gondii* less potent than *Neospora caninum*? *Parasitol. Res.* 108, 1235–1241.
- Williams, R.H., Morley, E.K., Hughes, J.M., Duncanson, P., Terry, R.S., Smith, J.E., Hide, G., 2005. High levels of congenital transmission of *Toxoplasma gondii* in longitudinal and cross-sectional studies on sheep farms provides evidence of vertical transmission in ovine hosts. *Parasitology* 130, 301–307.
- Work, K., Eriksen, L., Fennestad, K.L., Moller, T., Siim, J.C., 1970. Experimental toxoplasmosis in pregnant sows. I. Clinical, parasitological and serological observations. *Acta Pathol. Microbiol. Scand. B: Microbiol. Immunol.* 78, 129–139.
- Work, T.M., Dagenais, J., Rameyer, R., Breeden, R., 2015. Mortality patterns in endangered Hawaiian geese (Nene; *Branta sandvicensis*). *J. Wildl. Dis.* 51, 688–695.
- Wyss, R., Sager, H., Müller, N., Inderbitzin, F., König, M., Audige, L., Gottstein, B., 2000. Distribution of *Toxoplasma gondii* and *Neospora caninum* under aspects of meat hygiene. *Schweiz. Arch. Tierheilkd.* 142, 95–108.
- Xu, P., Song, X., Wang, W., Wang, F., Cao, L., Liu, Q., 2012. Seroprevalence of *Toxoplasma gondii* infection in chickens in Jinzhou, northeastern China. *J. Parasitol.* 98, 1300–1301.
- Xu, P., Li, X., Tang, F., Liu, Y.H., Kou, X., Zhao, M.L., Li, B., Guo, L., Liu, X.G., Zhao, Q., 2015. Seroprevalence and risk factors for *Toxoplasma gondii* in sheep and goats in Jinzhou, Northeastern China. *Trop. Biomed.* 32, 563–567.
- Yang, N., Mu, M.Y., Li, H.K., Long, M., He, J.B., 2012. Seroprevalence of *Toxoplasma gondii* infection in slaughtered chickens, ducks, and geese in Shenyang, northeastern China. *Parasit. Vectors* 5, 237.
- Yang, N., Mu, M.Y., Yuan, G.M., Zhang, G.X., Li, H.K., He, J.B., 2013. Seroprevalence of *Toxoplasma gondii* in slaughtered horses and donkeys in Liaoning province, northeastern China. *Parasit. Vectors* 6, 140.
- Yin, M.Y., Wang, J.L., Huang, S.Y., Qin, S.Y., Zhou, D.H., Liu, G.X., Tan, Q.D., Zhu, X.Q., 2015. Seroprevalence and risk factors of *Toxoplasma gondii* in Tibetan sheep in Gansu province, Northwestern China. *BMC Vet. Res.* 11, 41.
- Zhang, N., Wang, S., Wang, D., Li, C., Zhang, Z., Yao, Z., Li, T., Xie, Q., Liu, S., Zhang, H., 2016. Seroprevalence of *Toxoplasma gondii* infection and risk factors in domestic sheep in Henan province, central China. *Parasite* 23, 53.
- Zhou, M., Cao, S., Sevinc, F., Sevinc, M., Ceylan, O., Liu, M., Wang, G., Moumouni, P.F., Jirapattharasate, C., Suzuki, H., Nishikawa, Y., Xuan, X., 2017. Enzyme-linked immunosorbent assays using recombinant TgSAG2 and NcSAG1 to detect *Toxoplasma gondii* and *Neospora caninum*-specific antibodies in domestic animals in Turkey. *J. Vet. Med. Sci.* 78, 1877–1881.
- Zhu, J., Yin, J., Xiao, Y., Jiang, N., Ankarlev, J., Lindh, J., Chen, Q., 2008. A sero-epidemiological survey of *Toxoplasma gondii* infection in free-range and caged chickens in northeast China. *Vet. Parasitol.* 158, 360–363.