FIVE NEW LABDANE DITERPENE OXIDES FROM *EUPATORIUM JHANII**

ANTONIO G. GONZÁLEZ, JOSÉ M. ARTEAGA, JOSÉ L. BRETÓN and BRAULIO M. FRAGA

Department of Organic and Biochemistry, University of La Laguna,
Instituto de Productos Naturales Orgánicos, CSIC, Tenerife, Spain

(Received 14 June 1976)

Key Word Index—*Eupatorium jhanii*; Compositae; labdane diterpene oxides; jhanol; jhanidiol; jhanol acetate; jhanidiol-18-monoacetate; jhanidiol diacetate.

Abstract—Five new labdane diterpene oxides jhanol, jhanidiol, jhanol acetate, jhanidiol-18-monoacetate and jhanidiol diacetate were isolated from the aerial parts of *Eupatorium jhanii*.

INTRODUCTION

From species of *Eupatorium*, several sesquiterpene lactones with cytotoxic activity [1–67] and a number of interesting flavonoid compounds [7–11] have been isolated. However, there is not much information about the presence of diterpenoid compounds in this genus [12,13].

In the present communication, we report the isolation and structural determination of five hydroxylated derivatives of 8,13-epoxylabd-14-ene (manoyl oxide) from *Eupatorium jhanii*.

RESULTS

The first free alcohol isolated was jhanol, $C_{20}H_{34}O_2$, (1a). Spectral and chemical data showed it to be a primary alcohol: IR 3480 cm^{-1} , ^1H NMR two doublets, J 11 Hz at 3.14 and 3.45; it gave an acetate and was oxidized to the aldehyde 1c.

The presence of a vinyl group, suggested by the IR spectrum (bands at $3080, 1640, 980\text{ cm}^{-1}$), was confirmed by the ^1H NMR spectrum which exhibited the signals characteristic of this group: a set of three quartets at 4.95, 5.15 and 5.92 ppm [14].

The remaining oxygen atom must be part of an ether bridge connecting two fully substituted carbon atoms; the IR spectrum showed strong absorption at 1120 cm^{-1} while there was no ^1H NMR signal for a hydrogen geminal to the oxygen. This spectrum also showed two methyl singlets at 1.31 and 1.35 ppm (Me—C(0)) and two more at 0.80 and 0.88 ppm. Consequently jhanol is probably a compound belonging to the 8,13-epoxylabdane series.

This was proved by Huang–Minlon reduction of the aldehyde 1c, affording two substances identified as 8,13-epoxylabdane (2a) [15,16] and 8,13-epoxylabd-14-ene

(1d) [17,18]. Comparison of the methyl carbon-13 NMR signals of the jhanol acetate with those of manoyl oxide (2a), Table 1, showed that 1b is an 18-hydroxyditerpene. The C-18, C-19 and C-20 methyls of manoyl oxide [19,20] and also of other tri and tetracyclic diterpenes resonate at different field positions, near to 33, 20 and 15 ppm [21–23]. On the other hand, the ^{13}C -NMF (SFORD) spectrum of 1b showed only two quartets, at 17.80 and 15.23 ppm, and a new triplet at 72.5 ppm. The stereochemistry at C-4, depicted in 1b, was corroborated by the ^1H -NMR spectrum because both the methyl and oxymethylene signals occupy different field positions in the two possible structures [24,25]. The position of the aldehyde signal for 1c is typical of equatorial substituents [26].

Table 1. ^{13}C -NMR spectra of manoyl oxide (1d) jhanol acetate (1b) and jhanidiol-18-monoacetate (3b)

Carbon No	1d	1b	3b
1	39.12	38.61	79.14
2	18.62	17.20	28.89
3	42.23	35.93	34.04
4	33.32	36.56	36.22
5	56.55	50.64	49.26
6	20.03	19.95	19.73
7	43.40	43.01	42.77
8	75.09	74.89	74.80
9	55.60	56.05	55.86
10	37.11	37.02	42.91
11	51.41	15.42	18.49
12	35.73	35.93	35.30
13	73.25	73.28	73.28
14	148.07	148.04	147.96
15	110.21	110.23	110.28
16	28.71	78.52	28.81
17	25.58	25.43	25.69
18	33.40	72.99	72.49
19	21.39	17.80	16.91
20	15.41	15.83	11.71

* Part 33 in the series "Chemistry of the Compositae". For Part 32 see: A. G. González, J. Bermejo, J. L. Bretón, G. Martínez, B. Domínguez, J. M. Amaro, *J. Chem. Soc. Perkin I* 1976, 1663.

The most polar diterpene of this series, jhanidiol (**3a**), had the empirical formula $C_{20}H_{34}O_3$; its polarity and the 1H -NMR spectrum, which showed signals for protons under the hydroxyls at 3.14*d*, 3.45*d* and 3.41*t* ppm, suggested the presence of a primary and a secondary alcohol; according to the conditions, this compound gave the monoacetate **3b** or the diacetate **3c**. Comparison of the 1H -NMR resonances of **1a** and **3a** made it obvious

- 1**
 a $R = CH_2OH$
 b $R = CH_2OAC$
 c $R = CHO$
 d $R = Me$

- 3**
 a $R_1 = OH$ $R_2 = H$ $R_3 = CH_2OH$
 b $R_1 = OH$ $R_2 = H$ $R_3 = CH_2OAC$
 c $R_1 = OAC$ $R_2 = H$ $R_3 = CH_2OAC$
 d R_1 $R_2 = O$ $R_3 = CH_2OAC$
 e R_1 $R_2 = O$ $R_3 = CHO$
 f R_1 $R_2 = O$ $R_3 = COOH$

- 2**
 a $R_1 = R_2 = H$ $R_3 = CH_3$
 b $R_1 = R_2 = H$ $R_3 = CH_2OAC$
 c R_1 $R_2 = O$ $R_3 = CH_2OAC$

4

that the only difference between these compounds was the existence of an additional OH group in **3a**. Oxidation of monoacetate **3b** to the ketone **3d** and Huang-Minlon reduction yielded an alcohol identical with jhanol (**1a**).

Analyses of the ^{13}C -NMR spectrum of **3b** indicated the position and stereochemistry of oxymethine hydrogens. In the expoylabdane [19,20] and also in other series of diterpenes, C-2, C-6 and C-11 triplets were found near 20 ppm; the resonance signals of C-1, C-3, C-7 and C-12 appeared much lower in the field [21-23]. Comparison of the secondary carbon resonances of **1b** and **3b** suggested that one of the latter's carbons must be substituted. The paramagnetic shift undergone by C-2 (11.7) and C-10 (5.9), together with the diamagnetic one experienced by C-3 (11.9) and C-20 (4.12 ppm) clearly point to C-1 as the locus of the hydroxyl group with an equatorial orientation [27]. The 1H -NMR spectrum of **3c** corroborated this conclusion, showing the oxymethine resonance as a clear quartet at 4.53 ppm. Moreover, the spectra of all the oxidation products of **3a** had signals for the equatorial proton at C-11 [28].

Jhanidiol did not form an acetonide nor did the ketoacid **3f** show β -ketoacid properties. Bromination and dehydrobromination of ketodihydroacetate **2c** afforded the enone **4** which presented the characteristic

1H -NMR signals for a Δ^2 -1-one [29], in full accord with structure **3a**. Jhanol acetate (**1b**), jhanidiol-18-monoacetate (**3b**) and jhanidiol diacetate (**3c**) were also found as natural products in the plant.

EXPERIMENTAL

Mp's are uncorr. Optical activities were taken in $CHCl_3$, 1H -NMR spectra on a 90 MHz and ^{13}C -NMR on a 20 MHz

instrument in $CDCl_3$ with TMS as internal reference. Column and dry column chromatography was performed on Si gel.

Isolation of the diterpenes. Air-dried aerial part of the plant *Eupatorium jhanii* Robinson (4 kg) collected at El Delgadito (Mérida, Venezuela) in June, was finely cut and Soxhlet extracted several times with EtOH. Combined filtered ethanolic extracts were concentrated *in vacuo* and chromatographed on a Si gel column. Petrol, petrol-EtOAc and EtOAc eluted a mixture of diterpenes, triterpenes and flavones which was rechromatographed on dry column, yielding the following diterpenes, in the order of elution: jhanol acetate (**1b**), jhanidiol diacetate (**3c**), jhanidiol-18-monoacetate (**3b**), jhanol (**1a**), jhanidiol (**3a**).

Jhanol acetate (1b). **1b** (400 mg) was isolated as a liquid; IR $\nu_{CHCl_3}^{max} cm^{-1}$: 3080, 1720, 1640, 1120, 990, 920; NMR: δ 0.88 (6H, s, $C_{19}C_{20}$), 1.31 and 1.35 (each 3H, s, C-14 and C-17), 2.10 (3H, s), 3.68, 3.90 (each 1H, d, J 11 Hz, C-18), 4.97 (1H, q, J 11 Hz, J 2 Hz), 5.17 (1H, q, J 17 Hz, J 2 Hz), 5.94 (1H, q, J 11 Hz, J 17 Hz); MS: m/e (%) 348 (1) (M^+), 333, 273, 255, 191, 149 (100), 135. Identical to that obtained from jhanol by standard acetylation.

Jhanidiol acetate (3c). **3c** (500 mg) was isolated as a liquid; IR $\nu_{CHCl_3}^{max} cm^{-1}$: 3080, 1720, 1110, 980, 920; NMR: δ 0.80 (3H, s), 0.96 (3H, s), 1.20 (3H, s), 1.26 (3H, s), 1.96 (3H, s), 2.02 (3H, s), 3.67, 3.84 (each 1H, d, J 11 Hz), 4.53 (1H, q, J 9 and 5 Hz), 4.94 (1H, q, J 11 and 2 Hz), 5.12 (1H, q, J 17 and 2 Hz), 5.86 (1H, q, J 17 and 11 Hz). Identical to the product obtained by acetylation of jhanidiol. MS: m/e (%) 391 (M^+ -15), 331, 271, 253, 223, 201, 189, 176, 174, 149 (100), 133.

Jhanidiol-18-monoacetate (3b). **3b** (1200 mg) was isolated as a liquid; IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3330, 3080, 1740, 1640, 1240, 1120, 980, 920; NMR: δ 0.82, 0.88, 1.24, 1.29, 2.06 (all 3H, s), 3.38 (1H, q, $W_{1/2}$ 14 Hz), 3.62, 3.84 (each 1H, d, J 11 Hz), 4.92 (1H, q, J 11 and 2 Hz), 5.14 (1H, q, J 17 and 2 Hz), 5.80 (1H, q, J 17 and 11 Hz); MS: m/e (%) 364 (1) (M^+), 349 (100), 331, 313, 271, 253, 223, 201, 188, 149.

Jhanol (1a). **1a** (2.3 g) was isolated as a solid; mp 139–141° [α_D^{25} +27° (c 0.65)]; IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3480, 3080, 1640, 1120, 980, 930; NMR: δ 0.80, 0.88, 1.31, 1.35 (each 3H, s), 3.15, 3.47 (each 1H, d, J 11 Hz), 4.95 (1H, q, J 11 and 2 Hz), 5.15 (1H, q, J 17 and 2 Hz), 5.92 (1H, q, J 17 and 11 Hz); MS: m/e (%) 306 (1) (M^+), 291, (98), 273, 261, 255, 243, 208, 177 (100), 151, 149, 147, 135, 133, 123, 121. (Found: C78.30; H 10.89. $\text{C}_{20}\text{H}_{34}\text{O}_2$ requires: C 78.38; H 11.18%).

Jhanidiol (3a). **3a** (4.0 g) was isolated as a solid; mp 189° [α_D^{25} +90° (c 0.8)]; IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3460, 3080, 1640, 1120, 990, 920; NMR: δ 0.78, 0.92, 1.28, 1.34 (each 3H, s), 3.14, 3.45 (each 1H, d, J 11 Hz), 3.41 (1H, t, $W_{1/2}$ 14 Hz), 4.95 (1H, q, J 11 and 2 Hz), 5.15 (1H, q, J 17 and 2 Hz), 5.92 (1H, q, J 17 and 11 Hz); MS: m/e (%) 322 (0.5), (M^+), 307 (100), 289, 271, 259, 231, 224, 206, 193, 189, 175, 173, 161, 159, 149, 137. (Found: C 74.18; H 10.71. $\text{C}_{20}\text{H}_{34}\text{O}_2$ requires: C 74.49; H 10.63%).

Oxidation of jhanol. Pyridinium chlorochromate (1.5 g) was suspended in dry CH_2Cl_2 (20 ml) and jhanol (290 mg) in CH_2Cl_2 (15 ml) was added once to the stirred sol. After 1 hr, dry Et_2O (20 ml) was added and the supernatant decanted from the black gum formed. The insoluble residue was washed 3 $\times$ with 30 ml portions dry Et_2O . Combined CH_2Cl_2 -ether sol was passed through a short column of Celite and solvent removed by distillation, giving 8,13-epoxylabd-14-enal (210 mg) (**1c**); IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm^{-1} : 3080, 2700, 1720; NMR: δ 0.85, 1.04, 1.28, 1.31 (each 3H, s), 4.94 (1H, q, J 11 and 2 Hz), 5.12 (1H, q, J 17 and 2 Hz), 5.89 (1H, q, J 17 and 11 Hz) and 9.18 (1H, s).

Huang-Minlon reduction of 8,13-epoxylabd-18-al. $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$ was added to a soln of **1c** (210 mg) in diethylene glycol (14 ml) and the mixture refluxed for 90 min (temp. 130°). After the addition of KOH pellets (300 mg), refluxing continued for a further 45 min. Then the temp was raised to 210° by distillation of the excess water-hydrazine and the refluxing continued for another 3 hr. The usual work-up afforded a mixture of 2 products which were separated by dry column chromatography, using petrol- C_6H_6 (4:1) as eluent. 8,13-Epoxylabdane (dihydromanoyl oxide) (**2a**) (65 mg); isolated as a liquid [α_D^{25} +3° (c 1.1) Lit +5.9; NMR: δ 0.78, 0.80, 0.85 (t), 0.86, 1.18, 1.29 (each 3H, s), 8, 13-Epoxylabd-14-ene (manoyl oxide) (**1d**). **1d** (95 mg) was isolated as a solid; mp 26–28° [α_D^{25} +11° (c 2.0) Lit 22.5–26.5 [α_D^{25} +19°; NMR: δ 0.83 (6H, d), 0.90, 1.31, 1.33 (each 3H, s), 4.94 (1H, q, J 11 and 2 Hz), 5.16 (1H, q, J 17 and 2 Hz), 5.92 (1H, q, J 17 and 11 Hz); MS: m/e (%) 290 (1) (M^+), 275 (100), 257, 245, 192, 177, 149, 137.

Partial acetylation of jhanidiol. To a soln of **3a** (700 mg) in $\text{C}_5\text{H}_5\text{N}$ (2 ml), cold Ac_2O (3 ml) was added and the mixture was left for 10 min. The usual work-up gave a mixture of diacetate (70 mg), monoacetate (410 mg) and unaltered starting material, which was separated by dry column chromatography, using C_6H_6 - EtOAc (9:1) as eluent. The diacetate and 18-monoacetate of jhanidiol were identical to the natural products obtained.

Oxidation of jhanidiol monoacetate (3b). **3b** (410 mg) was oxidized as described above for **1a**. 18-acetate-1-keto-8,13-epoxylabd-14-ene (400 mg) was obtained; mp 87–89°; [α_D^{25} +70°]; IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3090, 1735, 1700, 1240, 1120, 990, 920, 910; NMR: δ 1.05, 1.21, 1.30, 1.35 (each 3H, s), 2.06 (3H, s), 2.8 (1H, m), 3.74, 3.98 (each 1H, d), 4.93 (1H, q), 5.13 (1H, q), 5.93 (1H, q); NMR: δ (benzene- d_6) 0.72, 0.92, 1.20, 1.36 (each 3H, s), 3.58, 3.76 (each 1H, d), 4.93 (1H, q), 5.26 (1H, q), 5.97 (1H, q); MS: m/e (%) 362 (1) (M^+), 347, 269, 224, 200, 191, 173, 167, (100).

Jhanol from jhanidiol. The above substance **3d** (300 mg) was reduced under Huang-Minlon conditions. Work up gave jhanol: mp, mixed mp and IR superimposable.

Hydrogenation of 3d. **3d** (400 mg) was dissolved in EtOAc

(20 ml) and hydrogenated for 1 hr over Pd/C (5%) at room temp and atm press. This gave **2c** (350 mg); NMR: δ 0.85 (3H, t, J 6 Hz), 1.04 (3H, s), 1.21 (6H, s) and 1.32 (3H, s).

Bromination of keto-18-acetoxy-8,13-epoxy-labdane. A soln of trimethylphenylammonium tribromide (400 mg) in dry THF (6 ml) was added in drops while stirring a soln of **2c** (350 mg) in the same solvent until the mixture turned yellow and it was then stirred at room temp for a further 30 min. The reaction mixture was poured into H_2O , extracted with CH_2Cl_2 and the solvent was evaporated, yielding a Br derivative (300 mg) which did not crystallize; IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm^{-1} : 1730, 1700, 1120, 960; NMR: δ 0.82 (3H, t, 6 Hz), 1.02 (3H, s), 1.18 (6H, s), 1.30 (3H, s), 2.03 (3H, s), 2.8 (1H, m), 3.64, 3.72 (each 1H, d), 4.12 (1H, m $W_{1/2}$ 27 Hz).

Dehydrobromination of 1-keto-2 β -bromo-18-acetoxy-18,13-epoxylabdane. A mixture of Li_2CO_3 (400 mg), LiBr (250 mg), DMF (15 ml) and C_6H_6 (6 ml) was heated until the H_2O and C_6H_6 had been eliminated. Then a soln of bromide (300 mg) in DMF (15 ml) was added and the reaction mixture refluxed for 1 hr under N_2 . The cooled soln was poured into H_2O , extracted with CH_2Cl_2 and washed with 10% aq HOAc, sat aq NaHCO_3 and H_2O . After the solvent was evaporated, the residue afforded 1-keto-18-acetoxy-8,13-epoxy-labd-2-ene (**4**); mp 84–88°; [α_D^{25} +46°]; UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm: 223 (c 22000); IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm^{-1} : 1730, 1700, 1680, 1120, 950; NMR: δ 0.84 (3H, t J 6 Hz), 1.02 (3H, s), 1.20 (6H, s), 1.32 (3H, s), 2.03 (3H, s), 2.75 (1H, m, $W_{1/2}$ 27 Hz), 3.71, 3.50 (each 1H, d, J 11 Hz), 5.82 and 6.32 (each 1H, d, J 11 Hz).

8,13-Epoxy-labd-14-en-1-keto-18-al (3e). **3a** (100 mg) was oxidized as described above for **1a**, giving **3e** (80 mg); mp 88–93°; [α_D^{25} +62°; M^+ 318; IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3080, 2700, 1720, 1700, 1640, 1260, 1130, 990, 960, 920, 895; NMR: δ 1.23, 1.27, 1.30, 1.35 (each 3H, s), 2.78 (1H, m), 4.92 (1H, q), 5.11 (1H, q), 5.86 (1H, q).

Oxidation of keto-aldehyde (3f). **3e** (100 mg) dissolved in Me_2CO (min quantity) was treated dropwise with Jones' reagent and left at room temp for 3 hr, upon which MeOH was added to destroy any excess reagent. The mixture was poured into water and extracted as usual, giving **3f**; mp 206–209°; [α_D^{25} +33° (c 0.57)]; IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm^{-1} : 3500–2500 (br) 1700, 1640, 1110; NMR: δ 1.17, 1.29 (each 3H, s), 4.90 (1H, q), 5.12 (1H, q), 5.88 (1H, q); MS: m/e (%) 319 (100) (M^+ -15), 256, 186, 173, 149.

Acknowledgements—The authors would like to thank Dr A. Morales, University of Mérida (Venezuela) for gathering the plant and for the sustained interest he has shown in the progress of this work.

REFERENCES

- Kupchan, S. M., Kelsey, J. E., Maruyama, M., Cassidy, J. M., Hemingway, S. C. and Knox, J. R. (1969) *J. Org. Chem.* **34**, 3876.
- Kupchan, S. M., Fujita, T., Maruyama, M. and Britton, R. W. (1973) *J. Org. Chem.* **38**, 1260.
- Herz, W. and Wahlberg, I. (1973) *J. Org. Chem.* **38**, 2485.
- Kupchan, S. M., Maruyama, M., Hemingway, R. J. Shibuya, S. and Fujita, T. (1973) *J. Org. Chem.* **38**, 2189.
- McPhail, A. T., Onan, R. D., Lee, K. H., Ibuka, T. and Huang, H. C. (1974) *Tetrahedron Letters* 3203.
- Herz, W. and Sharma, R. P. (1976) *Tetrahedron* **41**, 1015.
- Kupchan, S. M., Sigel, W., Knox, J. R. and Udayamurthy, M. S. (1969) *J. Org. Chem.* **34**, 1460.
- Wagner, H., Iyengar, M. A., Hörhammer, L. and Herz, W. (1972) *Phytochemistry* **11**, 1504.
- Wagner, H., Iyengar, M. A., Seligmann, O., Hörhammer, L. and Herz, W. (1972) *Phytochemistry* **11**, 1506.
- Herz, W., Gibaja, S., Bhat, S. V. and Srinivasan, A. (1972) *Phytochemistry* **11**, 2859.
- Talapatra, S. R., Bhar, D. S. and Talapatra, B. (1974) *Australian J. Chem.* **27**, 1137.

12. Hegnauer, R. (1964) *Chemotaxonomie der Pflanzen*, Vol. III, p. 482. Birkhauser, Basel.
13. Herz, W. and Sharma, R. P. (1976) *J. Org. Chem.* **41**, 1021.
14. Wenkert, E., Beak, P. and Grant, P. K. (1961) *Chem. Ind. (London)* 1574.
15. Hosking, J. R. and Brandt, C. W. (1934) *Ber Deut. Chem. Ges* **67**, 1173.
16. Hasen, A. J., Kimland, B., Almqvist, S. O. and Enzell, C. R. (1972) *Acta Chem. Scand.* **B26**, 832.
17. Hodges, R. and Reed, R. I. (1960) *Tetrahedron* **10**, 71.
18. Giles, J. A., Schumacher, J. N., Mims, S. S. and Bernasek, E. (1962) *Tetrahedron* **18**, 169.
19. Buckwalter, B. L., Burfitt, I. R., Nagel, A. A. and Wenkert, E. (1975) *Helv. Chim. Acta* **58**, 1567.
20. Almqvist, S. O., Enzell, C. R. and Wehrli, F. N. (1975) *Acta Chem. Scand.* **B29**, 695.
21. Carstenn-Lichterfelde, C., Pascual, C., Pons, J., Rabanal, R., Rodríguez, B. and Valverde, S. (1975) *Tetrahedron Letters* 3569.
22. Hanson, J. R., Sivers, M., Piozzi, F. and Savona, G. (1976) *J. Chem. Soc. Perkin I* 114.
23. Wahlberg, I., Almqvist, S. O., Nishida, I. and Enzell, C. R. (1975) *Acta Chem. Scand.* **B29**, 1047.
24. Wenkert, E., Afonso, A., Beak, P., Carney, R. V. J., Jeffs, P. N. and McChesney, J. D. (1965) *J. Org. Chem.* **30**, 713.
25. Gaudemar, A., Polonsky, J. and Wenkert, E. (1964) *Bull. Soc. Chim. France* 407.
26. King, T. J. and Yarley, J. P. (1961) *J. Chem. Soc.* 4308.
27. Eggert, H., Van Antwerp, G. F., Bhacca, N. S. and Djerassi, C. (1976) *J. Org. Chem.* **41**, 71.
28. Bhacca, N. S. and Williams, D. H. (1966) *Application of NMR Spectroscopy in Organic Chemistry*, p. 66. Holden Day, San Francisco.
29. Chan, W. R., Gibbs, J. A. and Taylor, D. R. (1973) *J. Chem. Soc. Perkin I* 1047.