JGR Atmospheres

RESEARCH ARTICLE

10.1029/2020JD033017

Key Points:

- Atmospheric Evaporative Demand mostly affect SPEI in dry periods
- There are spatial differences in the influence of the AED on drought assessment by means of the SPEI
- AED is not overestimating drought severity since it shows a comprehensive influence on dryness

Supporting Information:

- Supporting Information S1

Correspondence to:

S. M. Vicente-Serrano,
svicen@ipe.csic.es

Citation:

Tomas-Burguera, M., Vicente-Serrano, S. M., Peña-Angulo, D., Domínguez-Castro, F., Noguera, I., & El Kenawy, A. (2020). Global Characterization of the Varying Responses of the Standardized Precipitation Evapotranspiration Index to Atmospheric Evaporative Demand. *Journal of Geophysical Research: Atmospheres*, 125, e2020JD033017. <https://doi.org/10.1029/2020JD033017>

Received 28 APR 2020

Accepted 8 JUL 2020

Accepted article online 24 AUG 2020

Author Contributions:

Conceptualization: M. Tomas-Burguera, S. M. Vicente-Serrano

Formal analysis: M. Tomas-Burguera, S. M. Vicente-Serrano, D. Peña-Angulo

Methodology: M. Tomas-Burguera, S. M. Vicente-Serrano

Software: M. Tomas-Burguera, S. M. Vicente-Serrano, F. Domínguez-Castro, I. Noguera

Supervision: S. M. Vicente-Serrano

Writing - original draft: S. M. Vicente-Serrano

Writing - review & editing: M. Tomas-Burguera, S. M. Vicente-Serrano, D. Peña-Angulo, F. Domínguez-Castro, I. Noguera, A. El Kenawy

Global Characterization of the Varying Responses of the Standardized Precipitation Evapotranspiration Index to Atmospheric Evaporative Demand

M. Tomas-Burguera¹, S. M. Vicente-Serrano² , D. Peña-Angulo² , F. Domínguez-Castro^{3,4}, I. Noguera², and A. El Kenawy^{5,6}

¹Estación Experimental de Aula Dei, Consejo Superior de Investigaciones Científicas (EEAD-CSIC), Zaragoza, Spain,

²Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC), Zaragoza, Spain,

³Aragones Agency for Research and Development Researcher (ARAID), Zaragoza, Spain, ⁴Department of Geography, University of Zaragoza, Zaragoza, Spain, ⁵Department of Geography, Mansoura University, Mansoura, Egypt,

⁶Department of Geography, Sultan Qaboos University, Al Khoud, Oman

Abstract The standardized precipitation evapotranspiration index (SPEI) is one of the well-established drought metrics worldwide. It is simply computed using precipitation and atmospheric evaporative demand (AED) data. Although AED is considered a key driver of drought variability worldwide, it could have less impact on drought in specific regions and for particular times as a function of the magnitude of precipitation. Specifically, the influence of the AED might overestimate drought severity during both normal and humid periods, resulting in “false alarms” about drought impacts on physical and human environments. Here, we provided a global characterization of the sensitivity of the SPEI to changes of the AED. Results demonstrate that the contribution of AED to drought severity is largely impacted by the spatial and temporal variability of precipitation. Specifically, the impact of AED on drought severity was more pronounced during periods of low precipitation, compared to wet periods. Interestingly, drought severity in humid regions (as revealed by SPEI) also showed low sensitivity to AED under drier conditions. These results highlight the skill of SPEI in identifying the role of AED in drought evolution, especially in arid and semiarid regions whose climate is characterized typically by low precipitation. This advantage was also evident for humid environments, where SPEI did not overestimate drought severity due to the increased AED. These findings highlight the broader applicability of SPEI to accurately characterize drought severity worldwide.

1. Introduction

Drought is a complex natural hazard, which is impacted by a wide variety of climatic variables. Accordingly, it is important to unravel the contribution of these different drivers and assess their role in drought severity (Mukherjee et al., 2018; S. M. Vicente-Serrano, McVicar, et al., 2020). This assessment has strong implications for detecting trends of drought severity in both current and future climates. When interpreting drought trends using the standardized precipitation index (SPI) (Mckee et al., 1993), which employed precipitation data alone, there is often a low degree of uncertainty in evaluating drought trends in the present climate and future scenarios, as precipitation deficit is usually directly proportional to more severe and long-lasting drought events. Nevertheless, precipitation could not be seen as the only variable impacting drought evolution, especially under the current global warming conditions and even more for future climate change scenarios. Due to air temperature rise, there is a stronger atmospheric vapor pressure deficit (W. Yuan et al., 2019). In the same context, relative humidity witnesses a decreasing trend over the majority of the continental areas worldwide (Willett et al., 2014), which is likely to be accelerated in the future, as suggested by climate model outputs (M. Byrne & O’Gorman, 2013; M. P. Byrne & O’Gorman, 2018). This situation will reinforce the atmospheric evaporative demand (AED) and thus it could increase drought severity (S. M. Vicente-Serrano, McVicar, et al., 2020).

The AED intensification during dry periods can increase levels of vegetation stress (Allen et al., 2015; Hartmann et al., 2018; A. P. Williams et al., 2013), higher evaporation rates from soil (Teuling et al., 2013) and water bodies (Friedrich et al., 2018), and accelerate hydrological drought downstream. Some drought

indices like the Palmer drought severity index (PDSI) (Palmer, 1965), the reconnaissance drought index (RDI) (Tsakiris et al., 2007), the standardized moisture anomaly index (SZI) (Zhang et al., 2019), the standard Palmer drought index (SPDI) (Ma et al., 2014), and the standardized precipitation evapotranspiration index (SPEI) (Beguería et al., 2014; S. M. Vicente-Serrano et al., 2010) take into account AED when quantifying drought severity.

Some studies have suggested that there is some degree of uncertainty in quantifying drought severity using climate drought indices that account for the role of AED in their formulations (Berg & Sheffield, 2018; Cook et al., 2014; Greve et al., 2019; Scheff, 2018; Yang et al., 2019). Overall, the debate on the role of AED on drought severity (B. Cook et al., 2018; Dai et al., 2018; Greve et al., 2019) originates mainly from the notion that the role of AED is complex, depending largely on the drought type (e.g., meteorological, hydrological, and environmental) and prevailing climatic conditions (S. M. Vicente-Serrano, McVicar, et al., 2020).

Some studies argue that AED is not a good metric of actual evapotranspiration (ET) because AED exceeds ET particularly in arid regions, leading to an overestimation of soil drying (Berg & Sheffield, 2018; Benjamin I. Cook et al., 2014). While this assumption seems to be feasible, the inclusion of AED in climate drought indices cannot be introduced as a substitute of ET. Rather, irrespective of ET rates, accounting for the role of AED in drought metrics has a strong physical meaning with real implications in the intensification of hydrological, agricultural and environmental droughts (S. M. Vicente-Serrano, McVicar, et al., 2020). In the same context, some studies suggest that AED could not be seen as a good metric of plant water demand. These studies attributed this feedback to the role of the atmospheric CO₂ fertilization (Abigail L. S. Swann, 2018) and the role of vapor pressure deficit on stomata conductance, which can diminish plant water demand, even with high AED (Konings et al., 2017; Miralles et al., 2019). As such, it is expected that the atmospheric CO₂ fertilization would increase plant water use efficiency (Milly & Dunne, 2016; Roderick et al., 2015; A. L. S. Swann et al., 2016), alleviate plant water demand (Abigail L. S. Swann, 2018), and reduce the stress corresponding to hydrological deficits (Greve et al., 2019). However, the impact of atmospheric CO₂ fertilization is still uncertain (Brodribb et al., 2020; B. Cook et al., 2018; S. M. Vicente-Serrano, McVicar, et al., 2020). An important aspect is that the role of atmospheric CO₂ could be negligible during low precipitation periods (Menezes-Silva et al., 2019; Morgan et al., 2004; Z. Xu et al., 2016), making environmental and agricultural drought aggravated by any increase in AED (Allen et al., 2015; McDowell et al., 2016; McDowell & Allen, 2015). In addition, AED could largely reduce surface water resources due to reinforced land evaporation (B. I. Cook et al., 2019; Zhang et al., 2018). Also, vegetation changes may increase water consumption, especially in locations characterized by denser coverage, higher leaf area index, and longer vegetation active periods (Frank et al., 2015; J. S. Mankin et al., 2018; Justin S. Mankin et al., 2019).

There are also important statistical issues that should be carefully considered when interpreting drought changes associated with AED. These concerns are mostly related to how drought metrics are calculated (S. M. Vicente-Serrano et al., 2019). Overall, the different assumptions about the role of AED in drought severity, as revealed by climate drought metrics, are still questionable and deserve much more attention for better interpretation and understanding of drought severity.

In the past decade, SPEI has been one of the most widely used drought metrics worldwide. It has been used for different applications, including drought risk assessment (e.g., Blauhut et al., 2015; Domínguez-Castro et al., 2019), analysis of drought evolution (e.g., Chen & Sun, 2015; Spinoni et al., 2014; Spinoni, Barbosa, Bucchignani, et al., 2019), assessment of drought impacts in different socioeconomic (e.g., Bachmair et al., 2015, 2016; Gu et al., 2020; Leng & Hall, 2019; Ribeiro et al., 2019; Von Uexkull et al., 2016; Zampieri et al., 2017), hydrological (e.g., Barker et al., 2016; Peña-Gallardo et al., 2019) and environmental systems (e.g., Babst et al., 2019; Bachmair et al., 2018; Fang et al., 2019; Knapp et al., 2015; Mitchell et al., 2014; Slette et al., 2019; Zimmermann et al., 2015), and characterization of drought in future scenarios (e.g., Chen & Sun, 2017; Dewes et al., 2017; Gu et al., 2020; Naumann et al., 2018; Spinoni, Barbosa, De Jager, et al., 2019).

A key aspect of SPEI formulation is that it responds equally to the role of precipitation and AED (Sergio M. Vicente-Serrano et al., 2015), making it a suitable metric to assess the response of drought severity to AED changes in regions with different climate characteristics. Based on this index, Vicente-Serrano et al. (2015)

and Vicente-Serrano, McVicar, et al. (2020) indicated that in general precipitation showed more impact on drought evolution than AED at the global scale. The only exception was found for water-limited regions, where AED may be more relevant in defining SPEI. It is suggested that the role of AED in hydrological and environmental drought severity is less significant during humid conditions and more pronounced during dry periods characterized by low precipitation and soil moisture deficit (S. M. Vicente-Serrano, McVicar, et al., 2020). Albeit with these earlier attempts to explore the response of drought severity to AED, a comprehensive assessment of the sensitivity of SPEI to changes in AED from a temporal perspective is still lacking and worth investigation. Such an assessment is important to determine whether drought severity, as revealed by SPEI, can be overestimated due to the possible role of AED. This assessment matters to both arid and humid regions worldwide to evaluate the suitability of SPEI to reliably capture the influence of AED on drought severity. A high sensitivity of this index to AED, especially during wet periods, may make the use of this index less robust for assessing drought severity.

The overall aims of this study are (i) to provide a global characterization of the sensitivity of SPEI to AED changes under different precipitation magnitudes, and (ii) to define the regions and periods in which this sensitivity is much stronger due to changes in precipitation intensities. Hereinafter, the terms “sensitivity of SPEI to AED” refers to the relative contribution of AED to the variability of SPEI.

2. Materials and Methods

Data of monthly precipitation and potential evapotranspiration (as a metric of AED) were retrieved from the Climate Research Unit (CRU) data set CRU TS v. 3.26 (Harris et al., 2014), at a spatial resolution of 0.5° for the period 1901–2017. To guarantee an adequate spatial representation of the global main precipitation regimes, our analysis was restricted to the global land areas between 65°N and 65°S. From a practical perspective, calculating drought indices in desert regions is meaningless (Wu et al., 2007), but we included these regions to establish a wider range of validation of the obtained results.

Climate balance (D), defined as the difference between precipitation and AED ($P-AED$), is standardized to obtain SPEI. In order to assess the sensitivity of SPEI to AED during different periods of the year, here the climate balance was calculated at three different time scales: monthly, seasonal, and annual. Herein, SPEI at 1-month time scale was computed to characterize drought at monthly time scale. Also, SPEI at 3-month time scale was quantified for the months of February, May, August and November to account for drought in winter, spring, summer, and autumn, respectively. Similarly, SPEI at 12-month time scale for December was considered as representative of annual changes in drought.

The role of AED in climate balance variability was assessed following the approach of Hobbins et al. (2012). Nonetheless, as our research focuses on a different variable, we adapted this approach to evaluate the contribution of AED and precipitation (P) to the variability of the climate balance (D):

$$D = P - AED \quad (1)$$

The variability of D can be expressed as

$$\sigma_D^2 \approx J^T C J \quad (2)$$

where J is the Jacobian matrix, with the two partial derivatives of D , and C is the variance-covariance matrix

$$J^T = \left[\frac{\partial D}{\partial P} \quad \frac{\partial D}{\partial AED} \right] \quad (3)$$

$$C = \begin{bmatrix} \sigma_P^2 & \sigma_{P,AED} \\ \sigma_{AED,P} & \sigma_{AED}^2 \end{bmatrix} \quad (4)$$

Thus,

$$\sigma_D^2 \approx \frac{\partial D^2}{\partial P} \sigma_P^2 + \frac{\partial D}{\partial P} \frac{\partial D}{\partial AED} \sigma_{P, AED} + \frac{\partial D^2}{\partial AED} \sigma_{AED}^2 + \frac{\partial D}{\partial AED} \frac{\partial D}{\partial P} \sigma_{AED, P} \quad (5)$$

In order to quantify the contribution of each variable (i.e., P and AED) to the variability of climate balance (D), we split this expression into two parts, where B_P represents the contribution of P , while B_{AED} denotes the contribution of AED .

$$B_P = \frac{\partial D^2}{\partial P} \sigma_P^2 + \frac{\partial D}{\partial P} \frac{\partial D}{\partial AED} \sigma_{P, AED} \quad (6)$$

$$B_{AED} = \frac{\partial D^2}{\partial AED} \sigma_{AED}^2 + \frac{\partial D}{\partial AED} \frac{\partial D}{\partial P} \sigma_{AED, P} \quad (7)$$

As D is only a subtraction, the partial derivatives are easy to obtain.

$$\frac{\partial D}{\partial P} = 1 \text{ and } \frac{\partial D}{\partial AED} = -1 \quad (8)$$

Hence,

$$B_P = \sigma_P^2 - \sigma_{P, AED} \quad (9)$$

$$B_{AED} = \sigma_{AED}^2 - \sigma_{AED, P} \quad (10)$$

To allow the direct comparison between the obtained results for different sites and time scales, we computed the relative values of B_{AED} (Equation 11). Following this procedure, our results were independent of the absolute values of the variance, which are expected to vary among sites and time scales.

$$\beta_{AED} = 100 \frac{|B_P|}{|B_P| + |B_{AED}|} \quad (11)$$

SPEI is based on standardization of D . Indeed, the standardization process changes the variance. However, our preference was to quantify the relative contribution of AED to the variability of D (i.e., β_{AED}). This also provides a metric of the relative contribution of AED to the variability of SPEI.

Herein, it is noteworthy indicating that our methodology was applied independently to the monthly, seasonal, and annual series. However, all precipitation and AED series were detrended prior to making the analysis to avoid any possible effects of trends presented in the series (via their impact on variance) on the results.

We analyzed the sensitivity of the SPEI considering different precipitation conditions. First, we applied the sensitivity analysis to the complete series (1901–2017). However, to identify whether this sensitivity can be changeable as a function of precipitation dryness, we also calculated the sensitivity, but for sampled years, seasons, and months. These samples were selected using precipitation thresholds of the 10th, 25th, and 50th centiles. These centiles refer to periods with different drying conditions.

3. Results and Discussion

3.1. Local Sensitivity

Figure 1 illustrates representative examples of the sensitivity of SPEI to AED in four regions with different climatic conditions. Results are presented at the monthly, seasonal and annual scales. This analysis considered SPEI at 1-, 3-, and 12-month time scales as representative of monthly, seasonal, and annual variability of drought, respectively. Also, differences in the sensitivity of SPEI to AED , as a function of precipitation magnitude, are presented at the monthly, seasonal, and annual scales. Overall, results demonstrate that in the Sahel SPEI is completely (100%) sensitive to AED during the long dry season (November–March). This suggests that the interannual variability of SPEI during this period of the year is determined exclusively by AED . During the rainy season (April–October), the sensitivity of SPEI to AED decreases, reaching its

Figure 1. Sensitivity of SPEI to AED calculated for four different regions worldwide at the (from left to right) monthly (first panels), seasonal (second panels), and annual (third panels) scales, considering different precipitation thresholds (complete precipitation series and centiles 50th, 25th, and 10th). Blue circles represent the average precipitation.

lowest values (<20%) during months of higher precipitation (i.e., July and August). This implies that the interannual variability of SPEI during the rainy season is mainly driven by precipitation variability. Notably, the importance of the AED in drought evolution increases during dry years. For example, the sensitivity of SPEI to AED in July and August during the years below the 50th centile of precipitation exceeded 20% and during the years below the 10th centile of precipitation; the sensitivity of the SPEI was around 40%.

Figure 1 reveals that there exists a strong variability in the sensitivity of SPEI to AED in response to the dominant climate characteristics. For example, in the humid equatorial regions (Congo), the sensitivity of SPEI to AED is practically negligible at the annual and seasonal scales. Exceptionally, the sensitivity was higher than 20% during the seasons below the 10th centile of precipitation, suggesting that the temporal variability of drought in these regions depends mainly on precipitation variability. However, AED seems to have a considerable role in drought variability (as revealed by SPEI values) during the most dry months (mainly summertime). In the same context, the sensitivity of SPEI to AED is expected to be low in temperate humid regions (as represented by Scotland). The sensitivity of SPEI to AED in this region was small considering the complete series (1901–2017) at the monthly, seasonal, and annual scales. In Scotland, the contribution of AED to SPEI variability was 5% at the annual scale and around 10% during summer months (June–August). Similarly, the sensitivity was also small (20%) during the years below the 10th centile of precipitation. AED showed stronger influence on SPEI only during the very dry summers. In the semiarid northeastern Spain, the sensitivity of SPEI to AED was below 20% at the annual scale. Again, the sensitivity increased during summertime, approaching 100% during the years below the 10th centile of precipitation. This extreme sensitivity might be expected given that precipitation amounts are very low during dry years, making AED a dominant factor controlling D . Thus, the sensitivity of SPEI to AED was more important during these dry years and typically more relevant during the driest and warmest season of the year (summer).

An inspection of Figure 1 suggests that SPEI is more sensitive to AED variability during periods of low precipitation, which is coherent with what is desired from a drought index like SPEI. Indeed, the influence of AED on hydrological and environmental systems is more relevant during low precipitation periods (S. M. Vicente-Serrano, McVicar, et al., 2020). However, in general, the temporal variability of SPEI is mostly driven by precipitation, which is undoubtedly the main controller of drought variability. Our results noted remarkable spatial differences in the sensitivity of SPEI to AED, which are largely driven by the magnitude and variability of precipitation and climate seasonality. These spatial differences are well identified at the global scale, being modulated by precipitation anomalies.

3.2. Global Sensitivity

Figure 2 depicts the global spatial distribution of the sensitivity of SPEI to AED at the 12-month (annual) time scale. Considering the complete series, only the driest areas of the world exhibited a remarkable sensitivity, close to 100% of sensitivity in deserts. This sensitivity was lower (50–100%) in the neighboring dry areas. In the rest of the world, the sensitivity of SPEI to AED was much smaller. Notably, sensitivity of the SPEI to the AED was close to 0% over the majority of South America, Africa, South Asia, and eastern portions of North America. These spatial patterns inform that SPEI values were mainly driven by precipitation and, apart from very dry regions worldwide, SPEI showed low sensitivity to AED. Also, this stresses that SPEI is sensitive to AED only when precipitation amounts decline substantially. The percentage of the world areas showing a high sensitivity of SPEI to AED increased during dry years (Figure 3). For example, an inspection of the 50th centile reveals that a smaller percentage of world areas exhibit a considerable sensitivity of SPEI to AED, compared to the 25th and the 10th centiles of the dry years. Only when considering the 10th and 25th centiles of precipitation, a noticeable influence of AED on SPEI was identified over large areas of South America, South Africa, central Asia, Australia, Europe, and North America. At the monthly and seasonal scales, important spatial differences were noted (Figures S1 to S16 in the supporting information), which can be directly linked to climate seasonality. However, results stress that the spatial patterns of SPEI sensitivity to AED depend largely on precipitation amount, irrespective of SPEI time scale and the period of the year (Figures S17 and S18 in the supporting information). Overall, the sensitivity to AED increases during low precipitation periods, although the world area showing high sensitivity increased (mainly driven by higher AED) during boreal summers.

Here, we should stress that it cannot be inferred that the general drought conditions in these areas are only driven by AED, as this is clearly demonstrated by the negligible sensitivity of SPEI to AED in the complete series. Rather, drought is mostly driven by precipitation variability. Nevertheless, it is suggested that AED effect is more relevant during years with low precipitation. Therefore, the severity of drought, as suggested by SPEI, would be accentuated or diminished during low-precipitation years following the behavior of AED. Simply, apart from world deserts, the interannual variability of SPEI is driven by precipitation variability. However, AED can be significantly relevant to drought quantification during years of low precipitation.

Figure 2. Spatial distribution of the sensitivity of SPEI to AED at the annual scale (SPEI at 12 month for December). The sensitivity was calculated for the complete series (bottom right) and the series of P-AED for different precipitation amounts (i.e., 10th [top left], 25th [top right], and 50th [bottom left]).

This is strongly coherent with what would be desirable in a metric that tries to account for the effect of AED on drought severity. This is typically the case for SPEI, given that our global assessment confirms that the role of AED is only relevant during periods of low precipitation and/or limited soil moisture. During humid periods, an increase of the aerodynamic (e.g., air temperature) and radiative (i.e., solar radiation) components of AED is favorable for photosynthetic activity and growth (Niu et al., 2008; White et al., 1999). As such, any increase of the AED during these periods would not expect to increase vegetation stress in natural ecosystems and crops. Rather, an increase of AED could have negative influences on water resources through direct evaporation from lakes and reservoirs (Friedrich et al., 2018). However, this feedback is not expected to be relevant in absence of climatic conditions favoring for meteorological and hydrological droughts. In contrast, during dry periods, the enhanced AED could reinforce water stress in natural plants and cultivations (Allen et al., 2010, 2015; A. P. Williams et al., 2013; Hartmann et al., 2018; McDowell et al., 2016). Similarly, it would also increase the severity of hydrological droughts through accelerating evaporation from soil moisture (Teuling et al., 2013), water courses, reservoirs and lakes (Martínez-Granados et al., 2011), or irrigated lands (S. M. Vicente-Serrano et al., 2017). Therefore, according to the results obtained here, SPEI

is highly suitable for identifying enhanced drought severity associated with increased AED during dry periods. This important aspect highlights the potential of using this drought metric to assess environmental and hydrological impacts of drought in different regions worldwide, irrespective of the dominant climatic conditions.

Figure 3. Density plot showing the sensitivity of SPEI to AED worldwide at the annual scale, considering the complete series and precipitation series corresponding to the 10th, 25th and 50th centiles of annual precipitation.

3.3. The Role of Mean Precipitation

In addition to differences in SPEI sensitivity to AED, as driven by the precipitation amount, this study also suggests coherent spatial patterns considering the average precipitation amount (refer to Figure 1). Specifically, the sensitivity of SPEI to AED was much lower in humid regions than in dry regions, irrespective of the investigated sample (i.e., either from the complete series or series of precipitation below the 10th, 25th, and 50th percentiles) and the temporal scale (i.e., monthly, seasonal, and annual). In this context, although some regions exhibited an increase of the sensitivity of SPEI to AED during periods of low precipitation, other regions showed less sensitivity of

Figure 4. Relationship between the mean annual precipitation and the sensitivity of SPEI to AED considering the annual series. (left to right) The plots consider the (fourth panel) complete series and the precipitation series during the years corresponding to the (first panel) 10th, (second panel) 25th, and (third panel) 50th of annual precipitation. Colors represent the point density (red: high density, blue: low density).

SPEI to AED, even during low-precipitation periods. Again, this informs that the sensitivity of SPEI to AED is driven largely by climatology of precipitation, regardless of SPEI time scale (Figures 4 and S19 to S22). When considering the entire series, we found an exponential relationship between the sensitivity of SPEI to AED and mean precipitation. This exponential relationship was also found for the periods of low precipitation. Interestingly, the frequency of regions that showed a low sensitivity of SPEI to AED in very humid regions ($>3,000 \text{ mm year}^{-1}$) did not change substantially when considering the years of low precipitation. In areas characterized by subhumid to very arid conditions ($0 \text{ to } 800 \text{ mm year}^{-1}$), we noted an increase in the frequency of areas showing high sensitivity of SPEI to AED during low-precipitation years. This implied that the majority of these regions exhibited a clear increase in the sensitivity of SPEI to AED during low-precipitation years. Focusing on the driest years (10th centile), most of these regions showed sensitivities higher than 50%. This pattern is clearly evident when exploring the sensitivity of SPEI to AED, as a function of different precipitation averages (Figures 5 and S23 to S38).

3.4. General Implications

In general, our analysis based on long-term climate data of precipitation and AED revealed that the sensitivity of SPEI to AED showed a strong temporal coherence. In particular, SPEI was only sensible to AED during low-precipitation periods, in which an increase of AED would undoubtedly intensify the severity of hydrological, agricultural and environmental droughts. Interestingly, this behavior is not spatially homogeneous, as dry to subhumid regions, which are characterized by frequent and even critical water deficits, experienced high sensitivity to AED during low-precipitation periods. In contrast, the most humid regions of the world showed weak sensitivity to AED even under low precipitation conditions. High-precipitation amounts in these humid regions can modulate the impacts of the interannual variability of AED on drought severity, although some influence can be found in dry years. This was clearly evident for the Congo site, where AED showed a relatively moderate role (15–35%) during dry seasons with very low precipitation (Figure 1). These regions can still be triggered by extreme droughts, mainly driven by low precipitation (e.g., the Amazon, J. A. Marengo & Espinoza, 2016; Jose A. Marengo et al., 2011; Equatorial Africa, Sorí et al., 2017; or Indonesia, Ummenhofer et al., 2013; Wooster et al., 2012). In the meantime, AED could reinforce vegetation stress and water deficits during these anomalous dry events.

The role of AED in drought severity (as revealed by SPEI) is robust, confirming that drought severity cannot be impacted by any possible overestimation due to the role of AED. This role is obviously evident during low-precipitation periods (S. M. Vicente-Serrano, McVicar, et al., 2020). Any uncertainties originating from the overestimation of AED contribution on SPEI could only be raised during high-precipitation periods (S. M. Vicente-Serrano, McVicar, et al., 2020). Nevertheless, this possible effect is meaningless, given that the role of AED is practically negligible during nondry periods.

Small changes in the values drought indices accounting for AED in their formulations can be expected for the long term (Jonathan Spinoni et al., 2019), mostly as a consequence of the global weak changes in precipitation during the last decades (Orlowsky & Seneviratne, 2013; J. Spinoni et al., 2014). Nevertheless,

Figure 5. Boxplots showing the sensitivity of SPEI to AED at the annual scale for the (top left) complete series (centile 100th) and the precipitation series corresponding to the (bottom right) 10th, (bottom left) 25th, and (top right) 50th of annual precipitation. The central solid line indicates the median. The whiskers represent the 10th and the 90th, while the 25th and the 75th are plotted as the vertical lines of the bounding boxes.

impact-based studies suggest that when precipitation deficits are recorded, an increased AED contributes to drought intensification. This was perfectly described by Breshears et al. (2005) who introduced for the first time the term “global-change-type drought” to refer to reinforced drought conditions as a consequence of global warming. This concept is consistent with what is dominantly observed; mainly the global increase of AED due to recent warming processes (S. M. Vicente-Serrano, McVicar, et al., 2020). Therefore, although we cannot affirm with confidence that AED affects drought trends quantified by climate indices like SPEI, we still believe that dryness can be reinforced by enhanced AED when precipitation deficits occur. This has been evident for many regions worldwide, including Europe in 2003 (García-Herrera et al., 2010) and 2017 (García-Herrera et al., 2019), and southern portions of North America over the past two decades (Udall & Overpeck, 2017; Xiao et al., 2018). Moreover, this would not be inconsistent with the general evolution of mean vegetation and hydrological metrics over the last decades worldwide (Scheff, 2018). Rather, it would be consistent with enhanced forest decay (Carnicer et al., 2011) and mortality episodes in response to severe drought events reinforced by higher AED (Allen et al., 2015; Anderegg et al., 2013). Herein, it should be stressed that the role of AED on intensification of drought would be independent of the different thermodynamic and dynamic mechanisms that drive AED variability (see further details in Vicente-Serrano, McVicar, et al., 2020).

Studies based on climate change projections reveal a reinforcement of future dryness conditions, as revealed by SPEI (Naumann et al., 2018; Jonathan Spinoni, Barbosa, Buchignani, et al., 2019; S. M. Vicente-Serrano, Quiring, et al., 2020). This trend would be stronger in arid and semiarid regions. Cook et al. (2014) suggested that this could be a possible sign of dryness overestimation caused by SPEI formulation, which continues responding to changes in AED although ET supply is limited. Nevertheless, we must state here that AED is not included in SPEI as a substitute of ET (Beguéría et al., 2014). Rather, it is included as a more comprehensive driver of drought severity considering different drought types (S. M. Vicente-Serrano, McVicar, et al., 2020). Here, we have demonstrated that AED mostly affects SPEI during periods of low precipitation. From a hydrological perspective, AED during low-precipitation periods would contribute to dry the soil until soil moisture is depleted (A. Park Williams et al., 2020), given that AED would be the upper limit of ET. AED would increase free water evaporation and water consumption by irrigated lands; all of these aspects

would increase hydrological drought severity. From agricultural and ecological perspectives, AED is a better proxy of vegetation stress than ET, making it an important input of any drought metric including SPEI. This is simply because under dry conditions, characterized by low soil moisture levels, ET will be limited by leaf stomata regulation (Chaves et al., 2016). As such, ET would be less sensitive to AED increases, which would enhance plant stress, with a subsequent reduction in leaf photosynthesis, carbohydrate assimilation, secondary growth and ultimately xylem embolism and plant mortality in the most critical cases. Therefore, the sensitivity of SPEI to AED during anomalous low-precipitation periods can give strong indications on the reinforced role of AED in determining the severity of environmental and hydrological drought conditions.

Other studies have suggested that soil moisture simulations by coupled land-atmosphere models would be better metrics of dryness than climate drought indices including SPEI (e.g., Berg & Sheffield, 2018; Greve et al., 2019). Irrespective of the wide spectrum of uncertainty introduced in this kind of models when reproducing the temporal variability of soil moisture observations (see Ford & Quiring, 2019; Stillman et al., 2016; Yuan & Quiring, 2017), we stress that the enhanced role of AED on plant dryness under low-precipitation periods would not be well-recorded if we only consider soil moisture but we do not assess the AED influence. Moreover, several studies agree that the projections of soil moisture and SPEI for future scenarios are almost identical (see Feng et al., 2017; Vicente-Serrano et al., 2019; Xu et al., 2019), and even comparable to those suggested by PDSI (Dai et al., 2018).

Importantly, we would like to highlight some issues related to the possible goodness of SPEI for drought assessment in future scenarios. First, the adequate sensitivity of SPEI to AED identified in this study would be independent of any possible uncertainties in AED evolution in the upcoming decades. Second, SPEI is capable of including the possible influence of CO₂ atmospheric concentrations, which could affect AED in the future (Milly & Dunne, 2016; Roderick et al., 2015) since it is not constrained by any AED metric. Specifically, although there exist uncertainties related to the possible role of CO₂ concentrations on AED (B. Cook et al., 2018; S. M. Vicente-Serrano, McVicar, et al., 2020), the theoretical capacity of SPEI to adequately record the impacts of AED during low-precipitation periods would be independent of this issue. Regardless of AED behavior in response to the impact of CO₂ concentrations, this would not affect the overall accuracy of SPEI in capturing the specific role of AED during low-precipitation periods. Indeed, any future scenarios of a decrease in precipitation over some periods of regions would correspond to an increase in the sensitivity of SPEI to AED. However, this would be closely related to the expected role of this variable in the context of reinforcing hydrological and environmental drought conditions.

4. Conclusions

This study provides a global characterization of the sensitivity of the SPEI to the AED. Results demonstrate the following:

1. Drought severity (as suggested by SPEI values) is mostly driven by precipitation variability in the majority of the land regions, with the exception of the most arid regions worldwide.
2. In arid regions, AED contributes significantly to drought evolution. Importantly, this study stresses that the sensitivity of SPEI to AED is modulated largely by precipitation amounts, with a more pronounced role of AED in drought behavior during periods of low precipitation.
3. SPEI did not overestimate the role of AED during normal and humid periods, in which the influence of AED on the severity of hydrological, agricultural, and environmental drought severity should be negligible.
4. Moreover, this influence is not spatially homogeneous worldwide, as humid regions showed less sensitivity of SPEI to AED, even during the driest periods.

Overall, these spatial and temporal patterns make SPEI a highly suitable and recommended drought metric to monitor and quantify current and future drought severity under different global climatic conditions.

Data Availability Statement

Data used in this study are available at this site (https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_3.26/).

Acknowledgments

This work was supported by the research projects CGL2017-82216-R and PCI2019-103631, financed by the Spanish Commission of Science and Technology and Fondo Europeo de Desarrollo Regional (FEDER); CROSSDRO project financed by the AXIS (Assessment of Cross(X)—sectoral climate Impacts and pathways for Sustainable transformation), JPI-Climate cofunded call of the European Commission and INDECIS which is part of ERA4CS, an ERA-NET initiated by JPI Climate, and funded by FORMAS (SE), DLR (DE), BMFWF (AT), IFD (DK), MINECO (ES), and ANR (FR) with cofunding by the European Union (Grant 690462). Dhais Peña-Angulo received a “Juan de la Cierva” postdoctoral contract (FJCI-2017-33652 from Spanish Ministry of Economy and Competitiveness, MEC).

References

- Allen, C. D., Breshears, D. D., & McDowell, N. G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*, 6(8), 129. <https://doi.org/10.1890/ES15-00203.1>
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., et al. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 259(4), 660–684. <https://doi.org/10.1016/j.foreco.2009.09.001>
- Anderegg, W. R. L., Kane, J. M., & Anderegg, L. D. L. (2013). Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change*, 3(1), 30–36. <https://doi.org/10.1038/nclimate1635>
- Babst, F., Bouriaud, O., Poulter, B., Trouet, V., Girardin, M. P., & Frank, D. C. (2019). Twentieth century redistribution in climatic drivers of global tree growth. *Science Advances*, 5(1), eaat4313. <https://doi.org/10.1126/sciadv.aat4313>
- Bachmair, S., Kohn, I., & Stahl, K. (2015). Exploring the link between drought indicators and impacts. *Natural Hazards and Earth System Sciences*, 15(6), 1381–1397. <https://doi.org/10.5194/nhess-15-1381-2015>
- Bachmair, S., Svensson, C., Hannaford, J., Barker, L. J., & Stahl, K. (2016). A quantitative analysis to objectively appraise drought indicators and model drought impacts. *Hydrology and Earth System Sciences*, 20(7), 2589–2609. <https://doi.org/10.5194/hess-20-2589-2016>
- Bachmair, S., Tanguy, M., Hannaford, J., & Stahl, K. (2018). How well do meteorological indicators represent agricultural and forest drought across Europe? *Environmental Research Letters*, 13(3), 034042. <https://doi.org/10.1088/1748-9326/aaafda>
- Barker, L. J., Hannaford, J., Chiveron, A., & Svensson, C. (2016). From meteorological to hydrological drought using standardised indicators. *Hydrology and Earth System Sciences*, 20(6), 2483–2505. <https://doi.org/10.5194/hess-20-2483-2016>
- Begueria, S., Vicente-Serrano, S. M., Reig, F., & Latorre, B. (2014). Standardized precipitation evapotranspiration index (SPEI) revisited: Parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *International Journal of Climatology*, 34(10), 3001–3023. <https://doi.org/10.1002/joc.3887>
- Berg, A., & Sheffield, J. (2018). Climate change and drought: The soil moisture perspective. *Current Climate Change Reports*, 4(2), 180–191. <https://doi.org/10.1007/s40641-018-0095-0>
- Blauhut, V., Gudmundsson, L., & Stahl, K. (2015). Towards pan-European drought risk maps: Quantifying the link between drought indices and reported drought impacts. *Environmental Research Letters*, 10(1), 014008. <https://doi.org/10.1088/1748-9326/10/1/014008>
- Breshears, D. D., Cobb, N. S., Rich, P. M., Price, K. P., Allen, C. D., Balice, R. G., et al. (2005). Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences of the United States of America*, 102(42), 15,144–15,148. <https://doi.org/10.1073/pnas.0505734102>
- Brodribb, T. J., Powers, J., Cochard, H., & Choat, B. (2020). Hanging by a thread? Forests and drought. *Science*, 368(6488), 261–266. <https://doi.org/10.1126/science.aat7631>
- Byrne, M., & O’Gorman, P. (2013). Link between land-ocean warming contrast and surface relative humidities in simulations with coupled climate models. *Geophysical Research Letters*, 40, 5223–5227. <https://doi.org/10.1002/grl.50971>
- Byrne, M. P., & O’Gorman, P. A. (2018). Trends in continental temperature and humidity directly linked to ocean warming. *Proceedings of the National Academy of Sciences of the United States of America*, 115(19), 4863–4868. <https://doi.org/10.1073/pnas.1722312115>
- Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G., & Peñuelas, J. (2011). Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proceedings of the National Academy of Sciences of the United States of America*, 108(4), 1474–1478. <https://doi.org/10.1073/pnas.1010070108>
- Chaves, M. M., Costa, J. M., Zarrouk, O., Pinheiro, C., Lopes, C. M., & Pereira, J. S. (2016). Controlling stomatal aperture in semi-arid regions—The dilemma of saving water or being cool? *Plant Science*, 251, 54–64. <https://doi.org/10.1016/j.plantsci.2016.06.015>
- Chen, H., & Sun, J. (2015). Changes in drought characteristics over China using the standardized precipitation evapotranspiration index. *Journal of Climate*, 28(13), 5430–5447. <https://doi.org/10.1175/JCLI-D-14-00707.1>
- Chen, H., & Sun, J. (2017). Characterizing present and future drought changes over eastern China: Drought changes over eastern China. *International Journal of Climatology*, 37. <https://doi.org/10.1002/joc.4987>
- Cook, B., Mankin, J., & Anchukaitis, K. (2018). Climate change and drought: From past to future. *Current Climate Change Reports*, 4(2), 164–179. <https://doi.org/10.1007/s40641-018-0093-2>
- Cook, B. I., Seager, R., Park Williams, A., Puma, M. J., McDermid, S., Kelley, M., & Nazarenko, L. (2019). Climate change amplification of natural drought variability: The historic mid-twentieth-century North American drought in a warmer world. *Journal of Climate*, 32(17), 5417–5436. <https://doi.org/10.1175/JCLI-D-18-0832.1>
- Cook, B. I., Smerdon, J. E., Seager, R., & Coats, S. (2014). Global warming and 21st century drying. *Climate Dynamics*, 43(9–10), 2607–2627. <https://doi.org/10.1007/s00382-014-2075-y>
- Dai, A., Zhao, T., & Chen, J. (2018). Climate change and drought: A precipitation and evaporation perspective. *Current Climate Change Reports*, 4(3), 301–312. <https://doi.org/10.1007/s40641-018-0101-6>
- Dewes, C. F., Rangwala, I., Barsugli, J. J., Hobbins, M. T., & Kumar, S. (2017). Drought risk assessment under climate change is sensitive to methodological choices for the estimation of evaporative demand. *PLoS ONE*, 12(3), e0174045. Retrieved from. <https://doi.org/10.1371/journal.pone.0174045>
- Domínguez-Castro, F., Vicente-Serrano, S. M., Tomás-Burguera, M., Peña-Gallardo, M., Begueria, S., El Kenawy, A., et al. (2019). High-spatial-resolution probability maps of drought duration and magnitude across Spain. *Natural Hazards and Earth System Sciences*, 19(3), 611–628. <https://doi.org/10.5194/nhess-19-611-2019>
- Fang, W., Huang, S., Huang, Q., Huang, G., Wang, H., Leng, G., et al. (2019). Probabilistic assessment of remote sensing-based terrestrial vegetation vulnerability to drought stress of the Loess Plateau in China. *Remote Sensing of Environment*, 232, 111290. <https://doi.org/10.1016/j.rse.2019.111290>
- Feng, S., Trnka, M., Hayes, M., & Zhang, Y. (2017). Why do different drought indices show distinct future drought risk outcomes in the U.S. Great Plains? *Journal of Climate*, 30(1), 265–278. <https://doi.org/10.1175/JCLI-D-15-0590.1>
- Ford, T. W., & Quiring, S. M. (2019). Comparison of contemporary in situ, model, and satellite remote sensing soil moisture with a focus on drought monitoring. *Water Resources Research* Retrieved from <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85062331732&doi=10.1029%2F2018WR024039&partnerID=40&md5=db3e33e53648f77f20113b8a6529e756>, 55, 1565–1582. <https://doi.org/10.1029/2018WR024039>
- Frank, D. C., Poulter, B., Saurer, M., Esper, J., Huntingford, C., Helle, G., et al. (2015). Water-use efficiency and transpiration across European forests during the Anthropocene. *Nature Climate Change*, 5(6), 579–583. <https://doi.org/10.1038/nclimate2614>
- Friedrich, K., Grossman, R. L., Huntington, J., Blanken, P. D., Lenters, J., Holman, K. D., et al. (2018). Reservoir evaporation in the Western United States. *Bulletin of the American Meteorological Society*, 99(1), 167–187. <https://doi.org/10.1175/BAMS-D-15-00224.1>

- García-Herrera, R., Díaz, J., Trigo, R. M., Luterbacher, J., & Fischer, E. M. (2010). A review of the European summer heat wave of 2003. *Critical Reviews in Environmental Science and Technology*, 40(4), 267–306. <https://doi.org/10.1080/10643380802238137>
- García-Herrera, R., Garrido-Pérez, J. M., Barriopedro, D., Ordóñez, C., Vicente-Serrano, S. M., Nieto, R., et al. (2019). The European 2016/2017 drought. *Journal of Climate*, 32(11), 3169–3187. <https://doi.org/10.1175/JCLI-D-18-0331.1>
- Greve, P., Roderick, M. L., Ukkola, A. M., & Wada, Y. (2019). The aridity index under global warming. *Environmental Research Letters*, 14(12), 124,006. <https://doi.org/10.1088/1748-9326/ab5046>
- Gu, L., Chen, J., Yin, J., Sullivan, S. C., Wang, H.-M., Guo, S., et al. (2020). Projected increases in magnitude and socioeconomic exposure of global droughts in 1.5 and 2°C warmer climates. *Hydrology and Earth System Sciences*, 24(1), 451–472. <https://doi.org/10.5194/hess-24-451-2020>
- Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of monthly climatic observations—The CRU TS3.10 dataset. *International Journal of Climatology*, 34(3), 623–642. <https://doi.org/10.1002/joc.3711>
- Hartmann, H., Moura, C. F., Anderegg, W. R. L., Ruehr, N. K., Salmon, Y., Allen, C. D., et al. (2018). Research frontiers for improving our understanding of drought-induced tree and forest mortality. *New Phytologist* Retrieved from <https://www2.scopus.com/inward/record.uri?eid=2-s2.0-85042551215&doi=10.1111%2Fnp.15048&partnerID=40&md5=07a674361e2bd1b7d0d2dea607ef4fa4>, 218(1), 15–28. <https://doi.org/10.1111/nph.15048>
- Hobbins, M., Wood, A., Streubel, D., & Werner, K. (2012). What drives the variability of evaporative demand across the conterminous United States? *Journal of Hydrometeorology*, 13(4), 1195–1214. <https://doi.org/10.1175/JHM-D-11-0101.1>
- Knapp, A. K., Carroll, C. J. W., Denton, E. M., La Pierre, K. J., Collins, S. L., & Smith, M. D. (2015). Differential sensitivity to regional-scale drought in six central US grasslands. *Oecologia*, 177(4), 949–957. <https://doi.org/10.1007/s00442-015-3233-6>
- Konings, A. G., Williams, A. P., & Gentine, P. (2017). Sensitivity of grassland productivity to aridity controlled by stomatal and xylem regulation. *Nature Geoscience*, 10(4), 284. Retrieved from—288. <https://doi.org/10.1038/ngeo2903>
- Leng, G., & Hall, J. (2019). Crop yield sensitivity of global major agricultural countries to droughts and the projected changes in the future. *Science of the Total Environment*, 654, 811–821. <https://doi.org/10.1016/j.scitotenv.2018.10.434>
- Ma, M., Ren, L., Yuan, F., Jiang, S., Liu, Y., Kong, H., & Gong, L. (2014). A new standardized Palmer drought index for hydro-meteorological use. *Hydrological Processes*, 28(23), 5645–5661. <https://doi.org/10.1002/hyp.10063>
- Mankin, J. S., Seager, R., Smerdon, J. E., Cook, B. I., Williams, A. P., & Horton, R. M. (2018). Blue water trade-offs with vegetation in a CO₂-enriched climate. *Geophysical Research Letters*, 45, 3115–3125. <https://doi.org/10.1002/2018GL077051>
- Mankin, J. S., Seager, R., Smerdon, J. E., Cook, B. I., & Williams, A. P. (2019). Mid-latitude freshwater availability reduced by projected vegetation responses to climate change. *Nature Geoscience*, 12(12), 983–988. <https://doi.org/10.1038/s41561-019-0480-x>
- Marengo, J. A., & Espinoza, J. C. (2016). Extreme seasonal droughts and floods in Amazonia: Causes, trends and impacts. *International Journal of Climatology*, 36(3), 1033–1050. <https://doi.org/10.1002/joc.4420>
- Marengo, J. A., Tomasella, J., Alves, L. M., Soares, W. R., & Rodriguez, D. A. (2011). The drought of 2010 in the context of historical droughts in the Amazon region. *Geophysical Research Letters*, 38, L12703. <https://doi.org/10.1029/2011GL047436>
- Martínez-Granados, D., Maestre-Valero, J. F., Calatrava, J., & Martínez-Alvarez, V. (2011). The economic impact of water evaporation losses from water reservoirs in the Segura Basin, SE Spain. *Water Resources Management*, 25(13), 3153–3175. <https://doi.org/10.1007/s11269-011-9850-x>
- McDowell, N. G., & Allen, C. D. (2015). Darcy's law predicts widespread forest mortality under climate warming. *Nature Climate Change*, 5(7), 669–672. <https://doi.org/10.1038/nclimate2641>
- McDowell, N. G., Williams, A. P., Xu, C., Pockman, W. T., Dickman, L. T., Sevanto, S., et al. (2016). Multi-scale predictions of massive conifer mortality due to chronic temperature rise. *Nature Climate Change*, 6(3), 295–300. <https://doi.org/10.1038/nclimate2873>
- McKee, T. B., Doesken, N. J., & Kleist, J. (1993). The relationship of drought frequency and duration to time scales. *Eighth Conference on Applied Climatology*, 17–22.
- Menezes-Silva, P. E., Loram-Lourenço, L., Alves, R. D. F. B., Sousa, L. F., Almeida, S. E. d. S., & Farnese, F. S. (2019). Different ways to die in a changing world: Consequences of climate change for tree species performance and survival through an ecophysiological perspective. *Ecology and Evolution*, 9(20), 11,979–11,999. <https://doi.org/10.1002/ece3.5663>
- Milly, P. C. D., & Dunne, K. A. (2016). Potential evapotranspiration and continental drying. *Nature Climate Change*, 6(10), 946–949. <https://doi.org/10.1038/nclimate3046>
- Miralles, D., Gentine, P., Seneviratne, S. I., & Teuling, A. J. (2019). Land-atmospheric feedbacks during droughts and heatwaves: State of the science and current challenges. *Annals of the New York Academy of Sciences*, 8, 469.
- Mitchell, P. J., O'Grady, A. P., Hayes, K. R., & Pinkard, E. A. (2014). Exposure of trees to drought-induced die-off is defined by a common climatic threshold across different vegetation types. *Ecology and Evolution*, 4(7), 1088–1101. <https://doi.org/10.1002/ece3.1008>
- Morgan, J. A., Pataki, D. E., Körner, C., Clark, H., Del Grosso, S. J., Grünzweig, J. M., et al. (2004). Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO₂. *Oecologia* Retrieved from, 140(1), 11–25. <https://www2.scopus.com/inward/record.uri?eid=2-s2.0-1942544440&doi=10.1007%2Fs00442-004-1550-2&partnerID=40&md5=25b872f89af465862363ae853ef598ad>, <https://doi.org/10.1007/s00442-004-1550-2>
- Mukherjee, S., Mishra, A., & Trenberth, K. E. (2018). Climate change and drought: A perspective on drought indices. *Current Climate Change Reports*, 4(2), 145–163. <https://doi.org/10.1007/s40641-018-0098-x>
- Naumann, G., Alfieri, L., Wyser, K., Mentaschi, L., Betts, R. A., Carrao, H., et al. (2018). Global changes in drought conditions under different levels of warming. *Geophysical Research Letters*, 45, 3285–3296. <https://doi.org/10.1002/2017GL076521>
- Niu, S., Wu, M., Han, Y., Xia, J., Li, L., & Wan, S. (2008). Water-mediated responses of ecosystem carbon fluxes to climatic change in a temperate steppe. *New Phytologist*, 209–219. <https://doi.org/10.1111/j.1469-8137.2007.02237.x>
- Orlowsky, B., & Seneviratne, S. I. (2013). Elusive drought: Uncertainty in observed trends and short-and long-term CMIP5 projections. *Hydrology and Earth System Sciences*, 17(5), 1765–1781. <https://doi.org/10.5194/hess-17-1765-2013>
- Palmer, W. C. (1965). *Meteorological drought* (Research paper no. 45, pp. 1–58). Washington, DC: U.S. Department of Commerce Weather Bureau.
- Peña-Gallardo, M., Vicente-Serrano, S. M., Hannaford, J., Lorenzo-Lacruz, J., Svoboda, M., Domínguez-Castro, F., et al. (2019). Complex influences of meteorological drought time-scales on hydrological droughts in natural basins of the contiguous united states. *Journal of Hydrology*, 568, 611–625. <https://doi.org/10.1016/j.jhydrol.2018.11.026>
- Ribeiro, A. F. S., Russo, A., Gouveia, C. M., & Páscoa, P. (2019). Modelling drought-related yield losses in Iberia using remote sensing and multiscalar indices. *Theoretical and Applied Climatology*, 136(1–2), 203–220. <https://doi.org/10.1007/s00704-018-2478-5>
- Roderick, M. L., Greve, P., & Farquhar, G. D. (2015). On the assessment of aridity with changes in atmospheric CO₂. *Water Resources Research*, 51, 5450–5463. <https://doi.org/10.1002/2015WR017031>

- Scheff, J. (2018). Drought indices, drought impacts, CO₂, and warming: A historical and geologic perspective. *Current Climate Change Reports*, 4(2), 202–209. <https://doi.org/10.1007/s40641-018-0094-1>
- Slette, I. J., Post, A. K., Awad, M., Even, T., Punzalan, A., Williams, S., et al. (2019). How ecologists define drought, and why we should do better. *Global Change Biology*, 25(10), 3193–3200. <https://doi.org/10.1111/gcb.14747>
- Sori, R., Nieto, R., Vicente-Serrano, S. M., Drumond, A., & Gimeno, L. (2017). A Lagrangian perspective of the hydrological cycle in the Congo River basin. *Earth System Dynamics*, 8(3), 653–675. <https://doi.org/10.5194/esd-8-653-2017>
- Spinoni, J., Naumann, G., Carrao, H., Barbosa, P., & Vogt, J. (2014). World drought frequency, duration, and severity for 1951–2010. *International Journal of Climatology*, 34(8), 2792–2804. <https://doi.org/10.1002/joc.3875>
- Spinoni, J., Barbosa, P., Buccignani, E., Cassano, J., Cavazos, T., Christensen, J. H., et al. (2019). Future global meteorological drought hot spots: A study based on CORDEX data. *Journal of Climate*, 33(9), 3635–3661. <https://doi.org/10.1175/JCLI-D-19-0084.1>
- Spinoni, J., Barbosa, P., De Jager, A., McCormick, N., Naumann, G., Vogt, J. V., et al. (2019). A new global database of meteorological drought events from 1951 to 2016. *Journal of Hydrology: Regional Studies*, 22, 100593. <https://doi.org/10.1016/J.EJRH.2019.100593>
- Stillman, S., Zeng, X., & Bosilovich, M. G. (2016). Evaluation of 22 precipitation and 23 soil moisture products over a semiarid area in southeastern Arizona. *Journal of Hydrometeorology*, 17(1), 211–230. <https://doi.org/10.1175/JHM-D-15-0007.1>
- Swann, A. L. S., Hoffman, F. M., Koven, C. D., & Randerson, J. T. (2016). Plant responses to increasing CO₂ reduce estimates of climate impacts on drought severity. *Proceedings of the National Academy of Sciences of the United States of America*, 113(36), 10,019–10,024. <https://doi.org/10.1073/pnas.1604581113>
- Swann, A. L. S. (2018). Plants and drought in a changing climate. *Current Climate Change Reports*, 4(2), 192–201. <https://doi.org/10.1007/s40641-018-0097-y>
- Teuling, A. J., Van Loon, A. F., Seneviratne, S. I., Lehner, I., Aubinet, M., Heinesch, B., et al. (2013). Evapotranspiration amplifies European summer drought. *Geophysical Research Letters*, 40, 2071–2075. <https://doi.org/10.1002/grl.50495>
- Tsakiris, G., Pangalou, D., & Vangelis, H. (2007). Regional drought assessment based on the reconnaissance drought index (RDI). *Water Resources Management*, 21(5), 821–833. <https://doi.org/10.1007/s11269-006-9105-4>
- Udall, B., & Overpeck, J. (2017). The twenty-first century Colorado River hot drought and implications for the future. *Water Resources Research*, 53, 2404–2418. <https://doi.org/10.1002/2016WR019638>
- Ummenhofer, C. C., D'Arrigo, R. D., Anchukaitis, K. J., Buckley, B. M., & Cook, E. R. (2013). Links between Indo-Pacific climate variability and drought in the monsoon Asia Drought Atlas. *Climate Dynamics*, 40(5–6), 1319–1334. <https://doi.org/10.1007/s00382-012-1458-1>
- Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscale drought index sensitive to global warming: The standardized precipitation evapotranspiration index. *Journal of Climate*, 23(7), 1696–1718. <https://doi.org/10.1175/2009JCLI2909.1>
- Vicente-Serrano, S. M., McVicar, T. R., Miralles, D. G., Yang, Y., & Tomas-Burguera, M. (2020). Unraveling the influence of atmospheric evaporative demand on drought and its response to climate change. *Wiley Interdisciplinary Reviews: Climate Change*, 11(2). <https://doi.org/10.1002/wcc.632>
- Vicente-Serrano, S. M., Quiring, S., Peña-Gallardo, M., Domínguez-castro, F., & Yuan, S. (2020). A review of environmental droughts: Increased risk under global warming? *Earth Science Reviews*, 201, 102953. <https://doi.org/10.1016/j.earscirev.2019.102953>
- Vicente-Serrano, S. M., Zabalza-Martínez, J., Borràs, G., López-Moreno, J. I., Pla, E., Pascual, D., et al. (2017). Effect of reservoirs on streamflow and river regimes in a heavily regulated river basin of Northeast Spain. *Catena*, 149, 727–741. <https://doi.org/10.1016/j.catena.2016.03.042>
- Vicente-Serrano, S. M. M., Domínguez-castro, F., McVicar, T. R., Tomas-Burguera, M., Peña-gallardo, M., Noguera, I., et al. (2019). Global characterization of hydrological and meteorological droughts under future climate change: The importance of time-scales, vegetation-CO₂ feedbacks and changes to distribution functions. *International Journal of Climatology*, 40(5), 2557–2567. <https://doi.org/10.1002/joc.6350>
- Vicente-Serrano, S. M., Van der Schrier, G., Beguería, S., Azorin-Molina, C., & Lopez-Moreno, J.-I. (2015). Contribution of precipitation and reference evapotranspiration to drought indices under different climates. *Journal of Hydrology*, 526, 42–54. <https://doi.org/10.1016/j.jhydrol.2014.11.025>
- Von Uexkull, N., Croicu, M., Fjelde, H., & Buhaug, H. (2016). Civil conflict sensitivity to growing-season drought. *Proceedings of the National Academy of Sciences of the United States of America*, 113(44), 12,391–12,396. <https://doi.org/10.1073/pnas.1607542113>
- White, M. A., Running, S. W., & Thornton, P. E. (1999). The impact of growing-season length variability on carbon assimilation and evapotranspiration over 88 years in the eastern US deciduous forest. *International Journal of Biometeorology*, 42(3), 139–145. <https://doi.org/10.1007/s004840050097>
- Willett, K. M., Dunn, R. J. H., Thorne, P. W., Bell, S., De Podesta, M., Parker, D. E., et al. (2014). HadISDH land surface multi-variable humidity and temperature record for climate monitoring. *Climate of the Past*, 10(6), 1983–2006. <https://doi.org/10.5194/cp-10-1983-2014>
- Williams, A. P., Cook, E. R., Smerdon, J. E., Cook, B. I., Abatzoglou, J. T., Bolles, K., et al. (2020). Large contribution from anthropogenic warming to an emerging north American megadrought. *Science*, 368(6488), 314–318. <https://doi.org/10.1126/science.aaz9600>
- Williams, A. P., Allen, C. D., Macalady, A. K., Griffin, D., Woodhouse, C. A., Meko, D. M., et al. (2013). Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change*, 3(3), 292–297. <https://doi.org/10.1038/nclimate1693>
- Wooster, M. J., Perry, G. L. W., & Zoumas, A. (2012). Fire, drought and El Niño relationships on Borneo (Southeast Asia) in the pre-MODIS era (1980–2000). *Biogeosciences*, 9(1), 317–340. <https://doi.org/10.5194/bg-9-317-2012>
- Wu, H., Svoboda, M. D., Hayes, M. J., Wilhite, D. A., & Wen, F. (2007). Appropriate application of the standardized precipitation index in arid locations and dry seasons. *International Journal of Climatology*, 27(1), 65–79. <https://doi.org/10.1002/joc.1371>
- Xiao, M., Udall, B., & Lettenmaier, D. P. (2018). On the causes of declining Colorado River streamflows. *Water Resources Research*, 54, 6739–6756. <https://doi.org/10.1029/2018WR023153>
- Xu, L., Chen, N., & Zhang, X. (2019). Global drought trends under 1.5 and 2 °C warming. *International Journal of Climatology*, 39(4), 2375–2385. <https://doi.org/10.1002/joc.5958>
- Xu, Z., Jiang, Y., Jia, B., & Zhou, G. (2016). Elevated-CO₂ response of stomata and its dependence on environmental factors. *Frontiers in Plant Science*. Retrieved from <https://www2.scopus.com/inward/record.uri?eid=2-s2.0-84969581208&doi=10.3389%2Ffpls.2016.00657&partnerID=40&md5=16b54ab73d76a63032f7ed77b2803c58>
- Yang, Y., Roderick, M. L., Zhang, S., McVicar, T. R., & Donohue, R. J. (2019). Hydrologic implications of vegetation response to elevated CO₂ in climate projections. *Nature Climate Change*, 9(1), 44–48. <https://doi.org/10.1038/s41558-018-0361-0>
- Yuan, S., & Quiring, S. M. (2017). Evaluation of soil moisture in CMIP5 simulations over the contiguous United States using in situ and satellite observations. *Hydrology and Earth System Sciences*, 21(4), 2203–2218. <https://doi.org/10.5194/hess-21-2203-2017>

- Yuan, W., Zheng, Y., Piao, S., Ciais, P., Lombardozzi, D., Wang, Y., et al. (2019). Increased atmospheric vapor pressure deficit reduces global vegetation growth. *Science Advances*. Retrieved from <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85070882807&doi=10.1126%2Fsciadv.aax1396&partnerID=40&md5=accc4ceca4db1b9313b80ffb5f57dfec>, 5(8), eaax1396. <https://doi.org/10.1126/sciadv.aax1396>
- Zampieri, M., Ceglar, A., Dentener, F., & Toreti, A. (2017). Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales. *Environmental Research Letters*, 12(6), 064008. <https://doi.org/10.1088/1748-9326/aa723b>
- Zhang, B., Xia, Y., Huning, L. S., Wei, J., Wang, G., & AghaKouchak, A. (2019). A framework for global multicategory and multiscalar drought characterization accounting for snow processes. *Water Resources Research*, 55(11), 9258–9278. <https://doi.org/10.1029/2019wr025529>
- Zhang, Y., Yu, Z., & Niu, H. (2018). Standardized precipitation evapotranspiration index is highly correlated with total water storage over China under future climate scenarios. *Atmospheric Environment*, 194, 123–133. <https://doi.org/10.1016/j.atmosenv.2018.09.028>
- Zimmermann, J., Hauck, M., Dulamsuren, C., & Leuschner, C. (2015). Climate warming-related growth decline affects *Fagus sylvatica*, but not other broad-leaved tree species in central European mixed forests. *Ecosystems*, 18(4), 560–572. <https://doi.org/10.1007/s10021-015-9849-x>