**Genomic insights into the origin of trans-Mediterranean disjunct
distributions: The case of the saltmarsh band-winged
grasshopper (*Mioscirtus wagneri*)**

Running title: Inferring the origin of disjunct distributions

Víctor Noguerales^{1,2,3,*}, Pedro J. Cordero^{2,4}, L. Lacey Knowles³ and Joaquín Ortego⁵

¹ Department of Biological Sciences, University of Cyprus. PO Box 20537, Nicosia 1678,
Cyprus

² Genetic and Cultural Biodiversity Research Group, Instituto de Investigación en Recursos
Cinegéticos (IREC, CSIC-UCLM-JCCM). Ronda de Toledo 12, E-13071 Ciudad Real,
Spain

³ Department of Ecology and Evolutionary Biology, Museum of Zoology, University of
Michigan. 1109 Geddes Avenue, Ann Arbor, MI, 48109-1079, USA

⁴ Escuela Técnica Superior de Ingenieros Agrónomos (ETSIA), Universidad de Castilla-La
Mancha (UCLM), Ronda de Calatrava 7, E-13071 Ciudad Real, Spain

⁵ Department of Integrative Ecology, Estación Biológica de Doñana (EBD-CSIC). Avda.
Américo Vespucio 26, E-41092 Sevilla, Spain

*** CORRESPONDING AUTHOR**

Víctor Noguerales. Email: *noguerales.victor@ucy.ac.cy*

PRESENT ADDRESS

Víctor Noguerales. Island Ecology and Evolution Research Group, Institute of Natural Products and Agrobiology (IPNA-CSIC). Astrofísico Francisco Sánchez 3, E-38206 San Cristóbal de La Laguna, Tenerife, Spain. Email: *victor.noguerales@csic.es*

ACKNOWLEDGEMENTS

We are grateful to the people at the Knowles Lab for their advice during the preparation of genomic libraries and valuable help in data analysis. We also wish to thank José Miguel Aparicio and Nabil Amor for their help during field sampling and two anonymous referees for their constructive and valuable comments on an earlier version of the manuscript. Centro de Supercomputación de Galicia (CESGA) and Doñana's Singular Scientific-Technical Infrastructure (ICTS-RBD) provided access to computer resources. This work was funded by the Spanish Ministry of Economy and Competitiveness and the European Regional Development Fund (grants CGL2011-25053, CGL2016-80742-R, and CGL2017-83433-P).

Abstract

Aim: Two main biogeographic hypotheses have been proposed to explain the Mediterranean-Turanian disjunct distributions exhibited by numerous steppe-dwelling organisms, namely (i) dispersal during the Messinian salinity crisis (~5.96-5.33 Ma) followed by range fragmentation and vicariance, and (ii) Pleistocene colonization and recent processes of population subdivision (<2 Ma). Despite the two hypotheses postulate the role of climatic alterations and changes in landmass configuration on determining such disjunct distributions, estimates of the timing of lineage diversification have not been complemented so far with spatially-explicit tests providing independent evidence on the proximate processes underlying geographical patterns of population genetic connectivity/fragmentation.

Location: Mediterranean-Turanian region

Taxon: Saltmarsh band-winged grasshopper (*Mioscirtus wagneri*)

Methods: We integrate different sources of genetic (mtDNA and ddRADseq) and spatial information (configuration of emerged lands and niche modelling) to evaluate competing hypotheses of lineage diversification in the saltmarsh band-winged grasshopper, a halophile species showing a classical Mediterranean-Turanian disjunct distribution.

Results: Phylogenomic analyses reveal the presence of two North African cryptic lineages and support that trans-Mediterranean populations of the species diverged in the Pleistocene, with evidence of post-Messinian permeability of the Strait of Gibraltar to gene flow likely associated with sea level drops during glacial periods. Accordingly, spatial patterns of genetic differentiation are best explained by a scenario of population connectivity defined by the configuration of emerged landmasses and environmentally suitable habitats during glacial periods, a time when effective population sizes of the species peaked as inferred by genomic-based demographic reconstructions.

63 **Main conclusions:** Our results support post-Messinian colonization and Pleistocene
64 diversification as the biogeographic scenario best explaining the trans-Mediterranean
65 disjunct distributions of halophilous organisms.

66

67 **KEYWORDS:** bathymetry, biogeographic scenarios, ddRADseq, disjunct distributions,
68 Messinian, phylogenomics, Pleistocene

INTRODUCTION

Inferring the processes structuring genetic variation, through their effects on genetic drift and gene flow, and understanding their evolutionary consequences remain a central focus of both population genetics and phylogeography (Avice, 2009; Habel et al., 2015). However, despite large overlap in their theoretical bodies, these disciplines have tended to evolve independently. In part, this was due to the contrasting spatiotemporal scales at which the main evolutionary processes that they deal with occur and whose resolution has traditionally required the usage of different molecular markers (e.g., nuclear microsatellite *vs.* mitochondrial sequence markers; Wang, 2010). This idiosyncrasy has in part hampered our capacity to understand how the microevolutionary processes operating at population level ultimately drive speciation and lineage divergence at broader geographic and temporal scales (Papadopoulou & Knowles, 2017). Currently, the advent of high-throughput sequencing technologies and the possibility of generating vast genomic datasets has opened the door to investigate patterns of genetic variation at a wide range of evolutionary scales, blurring the boundaries of population genetics, phylogeography and phylogenetics (Rissler, 2016).

Applications of genomic data, especially when viewed through a statistical phylogeographic framework (Knowles, 2009), are essential to studies of the divergence histories of organisms across geologically dynamic and environmentally heterogeneous landscapes, such as the Mediterranean. The modern Mediterranean basin is the legacy of complex climate and geological dynamics that have taken place since the Late Miocene (Blondel & Aronson, 1999). This includes the Messinian salinity crisis (MSC, ~5.96-5.33 Ma), a geological episode during which the Mediterranean-Atlantic seaways progressively closed, leading to a dramatic sea-level drawdown (~1,500 m) in the Mediterranean Sea and its partial desiccation (Krijgsman, Hilgen, Raffi, Sierro, & Wilson, 1999). The resulting land

bridges connecting regions from southern Europe and northern Africa provided a favorable geographic setting for the expansion of terrestrial biotas throughout the Mediterranean region, especially for dry-adapted and steppe-dwelling taxa that benefited by the increased availability of open landscapes and arid environments (García-Alix et al., 2016). The reopening of the Atlantic-Mediterranean marine connection (Strait of Gibraltar) and refilling of the Mediterranean basin (~5.33 Ma; García-Castellanos et al., 2009) led to vicariant events in numerous organism groups that had previously expanded their distributions during the MSC (Chueca, Madeira, & Gómez-Moliner, 2015). The long-term persistence of populations in relict patches of optimal habitat since the Messinian has been often invoked as the most plausible explanation for the current disjunct distribution of many steppe or halophilous taxa with populations at the western- and easternmost portions of the Mediterranean basin (Ribera & Blasco-Zumeta, 1998). However, an alternative biogeographic hypothesis proposes that the origin of disjunct distributions in some trans-Mediterranean organisms reflect their ability to track existing suitable habitats in more recent times (i.e., post-Messinian; Allegrucci, Trucchi, & Sbordoni, 2011). According to this hypothesis, disjunctions arose by colonization events and subsequent isolation processes linked to Quaternary climate oscillations (Kadereit & Yaprak, 2008). Sea-level drawdown (~125 m; Litcher et al., 2010) and expansions of steppe-like environments (Kajtoch et al., 2016; Kirschner et al., 2020) during Pleistocene glacial periods might have facilitated the dispersal of steppe species across seaways and favorable habitats and, ultimately, led to the colonization of geographically distant regions that have remained isolated since then (Ortiz, Tremetsberger, Talavera, Stuessy, & García-Castaño, 2007). For example, the Strait of Gibraltar, where the distance between the Iberian Peninsula and Africa shortened to a few kilometers during the Pleistocene coldest stages, which might have favored the exchange of fauna and flora between the two continents (Graciá et al., 2013). However, the east-west vicariant distributions of several Mediterranean steppe-dwelling organisms have often been attributed to pre-Quaternary migration events linked to the MSC without proper evaluation of alternative

phylogeographic hypotheses (Ortego, Bonal, Cordero, & Aparicio, 2009; Ribera & Blasco-Zumeta, 1998). Clearly important insights can be gained from integrating precise information on the timing of population- and lineage-level divergence, detailed reconstructions of the demographic fate of the populations, and formal tests of alternative spatially-explicit scenarios of gene flow representing expectations of population connectivity under contrasting biogeographical hypotheses (Papadopoulou & Knowles, 2015).

Here, we apply an integrative and hypothesis testing framework to distinguish between competing hypotheses about the Mediterranean-Turanian disjunct distribution of the saltmarsh band-winged grasshopper, *Mioscirtus wagneri* (Eversmann, 1859) (Orthoptera: Acrididae). This species is the only representative of the monotypic genus *Mioscirtus* Saussure, 1888 (Cigliano, Braun, Eades, & Otte, 2020) and its vast range spans from the Iberian Peninsula to central Asia (Cordero, Llorente, & Aparicio, 2007). In the Mediterranean region it forms highly fragmented populations in Iberia, northwestern Africa (Morocco, Algeria and Tunisia) and the Middle East (Turkey, Israel, Palestine and Jordan), being absent from France, Balkan and Italian Peninsulas, northeastern Africa (Egypt and Libya), and all islands with the exception of Cyprus (Cigliano et al., 2020; Cordero et al., 2007; Figure 1). The saltmarsh band-winged grasshopper is a specialist species intimately associated to certain halophilic plants (*Suaeda* sp.) from the family Amaranthaceae on which it depends for feeding and shelter (Cordero et al., 2007). This restricted ecological requirement limits the distribution of the species to saline lowlands, including coastal marshes and inland endorheic lagoons and steppes. Three subspecies of the saltmarsh band-winged grasshopper have been described so far based on subtle phenotypic variations and differences in body size (Cigliano et al., 2020): *M. w. wagneri* (Eversmann, 1859) (type locality: Asia-Temperate, Middle Asia, Turkmenistan), *M. w. rogenhoferi* (Saussure, 1888) (type locality: Asia-Temperate, Western Asia, Iraq), and *M. w. maghrebi* Fernandes, 1968 (type locality: Africa, Northern Africa, Algeria). It has been hypothesized that the diversification of the saltmarsh

band-winged grasshopper in the Mediterranean region could be linked to the MSC. This would suggest that its present-day disjunct distribution may constitute the relicts of a wider range during the Late Tertiary that has since experienced gradual fragmentation (Ortego et al., 2009). However, the tempo and mode of divergence between the different populations and putative subspecies remain unresolved due to the lack of a comprehensive approach covering the entire Mediterranean distribution range of the species and formal testing of alternative biogeographical hypotheses.

In this study we focus on the trans-Mediterranean distribution range of the saltmarsh band-winged grasshopper, including populations separated >4,000 km that encompass the Mediterranean and Irano-Turanian regions and comprise the three putative subspecies (Cigliano et al., 2020) (Figure 1). Specifically, we integrate different sources of genetic (mtDNA and ddRADSeq) and spatial information (configuration of emerged lands and niche modelling) to evaluate competing phylogeographic hypotheses and determine whether lineage diversification is a consequence of dispersal during the MSC followed by range fragmentation and vicariance (~5.33 Ma) (Ribera & Blasco-Zumeta, 1998) or, alternatively, resulted from post-Messinian colonization and Pleistocene population subdivision (<2 Ma) (Graciá et al., 2013). First, we used coalescent-based methods to reconstruct the phylogenetic relationships among populations, estimate the timing of lineage divergence, and elucidate whether these are congruent with either Messinian or Pleistocene genetic fragmentation. Second, we integrated bathymetric information and environmental niche modelling (ENM) to determine the spatial configuration of emerged landmasses and climatically suitable areas at different time periods (Messinian, Pleistocene and present day) and generate alternative scenarios of population connectivity, the fit of which to our genetic data we evaluate based on the timing of population divergence estimated in the previous step. Finally, we examined the geographical distribution of genetic variation and reconstructed changes in effective population size through time in order to infer the

demographic responses of the different populations to past climate oscillations, to identify pulses of population expansion/contraction and to link them with the historical processes underlying genetic fragmentation and lineage formation.

MATERIALS AND METHODS

Sample collection

We collected 180 individuals from 20 populations of the saltmarsh band-winged grasshopper distributed throughout its Mediterranean distribution range, including populations from Spain, Portugal, Morocco, Tunisia, Turkey and Jordan (Figure 1; Table S1).

Genetic data

We extracted DNA and amplified and sequenced fragments of the 12S and 16S rRNA mitochondrial genes as detailed in González-Serna, Ortego & Cordero (2018). After sequence alignment, trimming and editing, gene fragments had 395 bp for 12S and 460 bp for 16S. From the 180 specimens, we selected 7-9 individuals per population (144 samples in total; Table S1) to be processed into three genomic libraries following the double-digestion restriction-fragment-based procedure (ddRADseq) described in Peterson, Weber, Jay, Fisher & Hoekstra (2012), with minor modifications detailed in Noguerales, Cordero & Ortego (2018). Each library was sequenced in a single-read 101-bp lane on an Illumina HiSeq2500 platform at The Centre for Applied Genomics (Toronto, Canada). Raw sequences were demultiplexed and preprocessed using STACKS 1.35 (Catchen, Hohenlohe, Bassham, Amores, & Cresko, 2013) and assembled using PYRAD 3.0.66 (Eaton, 2014). Appendix S1 provides all details on genomic data assembling and filtering.

Phylogenetic analyses: mtDNA data

We used mitochondrial DNA data and BEAST 1.8.0 (Drummond, Suchard, Xie, & Rambaut, 2012) to reconstruct the phylogenetic relationships among the different populations and estimate the timing of lineage divergence. We applied a HKY+I model of sequence evolution for both gene fragments as determined in JMODELTEST2 (Darriba, Taboada, Doallo, & Posada, 2012). We assumed a normal distributed substitution rate of 0.049 (SD = 0.0008) per site per million years for the 16S rRNA gene fragment (Papadopoulou, Anastasiou, & Vogler, 2010) and applied a continuous-time Markov chain (CTMCs) model for the 12S rRNA, as no explicit priors for the substitution rate for this gene fragment are available (e.g., González-Serna et al., 2018). A strict clock and a constant demographic model was used for phylogenetic reconstructions as determined by model testing using a generalized stepping-stone sampling approach (Baele, Lemey, & Suchard, 2016). We ran the analysis with two independent MCMC chains of 100 million generations each, sampling every 10,000 generations, and discarding the first 10% as burn-in.

Phylogenetic analyses: Genome-wide nuclear data

First, we reconstructed the phylogenetic relationships among populations using genome-wide SNP data and the coalescent-based method implemented in SVDQUARTETS (Chifman & Kubatko, 2014). We ran SVDQUARTETS exhaustively evaluating all possible quartets and performing nonparametric bootstrapping with 100 replicates for quantifying uncertainty in relationships. Second, we used BPP 4.0 to estimate the timing of lineage and population divergence (Flouri, Jiao, Rannala, & Yang, 2018). The phylogenetic tree inferred using SVDQUARTETS was fit as the fixed topology in BPP analyses (option A00). The *.loci* file from PYRAD was edited, converted into a BPP input file and filtered using custom R scripts (J-P. Huang, <https://github.com/airbugs/>; Huang, Hill, Ortego, & Knowles, 2020). Due to high computational burden, branch length estimation was inferred using five datasets consisting of

100, 200, 300, 400 and 500 randomly chosen loci (out of the 1,518 variable loci recovered) to confirm the consistency of the results (e.g., Huang et al., 2020). We applied an automatic adjustment of fine-tune parameters, allowing swapping rates to range between 0.30 and 0.70, and set the diploid option to indicate that the input sequences are unphased (for further details on the diploid option in BPP, see Flouri et al., 2018). Analyses were run for 500,000 generations, sampling every 10 generations, after a conservative burn in of 500,000 generations.

Population genetic structure

We inferred population genetic structure using genome-wide SNP data and the model-based clustering algorithm implemented in FASTSTRUCTURE 1.0 (Raj, Stephens, & Pritchard, 2014). We performed 10 independent runs for each of the different possible K genetic clusters (from $K = 1$ to $K = 10$) using the simple prior and a convergence criterion of 1×10^{-7} . We assessed the number of genetic clusters that best describes our data by estimating the metrics $K_{\emptyset}^*$, the value of K that maximizes log-marginal likelihood lower bound (LLBO) of the data, and K_{ϵ}^* , the smallest number of model components explaining at least 99% of cumulative ancestry contribution in our sample (Raj et al., 2014).

Landscape genetic analyses

We implemented a spatially-explicit approach based on circuit theory to model gene flow under alternative hypothetical scenarios of contemporary and historical population connectivity (McRae, 2006). Specifically, we used CIRCUITSCAPE 4.0.5 (McRae & Beier, 2007) to calculate resistance distance matrices between all pairs of populations based on isolation by resistance (IBR) scenarios defined according to (i) the spatial configuration of emerged lands during the present-day, last glacial maximum (LGM), and Messinian (IBR_{GEO};

see Appendix S2 for details) and (ii) the spatial configuration of climatically suitable habitats as inferred from projections of an environmental niche model (ENM) to present-day and LGM bioclimatic conditions (IBR_{CLI}; see Appendix S3 for details). In order to identify the resistance value for sea water that best explains observed estimates of genetic differentiation, we explored a range of resistance values (10-1,000,000) for this landscape feature plus a scenario that considered it as an impassable barrier to dispersal. For IBR_{GEO} scenarios, pixels coded as “emerged land” were assigned a resistance value of 1. For IBR_{CLI} scenarios, logistic environmental suitability scores (x) yielded by projections of the ENM were transformed as $1-x$, so that smaller pixel values offer lower resistance to gene flow. Additionally, we tested a null model of isolation-by-distance (IBD_{NULL}) by generating a completely “flat landscape” scenario based on a map in which all pixels (including sea water and emerged lands) have a fixed resistance (=1) value. We used multiple matrix regressions with randomization (MMRR; Wang, 2013) to test all resistance matrices against matrices of population genetic differentiation (F_{ST}) calculated in ARLEQUIN 3.5 (Excoffier & Lischer, 2010) for mtDNA and genome-wide SNP data.

Genetic diversity

We analyzed spatial clines of genetic diversity (π , nucleotide diversity) estimated for both mtDNA and genome-wide nuclear data using DNASP 5.10/6.12 (Rozas et al., 2017). We tested the association between genetic diversity (π) and geography (latitude and longitude) using generalized linear models (GLMs) as implemented in the R package ‘lm4’. GLMs were constructed using a Gaussian error distribution, an identity link function and a weighted least square (WLS) method, where weight equals the sample size for each population (Bates, Maechler, Bolker, & Walker, 2015).

Demographic inference

We inferred the demographic history of each population using STAIRWAY PLOT 2.0, which implements a flexible multi-epoch demographic model to estimate changes in effective population size (N_e) over time using the site frequency spectrum (SFS) (Liu & Fu, 2015). We used the script *easySFS.py* (I. Overcast, <https://github.com/isaacovercast/easySFS>) to calculate a folded SFS for each population. To avoid the effects of linkage disequilibrium and remove all missing data for the calculation of the SFS, we considered a single SNP per locus and retained only loci present in ~66% of individuals. Populations with <7 individuals (PHU and MER) were excluded from STAIRWAY PLOT analyses. Demographic reconstructions in STAIRWAY PLOT were run fitting a mutation rate of 2.8×10^{-9} per site per generation (Keightley, Ness, Halligan, & Haddrill, 2014), considering a one-year generation time (Cordero et al., 2007), and performing 200 bootstrap replicates to estimate 95% confidence intervals.

RESULTS

Genetic data: mtDNA data

We found 16 and 21 unique haplotypes for 12S and 16S gene fragments, respectively (Table S1). In particular, 11 and 14 haplotypes for 12S and 16S gene fragments, respectively, were exclusively found in a single population (Table S1). The remaining haplotypes were shared among individuals belonging to nearby populations located within the same geographic region (Table S1). Remarkably, individuals from Iberian (TIN) and African (HOC) populations located at both sides of the Strait of Gibraltar shared the same 12S and 16S haplotypes (Table S1).

Genetic data: Genome-wide nuclear data

Four individuals (one individual from PHU and three individuals from MER) recovered a very low number of reads (<10,000) and were discarded from subsequent analyses. After filtering and assembly steps in PYRAD, each remaining individual ($n = 140$) retained on average 1.83 million sequence reads (SD = 0.66 M), which represents 92.50% (SD = 2.29%) of their initial number of reads (Figure S1). On average, we recovered 49,415 loci (SD = 9,305; range = 23,170-79,257) per individual.

Phylogenetic analyses

Phylogenetic inference based on both mitochondrial (BEAST) and genome-wide nuclear (SVDQUARTETS) data consistently supported two highly divergent lineages only represented in the populations located in western Morocco (ZIM) and southern Tunisia (KEB) (Figure 2). The remaining populations were grouped into four clades corresponding to the Middle East, Africa along with southwestern Iberia, northeastern Iberia, and central-southeastern Iberia populations (Figure 2). All analyses showed consistently that Iberian populations were not reciprocally monophyletic, as populations from southwestern Iberia (CAS and TIN) were placed within the northern Africa clade. We found a discordance between mitochondrial and nuclear genealogies in the phylogenetic position of the northeastern Iberian clade (Ebro Valley): mtDNA data supported populations from northeastern Iberia as the most external lineage of the trans-Mediterranean clade, whereas nuclear data placed them as a sister lineage to the central-southeastern Iberian clade (Figure 2).

Divergence time estimation

Estimates of divergence time based on BEAST analyses for mtDNA revealed that the ZIM+KEB clade and the remaining lineages diverged from a shared common ancestor during the Pliocene (~4.5 Ma; Figure 2). According to BPP analyses based on genome-wide nuclear

data, the most ancient split was estimated around 0.01032 τ units (Figure 3). Assuming a genomic mutation rate of 2.8×10^{-9} per site per generation (Keightley et al., 2014) and a one-year generation time (Cordero et al., 2007), the oldest diversification event may have taken place during the Early Pleistocene, around 1.84 Ma ($\tau = 2\mu t$, being μ the mutation rate and t the divergence time). The subsequent split between ZIM and KEB populations was dated to occur during the Early Pleistocene (mtDNA data: ~ 1.7 Ma; genome-wide nuclear data: ~ 1.08 Ma, 0.00608 τ units). The diversification of the remaining clades triggered around the Middle Pleistocene (mtDNA data: ~ 1.0 Ma; genome-wide nuclear data: ~ 0.21 Ma, 0.00117 τ units) and finished with the split of northern Morocco and southwestern Iberia populations in the Late Pleistocene (mtDNA data: ~ 100 ka; genome-wide nuclear data: ~ 47.5 ka, 0.00026 τ units) (Figure 3). Inferences from BPP were consistent across analyses based on datasets considering a different number of loci (Figure S2).

Population genetic structure

Results from FASTSTRUCTURE analyses indicated $K = 6$ as the most likely number of genetic clusters ($K_{\theta}^c = 6$; $K_{\epsilon} = 6$). The inferred genetic groups were consistent with phylogenetic inferences (Figure 2b). The highly divergent lineages from ZIM and KEB populations of North Africa were included in two distinct genetic clusters (Figure 2b). The other four genetic clusters corresponded to the rest of North African populations plus southwestern Iberian populations, and populations from northeastern Iberia, central-southeastern Iberia, and the Middle East (Figure 2b).

Landscape genetic analyses

Pairwise F_{ST} matrices calculated for mitochondrial and genome-wide nuclear data (Table S2) were significantly correlated (Mantel test, $r = 0.581$, $p < 0.001$). Resistance distance matrices

estimated under most scenarios were positively correlated with population genetic differentiation (F_{ST}) at both nuclear and mitochondrial levels (Table S3). However, multiple matrix regression with randomization (MMRR) analyses consistently supported that the scenario of population connectivity based on the spatial configuration of emerged lands and climatically suitable areas during the LGM ($IBR_{CLI-LGM}$) was the best fit to our data and the only one retained into the final model (Table 1). This result was consistent across both mitochondrial and genome-wide nuclear data and when including and excluding from the analyses the highly divergent ZIM and KEB populations (Table 1).

Genetic diversity

Mitochondrial and genome-wide nuclear genetic diversity were not significantly correlated (Pearson's $r = -0.127$, $p = 0.592$). The highest levels of mitochondrial genetic diversity were found in ZIM and KEB populations followed by the three populations from northeastern Iberia, whereas several populations from central, southeastern and southwestern Iberia, northern Morocco and Jordan exhibited no mitochondrial genetic variation ($\pi = 0$; Table S1). Accordingly, we did not find any significant relationship between mitochondrial genetic diversity and longitude and latitude either analyzing all populations (longitude, $t = -0.360$, $p = 0.723$; latitude, $t = 0.030$, $p = 0.977$) or excluding ZIM and KEB populations (longitude, $t = 0.179$, $p = 0.860$; latitude, $t = 1.919$, $p = 0.074$). With regards to genome-wide nuclear data, populations from the Middle East (TUZ and DEA) showed the highest levels of genetic diversity, whereas populations from central and northeastern Iberia presented the lowest estimates (Figure 1; Table S1). Nuclear genetic diversity was positively correlated with longitude ($\beta = 1.19 \times 10^{-4}$, $t = 8.408$, $p < 0.001$) and negatively correlated with latitude ($\beta = -4.11 \times 10^{-4}$, $t = -7.407$, $p < 0.001$). Analyses excluding the highly divergent populations ZIM and KEB yielded analogous results (longitude: $\beta = 1.16 \times 10^{-4}$, $t = 8.981$, $p < 0.001$; latitude: $\beta = -4.74 \times 10^{-4}$, $t = -8.485$, $p < 0.001$).

377

378 **Demographic inference**

379 STAIRWAY PLOT analyses revealed strong declines in N_e for all populations after the end of the
380 last glacial period (~10-20 ka; Figure 4). Phylogenetically closer populations shared more
381 similar demographic histories in terms of the trends and magnitude of population size
382 changes. In line with analyses of nuclear genetic diversity, populations from central,
383 southeastern and northeastern Iberian Peninsula exhibited on average the lowest historical
384 estimates of N_e , Middle East populations presented the highest estimates, and populations
385 from northern Africa and southwestern Iberia had intermediate values (Figure 4).

386

387 **DISCUSSION**

388

389 Our phylogenetic analyses and the estimated timing of lineage splits, complemented with
390 evidence provided by spatially-explicit testing of alternative scenarios of gene flow, support a
391 post-Messinian colonization and Pleistocene genetic fragmentation as the most plausible
392 biogeographic hypothesis to explain the trans-Mediterranean disjunct distribution of the
393 saltmarsh band-winged grasshopper. Intriguingly, our analyses revealed that the species
394 exhibits a complex phylogeographic structure (Figure 2), with three main cladogenetic events
395 and a deep genetic split between parapatric northern Africa lineages that indicate cryptic
396 processes of allopatric divergence and call upon a taxonomic revision of this monotypic
397 genus (Cigliano et al., 2020). Shared genetic lineages between southwestern Iberia and the
398 Maghreb region support the recurrent permeability to gene flow of the Strait of Gibraltar
399 during the Pleistocene and point to the important role of sea-level fluctuations linked to
400 glacial cycles in promoting biotic exchanges between Europe and Africa.

401

Ancient cryptic lineages in the Maghreb region

One of the major questions in biogeography research on the western Palearctic realm is elucidating the processes underlying the disjunct distributions of many Mediterranean organisms (Sanmartín, 2003). We found that genetic variation in the saltmarsh band-winged grasshopper is organized into three main evolutionary lineages whose divergence most likely took place after the end of the Messinian age (Figures 2-3). The first two sister lineages are distributed in western Morocco (ZIM) and southern Tunisia (KEB), regions separated >1,600 km apart from each other. These lineages present a parapatric distribution with respect to their geographically closest populations that belong to the third and more widespread trans-Mediterranean clade. This intriguing phylogeographic structure contrasts with the prevailing biogeographic pattern in the Maghreb, where the foremost genetic discontinuities have been linked to the Moulouya River (northern Morocco) and the Kabylia region (northern Algeria) in numerous terrestrial organisms (e.g., Beddek et al., 2018). The genetic discontinuities and disjunct distributions of these lineages likely represent processes of allopatric isolation driven by inland landscape changes occurring along the Pliocene and Pleistocene throughout the Maghreb region. Since the Early Pliocene, the Mediterranean region experienced a progressive aridification-cooling climatic shift that led to the expansion of steppe formations (Jiménez-Moreno, Fauquette, & Suc, 2010). This shift to a drier climate regime was particularly notable in North Africa (Griffin, 2002), which could have facilitated the establishment and expansion of halophilous vegetation along vast endorheic lacustrine areas (i.e., sabkhas and chotts) located across the Maghreb and whose development dates back to the Late Messinian associated to local tectonic dynamics (Capella et al., 2018). These geomorphological and climate alterations could have synergistically contributed to create refugial areas for the species in northern Africa, leading to long-term isolation processes that might have impeded gene flow (e.g., via the evolution of reproductive isolation or lack of

secondary contact) with its parapatric sister lineage widely distributed across the Mediterranean.

Pleistocene divergence throughout the Mediterranean region

Focusing on the trans-Mediterranean clade, our phylogenetic inferences support an east-to-west divergence that led to the formation of three main lineages distributed in the Middle East, northern Africa and Iberian Peninsula with a distribution gap in Egypt and Libya. As consistently estimated on the basis of both mitochondrial and nuclear data (Figures 2-3), the split of these clades began around the Middle Pleistocene (~ 1.0 - 0.21 Ma) and continued until the Late Pleistocene (<100 ka). These findings are concordant with phylogeographic inferences (i.e., Quaternary westward expansion) for other trans-Mediterranean halophilic and steppe-dwelling taxa (Escudero, Vargas, Arens, Ouborg, & Luceño, 2010; Kadereit & Yaprak, 2008; Pérez-Collazos, Sánchez-Gómez, Jiménez, & Catalán, 2009; Weising & Freitag, 2007). Our analyses also revealed that the three main lineages previously described within the Iberian Peninsula on the basis of mitochondrial (Ortego et al., 2009) and nuclear microsatellite (Ortego, Aguirre, & Cordero, 2010) data are not reciprocally monophyletic (Figures 2-3), with populations from southwestern Iberia being nested within the northern Africa clade (see Husemann, Schmitt, Zachos, Ulrich, & Habel, 2014). In turn, we found a mitonuclear discordance in the phylogenetic position of the northeastern Iberian clade, which might have been caused by the interplay between historical secondary contact and different evolutionary and demographic processes (see Toews & Brelsford, 2012). These results point to two independent colonization events of the Iberian Peninsula during the Pleistocene followed by allopatric divergence from their respective African ancestors. Thus, our results strongly support that the Strait of Gibraltar was a permeable barrier to dispersal, probably in association with sea-level changes linked to Quaternary climatic oscillations (Graciá et al., 2013). The sea-level during Pleistocene glacial periods has been estimated to be

approximately 125 meters lower than at present in the western Mediterranean region (Rohling et al., 2017), a fact that would have reduced the distance between European and African coasts to less than 5-7 km at the Camariñal Sill area (Lujan et al., 2011). This contributed to the emergence of small islands and shoals facilitating “steeping-stone” dispersal and the sporadic exchange of terrestrial faunas across the Strait of Gibraltar (Cosson et al., 2005; see also Ortiz et al., 2007). Accordingly, our landscape genetic analyses supported that genetic differentiation among populations is best explained by the configuration of shorelines and suitable habitats during Pleistocene glacial periods and when sea surface is modelled to offer much more resistance to movement than emerged lands but without acting as impassable barrier to gene flow (Tables 1 and S3).

Genetic diversity and past demographic history

Demographic reconstructions revealed that all analyzed populations have experienced population size declines after the LGM coinciding with the beginning of the current interglacial stage (Figure 4), a period during which climatic conditions likely resulted in the progressive shrink of suitable open habitats for the species (Weising & Freitag, 2007). Palynological studies have evidenced that extensive areas of the Mediterranean region were vegetated by steppe-like formations during glacial periods that became progressively replaced by temperate forests during interglacial stages (Carrión et al., 2012; Sánchez-Goñi, Eynaud, Turon, & Shackleton, 1999). Consequently, the confinement of the saltmarsh band-winged grasshopper in refuges of suitable habitat during unfavorable periods may have led to processes of allopatric divergence along the Quaternary as reported for steppe-like and halophytic species presenting similar environmental requirements (Kajtoch et al., 2016; Kirschner et al., 2020). Demographic inferences based on genomic data contrast with the predictions of our ENM, which suggests that the extent of climatically suitable habitats for the species have tended to increase since the LGM (Figure 1f). This might reflect the difficulty of

climate-based niche modelling to identify refugial areas in highly specialized species tightly linked to scattered and small-size habitat patches (i.e., salt-marshes, sabkhas and chotts) for which spatially-explicit information about their past distribution and connectivity is not currently available (González-Serna, Cordero, & Ortego, 2019). The very limited dispersal capacity of the saltmarsh band-winged grasshopper documented in previous studies (Ortego, García-Navas, Nogueras, & Cordero, 2015) points to postglacial contraction of suitable habitats and regional extinction in northeastern Africa, rather than long-distance dispersal, as the most likely cause of the species' distribution gap in Egypt and Libya (Cigliano et al., 2020). Accordingly, our data showed a general pattern of genetic erosion, the total lack of genetic variation at mitochondrial level in many populations, and an east-to-west and south-to-north decline of nuclear genetic diversity (Figure 1b; see also Escudero et al., 2010). This pattern might reflect the strong fragmentation of most contemporary populations, historical declines revealed by demographic reconstructions, and founder events resulted from serial colonization from the putatively ancestral range located in the Middle East (Pérez-Collazos et al., 2009).

Taxonomic implications

Our study revealed cryptic diversity within the saltmarsh band-winged grasshopper, as well as incongruence between current subspecies designations and the inferred phylogenetic relationships (Cigliano et al., 2020). For example, the populations ZIM and KEB, putatively belonging to the subspecies *M. w. maghrebi*, accumulated more genetic divergence than the remaining populations assigned to the three subspecies (Figure 2). This finding highlights the need for evaluating the taxonomic status of these two North Africa populations as potentially distinct taxa, which would ideally require additional population sampling, the identification of potential contact zones with the other Maghrebian lineage to determine whether the two are reproductively isolated, and comprehensive morphological analyses that consider diagnostic

phenotypic traits (e.g., Huang et al., 2020; Nogueras et al., 2018). Our phylogenetic analyses also shed light on the controversial taxonomic status of Iberian populations (Cordero et al., 2007), supporting the existence of two putative subspecies: *M. w. maghrebi* in southwestern Iberia (Fernandes, 1968) and *M. w. wagneri* in central, southeastern and northeastern Iberia (Cordero et al., 2007). Finally, our study clarifies the phylogenetic position of Turkish populations, where subspecies *M. w. wagneri* and *M. w. rogenhoferi* have been indistinctly considered in the literature (Naskrecki & Ünal, 1995; Ünal, 2011). In this sense, our analyses supported the close phylogenetic relationship between Turkish and Jordanian populations (Figures 2), which is coherent from a geographical point of view and agrees with the assignment of Middle East populations to the subspecies *M. w. rogenhoferi*, as reported in most Orthoptera inventories from the region (Katbeh-Bader, 2001; Naskrecki & Ünal, 1995).

CONCLUSIONS

Our study demonstrates that the disjunct distribution of the most widespread lineage of the saltmarsh band-winged grasshopper across the Mediterranean basin was shaped by its capacity to track and colonize suitable habitats during the Pleistocene (Graciá et al., 2013), rather than a consequence of fragmentation and long-term persistence of relict populations after range expansions linked to the MSC (Ribera & Blasco-Zumeta, 1998). The cohesiveness of Moroccan and southern Iberia populations evidences the post-Messinian permeability of the Strait of Gibraltar, while the presence of highly divergent cryptic lineages in northern Africa highlights the pivotal role of the Maghreb as a diversification area during the Pliocene-Pleistocene (Husemann et al., 2014). Overall, our study emphasizes the power of combining phylogenetic inference with spatially-explicit testing of alternative

biogeographical scenarios to unravel hidden diversification patterns and gain insights into the proximate processes underlying the origin of disjunct distributions.

DATA AVAILABILITY STATEMENT

Raw Illumina reads have been deposited at the NCBI Sequence Read Archive (SRA) under BioProject PRJNA663622. Mitochondrial DNA sequences for the 12S and 16S gene fragments are deposited in GenBank with accession numbers MW018492-MW018671 and MW018175-MW018354, respectively. Input files for all analyses are available for download from the Dryad Digital Repository (<https://doi.org/10.5061/dryad.qfttdz0fk>).

ORCID

Víctor Noguerales <https://orcid.org/0000-0003-3185-778X>

Pedro J. Cordero <https://orcid.org/0000-0002-1371-8009>

Lacey L. Knowles <https://orcid.org/0000-0002-6567-4853>

Joaquín Ortego <https://orcid.org/0000-0003-2709-429X>

REFERENCES

- Allegretti, G., Trucchi, E., & Sbordoni, V. (2011). Tempo and mode of species diversification in *Dolichopoda* cave crickets (Orthoptera, Rhaphidophoridae). *Molecular Phylogenetics and Evolution*, 60, 108-121.
- Avice, J. C. (2009). Phylogeography: Retrospect and prospect. *Journal of Biogeography*, 36, 3-15.
- Baele, G., Lemey, P., & Suchard, M. A. (2016). Genealogical working distributions for Bayesian model testing with phylogenetic uncertainty. *Systematic Biology*, 65, 250-264.
- Bates, D., Maechler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using *lme4*. *Journal of Statistical Software*, 67, 1-48.
- Beddek, M., Zenboudji-Beddek, S., Geniez, P., Fathalla, R., Sourouille, P., Arnal, V., . . . Crochet, P. A. (2018). Comparative phylogeography of amphibians and reptiles in Algeria suggests common causes for the east-west phylogeographic breaks in the Maghreb. *PLoS ONE*, 13, e0201218.
- Capella, W., Barhoun, N., Flecker, R., Hilgen, F. J., Kouwenhoven, T., Matenco, L. C., . . . Krijgsman, W. (2018). Palaeogeographic evolution of the late Miocene Rifian Corridor (Morocco): Reconstructions from surface and subsurface data. *Earth-Science Reviews*, 180, 37-59.
- Carrión, J.S., Fernández, S., Fierro, E., López-Merino, L., Munuera, M., 2012. Paleoflora y paleovegetación de la Península Ibérica e Islas Baleares: Plioceno-Cuaternario. Murcia, Spain: Ministerio de Economía y Competitividad.
- Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). STACKS: an analysis tool set for population genomics. *Molecular Ecology*, 22, 3124-3140.
- Chifman, J., & Kubatko, L. (2014). Quartet inference from SNP data under the coalescent model. *Bioinformatics*, 30, 3317-3324.

- 570 Cigliano, M.M., Braun, H., Eades, D.C., & Otte, D. (2020). Orthoptera Species File (OSF).
 571 <http://orthoptera.speciesfile.org> (accessed 10 March 2020).
- 572 Chueca, L. J., Madeira, M. J., & Gómez-Moliner, B. J. (2015). Biogeography of the land snail
 573 genus *Allognathus* (Helicidae): Middle Miocene colonization of the Balearic Islands.
 574 *Journal of Biogeography*, *42*, 1845-1857.
- 575 Cordero, P. J., Llorente, V., & Aparicio, J. M. (2007). New data on morphometrics,
 576 distribution and ecology of *Mioscirtus wagneri* (Kittary, 1859) (Orthoptera, Acrididae) in
 577 Spain: is *maghrebi* a well defined subspecies? *Graellsia*, *63*, 3-16.
- 578 Cosson, J. F., Hutterer, R., Libois, R., Sara, M., Taberlet, P., & Vogel, P. (2005).
 579 Phylogeographical footprints of the Strait of Gibraltar and Quaternary climatic
 580 fluctuations in the western Mediterranean: a case study with the greater white-toothed
 581 shrew, *Crocidura russula* (Mammalia: Soricidae). *Molecular Ecology*, *14*, 1151-1162.
- 582 Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). JMODELTEST 2: more models,
 583 new heuristics and parallel computing. *Nature Methods*, *9*, 772-772.
- 584 Drummond, A. J., Suchard, M. A., Xie, D., & Rambaut, A. (2012). Bayesian phylogenetics
 585 with BEAUTI and the BEAST 1.7. *Molecular Biology and Evolution*, *29*, 1969-1973.
- 586 Eaton, D. A. R. (2014). PYRAD: assembly of *de novo* RADseq loci for phylogenetic analyses.
 587 *Bioinformatics*, *30*, 1844-1849.
- 588 Escudero, M., Vargas, P., Arens, P., Ouborg, N. J., & Luceño, M. (2010). The east-west-
 589 north colonization history of the Mediterranean and Europe by the coastal plant *Carex*
 590 *extensa* (Cyperaceae). *Molecular Ecology*, *19*, 352-370.
- 591 Excoffier, L., & Lischer, H. E. L. (2010). ARLEQUIN suite ver 3.5: a new series of programs to
 592 perform population genetics analyses under Linux and Windows. *Molecular Ecology*
 593 *Resources*, *10*, 564-567.
- 594 Fernandes, J. A. (1968). A new subspecies of *Mioscirtus wagneri* Evers. *Arquivos do Museu*
 595 *Bocage, Segunda Série- Notas e suplementos*, *2*, 1-3.

- Flouri, T., Jiao, X., Rannala, B., & Yang, Z. (2018). Species tree inference with BPP using genomic sequences and the multispecies coalescent. *Molecular Biology and Evolution*, 35, 2585-2593.
- García-Alix, A., Minwer-Barakat, R., Martín-Suarez, E., Freudenthal, M., Aguirre, J., & Kaya, F. (2016). Updating the Europe-Africa small mammal exchange during the late Messinian. *Journal of Biogeography*, 43, 1336-1348.
- García-Castellanos, D., Estrada, F., Jiménez-Munt, I., Gorini, C., Fernández, M., Verges, J., & De Vicente, R. (2009). Catastrophic flood of the Mediterranean after the Messinian salinity crisis. *Nature*, 462, 778-U796.
- González-Serna, M. J., Cordero, P. J., & Ortego, J. (2019). Spatiotemporally explicit demographic modelling supports a joint effect of historical barriers to dispersal and contemporary landscape composition on structuring genomic variation in a red-listed grasshopper. *Molecular Ecology*, 28, 2155-2172.
- González-Serna, M. J., Ortego, J., & Cordero, P. J. (2018). A review of cross-backed grasshoppers of the genus *Dociostaurus* Fieber (Orthoptera: Acrididae) from the western Mediterranean: Insights from phylogenetic analyses and DNA-based species delimitation. *Systematic Entomology*, 43, 136-146.
- Graciá, E., Giménez, A., Anadón, J. D., Harris, D. J., Fritz, U., & Botella, F. (2013). The uncertainty of Late Pleistocene range expansions in the western Mediterranean: A case study of the colonization of south-eastern Spain by the spur-thighed tortoise, *Testudo graeca*. *Journal of Biogeography*, 40, 323-334.
- Griffin, D. L. (2002). Aridity and humidity: two aspects of the late Miocene climate of North Africa and the Mediterranean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 182, 65-91.
- Habel, J. C., Zachos, F. E., Dapporto, L., Rodder, D., Radespiel, U., Tellier, A., & Schmitt, T. (2015). Population genetics revisited - towards a multidisciplinary research field. *Biological Journal of the Linnean Society*, 115, 1-12.

- 623 Huang, J.-P., Hill, J., Ortego, J., & Knowles, L.L. (2020). Paraphyletic species no more –
 624 genomic data resolve a Pleistocene radiation and validate morphological species of the
 625 *Melanoplus scudderi* complex (Insecta: Orthoptera). *Systematic Entomology*, 45, 594-
 626 605.
- 627 Husemann, M., Schmitt, T., Zachos, F. E., Ulrich, W., & Habel, J. C. (2014). Palaeartic
 628 biogeography revisited: evidence for the existence of a North African refugium for
 629 Western Palaeartic biota. *Journal of Biogeography*, 41, 81-94.
- 630 Jiménez-Moreno, G., Fauquette, S., & Suc, J.-P. (2010). Miocene to Pliocene vegetation
 631 reconstruction and climate estimates in the Iberian Peninsula from pollen data. *Review*
 632 *of Palaeobotany and Palynology*, 162, 403-415.
- 633 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., . . .
 634 Wolff, E. W. (2007). Orbital and millennial Antarctic climate variability over the past
 635 800,000 years. *Science*, 317, 793-796.
- 636 Kadereit, G., & Yaprak, A.E. (2008). *Microcnemum coralloides* (Chenopodiaceae-
 637 Salicornioideae): an example of intraspecific East-West disjunctions in the
 638 Mediterranean region. *Anales del Jardín Botánico de Madrid*, 65: 415-426.
- 639 Katbeh-Bader, A. (2001). Acridoidae (Orthoptera) of Jordan. *Zoology in the Middle East*, 21,
 640 89-100.
- 641 Kajtoch, L., Cieslak, E., Varga, Z., Paul, W., Mazur, M. A., Sramko, G., & Kubisz, D. (2016).
 642 Phylogeographic patterns of steppe species in Eastern Central Europe: a review and the
 643 implications for conservation. *Biodiversity and Conservation*, 25, 2309-2339.
- 644 Keightley, P. D., Ness, R. W., Halligan, D. L., & Haddrill, P. R. (2014). Estimation of the
 645 spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib
 646 family. *Genetics*, 196, 313-320.
- 647 Kirschner, P., Závieská, E., Gamisch, A., Hilpold, A., Trucchi, E., Paun, O., . . . Schönschwetter,
 648 P. (2020). Long term isolation of European steppe outposts boots the biome's
 649 conservation value. *Nature Communications*, 11, 1968.

- 650 Knowles, L. L. (2009). Statistical phylogeography. *Annual Review of Ecology Evolution and*
651 *Systematics*, *40*, 593-612.
- 652 Krijgsman, W., Hilgen, F. J., Raffi, I., Sierro, F. J., & Wilson, D. S. (1999). Chronology,
653 causes and progression of the Messinian salinity crisis. *Nature*, *400*, 652-655.
- 654 Litcher, M., Zviely, D., Klein, M., & Sivan, D. (2010). Sea-level changes in the Mediterranean:
655 Past, present and future – a review. In A. Israel, R. Eina, & J. Seckbach (Eds.),
656 *Seaweeds and their role in globally changing environment. Cellular origin, life in extreme*
657 *habitats and astrobiology* (pp. 5-17). London, New York: Springer.
- 658 Liu, X., & Fu, Y.-X. (2015). Exploring population size changes using SNP frequency spectra.
659 *Nature Genetics*, *47*, 555-559.
- 660 Luján, M., Crespo-Blanc, A., & Comas, M. (2011). Morphology and structure of the
661 Camarinal Sill from high-resolution bathymetry: evidence of fault zones in the Gibraltar
662 Strait. *Geo-Marine Letters*, *31*, 163-174.
- 663 McRae, B. H. (2006). Isolation by resistance. *Evolution*, *60*, 1551-1561.
- 664 McRae, B. H., & Beier, P. (2007). Circuit theory predicts gene flow in plant and animal
665 populations. *Proceedings of the National Academy of Sciences of the United States of*
666 *America*, *104*, 19885-19890.
- 667 Naskrecki, P., & Ünal, M. (1995). The Orthoptera of Hatay Province, S. Turkey. *Beiträge zur*
668 *Entomologie*, *45*, 393-419.
- 669 Nogueras, V., Cordero, P. J., & Ortego, J. (2018). Integrating genomic and phenotypic data
670 to evaluate alternative phylogenetic and species delimitation hypotheses in a recent
671 evolutionary radiation of grasshoppers. *Molecular Ecology*, *27*, 1229-1244.
- 672 Ortego, J., Aguirre, M. P., & Cordero, P. J. (2010). Population genetics of *Mioscirtus wagneri*,
673 a grasshopper showing a highly fragmented distribution. *Molecular Ecology*, *19*, 472-
674 483.
- 675 Ortego, J., Bonal, R., Cordero, P. J., & Aparicio, J. M. (2009). Phylogeography of the Iberian
676 populations of *Mioscirtus wagneri* (Orthoptera: Acrididae), a specialized grasshopper

- 677 inhabiting highly fragmented hypersaline environments. *Biological Journal of the Linnean*
 678 *Society*, *97*, 623-633.
- 679 Ortego, J., García-Navas, V., Nogueras, V., & Cordero, P. J. (2015). Discordant patterns of
 680 genetic and phenotypic differentiation in five grasshopper species codistributed across a
 681 microreserve network. *Molecular Ecology*, *24*, 5796-5812.
- 682 Ortiz, M. A., Tremetsberger, K., Talavera, S., Stuessy, T., & García-Castaño, J. L. (2007).
 683 Population structure of *Hypochaeris salzmänniana* DC. (Asteraceae), an endemic
 684 species to the Atlantic coast on both sides of the Strait of Gibraltar, in relation to
 685 Quaternary sea level changes. *Molecular Ecology*, *16*, 541-552.
- 686 Papadopoulou, A., Anastasiou, I., & Vogler, A. P. (2010). Revisiting the insect mitochondrial
 687 molecular clock: the Mid-Aegean trench calibration. *Molecular Biology and Evolution*, *27*,
 688 1659-1672.
- 689 Papadopoulou, A., & Knowles, L. L. (2015). Species-specific responses to island connectivity
 690 cycles: refined models for testing phylogeographic concordance across a Mediterranean
 691 Pleistocene Aggregate Island Complex. *Molecular Ecology*, *24*, 4252-4268.
- 692 Papadopoulou, A., & Knowles, L. L. (2017). Linking micro- and macroevolutionary
 693 perspectives to evaluate the role of Quaternary sea-level oscillations in island
 694 diversification. *Evolution*, *71*, 2901-2917.
- 695 Pérez-Collazos, E., Sánchez-Gómez, P., Jiménez, J. F., & Catalán, P. (2009). The
 696 phylogeographical history of the Iberian steppe plant *Ferula loscosii* (Apiaceae): a test of
 697 the abundant-centre hypothesis. *Molecular Ecology*, *18*, 848-861.
- 698 Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double
 699 digest RADseq: an inexpensive method for *de novo* SNP discovery and genotyping in
 700 model and non-model species. *PLoS ONE*, *7*, e37135.
- 701 Raj, A., Stephens, M., & Pritchard, J. K. (2014). FASTSTRUCTURE: Variational inference of
 702 population structure in large SNP data sets. *Genetics*, *197*, 573-U207.

- Ribera, I., & Blasco-Zumeta, J. (1998). Biogeographical links between steppe insects in the Monegros region (Aragón, NE Spain), the eastern Mediterranean, and central Asia. *Journal of Biogeography*, 25, 969-986.
- Rissler, L. J. (2016). Union of phylogeography and landscape genetics. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 8079-8086.
- Rohling, E. J., Hibbert, F. D., Williams, F. H., Grant, K. M., Marino, G., Foster, G. L., . . . Yokoyama, Y. (2017). Differences between the last two glacial maxima and implications for ice-sheet, $\delta^{18}\text{O}$, and sea-level reconstructions. *Quaternary Science Reviews*, 176, 1-28.
- Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J. C., Guirao-Rico, S., Librado, P., Ramos-Onsins, S. E., & Sánchez-Gracia, A. (2017). DNASP 6: DNA sequence polymorphism analysis of large data sets. *Molecular Biology and Evolution*, 34, 3299-3302.
- Sánchez-Goñi, M. F., Eynaud, F., Turon, J. L., & Shackleton, N. J. (1999). High resolution palynological record off the Iberia margin: direct land-sea correlation for the Last Interglacial complex. *Earth and Planetary Science Letters*, 171: 123-137.
- Sanmartín, I. (2003). Dispersal vs. vicariance in the Mediterranean: historical biogeography of the Palearctic Pachydeminae (Coleoptera, Scarabaeoidea). *Journal of Biogeography*, 30, 1883-1897.
- Toews, D. P. L., & Brelsford, A. (2012). The biogeography of mitochondrial and nuclear discordance in animals. *Molecular Ecology*, 21, 3907-3930.
- Wang, I. J. (2010). Recognizing the temporal distinctions between landscape genetics and phylogeography. *Molecular Ecology*, 19, 2605-2608.
- Ünal, M. (2011). Turkish Orthoptera Site (TOS). <http://www.orthoptera-tr.org> (accessed 16 March 2020).
- Wang, I. J. (2013). Examining the full effects of landscape heterogeneity on spatial genetic variation: a multiple matrix regression approach for quantifying geographic and ecological isolation. *Evolution*, 67, 3403-3411.

730 Weising, K., & Freitag, H. (2007). Phylogeography of halophytes from European coastal and
731 inland habitats. *Zoologischer Anzeiger*, 246, 279-292.

732

733 BIOSKETCH

734 **Víctor Noguerales** is interested in understanding the ecological and evolutionary processes
735 shaping spatial patterns of biological diversity. His research harnesses the power of genomic
736 data to respond questions on how genetic variation arises and is maintained across different
737 levels of biological organization.

738

739 **Author contributions:** All authors conceived and designed the study and analyses. VN, PJC
740 and JO collected the samples. VN performed the lab work and analyzed the data guided by
741 JO, who performed the ENMs. VN wrote the manuscript with help of JO and inputs from PJC
742 and LLK.

TABLE 1 Multiple matrix regressions with randomization (MMRR) testing the relationship between mitochondrial (mtDNA) and nuclear (nDNA) genetic differentiation and distances calculated under alternative isolation-by-resistance (IBR) scenarios defined by the spatial configuration of emerged lands in the present (IBR_{GEO-CUR}), last glacial maximum (IBR_{GEO-LGM}) and Messinian (IBR_{GEO-MES}), and the spatial configuration of climatically suitable habitats in the present (IBR_{CLI-LGM}) and last glacial maximum (IBR_{CLI-LGM}). The IBD_{NULL} scenario refers to a layer in which all pixels had the same resistance value (=1). Each scenario initially considered a range of hypothetical resistance values offered by the sea water and only the one best fitting the data was included in final multivariate analyses presented in this table (see Table S3). Analyses were performed both including all populations and excluding the highly divergent ZIM and KEB populations.

Variable	All populations						Without ZIM and KEB					
	mtDNA			nDNA			mtDNA			nDNA		
	β	t	p	β	t	p	β	t	P	β	t	p
Explanatory terms												
Intercept	0.058			0.077			0.052			0.096		
IBR _{CLI-LGM}	0.531	8.714	0.001	0.410	6.662	0.001	0.552	8.057	0.001	0.356	4.977	0.001
Rejected terms												
IBD _{NULL}	0.222	2.269	0.119	0.114	1.373	0.477	0.203	1.734	0.283	0.068	0.552	0.607
IBR _{GEO-CUR}	-0.218	-0.569	0.737	-0.065	-	0.910	-	-	0.571	-0.709	-	0.220
					0.170		0.420	0.913			1.487	
IBR _{GEO-LGM}	0.135	0.266	0.868	0.009	0.010	0.997	-	-	0.799	-0.245	-	0.724
							0.241	0.412			0.402	
IBR _{GEO-MES}	0.226	2.092	0.204	0.130	1.601	0.467	0.202	1.652	0.310	0.052	0.407	0.715

IBR _{CLI-CUR}	-0.179	-0.565	0.769	0.081	0.167	0.927	-	-	0.519	0.029	0.195	0.877
							0.392	1.106				

751

FIGURE 1 (a) Genetic cluster membership and (b) genetic diversity of the study populations of saltmarsh band-winged grasshopper (*Mioscirtus wagneri*) as inferred using genome-wide nuclear data. Background maps display elevation and sea depth information. Panels c-d show the approximate spatial configuration of emerged lands during the maximum sea-level drop estimated for (c) the last glacial maximum (LGM; -125 m) and (d) the Messinian (-1,500 m). Panels e-f show the climate-based habitat suitability during the (e) present and (f) the LGM (~21 ka) as inferred by environmental niche modelling. Yellow dots indicate the location of sampling sites (panels c and d). Population codes as in Table S1.

761

762

FIGURE 2 (a, c) Phylogenetic relationships and (b) genetic structure of populations of the saltmarsh band-winged grasshopper (*Mioscirtus wagneri*). Panel (a) shows the maximum clade credibility tree for mitochondrial data (16S and 12S gene fragments) as inferred in BEAST. Estimates of divergence time for the main clades (median and lower and upper 95% highest posterior density, in brackets) and branch support values (*=1.0) are indicated. Panel (b) shows population genetic structure for the most likely number of clusters ($K = 6$) as inferred by FASTSTRUCTURE using genome-wide SNP data. Panel (c) shows the species-tree inferred by SVDQUARTETS using genome-wide SNP data. Bootstrapping support values are indicated on the nodes (*=100%). Putative membership of the populations to the different subspecies and population codes as in Table S1.

FIGURE 3 Divergence times estimated using BPP with a subset of 500 randomly chosen loci. The topology was fixed using the phylogenetic tree inferred using SVDQUARTETS. Bars on nodes indicate the 95% highest posterior densities (HPD) of the estimated divergence times. Background colors represent geological divisions as Early Pleistocene (EP, ~2.58-0.77 Ma), Middle Pleistocene (MP, ~0.77 Ma - 126 ka), Late Pleistocene (LP, ~126-11.7 ka) and Holocene (H, ~11.7 ka to present). Population codes as in Table S1.

FIGURE 4 Demographic history of the studied populations inferred using STAIRWAY PLOT. Lines represent the median estimate of the effective population size (N_e). Axes are logarithmically scaled and populations are grouped into the different panels according to the results of phylogenetic and clustering analyses. Upper panels represent temperature anomaly (ΔT °C) in the Late Quaternary as estimated from the EPICA (European Project for Ice Coring in Antarctica) Dome C ice core (Jouzel et al., 2007). Highlighted periods show the extent of the last glacial period (~110–11.7 ka), last glacial maximum (LGM: ~19–21 ka), and Holocene (~11.7 ka to present). Population codes as in Table S1.

- Supporting information -

Journal of Biogeography

Genomic insights into the origin of trans-Mediterranean disjunct distributions: The case of the saltmarsh band-winged grasshopper (*Mioscirtus wagneri*)

Running title: Inferring the origin of disjunct distributions

Víctor Nogueras^{1,2,3,*}, Pedro J. Cordero^{2,4}, L. Lacey Knowles³ and Joaquín Ortego⁵

¹ Department of Biological Sciences, University of Cyprus. PO Box 20537, Nicosia 1678, Cyprus

² Genetic and Cultural Biodiversity Research Group, Instituto de Investigación en Recursos Cinegéticos (IREC, CSIC-UCLM-JCCM). Ronda de Toledo 12, E-13071 Ciudad Real, Spain

³ Department of Ecology and Evolutionary Biology, Museum of Zoology, University of Michigan. 1109 Geddes Avenue, Ann Arbor, MI, 48109-1079, USA

⁴ Escuela Técnica Superior de Ingenieros Agrónomos (ETSIA), Universidad de Castilla-La Mancha (UCLM), Ronda de Calatrava 7, E-13071 Ciudad Real, Spain

⁵ Department of Integrative Ecology, Estación Biológica de Doñana (EBD-CSIC). Avda. Américo Vespucio 26, E-41092 Sevilla, Spain

*** CORRESPONDING AUTHOR**

Víctor Nogueras. Email: *nogueras.victor@ucy.ac.cy*

PRESENT ADDRESS

Víctor Nogueras. Island Ecology and Evolution Research Group, Institute of Natural Products and Agrobiology (IPNA-CSIC). Astrofísico Francisco Sánchez 3, E-38206 San Cristóbal de La Laguna, Tenerife, Spain. Email: *victor.nogueras@csic.es*

Appendix S1. Genomic data assembly and filtering.

Raw sequence reads were demultiplexed and quality-filtered using the *process_radtags* script within the STACKS 1.35 pipeline (Catchen, Hohenlohe, Bassham, Amores, & Cresko, 2013). Only reads with a Phred score > 10 (using a sliding window of 15%), no adaptor contamination, and that had an unambiguous barcode and restriction cut site were retained. Then, we checked the quality of raw sequences in FASTQC 0.11.5 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and trimmed them to 80 bp using TRIMMOMATIC 0.36 (Bolger, Lohse, & Usadel, 2014) in order to remove low-quality reads near the 3' ends. Reads retained after *process_radtags* were further quality-filtered using the program PYRAD 3.0.66 (Eaton, 2014) to convert base calls with a Phred score < 20 into *N*s and discard reads with > 2 *N*s (Figure S1). Afterwards, we used PYRAD to cluster retained reads within- and across samples considering a clustering threshold of sequence similarity of 85% ($W_{\text{clust}} = 0.85$). Clusters with a minimum coverage depth less than 5 ($d = 5$) and loci containing one or more heterozygous sites across more than 15% of individuals ($\text{maxSH} = 0.15$) and more than 10 polymorphic sites ($\text{maxSNPs} = 10$) were discarded (Eaton, 2014; Hipp et al., 2014). Final datasets used in all downstream analyses were generated in PYRAD by exporting loci present in at least the 50% of the samples ($\text{minCov} = 50\%$).

Appendix S2. Sea-level drop scenarios.

We modelled the contemporary and past (Pleistocene, Messinian) configuration of emerged lands using a digital terrain model incorporating fine-scale bathymetric data for the Mediterranean Sea available in the EMODNET BATHYMETRY portal (<http://www.emodnet-bathymetry.eu>). According to previous geological studies, we assumed a maximum sea-level drop of 125 m during Pleistocene

glaciations (Litcher, Zviely, Klein, & Sivan, 2010; Rohling et al., 2017) and 1,500 meters during the Messinian (Krijgsman, Hilgen, Raffi, Sierro, & Wilson, 1999; Roveri et al., 2014). A contemporary scenario was constructed considering the spatial configuration of current shoreline. In all scenarios, pixels were binary coded as either “emerged land” or “sea water”.

Appendix S3. Environmental niche modeling.

We constructed an environmental niche model (ENM) to predict climatically suitable habitats for the saltmarsh band-winged grasshopper in the present and during the LGM and test whether their spatial configuration explain patterns of genetic differentiation among populations. We built ENM by applying the maximum entropy algorithm implemented in MAXENT 3.3.3 (Phillips, Anderson, & Schapire, 2006; Phillips & Dudik, 2008) and using the present-day 19 bioclimatic variables from the WORLDCLIM dataset (<http://www.worldclim.org/>) interpolated to 30-arcsec resolution ($\sim 1 \text{ km}^2$) (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005). To generate environmental suitability maps during the LGM, contemporary species-climate relationships were projected onto the LGM bioclimatic conditions derived from the CCSM4 (Community Climate System Model) general atmospheric circulation model (Collins et al., 2006). We built ENM using occurrence data for our genotyped populations (Table S1), which allowed us to evaluate potential differences in the spatial distribution of environmental suitable habitats based on models constructed both including and excluding the highly divergent cryptic lineages from north Africa (populations ZIM and KEB). We used the R package ‘*ENMeval*’ (Muscarella et al., 2014) to conduct parameter tuning and determine the optimal feature class (FC) and regularization multiplier (RM) settings for MAXENT using a delete-one jackknife optimization approach, as recommended for small datasets (Muscarella et al., 2014; Shcheglovitova & Anderson, 2013; e.g., Wachter et al., 2016). We tested a total of 88 models of varying complexity by combining a range of regularization multipliers (RM)

(from 0 to 10 in increments of 1) with eight different feature classes (FC) combinations (L, LQ, LQP, H, T, LQH, LQHP, LQHPT, where L = linear, Q = quadratic, H = hinge, P = product and T = threshold) (Muscarella et al., 2014) and followed the approach detailed in González-Serna, Cordero, & Ortego (2019) for variable selection. We compared MAXENT models with different settings and evaluated their relative performance using the minimum training presence omission rate (OR_{MTP}) as the primary optimality criterion to protect from overfitting and the area under the curve of the receiver-operating characteristic plot on the testing data (AUC_{TEST}) as secondary criterion to maximize the discriminatory ability of the model (Wachter et al., 2016). An OR_{MTP} close to zero is indicative of a low degree of model overfitting (Radosavljevic & Anderson, 2014) whereas an AUC_{TEST} value close to one suggests a high discriminatory ability of the model (Peterson et al., 2011).

Parameter tuning performed using '*ENMeval*' showed that a threshold (T) feature class and a regularization multiplier of 1 increased the performance of the model in terms of OR_{MTP} and AUC_{TEST} statistics. After removing highly correlated variables and those with a zero percent contribution, ten bioclimatic layers were retained to build the final ENM (sorted by percent contribution, BIO12: 44.1 %; BIO1: 15.3 %; BIO7: 10.6 %; BIO14: 6.8 %; BIO4: 6.5 %; BIO3: 5.1 %; BIO9: 4.2 %; BIO8: 3.6 %; BIO19: 3.4 %; BIO15: 0.3 %). The moderately high AUC_{TEST} ($AUC_{TEST} = 0.85$) and low OR_{MTP} ($OR_{MTP} = 0.05$) estimates for the most supported model indicate that it has a reasonably high discriminatory power and a low degree of overfitting, respectively. ENMs built excluding the two occurrence points from the highly divergent populations ZIM and KEB yielded qualitatively identical results (results not shown).

REFERENCES

- Bolger, A. M., Lohse, M., & Usadel, B. (2014). TRIMMOMATIC: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*, 2114-2120.
- Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). STACKS: an analysis tool set for population genomics. *Molecular Ecology*, *22*, 3124-3140.
- Collins, W. D., Bitz, C. M., Blackmon, M. L., Bonan, G. B., Bretherton, C. S., Carton, J. A., . . . Smith, R. D. (2006). The Community Climate System Model version 3 (CCSM3). *Journal of Climate*, *19*, 2122-2143.
- Eaton, D. A. R. (2014). PYRAD: assembly of de novo RADseq loci for phylogenetic analyses. *Bioinformatics*, *30*, 1844-1849.
- González-Serna, M. J., Cordero, P. J., & Ortego, J. (2019). Spatiotemporally explicit demographic modelling supports a joint effect of historical barriers to dispersal and contemporary landscape composition on structuring genomic variation in a red-listed grasshopper. *Molecular Ecology*, *28*, 2155-2172.
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, *25*, 1965-1978.
- Hipp, A. L., Eaton, D. A. R., Cavender-Bares, J., Fitzek, E., Nipper, R., & Manos, P. S. (2014). A framework phylogeny of the American oak clade based on sequenced RAD data. *PLoS ONE*, *9*, e102272.
- Krijgsman, W., Hilgen, F. J., Raffi, I., Sierro, F. J., & Wilson, D. S. (1999). Chronology, causes and progression of the Messinian salinity crisis. *Nature*, *400*, 652-655.
- Litcher, M., Zviely, D., Klein, M., & Sivan, D. (2010). Sea-level changes in the Mediterranean: Past, present and future – a review. In A. Israel, R. Eina, & J. Seckbach (Eds.), *Seaweeds and their role in globally changing environment. Cellular origin, life in extreme habitats and astrobiology* (pp. 5-17). London, New York: Springer.
- Muscarella, R., Galante, P. J., Soley-Guardia, M., Boria, R. A., Kass, J. M., Uriarte, M., & Anderson, R. P. (2014). *ENMeval*: An R package for conducting spatially independent evaluations and estimating optimal model complexity for MAXENT ecological niche models. *Methods in Ecology and Evolution*, *5*, 1198-1205.
- Peterson, A.T, Soberón, J., Pearson, R.G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., & Araújo, M.B. (2011). *Ecological niches and geographic distributions. Monographs in Population Biology*. UK: Princeton University Press.
- Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, *190*, 231-259.

- Phillips, S. J., & Dudik, M. (2008). Modeling of species distributions with MAXENT: new extensions and a comprehensive evaluation. *Ecography*, *31*, 161-175.
- Radosavljevic, A., & Anderson, R. P. (2014). Making better MAXENT models of species distributions: complexity, overfitting and evaluation. *Journal of Biogeography*, *41*, 629-643.
- Rohling, E. J., Hibbert, F. D., Williams, F. H., Grant, K. M., Marino, G., Foster, G. L., . . . Yokoyama, Y. (2017). Differences between the last two glacial maxima and implications for ice-sheet, $\delta^{18}\text{O}$, and sea-level reconstructions. *Quaternary Science Reviews*, *176*, 1-28.
- Roveri, M., Flecker, R., Krijgsman, W., Lofi, J., Lugli, S., Manzi, V., . . . Stoica, M. (2014). The Messinian Salinity Crisis: Past and future of a great challenge for marine sciences. *Marine Geology*, *352*, 25-58.
- Shcheglovitova, M., & Anderson, R. P. (2013). Estimating optimal complexity for ecological niche models: A jackknife approach for species with small sample sizes. *Ecological Modelling*, *269*, 9-17.
- Wachter, G. A., Papadopoulou, A., Muster, C., Arthofer, W., Knowles, L. L., Steiner, F. M., & Schlick-Steiner, B. C. (2016). Glacial refugia, recolonization patterns and diversification forces in Alpine-endemic *Megabunus* harvestmen. *Molecular Ecology*, *25*, 2904-2919.

TABLE S1 Geographical location, putative subspecies membership, and number of analyzed individuals (n) for mitochondrial (mtDNA) and genome-wide nuclear (nDNA) data for each sampled population of the saltmarsh band-winged grasshopper (*Mioscirtus wagneri*). Haplotypes for 12S and 16S mitochondrial gene fragments (number of individuals with each haplotype in parentheses) and mtDNA and nDNA genetic diversity (π , nucleotide diversity) for each population are also indicated.

Population	Code	Country	Subspecies	Latitude	Longitude	n (mtDNA)	n (nDNA)	Haplotypes (12S)	Haplotypes (16S)	π ($\times 10^{-2}$) (mtDNA)	π ($\times 10^{-2}$) (nDNA)
Balsa la Salada	SAL	Spain	<i>M. w. wagneri</i>	41.50045 8	-0.733926	8	7	H1(7), H2(1)	H1(8)	0.088	0.054
Salada Grande	GRA	Spain	<i>M. w. wagneri</i>	41.04366 3	-0.199725	10	7	H1(1), H3(9)	H1(7), H2(3)	0.078	0.058
Pantano de Utxesa	UTX	Spain	<i>M. w. wagneri</i>	41.50385 3	0.548702	8	7	H3(8)	H1(5), H2(3)	0.063	0.068
Salobral de Ocaña	OCA	Spain	<i>M. w. wagneri</i>	39.98430 2	-3.635915	9	7	H4(7), H5(2)	H3(9)	0.046	0.087
Saladares de Huerta de Valdecarábanos	HUE	Spain	<i>M. w. wagneri</i>	39.83347 0	-3.617390	8	7	H4(8)	H3(8)	0.000	0.066
El Pedernoso	PED	Spain	<i>M. w. wagneri</i>	39.49183 4	-2.767505	9	7	H4(9)	H3(9)	0.000	0.044
Laguna de Peña Hueca	PHU	Spain	<i>M. w. wagneri</i>	39.51666 7	-3.333333	9	6	H4(9)	H3(9)	0.000	0.053
Las Tablas de Daimiel	DAI	Spain	<i>M. w. wagneri</i>	39.13684 7	-3.706739	10	7	H4(10)	H3(10)	0.000	0.033

Saladar	de	COR	Spain	<i>M. w. wagneri</i>	38.54381 4	-1.610672	9	7	H4(8), H6(1)	H3(7), H4(1), H5(1)	0.026	0.106	
Saladar	de	El	MAR	Spain	<i>M. w. wagneri</i>	37.64597 7	-2.600745	9	7	H4(9)	H3(9)	0.000	0.115
Castro Marim		CAS	Portugal	<i>M. w. wagneri</i> <i>maghrebi</i>	37.21379 9	-7.470991	9	7	H7(9)	H6(2), H7(7)	0.046	0.262	
Río Tinto		TIN	Spain	<i>M. w. wagneri</i> <i>maghrebi</i>	37.17461 5	-6.930724	8	7	H7(8)	H8(8)	0.000	0.310	
Al Hoceima		HOC	Morocco	<i>M. w. maghrebi</i>	35.19379 9	-3.864230	8	7	H7(8)	H8(8)	0.000	0.269	
Sebkha Zima		ZIM	Morocco	<i>M. w. maghrebi</i>	32.08040 7	-8.649739	9	7	H8(8), H9(1)	H9(7), H10(2),	0.091	0.278	
Boulaajoul		BOU	Morocco	<i>M. w. maghrebi</i>	32.89969 8	-4.970687	9	7	H7(1), H10(8)	H8(9)	0.026	0.384	
Merzouga		MER	Morocco	<i>M. w. maghrebi</i>	31.08103 9	-4.042237	10	4	H11(9), H12(1)	H11(2), H12(4), H13(4),	0.024	0.377	
Kairouan		KAI	Tunisia	<i>M. w. maghrebi</i>	35.68250 3	10.22462 4	10	8	H11(10)	H12(1), H13(7), H14(1), H15(1),	0.024	0.512	
Kebili		KEB	Tunisia	<i>M. w. maghrebi</i>	33.77542 8	9.434559	10	8	H13(9), H14(1)	H16(8), H17(1), H18(1)	0.094	0.340	
Tuz Gölü		TUZ	Turkey	<i>M. w. wagneri</i> <i>rogenhoferi</i>	38.70937 1	33.10047 6	9	8	H15(9)	H19(8), H20(1)	0.026	0.540	
Dead Sea		DEA	Jordan	<i>M. w. rogenhoferi</i>	31.81517 2	35.62892 0	9	8	H16(9)	H21(9)	0.000	0.923	

TABLE S2 Genetic differentiation (F_{ST}) between populations of the saltmarsh band-winged grasshopper (*Mioscirtus wagneri*) estimated using mitochondrial (above diagonal) and genome-wide nuclear (below diagonal) data. The significance of F_{ST} values was tested using Fisher's exact test after 10,000 permutations as implemented in ARLEQUIN 3.5. Significant pairwise F_{ST} values in bold. Population codes as in Table S1.

	SAL	GRA	UTX	OCA	HUE	PED	PHU	DAI	COR	MAR	CAS	TIN	HOC	ZIM	BOU	MER	KAI	KEB	TUZ	DEA
SAL	-	0.478	0.571	0.943	0.961	0.964	0.964	0.966	0.951	0.964	0.957	0.970	0.968	0.975	0.960	0.925	0.952	0.976	0.953	0.964
GRA	0.078	-	0.000	0.943	0.959	0.961	0.961	0.963	0.950	0.961	0.957	0.969	0.966	0.976	0.959	0.927	0.952	0.977	0.950	0.961
UTX	0.075	0.088	-	0.950	0.970	0.972	0.972	0.974	0.959	0.972	0.963	0.978	0.976	0.977	0.966	0.929	0.958	0.978	0.960	0.972
OCA	0.469	0.468	0.456	-	0.107	0.125	0.125	0.141	0.083	0.125	0.948	0.971	0.967	0.979	0.957	0.894	0.939	0.981	0.952	0.969
HUE	0.462	0.463	0.448	0.105	-	0.000	0.000	0.000	0.000	0.000	0.971	1.000	1.000	0.984	0.983	0.912	0.964	0.986	0.981	1.000
PED	0.512	0.513	0.492	0.155	0.130	-	0.000	0.000	0.000	0.000	0.973	1.000	1.000	0.985	0.984	0.917	0.966	0.987	0.982	1.000
PHU	0.436	0.439	0.418	0.170	0.142	0.158	-	0.000	0.000	0.000	0.973	1.000	1.000	0.985	0.984	0.917	0.966	0.987	0.982	1.000
DAI	0.457	0.459	0.446	0.154	0.130	0.135	0.091	-	0.012	0.000	0.975	1.000	1.000	0.985	0.985	0.921	0.968	0.988	0.983	1.000
COR	0.496	0.497	0.476	0.180	0.194	0.214	0.234	0.229	-	0.000	0.958	0.983	0.981	0.981	0.968	0.903	0.950	0.983	0.964	0.982
MAR	0.502	0.502	0.488	0.351	0.347	0.403	0.360	0.370	0.377	-	0.973	1.000	1.000	0.985	0.984	0.917	0.966	0.987	0.982	1.000
CAS	0.537	0.536	0.522	0.496	0.487	0.528	0.427	0.472	0.519	0.517	-	0.107	0.832	0.980	0.855	0.742	0.877	0.983	0.967	0.979
TIN	0.553	0.553	0.539	0.506	0.496	0.539	0.450	0.491	0.529	0.526	0.142	-	1.000	0.985	0.937	0.778	0.925	0.987	0.987	1.000
HOC	0.525	0.524	0.507	0.492	0.483	0.517	0.406	0.465	0.508	0.512	0.230	0.259	-	0.984	0.868	0.793	0.899	0.987	0.985	1.000
ZIM	0.866	0.866	0.857	0.854	0.846	0.875	0.787	0.824	0.872	0.869	0.737	0.777	0.675	-	0.982	0.970	0.980	0.943	0.980	0.985
BOU	0.496	0.498	0.481	0.474	0.463	0.503	0.369	0.438	0.496	0.492	0.198	0.241	0.131	0.582	-	0.821	0.898	0.984	0.973	0.986
MER	0.599	0.598	0.585	0.570	0.559	0.613	0.449	0.521	0.605	0.598	0.303	0.361	0.248	0.664	0.146	-	0.240	0.971	0.903	0.917
KAI	0.501	0.500	0.486	0.453	0.445	0.480	0.407	0.442	0.469	0.472	0.276	0.276	0.280	0.755	0.270	0.372	-	0.981	0.950	0.966
KEB	0.859	0.859	0.850	0.848	0.840	0.868	0.783	0.818	0.866	0.862	0.733	0.772	0.674	0.681	0.583	0.660	0.752	-	0.981	0.986
TUZ	0.563	0.564	0.548	0.546	0.538	0.571	0.452	0.511	0.564	0.562	0.378	0.420	0.324	0.636	0.258	0.326	0.379	0.633	-	0.973

DEA	0.508	0.509	0.490	0.484	0.475	0.505	0.405	0.459	0.496	0.497	0.323	0.353	0.279	0.661	0.242	0.332	0.295	0.658	0.168	-
-----	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	---

TABLE S3 Univariate matrix regressions with randomization testing the relationships between mitochondrial (mtDNA) and nuclear (nDNA) genetic differentiation and distances calculated under alternative isolation-by-resistance (IBR) scenarios. The coefficient of determination (R^2) and p -value are provided for each univariate model. The IBD_{NULL} scenario refers to a layer in which all pixels had the same resistance value (=1). For the IBR_{GEO} and IBR_{CLI} scenarios, we considered sea water as an impassable barrier to dispersal or assigned it a range of resistance values (1-1,000,000) in order to identify the resistance value for this landscape feature that best explain observed estimates of genetic differentiation. The IBR_{GEO} scenarios are based on the spatial configuration of emerged lands in the present (IBR_{GEO-CUR}), last glacial maximum (IBR_{GEO-LGM}) and Messinian (IBR_{GEO-MES}). The IBR_{CLI} scenarios are based on the spatial configuration of climatically suitable habitats in the present (IBR_{CLI-LGM}) and last glacial maximum (IBR_{CLI-LGM}). Analyses were performed both including all populations and excluding the highly divergent ZIM and KEB populations.

Scenario	Sea-water resistance value	All populations				Without ZIM and KEB			
		mtDNA		nDNA		mtDNA		nDNA	
		R^2	p	R^2	p	R^2	p	R^2	p
IBD _{NULL}	-	0.221	0.001	0.057	0.123	0.233	0.001	0.097	0.001
IBR _{GEO-CUR}	Barrier	0.182	0.002	0.185	0.001	0.192	0.004	0.121	0.003
IBR _{GEO-CUR}	10	0.242	0.001	0.050	0.155	0.257	0.001	0.082	0.004
IBR _{GEO-CUR}	10 ²	0.275	0.001	0.144	0.005	0.285	0.001	0.127	0.001
IBR _{GEO-CUR}	10 ³	0.201	0.001	0.184	0.001	0.211	0.002	0.124	0.001
IBR _{GEO-CUR}	10 ⁴	0.184	0.002	0.185	0.001	0.194	0.003	0.121	0.002
IBR _{GEO-CUR}	10 ⁵	0.182	0.003	0.185	0.001	0.192	0.002	0.121	0.003
IBR _{GEO-CUR}	10 ⁶	0.182	0.003	0.185	0.001	0.192	0.003	0.121	0.002
IBR _{GEO-LGM}	Barrier	0.184	0.001	0.187	0.001	0.190	0.003	0.106	0.003
IBR _{GEO-LGM}	10	0.239	0.001	0.065	0.108	0.249	0.001	0.093	0.002

IBR _{GEO-}	10 ²	0.286	0.001	0.166	0.004	0.293	0.001	0.136	0.010
LGM									
IBR _{GEO-}	10 ³	0.205	0.001	0.190	0.001	0.212	0.001	0.114	0.005
LGM									
IBR _{GEO-}	10 ⁴	0.186	0.003	0.188	0.001	0.192	0.004	0.107	0.002
LGM									
IBR _{GEO-}	10 ⁵	0.184	0.002	0.187	0.001	0.190	0.003	0.106	0.004
LGM									
IBR _{GEO-}	10 ⁶	0.184	0.001	0.187	0.001	0.190	0.003	0.106	0.005
LGM									
IBR _{GEO-}	Barrier	0.260	0.001	0.102	0.033	0.262	0.001	0.105	0.001
MES									
IBR _{GEO-}	10	0.237	0.001	0.069	0.094	0.243	0.001	0.087	0.001
MES									
IBR _{GEO-}	10 ²	0.253	0.001	0.092	0.052	0.256	0.001	0.100	0.002
MES									
IBR _{GEO-}	10 ³	0.259	0.001	0.101	0.046	0.262	0.001	0.104	0.003
MES									
IBR _{GEO-}	10 ⁴	0.260	0.001	0.102	0.040	0.262	0.001	0.105	0.001
MES									
IBR _{GEO-}	10 ⁵	0.260	0.001	0.102	0.043	0.262	0.001	0.105	0.001
MES									
IBR _{GEO-}	10 ⁶	0.260	0.001	0.102	0.032	0.262	0.001	0.105	0.001
MES									
IBR _{CLI-CUR}	Barrier	0.195	0.001	0.189	0.001	0.203	0.001	0.119	0.001
IBR _{CLI-CUR}	10	0.246	0.001	0.062	0.001	0.253	0.001	0.078	0.001
IBR _{CLI-CUR}	10 ²	0.272	0.001	0.165	0.001	0.277	0.001	0.119	0.001
IBR _{CLI-CUR}	10 ³	0.209	0.001	0.188	0.001	0.216	0.001	0.117	0.001
IBR _{CLI-CUR}	10 ⁴	0.197	0.001	0.189	0.001	0.204	0.001	0.119	0.001
IBR _{CLI-CUR}	10 ⁵	0.195	0.001	0.189	0.001	0.204	0.001	0.119	0.001
IBR _{CLI-CUR}	10 ⁶	0.195	0.001	0.189	0.001	0.203	0.001	0.119	0.001
IBR _{CLI-LGM}	Barrier	0.192	0.001	0.187	0.001	0.198	0.001	0.102	0.001
IBR _{CLI-LGM}	10	0.239	0.001	0.081	0.001	0.249	0.001	0.105	0.001
IBR _{CLI-LGM}	10 ²	0.288	0.001	0.180	0.001	0.301	0.001	0.141	0.001
IBR _{CLI-LGM}	10 ³	0.210	0.001	0.191	0.001	0.219	0.001	0.103	0.001
IBR _{CLI-LGM}	10 ⁴	0.194	0.001	0.187	0.001	0.200	0.001	0.103	0.001
IBR _{CLI-LGM}	10 ⁵	0.192	0.001	0.187	0.001	0.198	0.001	0.102	0.001
IBR _{CLI-LGM}	10 ⁶	0.192	0.001	0.187	0.001	0.198	0.001	0.102	0.001

FIGURE S1 Number of reads per individual before and after different quality filtering steps by STACKS and PYRAD. The cumulative stacked bars represent the total number of raw reads obtained for each individual. Within each bar, the red color represents the reads that were discarded by STACKS (*process_radtags*) due to low quality, adapter contamination or an ambiguous barcode. Orange color represents the reads that were subsequently discarded by PYRAD (*step 2*) after filtering out reads that did not comply with the quality criteria (reads with >2 sites with a Phred quality score < 20 were discarded). Finally, green color represents the total number of retained reads used to identify homologous loci during the subsequent steps performed in PYRAD. Dashed horizontal black line indicates the average number of reads across all individuals ($n = 140$). We do not show individuals that were discarded from analyses (one individual from PHU and three individuals from MER), as they recovered a very low number of reads (< 10,000). Population codes as Table S1.

FIGURE S2 Divergence times estimated using BPP with different subsets of loci (100, 200, 300 and 400 loci). The topology was fixed using the phylogenetic tree inferred using SVDQUARTETS. Bars on nodes indicate the 95% highest posterior densities (HPD) of the estimated divergence times. Population codes as in Table S1.

Figure 1 in high resolution (PNG. format)

Figure 2 in high resolution (PNG. format)

Figure 3 in high resolution (PNG. format)

Figure 4 in high resolution (PNG. format)