Analyzing publicly available videos about recreational fishing reveals key ecological and social insights: A case study about groupers in the Mediterranean Sea

Valerio Sbragaglia, Salvatore Coco, Ricardo A. Correia, Marta Coll, Robert Arlinghaus

PII: S0048-9697(20)36201-X

DOI: <https://doi.org/10.1016/j.scitotenv.2020.142672>

Reference: STOTEN 142672

To appear in: *Science of the Total Environment*

Received date: 23 June 2020

Revised date: 2 September 2020

Accepted date: 24 September 2020

Please cite this article as: V. Sbragaglia, S. Coco, R.A. Correia, et al., Analyzing publicly available videos about recreational fishing reveals key ecological and social insights: A case study about groupers in the Mediterranean Sea, *Science of the Total Environment* (2020), <https://doi.org/10.1016/j.scitotenv.2020.142672>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Analyzing publicly available videos about recreational fishing reveals key ecological and social insights: a case study about groupers in the Mediterranean Sea

Valerio Sbragaglia^{1,2,*}, Salvatore Coco³, Ricardo A. Correia^{4,5,6,7}, Marta Coll¹, Robert Arlinghaus^{2,8}

¹ Department of Marine Renewable Resources, Institute of Marine Science (ICM-CSIC), Passeig Marítim de la Barceloneta 37-49, 08003, Barcelona, Spain

² Department of Biology and Ecology of Fishes, Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Müggelseedamm 310, 12587 Berlin, Germany

³ School of Biosciences and Veterinary Medicine, University of Camerino, Viale Circonvallazione 93-95, 62024 Matelica, Italy

⁴ Helsinki Lab of Interdisciplinary Conservation Science (HELICS), Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

⁵ Helsinki Institute for Sustainability Science (HELSUS), Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

⁶ DBIO & CESAM – Centre for Environmental and Marine Studies, University of Aveiro, Aveiro, Portugal

⁷ Instituto de Ciências Biológicas e da Saúde, Universidade Federal de Alagoas, Campus A. C. Simões, Avenida Aurival Melo Mota, Tabuleiro dos Martins, Maceió, Alagoas, Brasil

⁸ Division of Integrative Fisheries Management, Department of Crop and Animal Sciences, Faculty of Life Sciences, Humboldt-Universität zu Berlin, Philippstrasse 13, Haus 7, 10115 Berlin, Germany

* Corresponding author email: valeriosbra@gmail.com

ABSTRACT

iEcology and conservation culturomics are two emerging research approaches that rely on digital data for studying ecological patterns and human-nature interactions. We applied data mining of videos published on YouTube related to recreational fishing of four species of groupers (family: Epinephelidae) in Italy between 2011 and 2017 to learn whether digital user-supplied data help uncover key spatio-temporal ecological patterns characteristic of the studied species. Our results support an ontogenetic deepening of the dusky grouper (*Epinephelus marginatus*) as revealed by a positive relationship between body mass and depth of captures declared in spearfishing videos. In addition, the data support a northward expansion of the white grouper (*Epinephelus aeneus*) because the average latitude associated to the catch was found to be positively correlated with the years when the videos were uploaded on YouTube. Furthermore, the georeferenced data about the white grouper filled a knowledge gap in a well-established international occurrence records dataset. The approach presented here could help mitigating data deficiencies and inform about harvesting patterns shown by recreational anglers and spearfishers. Our work illustrates the value of digital data associated with recreational fishing for advancing fish and fisheries research. The approach can be broadened to larger spatial and temporal scales, and to different species, contributing to a better understanding of macroecological patterns, assessment and conservation of exploited species, and monitoring of recreational fisheries.

KEYWORDS: depth refuge hypothesis, ontogenetic deepening, meridionalization, northward expansion, angling, spearfishing

INTRODUCTION

Scientists increasingly recognize the value of data gathered from non-traditional user supplied sources, such as fishers' local ecological knowledge, as a complement to scientific data for advancing marine ecology, resource conservation and management (e.g., Azzurro et al., 2019; Sáenz-Arroyo et al., 2005). In recent decades, many aspects of human culture, knowledge, and social interactions are being regularly recorded online in digital format (Reed, 2018). This has fostered the emergence of new research areas: iEcology (Jaric et al., 2020) and conservation culturomics (Ladle et al., 2016). Although both research areas applied quantitative analysis of large bodies of digital data, iEcology aims to characterize ecological patterns and processes (e.g., species occurrences, distributional range shifts; Jaric et al., 2020) using data generated for other purposes, while conservation culturomics aims to characterize and understand contemporary problems in conservation by looking at them from the perspective of human-nature interactions (e.g., attitudes of stakeholders, human behavior in the context of resource exploitation; Ladle et al., 2016). The systematic analysis of digital data in both research areas constitutes a promising research approach, which can also contribute to fisheries science and may allow examining specific ecological hypotheses.

iEcology and conservation culturomics are particularly interesting in the context of recreational fisheries because of the many people that are involved in this activity, of which a sizable fraction uses digital platforms on the internet. New monitoring and assessment methods tailored towards recreational fishing have been developed to collect and analyze data from digital platforms (e.g., Carter et al., 2015; Martin et al., 2014; Monkman et al., 2018b; Sbragaglia et al., 2019a; Venturelli et al., 2017). For example, digital information of catches by recreational fishers constitutes a

widespread spatio-temporal network of samples that - if appropriately analyzed - can provide an unprecedented body of information, especially for aquatic environments where sampling is constrained across time and space. In this context, data mining on YouTube is of particular interest because YouTube represents a worldwide dynamic cultural system (Burgess and Green, 2018), which is embedded into the recreational fishers' culture (Sbragaglia et al., 2019a). Recreational fishers often share videos of their catches or memorable fishing trips together with other important information such as the mass of the fish, the depth of capture, and the place of capture. Mining data about recreational fishing from YouTube can foster new opportunities for comparative studies among recreational fishing modalities, such as angling and spearfishing (Sbragaglia et al., 2019a), which can support marine science, resource conservation and management.

Groupers (family: Epinephelidae) are iconic and emblematic top predators of coral and rocky reefs around the world (Hackradt et al., 2014; Heemstra, 1993; Sadovy de Mitcheson et al., 2013). Groupers can have a keystone role from an ecological (Libralato et al., 2006; Valls et al., 2015) and cultural (Garibaldi and Turner, 2004) perspective. Several species of groupers are exploited in commercial (Sadovy de Mitcheson et al., 2013) and recreational (Giglio et al., 2017) fisheries. Simultaneously, groupers are also charismatic and important species for recreational divers (Giglio et al., 2015; Gill et al., 2015). Since 1998, the International Union for Conservation of Nature (IUCN) established the grouper and wrasse specialist group due to growing concern for these species' conservation status. The global results of the IUCN Red List assessment indicated that among 163 grouper species evaluated, 42 species show a risk of extinction in the wild or were Near Threatened (NT), and 50 species lacked

sufficient data to perform an accurate assessment (i.e., they were Data Deficient, DD; Luiz et al., 2016; Sadovy de Mitcheson et al., 2013). Poor management of grouper fisheries can have substantial ecological, social and economic implications for coastal communities worldwide (Luiz et al., 2016). While previous studies have indicated that fisher's knowledge can contribute to assessment of species at risk, such as groupers (e.g., Sáenz-Arroyo et al., 2005), the use of digital data for similar purposes remains underexplored.

Here, we focus on the analysis of YouTube videos featuring recreational fishing of four target species of groupers in Italy as a case study for the Mediterranean Sea, which is one of the areas of the world where several Data Deficient grouper species are predicted to be threatened (Luiz et al., 2016). We explore the potential application of an iEcology approach to characterize spatio-temporal ecological patterns, mitigate data deficiencies, and inform about recreational harvest patterns. The first target species is the dusky grouper (*Epinephelus marginatus*; see also Table 1), an emblematic species for conservation in the Mediterranean Sea and a common target by many recreational fishing modalities, especially spearfishing (e.g., Dedeu et al., 2019). It has been widely accepted that selective spearfishing pressure on large individuals in shallow waters is the major driver of the ontogenetic deepening (i.e., the pattern that older and larger fish are found in deeper waters compared with smaller and younger individuals that remain shallower) of the dusky grouper (García-Rubies et al., 2013; Harmelin and Harmelin-Vivien, 1999; Reñones et al., 1999). Ontogenetic deepening can occur for natural reasons of habitat preferences or may also be triggered by size-selective fishing (Audzijonyte and Pecl, 2018; Frank et al., 2018). Specifically, the “depth refuge” hypothesis predicts that larger individuals - the main

target of spearfishers - will search for protection in deeper water where spearfishers cannot operate (Lindfield et al., 2014). However, the role of harvesting in driving ontogenetic deepening in exploited stocks is still controversially debated (Audzijonyte and Pecl, 2018). This macroecological pattern – also known as Heincke’s law (Cushing, 1975; Heincke, 1913) - has been widely described, but whether it is entirely caused by fishing exploitation, climate change, life-history adaptations, habitat preferences or a combination of all these factors is still largely unknown (Audzijonyte and Pecl, 2018; Baudron et al., 2019; Frank et al., 2018; Frank et al., 2019). If spearfishing harvesting is the most important driver of the ontogenetic deepening of the dusky grouper, videos posted on YouTube should document a positive relationship between depth and body mass of the catch associated with an increasing depth of catch across years. Such time-dependent relationships would strongly suggest an arms race between recreational spearfishers and groupers characterized by larger individuals seeking refuge from spearfishers in deeper water, and spearfishers systematically increasing their operating depth for chasing them.

The second target species of relevance in the Mediterranean is the white grouper (*Epinephelus aeneus*; see also Table 1), which is mainly targeted by recreational angling, especially trolling (Giovos et al., 2018), rather than spearfishing. The white grouper was considered to be absent in central Italy, but a specimen has been captured in January 2014 off Monaco (Pollard et al., 2018), and two other specimens have been recently caught off Corsica, France (Ruitort, 2012). Moreover, the species is also expanding northward in the Atlantic Ocean (Bañón et al., 2020). These records, together with other fishery-dependent data and anecdotal reports from recreational fishers, suggest that the species may be extending its spatial

distribution to the northern parts of the Mediterranean Sea (ĐoĐo et al., 2016; Dulčić et al., 2006; Glamuzina et al., 2000). If that is the case, records posted on YouTube by recreational fishers should document this pattern.

The last two target species studied here are the goldblotch grouper (*Epinephelus costae*) and the dogtooth grouper (*Epinephelus caninus*). The goldblotch grouper is mainly targeted by recreational trolling and spearfishing (Giovos et al., 2018), while the dogtooth grouper, which lives in deeper water than the other species studied here (Table 1), is mainly targeted by anglers due to the depth limitation of spearfishing. Both species were declared “Data Deficient” in the last Mediterranean IUCN report (Francour and Pollard, 2018a; Francour and Pollard, 2018b; Table 1), therefore we lack knowledge on their population trend data and, furthermore, their importance for recreational fishing in the region remain uncertain.

We first characterized the annual rhythms in the upload patterns of videos by recreational anglers and spearfishers, which could indicate differences in catchability of groupers throughout the year as previously suggested for recreational fishing of the common dentex, *Dentex dentex* (Sbragaglia et al., 2019a). Moreover, we documented the declared mass and depth of the groupers showed in the videos to reveal possible differences in the harvesting patterns between recreational anglers and recreational spearfishers (Sbragaglia et al., 2019a). Next, we tackled the following ecological questions:

- (i) Does the declared mass of the dusky grouper increase with the depth in the spearfishing videos supporting an ontogenetic deepening?
- (ii) Does the declared depth of the dusky grouper increase across years in the spearfishing videos supporting an arms race between groupers and spearfishers?

(iii) Does the geographical pattern of the location declared in the videos support a northward expansion of the white grouper?

MATERIALS AND METHODS

Ethical aspects

The data we mined from YouTube are publicly available. However, we followed the framework presented by Monkman et al. (2018a) on the ethics of using social media in fisheries research and the recent guidelines presented by Di Minin E. et al. (*in press*) for ensuring data privacy concerns are fully considered and the European Union's (EU) General Data Protection Regulation (GDPR) is complied with when using social media data. Specifically, we minimized the data by discarding all but required information and pseudonymised them by replacing IDs (e.g., channel title, channel ID). Finally, we kept all data related to personal information in one dataset, while the rest of data presented in the paper were stored in another dataset.

Study system and fishing forms examined

We explored recreational fishing of groupers in Italy. We mined data from 2011 to 2017 by using a systematic analysis to collect quantitative and semi-quantitative data on harvesting patterns from YouTube videos (Correia et al., *in press*). We automatically retrieved the metadata of videos published concerning the species of interest and sorted them into two groups: one related to captures by recreational angling and the other one related to captures by recreational spearfishing. Spearfishing was defined as underwater fishing practiced by the exclusive use of free-diving techniques and a

speargun (e.g., Sbragaglia et al., 2018); angling was defined as hook-and-line fishing from either the coastline or from a boat with natural baits or artificial lures.

Data mining and cross check of automatic identification

We collected the data using the YouTube Data API (v3), following the steps reported in a previous study (Sbragaglia et al., 2019a). First, we extracted the data from YouTube's API in October 2018 using one single keyword represented by the common name of groupers in Italian ("*cernia*"). Although the use of both vernacular and scientific names is suggested (Correia et al., 2018), the use of the common name "*cernia*" allows identifying videos related to the target species of our study: dusky, white, goldblotch, and dogtooth groupers. We compiled a raw dataset with the title and descriptions of videos.

In a second step, we automatically searched the title and description of each video for specific keywords that were already used in a previous study (Sbragaglia et al., 2019a). The keywords were subdivided into two groups with the aim to sort the videos regarding recreational angling and recreational spearfishing. Finally, we stored the results in a dataset that was subsequently manually cross checked.

We excluded the videos that were: (i) not related to the target species; (ii) not showing the catch of the target species (i.e., catch and release or not shooting while spearfishing); (iii) not related to the target country; and (iv) duplicates of previously published videos. Then, we applied a manual cross check of the automatic classification to identify the occurrence of false negatives (i.e., target videos previously not recognized following the keywords), false positives (i.e., videos erroneously attributed to one of the two groups) and mismatched categorizations (i.e., videos

erroneously attributed to one fisher group instead of the other). Finally, during the manual cross check, we explored the content of videos and annotated the species, the mass of the fish, and the depth at which the fish was captured by screening the title or description of the videos as well as the video footage itself. We also annotated the location of the videos related to the white grouper. We run all the analyses related to data mining in R (<https://www.r-project.org/>; version 3.5.0) with the additional package “*jsonlite*” (Ooms, 2014), “*lubridate*” (Grolemund and Wickham, 2011), and “*curl*” (<https://cran.r-project.org/web/packages/curl/index.html>).

Data analysis

We estimated annual periodicity of the upload patterns of videos for each of the four target species and each group (angling and spearfishing) by using RAIN (rhythmicity analysis incorporating nonparametric methods). This method is a robust non-parametric method for the detection of rhythms in data that can detect arbitrary oscillations (Thaben and Westermark, 2014). We estimated differences in declared mass and depth between angling and spearfishing using: (i) a two-sample unpaired Welch’s t-test (a parametric test that better controls Type 1 error when the assumption of homogeneity of variance is not met; Delacre et al., 2017), followed by the estimation of Cohen’s *d* effect sizes (Cohen, 1988); or (ii) the non-parametric Mann–Whitney *U* test followed by the estimation of Vargha and Delaney’s *A* effect sizes (Vargha and Delaney, 2000), which accounted for cases where the distribution of response variable did not allow the use of a parametric statistical test. Next, the relationship between declared depth and body mass for spearfishing videos related to the dusky grouper (question i) was estimated using a linear regression model; while

the correlation between the declared depth and years for the spearfishing videos related to the dusky grouper (question *ii*) as well as the correlation between latitude and years for the white grouper (question *iii*) were estimated using the Kendall's rank correlation coefficient (r_t) because data did not meet the assumptions to use parametric statistical tests. In the cases in which we used parametric tests, response variables were transformed by finding the exponent (λ), as a power transformation producing a normally distributed response variable. We assessed model fits by checking the plot of the residuals vs. the fitted values. In all cases we used a 95% confidence interval. We run all the analyses related to data mining in R (<https://www.r-project.org/>; version 3.5.0) with the additional package "*rain*" (Thaben and Westermarck, 2014); "*rcompanion*" (<https://CRAN.R-project.org/package=rcompanion>); "*lsr*" (<https://cran.r-project.org/web/packages/lsr>); "*effsize*" (<https://cran.r-project.org/web/packages/effsize>).

RESULTS

We identified a total of 2097 videos published between 2011 and 2017: 1714 (82%) videos were related to spearfishing and 383 (18%) related to angling. We sorted the videos identified among the four target species: 1670 related to the dusky grouper (80%); 262 related to the white grouper (12%); 100 related to goldblotch grouper (6%); and 32 related to dogtooth grouper (1%). We were not able to assign a species to the groupers fished in 33 videos (1%; Table 2).

Dusky grouper

Among the 1670 videos identified for the dusky grouper (Fig. 1), 171 were related to angling (10%), while 1499 were related to spearfishing (90%). The upload of spearfishing videos indicated a significant ($p < 0.001$) annual periodicity with a peak in August (Fig. 2A), while angling videos did not show significant annual periodicity ($p = 0.976$). We retrieved the body mass declared in a total of 460 videos (34 for angling and 426 for spearfishing). The declared body mass was significantly ($t_{41.97} = 6.46$; $p < 0.001$; $d = 0.94$) higher in angling videos (mean = 13.6 kg; SD = 6.1 kg) than in spearfishing ones (mean = 8.4 kg; SD = 5.3 kg; Fig. 2B). We retrieved the declared depth of dusky grouper capture in a total of 155 videos (6 for angling and 149 for spearfishing). The declared depth showed that values for angling videos were significantly ($U = 891$; $p < 0.001$; $A = 0.99$) higher (median = 67 m) than those for spearfishing videos (median = 24 m; Fig. 2C). Finally, we retrieved a total of 75 videos of spearfishing related to the dusky grouper where we were able to assess both depth and body mass of the specimen shown in the video. We found a significant positive correlation between declared depth and body mass ($R = 0.17$; $F_{1,73} = 15.3$; $p < 0.001$), indicating that larger dusky grouper captures tended to occur in deeper sites (Fig. 2D). The declared depth for the spearfishing videos was not significantly correlated with years ($r_t = -0.08$; $p = 0.146$; $N = 149$; Fig. 2E), and thus we did not find evidence of a deepening of spearfishing efforts in recent years.

White grouper

Among the 262 videos identified for the white grouper, 136 were related to angling (52%), while 126 were related to spearfishing (48%). The upload patterns of both groups did not indicate significant annual periodicity (angling: $p = 0.988$; spearfishing: p

= 0.999; Fig. 3A). We retrieved the declared body mass of the white grouper in a total of 67 videos (42 for angling and 25 for spearfishing). The declared body mass was significantly ($t_{42.69} = 4.66$; $p < 0.001$; $d = 1.24$) higher in angling videos (mean = 11.3 kg; SD = 5.5 kg) than in spearfishing ones (mean = 6.6 kg; SD = 4.2 kg; Fig. 3B). We retrieved the declared depth of capture for white groupers in a total of 21 videos (6 for angling and 15 for spearfishing). The declared depths in angling videos were significantly ($U = 80$; $p < 0.01$; $A = 0.89$) higher (median = 44 m) than those for spearfishing videos (median = 29 m; Fig. 3C). Finally, we retrieved the location of fishing effort in a total of 64 videos (44 for angling and 20 for spearfishing) and found a significant positive correlation between latitude and years ($r_t = 0.25$; $p < 0.01$), which suggests a northward shift in fishing location, from 2011 to 2017 (Fig. 3D). The location of the videos was mostly from the South of Italy (Fig. 3E).

Goldblotch grouper

Among the 100 videos identified for the goldblotch grouper, 25 were related to angling (25%), while 75 to spearfishing (75%). The upload patterns of both groups did not indicate significant annual periodicity (angling: $p = 1$; spearfishing: $p = 0.999$; Fig. 4A). We retrieved the body mass declared for the goldblotch grouper in a total of 15 videos (1 for angling and 14 for spearfishing). Only one angling video declared the body mass for this species – 11 kg – but this mass is larger than any among those retrieved for spearfishing videos (mean = 4.6 kg; SD = 1.7 kg; Fig. 4B). Finally, we retrieved the depth declared for the capture of the goldblotch grouper in a total of 12 videos (2 for angling and 10 for spearfishing). The depth declared in the two videos retrieved for angling

was 32 and 49 m, which is within the range of depth declared for spearfishing videos (mean = 35 m; SD = 11 m; Fig. 4C).

Dogtooth grouper

Among the 32 videos identified for the dogtooth grouper, 31 related to angling (97%) and only 1 to spearfishing (3%). The upload pattern across years did not show significant annual periodicity (angling: $p = 1$; Fig. 5A). We retrieved the body mass declared of the dogtooth grouper in a total of 12 videos (all of them for angling). The declared mass ranged from 15 to 60 kg (mean = 36 kg; SD = 14 kg; Fig. 5B). Finally, we retrieved the depth declared for the capture of the dogtooth grouper in a total of 5 videos (all of them for angling). The depth ranged from 61 to 443 m (Fig. 5C).

DISCUSSION

We show how data mining of recreational fishing on YouTube can provide large volumes of data that are useful to characterize spatio-temporal ecological patterns, mitigate data deficiencies and inform about recreational harvesting patterns. Our results support two ecological hypotheses possibly related to human disturbance and climate change effects on groupers in the Mediterranean Sea. First, our results showed that the declared body mass of the dusky grouper was positively correlated to the declared depth in recreational spearfishing videos; this may support the depth refuge hypothesis (Lindfield et al., 2014), but could also be explained by a natural ontogenetic deepening of the species where larger fish use deeper habitats (Audzijonyte and Pecl, 2018). Second, our work contributes to a large body of studies documenting a poleward expansion of fish species in response to climate change (Burrows et al.,

2011; Cheung et al., 2013) by showing a positive correlation between latitude and the years in those videos in which the capture of white groupers have been documented. Our results also confirm the results of a previous study on recreational fishing of the common dentex on YouTube (Sbragaglia et al., 2019a), by highlighting different harvesting patterns of groupers between recreational anglers and recreational spearfishers.

The depth refuge hypothesis in the dusky grouper

We found support for an ontogenetic deepening of the dusky grouper by showing that the depth and the mass of the captured dusky grouper were positively related in spearfishing videos. Such pattern can be interpreted according to the depth refuge hypothesis, which was supported using fishery-independent data (i.e., baited remote underwater stereo-video systems) for SCUBA spearfishing on coral reef fish populations in the southern Mariana Islands (Lindfield et al., 2014). In particular, the authors found greater lengths of scarines and acanthurids in deeper waters (Lindfield et al., 2014). However, the declared depth in spearfishing videos of the dusky grouper did not increase over years, which suggests that spearfishers are not increasing their operating depth over the time period studied here (2011-2017). If fishing pressure for larger fish would be the most important driving force of the observed pattern, we might expect that the information associated to the videos would indicate a deeper fishing effort in response to ontogenetic deepening. However, this was not the case, and thus we have no support that the ontogenetic deepening is caused by size-selective harvesting over the time frame analyzed here.

There are three additional mechanisms that could explain the ontogenetic deepening of the dusky grouper that are not directly related to spearfishing harvesting: (i) small- and medium-sized individuals could have more pronounced preference for shallow waters than large individuals (e.g., Harmelin and Harmelin-Vivien, 1999); (ii) small- and medium-sized individuals could have a higher mortality rate in deeper water (Audzijonyte and Pecl, 2018), which in part can be caused by cannibalism (Concini et al., 2015), and (iii) larger individual could move to deeper and cooler waters due to a growth/metabolism trade-off, which predicts that larger individuals allocate more energy to reproduction than smaller ones and therefore might move to cooler waters to decrease their metabolic costs and thereby increase their fitness (Audzijonyte and Pecl, 2018; Frank et al., 2018; Macpherson and Duarte, 1991). Considering that the surface water temperature of the north-western Mediterranean Sea is increasing (Bianchi et al., 2018; Lejeusne et al., 2010), the growth/metabolism trade-off could indeed be one of the drivers of the ontogenetic deepening of the dusky grouper.

In this context, selective harvesting of larger individuals - a common scenario in many fisheries worldwide - could act at different levels on the ontogenetic deepening. First, as highlighted by (Frank et al., 2018), intensive and selective harvesting of larger individuals may reduce the number of older and larger individuals from highly-exploited shallow water compared to less-exploited deeper waters. Second, the evolutionary effects associated to selective harvesting of larger individuals could confound life-history effects (Frank et al., 2018). For example, fisheries-induced evolution of life history could affect the growth/metabolism trade-off by attenuating the ontogenetic deepening in response to climate change. Indeed, fisheries-induced

evolution may foster the evolution of a fast life history that is characterized by elevated reproductive investment, reduced age and size at maturation and reduced post maturation growth and longevity (Heino et al., 2015). Therefore, individuals could move to deeper and cooler waters at a smaller size and age because of anticipated maturation and increase reproductive investment, attenuating the ontogenetic deepening. A further mechanism associated to intense and selective-harvesting of larger individuals is related to behavior, but escaped the attention of the recent discussion on this topic (Audzijonyte and Pecl, 2018; Frank et al., 2018). Experimental selective harvesting of larger individuals has suggested that an evolutionary decrease of risk-taking behavior could happen (Sbragaglia et al., 2019b), which in turn could increase the prevalence of shy fish (i.e., those less prone to take risks) in deep waters where there is less anthropogenic disturbance compared to shallow waters (deep reef refugia concept; Bongaerts et al., 2010). Such prediction is not only expected for spearfishing harvesting, which has been shown to increase fish wariness (Januchowski-Hartley et al., 2011; Samis et al., 2019; Sbragaglia et al., 2018), but also in other recreational and commercial fisheries triggering a “timidity syndrome” in response to intensive harvesting (Arninghaus et al., 2017 and references therein). Moreover, it must be considered that the average depth of spearfishing in free diving is reported to be around 18-25 m (FIPSAS, 2002). Although in recent decades deep spearfishing at depth of 40-45 m has become more common, only a minority of spearfishers can operate at such extreme depth, for example only 10% of spearfishers operate at more than 25 m in the canary islands (Martín-Sosa, 2019). In summary, our results suggest that the ontogenetic deepening of the dusky grouper could not solely determine by spearfishing harvesting and the mechanisms explained above could contribute in

driving it. The relative contribution of the different mechanisms is still unknown and represents an intriguing question for future research in many fisheries worldwide (Audzijonyte and Pecl, 2018).

Northward expansion in the white grouper

Our results support a northward expansion of the white grouper during the time period that we studied (2011-2017). We found a positive relationship between latitude at which the white grouper catches were declared in the videos and the years of study. Our results agreed with previous published records using different data sources that documented an ongoing northward expansion of the white grouper. For example, two specimens of white grouper were captured in the northern Adriatic Sea, which represented the northernmost occurrence of the white grouper in the Mediterranean Sea (ĐoĐo et al., 2016; Dulčić et al., 2006). Our study shows a retrospective quantitative increase of the latitude across years related to the south Tyrrhenian and Ionian Sea. Although our study is representative of only those recreational fishers sharing their catches on YouTube (possible associated biases are discussed below), it demonstrates that recreational fishers represent a widespread network of observers of ongoing changes of marine biota that can be quantitatively measured using data mining on social media. Indeed, the georeferenced records reported here about the white grouper filled a geographical gap in the occurrence records dataset of the Global Biodiversity Information Facility (GBIF; Fig. S1), which is an international network and research infrastructure funded by the world's governments and aimed at providing open access biodiversity data. As such, we argue that recreational fishers could represent an important monitoring tool for climate change effects on marine biota as

recently demonstrated by using their local ecological knowledge in the context of meridionalization of the Mediterranean Sea (Sbragaglia et al., 2020).

Mitigation of data deficiency in goldblotch and dogtooth groupers

The IUCN status of Data Deficiency refers to lack of knowledge on distribution or population trend and implies that more information is required for the assessment of extinction risk (IUCN, 2012). Our study showed that publicly available videos about recreational fishing can provide additional and retrospective information and therefore can be a valuable additional tool for future assessment of species that are Data Deficient. Moreover, our results showed that the goldblotch grouper seems to be targeted preferentially by spearfishers compared to anglers. Information on Mediterranean recreational fishing of goldblotch grouper could be scattered especially because it can be confounded with juveniles of the dogtooth grouper or the mottled grouper, *Mycteroperca rubra* (Cotil et al., 2004). A systematic analysis of publicly available videos could overcome this limitation. As regarding the dogtooth grouper, it is preferentially targeted by recreational anglers because of clear depth-limitation of spearfishing. We documented a maximum declared mass that is close to the published records (Francour and Pollard, 2018a; Morales-Nin et al., 2005). Although recreational anglers may probably inflate the mass of the fish declared in the videos for increasing social engagement on YouTube, as previously suggested for recreational spearfishers (Sbragaglia et al., 2019a), our study suggests that large-sized species can be vulnerable to deep-sea angling. In this context, harvesting of dogtooth grouper is of special interest for Mediterranean deep-sea ecosystems because it is the grouper species with the deepest depth range and the maximum size and age (up to 60 kg and age of more

than 50 years; Morales-Nin et al., 2005). The dogtooth grouper could be considered a keystone species (sensu Valls et al., 2015), and harvesting of large-sized individuals from the deep sea could have strong cascading effects on vulnerable ecosystems inhabited by this species such as deep coral reefs and seamounts (Canese and Bava, 2015; Francour and Pollard, 2018a). In general, harvesting of keystone predators may remove top-down control on ecosystems triggering trophic cascades (Frank et al., 2005) and alternative stable states (Palkovacs et al., 2018). This could be particularly important in the assessment of IUCN status of the dogtooth grouper because habitat alteration can be used to indirectly assign a specific threat category (IUCN, 2012).

Annual rhythms of capture

We only detected annual rhythms in the upload pattern of the spearfishing videos related to the dusky grouper but not to the other three species. These results may indicate – assuming that the videos are uploaded within few days after the fishing trip – a seasonal change in the catchability of the dusky grouper species with peaks in August. This may be related to a seasonal depth migration of the species that becomes more accessible to spearfishers in shallow waters during the summer as suggested in a similar study with the common dentex, *Dentex dentex* (Sbragaglia et al., 2019a). Indeed, in the Western Mediterranean Sea the dusky grouper performs seasonal spawning migration moving to shallow water during the summer as documented by direct observations (Zabala et al., 1997) and telemetry data (Koeck et al., 2014). Such seasonal migrations seem to usually occur in August (Zabala et al., 1997), which is also the month in which we detected the peak of videos upload, supporting our interpretation. Additionally, spearfishers may show seasonal patterns of fishing effort

(e.g., peaks in August could be associated to vacation times and more free time to go fishing) or annual periodicity in the operating depth. In the latter case, Northern Mediterranean spearfishers wear a thinner wetsuit in the summer than in winter to cope with changes of water temperature, which implies the use of less weight during the summer that subsequently increase the diving performance and hence the operating depth. Our results are in contrast to what was shown in a recent study using local ecological knowledge of eastern Mediterranean recreational fishers, where the fishing pressure on groupers was greater in spring and autumn compared to the other seasons (Mavruk et al., 2018). Such discrepancies could be related to the small sample size used in that study ($N = 10$), to real differences between the two areas or to complementary data provided by the two methods (local ecological knowledge and data mining on YouTube; see below for further discussion).

Declared body mass and depth

We document that the declared body mass and depth in angling videos is greater than in spearfishing videos for both the dusky and white groupers. These results are in accordance to what was previously shown by mining data on recreational fishing of common dentex (Sbragaglia et al., 2019a). In particular, catching a grouper (or a common dentex) with spearfishing techniques requires more skills and usually long free diving at considerable depths and elevated personal investment of energy relative to angling (Sbragaglia et al., 2019a). Therefore, spearfishers are probably more stimulated to post videos than anglers, including those showing the catch of relatively small specimens. This could explain the differences in number of videos uploaded and the differences in declared mass we observed in our study.

Our results partially disagree with local ecological knowledge of fishers in eastern Mediterranean Sea, which indicated that the maximum size of the dusky grouper was smaller for anglers compared to other fishing modalities including spearfishing (Mavruk et al., 2018), while the maximum declared body mass we recorded was slightly larger for anglers (30 kg) than spearfishers (27.5 kg). The same study reported the opposite for the white grouper (i.e., smaller maximum size for spearfishing than other fishing techniques, including angling; Mavruk et al., 2018), which agreed with our results (angling max declared mass = 28 kg; spearfishers maximum declared body mass = 17 kg). Finally, Mavruk et al. (2018) also reported that the average size of the white grouper was smaller for angling compared to other fishing techniques, including spearfishing, which disagreed with our results where we recorded a greater average body mass for angling (11.3 kg) with respect to spearfishing (6.6 kg). As discussed above in the context of annual rhythmicity, the observed discrepancies with the study by Mavruk et al. (2018) could be related to several factors including complementary data provided by the two methods. For example, Mavruk et al. (2018) reported that recreational anglers and recreational spearfishers usually fish in shallow waters. Although this is true for spearfishing for obvious depth-related operating constraints, our data show that recreational anglers can operate at greater depth.

Challenges and limitations

Our results provide novel insights into ecological aspects and monitoring of groupers targeted by recreational fishing in the Mediterranean Sea, but limitations must be considered. First of all, the availability and representativeness of YouTube digital data

depends on accessibility to the internet and video recording technology, which are strongly influenced by demography, socio-cultural and economic aspects (Correia et al., *in press*). Moreover, YouTube is a dynamic cultural system where algorithms for data management often change and users can modify or delete information. Importantly, mining data from YouTube is unlikely to be representative of the whole population of recreational fishers and those fishers posting videos on YouTube probably represent the most avid individuals (i.e., those recreational fishers that spend much of their time fishing; Griffiths et al., 2013; Rocklin et al., 2014) as more specialized anglers usually have larger degree of media interactions (Ditton et al., 1992). Therefore, it is very likely that our data represent catches of the most efficient and skilled recreational fishers.

Furthermore, we extracted data from YouTube's API using one single keyword represented by the common name of groupers in Italian ("*cernia*"). This could have limited the amount of videos that are showing recreational catches of groupers without explicitly using the name in the title, description and tags of the videos. For example, several recreational fishers post compilations of catches that are difficult to identify with the methodology applied here, but could contain valuable data regarding groupers. This could have contributed to the overrepresentation of data related to the dusky grouper. In summary, the results presented here are not an absolute representation of recreational fisheries of groupers, but they represent a complementary contribution to existing knowledge (see also Fig. S1). Future challenges are represented by making data mining on social media more robust with the use of machine learning approaches (Roll et al., 2018; Toivonen et al., 2019) and by disentangling the proportion of recreational fishers posting their catches on social

media and how much they differ from a representative population of recreational fishers.

Implications

Despite the limitations highlighted above, the approach we presented in our study is expected to strongly contribute to conservation and management of aquatic ecosystems in the coming years (Jaric et al., 2020; Ladle et al., 2016). One of the main reasons is that recreational fishers are more than 5 times the number of commercial capture fishers (Arlinghaus et al., 2019; FAO, 2018); for example, it has been estimated that Mediterranean marine recreational fishers are at about 8.7 million (Hyder et al., 2018). Therefore, recreational fishers can provide a unique network of detailed spatial and temporal representation of macroecological patterns. In particular the use of YouTube as data source will certainly support these developments; in fact YouTube is the second most visited website in the world (2 billion users) with 1 billion hours of videos played every day in 80 different languages (YouTube, 2020).

Credit author statement

Valerio Sbragaglia: Conceptualization; Formal analysis; Funding acquisition; Investigation; Visualization; Writing - original draft. **Salvatore Coco:** Data curation; Validation; Writing - review & editing. **Ricardo A. Correia:** Methodology; Writing - review & editing. **Marta Coll:** Writing - review & editing. **Robert Arlinghaus:** Writing - review & editing.

Acknowledgments

VS is supported by a “*Juan de la Cierva Incorporación*” research fellowship (IJC2018-035389-I) granted by the Spanish Ministry of Science and Innovation. RC was partly supported by funding from the Helsinki Institute of Sustainability Science (HELSUS) and the University of Helsinki. RA received funding through the European Union (European Maritime and Fisheries Fund) and the State of Mecklenburg-Vorpommern (Germany) (Grant MV-I.18-LM-004, B 730117000069), and the German Federal Ministry of Education and Research (Grants 01LC1826E and 033W046A).

REFERENCES

- Arlinghaus R, Abbott JK, Fenichel EP, Carpenter SR, Hunt LM, Alós J, et al. Opinion: Governing the recreational dimension of global fisheries. *Proceedings of the National Academy of Sciences* 2019; 116: 5209-5213.
- Arlinghaus R, Laskowski KL, Alós J, Klefoth T, Monk CT, Nakayama S, et al. Passive gear-induced timidity syndrome in wild fish populations and its potential ecological and managerial implications. *Fish and Fisheries* 2017; 18: 360-373.
- Audzijonyte A, Pecl GT. Deep impact of fisheries. *Nature Ecology & Evolution* 2018; 2: 1348-1349.
- Azzurro E, Sbragaglia V, Cerri J, Bariche M, Bolognini L, Ben Souissi J, et al. Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological knowledge. *Global Change Biology* 2019; 25: 2779-2792.
- Bañón R, de Carlos A, Alonso-Fernández A, Ramos F, Baldó F. Apparently contradictory routes in the expansion of two fish species in the Eastern Atlantic. *Journal of Fish Biology* 2020; 96: 1051-1054.
- Baudron AR, Pecl G, Gardner C, Fernandes PG, Audzijonyte A. Ontogenetic deepening of Northeast Atlantic fish stocks is not driven by fishing exploitation. *Proceedings of the National Academy of Sciences* 2019; 116: 2397-2402.
- Bianchi CN, Caroli F, Guidetti P, Morri C. Seawater warming at the northern reach for southern species: Gulf of Genoa, NW Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* 2018; 98: 1-12.
- Bongaerts P, Ridgway T, Sampayo EM, Hughes-Guldborg O. Assessing the 'deep reef refugia' hypothesis: focus on Caribbean reefs. *Coral Reefs* 2010; 29: 309-327.
- Burgess J, Green J. YouTube: Online video and participatory culture: John Wiley & Sons, 2018.
- Burrows MT, Schoeman DS, Buckley LB, Moore P, Poloczanska ES, Brander KM, et al. The pace of shifting climate in marine and terrestrial ecosystems. *Science* 2011; 334: 652-655.
- Canese S, Bava S. The decline of top predators in deep coral reefs. 1st Mediterranean symposium on the conservation of dark habitats (Portorož, Slovenia, 31 October 2014), 2015, pp. 67.
- Carter DW, Crosson S, Liese C. Nowcasting intraseasonal recreational fishing harvest with internet search volume. *Plos One* 2015; 10: e0137752.
- Cheung WW, Watson P, Pauly D. Signature of ocean warming in global fisheries catch. *Nature* 2013; 497: 365.
- Cheung WWL, Pitcher TJ, Pauly D. A fuzzy logic expert system to estimate intrinsic extinction vulnerabilities of marine fishes to fishing. *Biological Conservation* 2005; 124: 97-111.
- Cohen J. The effect size index: d. *Statistical power analysis for the behavioral sciences* 1988; 2: 284-288.
- Coll J, Linde M, García-Rubies A, Riera F, Grau AM. Spear fishing in the Balearic Islands (west central Mediterranean): species affected and catch evolution during the period 1975–2001. *Fisheries Research* 2004; 70: 97-111.
- Condini MV, Hoetinghaus DJ, Garcia AM. Trophic ecology of dusky grouper *Epinephelus marginatus* (Actinopterygii, Epinephelidae) in littoral and neritic habitats of southern Brazil as elucidated by stomach contents and stable isotope analyses. *Hydrobiologia* 2015; 743: 109-125.
- Correia RA, Ladle R, Jarić I, Malhado ACM, Mittermeier JC, Roll U, et al. Digital data sources and methods for conservation culturomics. *Conservation Biology in press*.
- Cushing DH. *Marine ecology and fisheries*: CUP Archive, 1975.

- Dedeu AL, Boada J, Gordo A. The first estimates of species compositions of Spanish marine recreational fishing reveal the activity's inner and geographical variability. *Fisheries Research* 2019; 216: 65-73.
- Delacre M, Lakens D, Leys C. Why psychologists should by default use Welch's t-test instead of Student's t-test. *International Review of Social Psychology* 2017; 30.
- Di Minin E., Fink C., Hausmann A., Kremer J., R. K. How to address data privacy concerns when using social media data in conservation science. *Conservation Biology* *in press*.
- Ditton RB, Loomis DK, Choi S. Recreation specialization: Re-conceptualization from a social worlds perspective. *Journal of Leisure Research* 1992; 24: 33-51.
- ĐoĐo Z, Antolovic N, Dulčić J. New record of white grouper *epinephelus aeneus* (osteichthyes: Serranidae) in croatian adriatic waters/nuova segnalazione della cernia bianca *epinephelus aeneus* (osteichthyes: Serranidae) in acque adriatiche croate. *Annales: Series Historia Naturalis*. 26. Scientific and Research Center of the Republic of Slovenia, 2016, pp. 37.
- Dulčić J, Tutman P, Caleta M. Northernmost occurrence of the white grouper, *Epinephelus aeneus* (Perciformes: Serranidae), in the Mediterranean area. *Acta Ichthyologica et Piscatoria* 2006; 1.
- FAO. The State of World Fisheries and Aquaculture (SOFIA) - Meeting the Sustainable Development Goals. Food and Agriculture Organization, Rome, Italy, (p. xiii + 210), 2018.
- FIPSAS. Libro bianco; Le gare di pesca in apnea. Comitato d. Settore Attività Subacquee, 2002.
- Francour P, Pollard DA. *Epinephelus caninus*. The IUCN Red List of Threatened Species 2018: e.T132796A100463319. Downloaded on 14 April 2020. 2018a.
- Francour P, Pollard DA. *Epinephelus costae*. The IUCN Red List of Threatened Species 2018: e.T132762A100464786 Downloaded on 14 April 2020. 2018b.
- Frank KT, Petrie B, Choi JS, Leggett WC. Trophic cascades in a formerly cod-dominated ecosystem. *Science* 2005; 308: 1621-1623.
- Frank KT, Petrie B, Leggett WC, Boyce DG. Exploitation drives an ontogenetic-like deepening in marine fish. *Proceedings of the National Academy of Sciences* 2018; 115: 6422-6427.
- Frank KT, Petrie B, Leggett WC, Boyce DG. Reply to Baudron et al.: Fishing matters: Age-specific deepening is driven by exploitation. *Proceedings of the National Academy of Sciences* 2019; 116: 2393-2394.
- García-Rubies A, Hereu B, Zabala M. Long-term recovery patterns and limited spillover of large predatory fish in a Mediterranean MPA. *Plos One* 2013; 8: e73922.
- Garibaldi A, Turner N. Cultural keystone species implications for ecological conservation and restoration. *Ecology and Society* 2004; 9: 1.
- Giglio VJ, Bender MG, Zupelini C, Ferreira CEL. The end of the line? Rapid depletion of a large-sized grouper through spearfishing in a subtropical marginal reef. *Perspectives in Ecology and Conservation* 2017; 15: 115-118.
- Giglio VJ, Luiz OJ, Schiavetti A. Marine life preferences and perceptions among recreational divers in Brazilian coral reefs. *Tourism Management* 2015; 51: 49-57.
- Gill DA, Schuhmann PW, Oxenford HA. Recreational diver preferences for reef fish attributes: Economic implications of future change. *Ecological Economics* 2015; 111: 48-57.
- Giovas I, Keramidas I, Antoniou C, Deidun A, Font T, Kleitou P, et al. Identifying recreational fisheries in the Mediterranean Sea through social media. *Fisheries Management and Ecology* 2018; 25: 287-295.
- Glamuzina B, Tutman P, Geffen JA, Kožul V, Boško S. First record of white grouper, *Epinephelus aeneus* (Serranidae) in the south eastern Adriatic. *Cybiu* 2000; 24: 306-308.
- Griffiths SP, Zischke MT, Tonks ML, Pepperell JG, Tickell S. Efficacy of novel sampling approaches for surveying specialised recreational fisheries. *Reviews in Fish Biology and Fisheries* 2013; 23: 395-413.

- Grolemund G, Wickham H. Dates and times made easy with lubridate. *Journal of Statistical Software* 2011; 40: 1-25.
- Hackradt CW, García-Charton JA, Harmelin-Vivien M, Pérez-Ruzafa Á, Le Diréach L, Bayle-Sempere J, et al. Response of rocky reef top predators (Serranidae: Epinephelinae) in and around marine protected areas in the western Mediterranean Sea. *Plos One* 2014; 9: e98206.
- Harmelin J-G, Harmelin-Vivien M. A review on habitat, diet and growth of the dusky grouper *Epinephelus marginatus* (Lowe, 1834). *Marine Life* 1999; 9: 11-20.
- Heemstra PC. Groupers of the world (Family Serranidae, Subfamily Epinephelinae). An annotated and illustrated catalogue of the grouper, rockcod, hind, coral grouper and lyretail species known to date. *FAO species catalogue* 1993; 16.
- Heincke F. Investigations on the plaice. General report 1. Rapports et procès verbaux des réunions (International Council for the Exploration of the Sea, Copenhagen) 1913; Vol 17.
- Heino M, Díaz Pauli B, Dieckmann U. Fisheries-induced evolution. *Annual Review of Ecology, Evolution and Systematics* 2015; 46: 461-480.
- Hyder K, Weltersbach MS, Armstrong M, Ferter K, Townhill B, Andersen A, et al. Recreational sea fishing in Europe in a global context-Participation rates, fishing effort, expenditure, and implications for monitoring and assessment. *Fish and Fisheries* 2018; 19: 225-243.
- IUCN. IUCN Red List Categories and Criteria: Version 3.1. Second Edition. Gland, Switzerland and Cambridge, UK: IUCN. iv + 32pp 2012.
- Januchowski-Hartley FA, Graham NAJ, Feary DA, Morove T, Cinner JE. Fear of fishers: Human predation explains behavioral changes in coral reef fishes. *Plos One* 2011; 6: e22761.
- Jaric I, Correia RA, Brook BW, Buettel JC, Courchamp F, Di Minin E, et al. iEcology: Harnessing large online resources to generate ecological insights. *Trends in Ecology and Evolution* 2020; 35: 630-639.
- Koeck B, Pastor J, Saragoni G, Dalias M, Payrol J, Lenfant P. Diel and seasonal movement pattern of the dusky grouper *Epinephelus marginatus* inside a marine reserve. *Marine environmental research* 2014; 91: 38-47.
- Ladle RJ, Correia RA, Do Y, Joo GJ, Machado AC, Proulx R, et al. Conservation culturomics. *Frontiers in Ecology and the Environment* 2016; 14: 269-275.
- Lejeusne C, Chevaldonné P, Pétigot-Martini C, Boudouresque CF, Pérez T. Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends in Ecology & Evolution* 2010; 25: 250-260.
- Libralato S, Christensen V, Pauly D. A method for identifying keystone species in food web models. *Ecological Modelling* 2006; 195: 153-171.
- Lindfield SJ, McIlwain JL, Harvey ES. Depth refuge and the impacts of SCUBA spearfishing on coral reef fishes. *Plos One* 2014; 9: e92628.
- Luiz OJ, Woods RM, Madin EMP, Madin JS. Predicting IUCN extinction risk categories for the world's data deficient groupers (Teleostei: Epinephelidae). *Conservation Letters* 2016; 9: 342-350.
- Macpherson E, Duarte CM. Bathymetric trends in demersal fish size: is there a general relationship? *Marine Ecology Progress Series* 1991; 71: 103-112.
- Martín-Sosa P. Spearfishing in The Canary Islands: is the devil as black as it seems to be? *Scientia insularum* 2019; 2: 9-36.
- Martin DR, Chizinski CJ, Eskridge KM, Pope KL. Using posts to an online social network to assess fishing effort. *Fisheries Research* 2014; 157: 24-27.
- Mavruk S, Saygu İ, Bengil F, Alan V, Azzurro E. Grouper fishery in the Northeastern Mediterranean: An assessment based on interviews on resource users. *Marine Policy* 2018; 87: 141-148.
- Monkman GG, Kaiser M, Hyder K. The ethics of using social media in fisheries research. *Reviews in Fisheries Science & Aquaculture* 2018a; 26: 235-242.

- Monkman GG, Kaiser MJ, Hyder K. Text and data mining of social media to map wildlife recreation activity. *Biological Conservation* 2018b; 228: 89-99.
- Morales-Nin B, Bauza D, Grau AM. Otolith characteristics and age estimation of an exceptionally old dogtooth grouper (*Epinephelus caninus*) captured off Majorca Island (western Mediterranean). *Cybium* 2005; 29: 100-102.
- Ooms J. The jsonlite package: A practical and consistent mapping between json data and r objects. *arXiv preprint arXiv:1403.2805* 2014.
- Palkovacs EP, Moritsch MM, Contolini GM, Pelletier F. Ecology of harvest-driven trait changes and implications for ecosystem management. *Frontiers in Ecology and the Environment* 2018; 16: 20-28.
- Pollard DA, Francour P, Fennessy S. *Epinephelus aeneus*. The IUCN Red List of Threatened Species 2018 2018; e.T132722A100459597. Downloaded on 24 July 2019.
- Reed TV. Digitized lives: Culture, power and social change in the internet era: London, Routledge, 2018.
- Reñones O, Goñi R, Pozo M, Deudero S, Moranta J. Effects of protection on the demographic structure and abundance of *Epinephelus marginatus* (Lowe, 1834). Evidence from Cabrera Archipelago National Park (West-central Mediterranean). *Marine Life* 1999; 9.
- Rocklin D, Levrel H, Drogou M, Herfaut J, Véron G. Combining telephone surveys and fishing catches self-report: The French sea bass recreational fishery assessment. *Plos One* 2014; 9: e87271.
- Roll U, Correia RA, Berger-Tal O. Using machine learning to disentangle homonyms in large text corpora. *Conservation Biology* 2018; 32: 716-724.
- Ruitort J. Premier signalement d'*Epinephelus aeneus* (Geoffroy st. Hilaire, 1817)(Perciformes, serranidae) dans les eaux méditerranéennes françaises. *Bulletin de la Société des Sciences Historiques et Naturelles de la Corse* 2012: 738-739.
- Sadovy de Mitcheson Y, Craig MT, Bertoni AA, Carpenter KE, Cheung WWL, Choat JH, et al. Fishing groupers towards extinction: a global assessment of threats and extinction risks in a billion dollar fishery. *Fish and Fisheries* 2013; 14: 119-136.
- Sáenz-Arroyo A, Roberts CM, Torre J, Cañño-Olvera M. Using fishers' anecdotes, naturalists' observations and grey literature to reassess marine species at risk: the case of the Gulf grouper in the Gulf of California, Mexico. *Fish and Fisheries* 2005; 6: 121-133.
- Samia DSM, Bessa E, Blumstein DT, Nunes JACC, Azzurro E, Morroni L, et al. A meta-analysis of fish behavioural reaction to underwater human presence. *Fish and Fisheries* 2019; 20: 817-829.
- Sbragaglia V, Cerri J, Bolognini L, Dragičević B, Dulčić J, Grati F, et al. Local ecological knowledge of recreational fishers reveals different meridionalization dynamics of two Mediterranean subregions. *Marine Ecology Progress Series* 2020; 634: 147-157.
- Sbragaglia V, Correia RA, Coco S, Arlinghaus R. Data mining on YouTube reveals fisher group-specific harvesting patterns and social engagement in recreational anglers and spearfishers. *ICES Journal of Marine Science* 2019a; fsz100.
- Sbragaglia V, López-Olmeda JF, Frigato E, Bertolucci C, Arlinghaus R. Fisheries-induced evolution of the circadian system and collective personality traits. *bioRxiv* 2019b: 622043.
- Sbragaglia V, Morroni L, Bramanti L, Weitzmann B, Arlinghaus R, Azzurro E. Spearfishing modulates flight initiation distance of fishes: the effects of protection, individual size, and bearing a speargun. *ICES Journal of Marine Science* 2018; 75: 1779-1789.
- Thaben PF, Westermark PO. Detecting rhythms in time series with RAIN. *J Biol Rhythms* 2014; 29: 391-400.
- Toivonen T, Heikinheimo V, Fink C, Hausmann A, Hiippala T, Järvi O, et al. Social media data for conservation science: A methodological overview. *Biological Conservation* 2019; 233: 298-315.

- Valls A, Coll M, Christensen V. Keystone species: toward an operational concept for marine biodiversity conservation. *Ecological Monographs* 2015; 85: 29-47.
- Vargha A, Delaney HD. A critique and improvement of the CL common language effect size statistics of McGraw and Wong. *Journal of Educational and Behavioral Statistics* 2000; 25: 101-132.
- Venturelli PA, Hyder K, Skov C. Angler apps as a source of recreational fisheries data: opportunities, challenges and proposed standards. *Fish and Fisheries* 2017; 18: 578-595.
- YouTube. <https://www.youtube.com/about/press> (Last accessed on 21 April 2020), 2020.
- Zabala M, García-Rubies A, Louisy P, Sala E. Spawning behaviour of the mediterranean dusky grouper *Epinephelus marginatus* (Lowe, 1834)(Pisces, Serranidae) in the Medes islands marine reserve (NW Mediterranean; Spain). *Scientia Marina* 1997; 61: 65-77.

Journal Pre-proof

Table 1 – Summary of information related to the four target species of groupers. Depth range and conservation statuses are those reported by the IUCN red list Mediterranean assessment (<https://www.iucnredlist.org>). The intrinsic vulnerability index is extracted by FishBase (<http://www.fishbase.org>). Such index is calculated using a fuzzy logic expert system according to species life history and ecological characteristics associated to fishing vulnerability (the index values ranging from 1 to 100, with 100 being the most vulnerable; Cheung et al., 2005).

Species	Depth range (m)	Intrinsic vulnerability index	IUCN status (Mediterranean)
Dusky grouper (<i>E. marginatus</i>)	8-300	72/100	Endangered (EN)
White grouper (<i>E. aeneus</i>)	0-200	52/100	Near Threatened (NT)
Goldblotch grouper (<i>E. costae</i>)	20-80	66/100	Data Deficient (DD)
Dogtooth grouper (<i>E. caninus</i>)	30-400	87/100	Data Deficient (DD)

Table 2 – Number of videos related to recreational fishing of groupers mined on YouTube between 2011 and 2017. The different species with their common and scientific name are reported together with the number of videos mined for each recreational fishing modality.

Species	Modalities		Total
	Angling	Spearfishing	
Dusky grouper (<i>E. marginatus</i>)	171	1499	1670
White grouper (<i>E. aeneus</i>)	136	126	262
Goldblotch grouper (<i>E. costae</i>)	25	75	100
Dogtooth grouper (<i>E. caninus</i>)	31	1	32
Not identified	20	13	33

Figure 1 – The annual number of videos related to recreational fishing of groupers mined on YouTube between 2011 and 2017 in Italy. The annual number of videos is reported for each species according to the recreational fishing modality (angling: red triangles; spearfishing: blue circles) together with the videos where the species was not identified. The images of the groupers are adapted from Heemstra (1993).

Figure 2 – The summary of the data for the videos related to recreational fisheries of the dusky grouper (*Epinephelus marginatus*) between 2011 and 2017 in Italy: (A) The number of uploaded videos for each month (angling: N=171; spearfishing: N=1499); (B) the declared mass (kg) for recreational anglers (N=34) and recreational spearfishers (N=426); (C) the declared depth (m) for recreational anglers (N=6) and recreational spearfishers (N=149); (D) the linear correlation and 95% confidence interval between declared mass (kg) and declared depth (m) for recreational spearfishing (N=75); (E) the declared depth (m) for each year (small circles) together with the mean (large circles) and standard deviation (vertical lines) for recreational spearfishing (N=149). The black horizontal line represents significant differences between groups (***: $p < 0.001$). The image of the grouper is adapted from Heemstra (1993).

Figure 3 – The summary of the data for the videos related to recreational fisheries of the white grouper (*Epinephelus aeneus*) between 2011 and 2017 in Italy: (A) The number of uploaded videos for each month (angling: N=136; spearfishing: N=126); (B) the declared mass (kg) for recreational anglers (N=42) and recreational spearfishers (N=25); (C) the declared depth (m) for recreational anglers (N=6) and recreational spearfishers (N=15); (D) the latitude (small circles) of the location retrieved from the videos for each year together with the mean (large circles) and standard deviation (vertical lines) for recreational anglers and spearfishers (N=64); (E) the geographical distribution of the videos according to the information retrieved in their title, description as well as in the video itself. Red triangles represent recreational angling, while blue circles represent recreational spearfishing. The black horizontal line represents significant differences between groups (***: $p < 0.001$). The image of the grouper is adapted from Heemstra (1993).

Figure 4 – The summary of the data for the videos related to recreational fisheries of the goldblotch grouper (*Epinephelus costae*) between 2011 and 2017 in Italy: (A) The number of uploaded videos for each month (angling: N=25; spearfishing: N=75); (B) the declared mass (kg) for recreational anglers (N=1) and recreational spearfishers (N=14); (C) the declared depth (m) for recreational anglers (N=2) and recreational spearfishers (N=10). The image of the grouper is adapted from Heemstra (1993).

Figure 5 – The summary of the data for the videos related to recreational fisheries of the dogtooth grouper (*Epinephelus caninus*) between 2011 and 2017 in Italy: (A) The number of uploaded videos for each month (angling: N=31; spearfishing: N=1); (B) the declared mass (kg) for recreational anglers (N=12); (C) the declared depth (m) for recreational anglers (N=5). The image of the grouper is adapted from Heemstra (1993).

SUPPLEMENTARY INFORMATION

Figure S1 – Representation of the Mediterranean Sea showing all the occurrence records of the white grouper (*Epinephelus aeneus*) available in the Global Biodiversity Information Facility (green circles, N = 122; GBIF.org. (25 August 2020) GBIF Occurrence Download: <https://doi.org/10.15468/dl.bdmkpu>) together with the georeferenced catches documented here using data mining on YouTube between 2011 and 2017 (purple circles, N = 64; see also Fig. 3E for more details).

Credit author statement

Valerio Sbragaglia: Conceptualization; Formal analysis; Funding acquisition; Investigation; Visualization; Writing - original draft. **Salvatore Coco:** Data curation; Validation; Writing - review & editing. **Ricardo A. Correia:** Methodology; Writing - review & editing. **Marta Coll:** Writing - review & editing. **Robert Arlinghaus:** Writing - review & editing.

Journal Pre-proof

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Graphical abstract

Highlights

- Digital data can provide novel ecological and environmental insights
- Recreational fishers represent a widespread network of observers of natural phenomena
- Recreational fishers share digital information about catches on social media
- Digital data about recreational fishing can be systematically mined on social media
- We show that such data can be used to explore several macroecological patterns

Journal Pre-proof