

1 **Title: Analysis of lipid composition reveals mechanisms of ethanol tolerance in the model yeast**
2 ***Saccharomyces cerevisiae***

3
4 Lairón-Peris M,^{a*} Routledge SJ,^{b*} Linney JA,^b Alonso-del-Real J,^a Spickett, C.M.^b Pitt AR,^{b,c}
5 Guillamon JM,^a Barrio E,^{a,d} Goddard AD^{#,b} Querol A^{#a}

6
7 ^aFood Biotechnology Department, Institute of Agrochemistry and Food Technology, CSIC, Valencia,
8 Spain

9 ^bCollege of Health and Life Sciences, Aston University, Birmingham, UK

10 ^cManchester Institute of Biotechnology and Department of Chemistry, University of Manchester,
11 Manchester, UK

12 ^dGenetics Department, University of Valencia, Valencia, Spain

13

14 Running head: Membrane lipid composition in *S. cerevisiae* strains

15 # These authors contributed equally. Address correspondence to Alan Goddard,

16 a.goddard@aston.ac.uk Amparo Querol, aquerol@iata.csic.es

17 * María Lairón-Peris and Sarah J. Routledge contributed equally to this work. Author order was
18 determined on the basis of alphabetical order.

19

20 **Abstract**

21 *Saccharomyces cerevisiae* is an important unicellular yeast species within the biotechnological and
22 food and beverage industries. A significant application of this species is the production of ethanol,
23 where concentrations are limited by cellular toxicity, often at the level of the cell membrane. Here,
24 we characterize 61 *S. cerevisiae* strains for ethanol tolerance and further analyse five representatives
25 with varying ethanol tolerances. The most tolerant strain, AJ4, was dominant in co-culture at 0% and
26 10% ethanol. Unexpectedly, although it does not have the highest NIC or MIC, MY29 was the
27 dominant strain in co-culture at 6% ethanol, which may be linked to differences in its basal lipidome.
28 Whilst relatively few lipidomic differences were observed between strains, a significantly higher PE
29 concentration was observed in the least tolerant strain, MY26, at 0% and 6% ethanol compared to the
30 other strains that became more similar at 10%, indicating potential involvement of this lipid with
31 ethanol sensitivity. Our findings reveal that AJ4 is best able to adapt its membrane to become more
32 fluid in the presence of ethanol and lipid extracts from AJ4 also form the most permeable membranes.
33 Furthermore, MY26 is least able to modulate fluidity in response to ethanol and membranes formed
34 from extracted lipids are least leaky at physiological ethanol concentrations. Overall, these results

35 reveal a potential mechanism of ethanol tolerance and suggests a limited set of membrane
36 compositions that diverse yeast species use to achieve this.

37

38 **Importance**

39 Many microbial processes are not implemented at the industrial level because the product yield is
40 poorer and more expensive than can be achieved by chemical synthesis. It is well established that
41 microbes show stress responses during bioprocessing, and one reason for poor product output from
42 cell factories is production conditions that are ultimately toxic to the cells. During fermentative
43 processes, yeast cells encounter culture media with high sugar content, which is later transformed
44 into high ethanol concentrations. Thus, ethanol toxicity is one of the major stresses in traditional and
45 more recent biotechnological processes. We have performed a multilayer phenotypic and lipidomic
46 characterization of a large number of industrial and environmental strains of *Saccharomyces* to
47 identify key resistant and non-resistant isolates for future applications.

48

49 **Keywords:** ethanol, *S. cerevisiae*, membrane properties

50

51 **Introduction**

52 *Saccharomyces cerevisiae* is a unicellular eukaryotic microorganism that has been employed as a
53 model organism to study diverse relevant phenomena in biology at molecular level (1). Due to its
54 high fermentative capability, it is also widely used in the biotechnology field for the performance of
55 industrial fermentations of products such as wine, beer or bread (2) or traditional Latin American
56 beverages like pulque, masato, chicha, tequila, or cachaça (3–7). *S. cerevisiae* also has a relevant role
57 in bioethanol production (8). *S. cerevisiae* has been isolated from different sources and environments
58 all over the world, including fruits, soils, cactus, insects, oak, and cork tree barks (9, 10). The
59 physiological and genetic diversity among the *Saccharomyces* genus is high, due to their colonization
60 of different environments; the most studied species are those associated with industrial processes of
61 economic importance as wine production (11–17), cider (18) and beer (11). *Saccharomyces* yeasts
62 that have been selected to carry out these fermentations in a controlled manner show particular
63 characteristics, as selective pressures imposed by the fermentative environment, such as low pH and
64 the high ethanol levels in the media, favor yeasts with the most efficient fermentative catabolism,
65 particularly *S. cerevisiae* strains, but there are species in the *Saccharomyces* genus which are also
66 found spontaneously in these fermentation products including *S. uvarum*. Depending on the
67 fermentation process, other factors apart from alcohol concentration, as temperature, can be
68 considered stress factors (19–21).

69 Ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) is a small molecule containing a methyl group and a hydroxyl group and
70 consequently it is soluble in both aqueous and lipidic phases. Because of these properties, it can
71 penetrate inside cells, which generates important stresses; incorporation into the cell membrane can
72 increase fluidity, which is a fundamental driver of membrane properties (22, 23).

73 This fluidity change induces a loss of membrane integrity, becoming more permeable (24). Ethanol
74 causes other detrimental effects to the cells, including alterations on mitochondrial structure, reducing
75 ATP levels and respiratory frequency and favoring acetaldehyde and reactive oxygen species (ROS)
76 generation, which can cause lipid peroxidation, DNA damage and oxidative stress (25, 26). As a
77 consequence, a notable reduction in cellular viability occurs. Cell membranes are composed of lipids
78 (mainly phospholipids and sterols, but also sphingolipids and glycolipids) and proteins. Membrane
79 lipids are amphipathic, possessing hydrophobic (apolar) and hydrophilic (polar) regions. Embedded
80 membrane proteins are strongly associated with the apolar core of the bilayer and peripheral proteins
81 are more loosely associated with the membrane via several mechanisms. A key factor contributing to
82 membrane fluidity is the fatty acids and sterol composition of the membrane (27).

83 The molecular structure of ethanol allows passive diffusion across the membrane and likely
84 incorporation into the bilayer structure (28). When this happens, van der Waals attractive forces
85 decrease, increasing membrane fluidity (29). Using fluorescence anisotropy studies a direct
86 relationship between plasma membrane fluidity and ethanol concentration has been reported (30, 31).
87 This increase in fluidity, together with the loss of structural integrity previously mentioned, result in
88 loss of various intracellular components including amino acids and ions (24), producing alterations
89 in a cellular homeostasis.

90 The alterations in membrane properties are fundamental in the mechanism of ethanol toxicity but the
91 physical changes that the membrane structure undergoes as a result of ethanol presence in the media
92 have not been completely described. It is widely accepted that ethanol is intercalated in lipidic heads
93 of the membrane, with the OH group of the ethanol associated with the phosphate group of the lipidic
94 heads and the hydrophobic tails aligned with the hydrophobic core of the membrane. When this
95 interaction takes place, ethanol molecules substitute interfacial water molecules, generating lateral
96 spaces between polar heads, and, as a consequence, spaces in the hydrophobic core (32). These gaps
97 result in unfavorable energy, so the system tries to minimize it by creating an interdigitated phase.
98 This modification in the membrane causes a decrease in its thickness of at least 25% (33, 34) and as
99 a consequence of this thinning, alterations in membrane protein structure and function can occur,
100 leading to cellular inactivation during the fermentation process (35).

101 It has been demonstrated that membrane thickness affects membrane protein functionality, in which
102 maximum activity takes place with a defined thickness (36, 37). If this thickness changes, exposure
103 of hydrophobic amino acid residues in integral membrane proteins can take place, resulting in a

104 phenomenon known as hydrophobic maladjustment (35), that can lead to aggregation of membrane
105 proteins to minimize the exposition of their hydrophobic parts in the aqueous media (38). Studies that
106 use membrane models formed by phosphatidylcholine and ergosterol that are exposed to different
107 ethanol concentrations have demonstrated that lipid composition protects the membrane because
108 interdigitated phase formation is delayed (39).

109 In Arroyo-López et al. (40) different *Saccharomyces* species were characterized for their ethanol
110 tolerance, identifying *S. cerevisiae* as the most ethanol tolerant. In the present work, we have selected
111 61 *S. cerevisiae* strains, from different origins and isolation sources. The purpose of this study was to
112 establish differences in the behavior of strains that represent the different *S. cerevisiae* groups, to
113 determine the most resistant ones, so they are better to perform industrial fermentations. With this
114 aim, we both monitored the growth in a liquid medium with different ethanol concentrations, using
115 absorbance measurements, and in a solid media, carrying out drop test analysis on ethanol plates.
116 Growth data were statistically analyzed for each of the *S. cerevisiae* strains and strains showing a
117 different behavior under ethanol stress were selected to conduct membrane studies that allow
118 correlations of lipid composition in yeast populations with responses to environmental stress such as
119 ethanol.

120

121 **Results**

122 **Ethanol tolerance of the strains in solid media.**

123 A total of 61 yeast strains belonging to *S. cerevisiae* were selected to assess ethanol tolerance. The
124 strains have been identified by sequencing of the D1/D2 26S, sequencing of the D1/D2 26S rRNA
125 gene was deposited in GenBank with the accession numbers MW559910-MW559970. All the strains
126 have been identified as *S. cerevisiae* with the exception of MY62 that is a *S. cerevisiae* strain
127 containing a limited amount of *S. kudriavzevii* genome. 21 are industrial strains and were selected for
128 their use in winemaking and 40 of them belong to the IATA-CSIC collection. The sources from which
129 these 40 strains were retrieved are diverse: agave, beer, bioethanol, chicha, cider, cocoa, honey water,
130 masato, sake, sugar cane, wine, natural wild strains, etc. *S. cerevisiae* yeast strains' ethanol tolerance
131 was first assessed in plates with GPY + different ethanol percentages. To observe the influence of
132 ethanol on these strains we performed four biological replicates of each strain growth in 6 different
133 media. One biological replicate for each of the strains and media can be seen in Fig. S1. With the
134 growth data of each of the strains and taking into account, the 4 replicates values of growth for each
135 strain, a heatmap with the growth data in ethanol was constructed (Fig. 1). This heatmap is
136 hierarchically clustered into two big clusters with different subclusters. The first cluster is made up
137 of the strains which are more tolerant of ethanol (a total number of 22 of the 61 strains) and another
138 one with the rest of the strains which show intermediate and low growth with this compound (39

139 strains). Among the first cluster, with the most tolerant strains, it is interesting that 19 of the 22 strains
140 belong to commercial wine strains. The other 3 strains which are included in this heatmap are AJ4, a
141 Lallemand commercial strain, which is also one of the most tolerant strains of all the screened ones,
142 MY48, a cachaça strain and MY43, a cider yeast strain.

143 The other cluster, with the 39 intermediate-low tolerant strains, appears to be divided into two
144 subclusters too. One of the subclusters is composed of MY33 and MY34, which are the less ethanol
145 tolerant strains, and belong to the sake group. It is interesting to note that in the other subcluster, there
146 are strains with different behaviors. As an example, strains MY46 (cachaça) and MY44 (cider) growth
147 in ethanol media are affected by low ethanol concentrations (ethanol percentage of 6%), but they can
148 grow (at a low rate) until 16% of ethanol is present in solid media. On the other hand, there are other
149 strains, such as MY37 (Masato) and MY22 (natural), whose growth is not affected until 10% of
150 ethanol is present in GPY solid media but in the next ethanol step (14%) they do not grow at all.

151

152 **Ethanol tolerance of the strains in liquid media.**

153 Ethanol tolerance of the set of *S. cerevisiae* strains was evaluated in minimal YNB liquid media at
154 28°C. Yeast growth was evaluated by OD₆₀₀ determination in microtiter plates containing this media
155 with different ethanol concentrations and for each strain, the area under the curve during these
156 growths was calculated. With the area under the curve reduction due to the addition of ethanol, NIC
157 (non-inhibitory concentration) and MIC (minimum inhibitory concentration) parameters were
158 calculated for 57 of the 61 strains. Not all of the 61 strains could be evaluated following this method:
159 the data obtained with flor strains MY28 and MY31 could not be used because these strains flocculate
160 and the data obtained with them are not reproducible. The data obtained with the strains MY55 and
161 MY56 were not used as they have problems growing in minimal media YNB. The complete list with
162 the NIC and MIC values for each one of the selected strains can be found in Table S1. Fig. 2 depicts
163 a graph representing these values for each one of the strains.

164

165 **Strain selection**

166 After performing the phenotypic characterization in ethanol of our collection of 61 strains, to further
167 characterize some representatives of the different behaviors we decided to select 5 of them as they
168 showed a range of tolerances: AJ4, MY3, MY14, MY26, and MY29. Fig. 3A shows the results of the
169 drop test in GPY+ethanol media of these 5 strains and Fig. 3B the NIC and MIC parameters of growth
170 in YNB liquid media+ethanol.

171 AJ4 shows high NIC and MIC values during YNB growth in liquid media, and in solid media in GPY
172 + ethanol it clusters amongst the most tolerant *S. cerevisiae* strains too. This strain is a Lallemand
173 commercial strain that has been reported as a highly tolerant ethanol strain (41). It has a high NIC

174 value $11.62\% \pm (0.33\%)$, which means that a high concentration of ethanol is needed to affect its
175 growth.

176 MY29, which is a flor strain isolated from sherry wine, is classified within the second cluster with
177 the strains that show an intermediate growth in GPY+ethanol in solid media. It grows well until 14%
178 ethanol; however, viability is reduced in 16% ethanol, and it is unable to grow at 18% ethanol.
179 Regarding the liquid assay in YNB+ethanol, its MIC value is amongst the highest MIC values of all
180 the strains ($15.41\% \pm 2.93\%$), but its NIC value ($7.5\% \pm 1.48\%$) can be classified as a medium-low
181 value. This result shows that MY29 is a *S. cerevisiae* strain whose behaviour can be classified as
182 intermediate in ethanol conditions. Moreover, MY29 is the most tolerant sherry wine strain of the
183 five strains analyzed.

184 MY26, which is an agave strain, is among the least tolerant strains in solid media and is also the strain
185 that shows the lowest growth among the three agave strains that we selected for our study. In liquid
186 media, its NIC value is also low, being affected by an ethanol concentration of $7.24\% \pm 0.77\%$ but
187 its MIC value is high ($15.34\% \pm 0.4\%$). This strain shows similar behavior in liquid media as MY29,
188 but in solid media, it proved to be less tolerant as it was not able to grow in 14% ethanol plates, and
189 MY26 could grow in this condition too.

190 MY3 and MY14 are commercial wine strains, which are classified in the cluster of the most tolerant
191 strains regarding their growth on ethanol plates. Nevertheless, MY14 appears to be affected by the
192 ethanol at low concentrations (NIC value of $6.787\% \pm 0.337\%$ and MIC value of $13.93\% \pm 0.91\%$)
193 and MY3 seems to start being affected by ethanol at higher concentrations but has a low range, as it
194 has a low MIC value (NIC $8.89\% \pm 1.26\%$ and MIC $12.97\% \pm 0.13\%$).

195

196 **Competition fermentations**

197 These five strains, AJ4, MY3, MY14, MY26, and MY29 were selected for their different behavior
198 regarding ethanol susceptibility. They were inoculated into mixed culture fermentations to assess the
199 correlation between ethanol tolerance and competition capacity under different ethanol
200 concentrations (0%, 6%, and 10%). As one GPY fermentation would be insufficient for observing
201 domination of the culture by one single strain, we followed a method in which we inoculated a sample
202 of the culture after sugar depletion into new fresh media with the corresponding ethanol
203 concentration.

204 After the tenth pass, AJ4 completely dominated the 0% and the 10% fermentations. However, in 6%
205 fermentations, MY29 strain completely dominated one of the three replicate fermentations and clearly
206 dominated the other two. The other 2 strains which are present in this 6 % fermentation when sugar
207 is depleted are AJ4 and MY14, although in low proportion. Neither MY3 nor MY26 colonies were
208 found in any of the fermentation (Fig. 4). AJ4 dominating high ethanol concentration cultures was

209 quite an expected result regarding its ethanol tolerance determined in the present work. However, it
210 does not seem clear why MY29 dominates 6% ethanol cultures, given its moderate tolerance
211 compared to other strains such as AJ4, MY3 or even MY14. Here, probably, complex interaction
212 among strains play an important role in domination, which has been studied previously for another
213 set of strains (42), and demonstrated to be of importance together with growth capacity under the
214 studied media conditions (43).

215

216 **Lipid composition and membrane properties**

217 Several studies have demonstrated that yeasts can adapt their membrane composition in response to
218 ethanol stress (44–46). To better understand the effects of ethanol upon the yeast strains, we
219 investigated the properties of the membranes in the presence and absence of ethanol. We determined
220 the total lipid composition of each of the strains by mass spectrometry (Table S2 and S3). The number
221 of species identified for major lipid classes for strains grown in media containing 0% or 6% ethanol
222 is shown in Fig. 5. For the strains grown in the absence of ethanol, for ceramide 1-phosphates (CerP),
223 there were significantly fewer species observed in MY29 (109.6 ± 6.61) compared to AJ4 and MY3
224 (128.2 ± 1.49 and 130 ± 0.55), where $P < 0.01$ (two-way anova and Tukey's multiple comparisons
225 test) and MY14 (126.6 ± 1.86) where $P < 0.05$. For cardiolipin species (CL), there were significantly
226 fewer observed in AJ4 and MY3 (3.0 ± 0.45 and 3.0 ± 0.31); ($P < 0.01$), and MY14 and MY26 (4.2
227 ± 1.3 and 4.0 ± 0.55); ($P < 0.05$) when compared to MY29 (9.67 ± 1.8). There were fewer
228 diacylglycerols observed in MY29 compared to MY3 (180.2 ± 1.93 and 193.0 ± 1.41); ($P < 0.05$).
229 For glycerophosphatidicacid (GPA) species, there were significantly fewer species identified for
230 MY29 (126.4 ± 15.17) compared to AJ4 (178.0 ± 2.28 ; $P < 0.0001$), MY3 (175.0 ± 1.05 ; $P < 0.001$),
231 MY14 (170.4 ± 5.30 ; $P < 0.001$), and MY26 (167.8 ± 6.67 ; $P < 0.01$). There were also fewer
232 glycerophosphatidylethanolamine GPEth species identified for MY29 compared to each of the strains
233 ($P < 0.01$ in each case) (259.6 ± 3.2 AJ4; 258.4 ± 1.36 MY3; 254.8 ± 2.85 MY14; 252.4 ± 3.26 MY26
234 and 186.2 ± 35.034 for MY29). For glycerophosphoserine species (GPSer), there were fewer species
235 in MY29 (120.0 ± 12.99) compared to AJ4 and MY3 (157.6 ± 2.50 and 159 ± 1.41 ; $P < 0.001$), MY14
236 (151.6 ± 3.41 ; $P < 0.01$) and MY26 (147.4 ± 3.94 ; $P < 0.05$). Lastly, there were less monoacylglycerols
237 (MG) species observed in MY29 (19.0 ± 0.84) than for MY3 (24.6 ± 0.51 ; $P < 0.01$).

238

239 There were no significant differences observed between the species grown in the presence of 6%
240 ethanol; however, significant changes were seen between the 0% and 6% ethanol samples. For CL,
241 there were significantly fewer species observed for MY29 grown in 6% compared to 0% ethanol (3.0
242 ± 0.44 and 9.66 ± 1.80 ; $P < 0.01$). For DG, there were more species in 0% MY3 than 6% ($193.0 \pm$
243 1.41 and 178.4 ± 2.13 ; $P < 0.05$), for GPA there were significantly fewer species in MY29 at 0%

244 compared to 6% (126.4 ± 15.17 and 157.0 ± 4.03 ; $P < 0.05$), and for GPETH there were also
245 significantly fewer species in MY29 at 0% compared to 6% ethanol (186.2 ± 35.04 and 241.2 ± 1.82 ;
246 $P < 0.05$). There were significantly more MG species in MY3 at 0% (24.6 ± 0.51 and 20 ± 1.22 ; $P <$
247 0.05) and more TG species in MY3 at 0% compared to 6% ethanol (73.2 ± 1.39 and 66.6 ± 1.03 ; $P <$
248 0.01). Strikingly, MY29 seems to have the most different total lipid composition at 0% ethanol and
249 to remodel this most dramatically, in terms of species diversity, at 6%. However, at 6% ethanol,
250 species diversity in MY29 is similar to the other strains, perhaps indicating an optimal membrane
251 composition for ethanol tolerance.

252

253 Acyl chain length and saturation have been shown to be important factors in regulating membrane
254 fluidity and ethanol tolerance in yeast (44–46). We therefore investigated this for AJ4, MY3, MY14,
255 MY26 and MY29 strains in both 0% and 6% ethanol. While there were no significant changes in
256 average carbon length of the acyl chains for each of the strains grown in 0% compared to 6% ethanol
257 (Fig. S2), there were significant differences in saturation (Fig. 6). For the strains grown in 0% ethanol
258 (Fig. 6A), DG species contained a significantly lower percentage saturated acyl chains in MY29
259 compared to AJ4 (37.95 ± 0.35 and 40.22 ± 0.30 ; $P < 0.01$). There was a significantly higher
260 percentage of monounsaturated CL species in MY29 (30 ± 7.83) compared to AJ4 and MY3 (0 ± 0.0
261 in both cases; $P < 0.01$), and MY26 (3.33 ± 3.33 ; $P < 0.05$). For GPA, there was a significantly higher
262 percentage saturated chains in MY29 (34.51 ± 1.07) compared to MY14 (31.30 ± 0.88); $P < 0.05$. For
263 GPETH, there were more saturated chains in MY29 compared to AJ4, MY3, MY14, and MY26 (31.21
264 ± 3.79 ; 25.30 ± 0.24 ; 24.92 ± 0.16 ; 24.96 ± 0.26 ; 24.38 ± 0.26 ; $P < 0.05$ in each case). There was a
265 significantly greater number of saturated GPSer species in MY29 compared to MY26 (32.44 ± 1.70
266 and 29.24 ± 0.22 ; $P < 0.05$) and a lower number of monounsaturated species in MY29 (40.07 ± 2.20)
267 compared to MY3 and MY14 (45.11 ± 0.62 and 44.7 ± 0.59 ; $P < 0.05$). Lastly, there was a
268 significantly higher percentage of MG species containing two unsaturations in MY29 (10.59 ± 0.40)
269 compared to MY3 (8.14 ± 0.17) ($P < 0.05$). Once again, MY29 is the most different in terms of
270 saturated species at 0% ethanol and remodels its membrane to be more similar to the other strains at
271 6%.

272

273 There were no significant differences observed between strains for 6% ethanol samples (Fig. 6B), but
274 there were between strains grown in 0% compared to 6% ethanol. There was a significantly higher
275 percentage of saturated DG species for AJ4 at 0% than 6% ethanol (40.22 ± 0.30 and 38.08 ± 0.44),
276 and a lower percentage of monounsaturated species for AJ4 (32.80 ± 0.09 and 34.75 ± 0.38 ; $P <$
277 0.001) and MY3 (33.06 ± 0.21 and 34.54 ± 0.25 ; $P < 0.05$) at 0% compared to 6% ethanol. For
278 saturated GPETH species, there was a significantly higher percentage in 0% MY29 than 6% MY29

279 (31.21 ± 3.79 and 24.65 ± 0.26; P < 0.05), and significantly fewer monounsaturated species in 0%
280 MY29 compared to 6% (40.23 ± 0.55 and 41.94 ± 0.42; P < 0.05). There were significantly more
281 monounsaturated GPGro species in MY29 at 0% compared to 6% ethanol (19.12 ± 4.95 and 12.37 ±
282 1.05). In addition, there were significantly fewer monounsaturated GPSer species in 0% MY29 than
283 in 6% (40.07 ± 2.20 and 44.77 ± 0.23). Lastly, for TG species, there were significantly more saturated
284 species in MY14 at 0% ethanol than in MY14 at 6% (35.94 ± 0.58 and 30.86 ± 1.16; P < 0.001), more
285 monounsaturated species in AJ4 6% (26.33 ± 0.503; P < 0.01), MY14 6% (6.24 ± 0.55; P < 0.01), and
286 MY26 6% (25.73 ± 0.26; P < 0.05) compared to the 0% samples (23.40 ± 0.64; 23.60 ± 0.40 and
287 23.55 ± 0.25 respectively), and fewer species containing two unsaturations in MY3 (26.50 ± 0.47; P
288 < 0.01) and MY14 at 0% (26.98 ± 0.55; P < 0.05) compared to 6% (29.43 ± 0.68 and 29.39 ± 0.48)
289 samples.

290

291 To assess variation in overall lipid unsaturation the unsaturation index (UI) was calculated at the lipid
292 level by lipid class for species identified in each strain at 0% and 6% ethanol (Table 2) using the
293 percentage of lipids weighted by the number of unsaturated bonds: UI = % with one unsaturation +
294 (2 x % with two unsaturations) + (3 x % with three unsaturations) + (4 x % with four unsaturations).
295 The UI for DG was significantly lower for AJ4 compared to MY29 at 0% ethanol (86.76 ± 0.64 and
296 90.03 ± 0.61, P < 0.01) and higher for GPEth species in the 0% AJ4, MY14, MY26 strains compared
297 to MY29 (108.72 ± 0.35, 108.72 ± 0.28, 109.36 ± 0.60 and 97.36 ± 7.13 respectively, where P < 0.05
298 in each case). The UI for MY29 at 0% was also significantly lower than at 6% ethanol (108.73 ± 0.92,
299 P < 0.05). Lastly, the UI for MG species at 0% ethanol was significantly lower for MY3 compared to
300 MY29 (73.30 ± 16.58 and 83.27 ± 18.95, P < 0.05), and the UI for MY29 at 0% ethanol was
301 significantly higher compared to 6% MY29 (83.27 ± 18.95 and 78.74 ± 1.52, P < 0.05).

302

303 Due to changes observed in phosphatidylethanolamine (PE) and phosphatidylserine (PS) species
304 diversity in Fig. 5, we undertook quantitative TLC analysis of these lipids. This showed significant
305 differences in the abundance of PE in MY26 grown in 0% ethanol (0.41 ± 0.02), where the abundance
306 was higher compared to AJ4 (0.03 ± 0.01; P < 0.0001), MY3 (0.08 ± 0.01; P < 0.0001), MY14 (0.17
307 ± 0.01; P < 0.0001) and MY29 (0.18 ± 0.04; P < 0.0001) grown in 0% ethanol as illustrated by Fig.
308 7. There was also a significantly greater abundance of PE in 6% MY26 (0.41 ± 0.05) compared to
309 6% AJ4 (0.08 ± 0.03; P < 0.05), MY3 (0.07 ± 0.02; P < 0.0001), MY14 (0.09 ± 0.01; P < 0.0001) and
310 MY29 (0.13 ± 0.01; P < 0.0001). In addition, there was a lower abundance of PE in MY26 at 10%
311 ethanol (0.20 ± 0.06) compared to MY26 at both 0% (0.41 ± 0.02) and 6% ethanol (0.41 ± 0.051); P
312 < 0.001). There was a significantly lower abundance of PS in AJ4 at 0% ethanol (0.06 ± 0.01)
313 compared to MY14 and MY29 (0.36 ± 0.06 and 0.30 ± 0.09; P < 0.01 and P < 0.05, respectively).

314 There was also a significantly lower abundance of PS in MY3 compared to MY14 at 0% ethanol
315 (0.09 ± 0.01 and 0.36 ± 0.06 ; $P < 0.05$). It is notable that MY26, the least tolerant strain, is the most
316 different at 0% and 6% ethanol, but has a similar composition to the other strains at 10%.

317

318 We next examined the effect of ethanol upon the fluidity of the yeast membranes as they grew in
319 cultures with and without ethanol. We utilized the fluorescent dye, Laurdan, which has been used to
320 study phase properties of membranes as it is sensitive to the polarity of the membrane environment
321 (47). GP (Generalized Polarization) values, which correlate inversely with fluidity, were calculated
322 at six timepoints during the growth of AJ4, MY3, MY14, MY26 and MY29 strains in GPY, GPY
323 containing 6% ethanol and GPY containing 10% ethanol. The assay suggests that the fluidity of the
324 yeast membranes decreases with culture time as shown by the increase in GP (Fig. 8). AJ4 and MY14
325 strains demonstrated large changes in fluidity when treated with 10% ethanol (AJ4 showed a GP value
326 change of -0.0002 ± 0.0009 at 10% and a GP value change of 0.0233 ± 0.0025 at 0% and MY14
327 showed a GP value change of -0.0101 ± 0.002 at 10% and a GP value change of 0.009 ± 0.002 at 0%)
328 ($P < 0.001$ and $P < 0.01$, respectively). MY29 also became significantly more fluid at 10% ethanol
329 (GP value change of -0.0016 ± 0.0011 at 10% and a GP value change of 0.0084 ± 0.0019 at 0%) ($P <$
330 0.05). However, these strains did not show any increases in fluidity with 6% ethanol. The other strains
331 showed no significant differences to fluidity with ethanol treatment. It is notable that the most tolerant
332 strains show the largest increases in membrane fluidity in response to ethanol exposure.

333

334 To examine membrane permeability, we investigated the integrity of liposomes composed of lipids
335 extracted from each of the strains and loaded with carboxyfluorescein (CF) dye. The liposomes were
336 challenged with increasing concentrations of exogenous ethanol, and the fluorescence increase from
337 CF dye release was measured. The data in Fig. 9 shows that the liposomes containing lipids extracted
338 from AJ4 demonstrated a significantly greater increase in fluorescence at high ethanol concentrations
339 than those composed of lipids from the other strains (ANOVA and Tukey's multiple comparisons test
340 (90.98 ± 4.29 fluorescence increase; $P < 0.001$). MY3 and MY26 liposomes were less "leaky" overall
341 (46.38 ± 2.97 and 47.41 ± 7.84 of fluorescence increase). This increase in fluorescence indicates
342 increased "leakiness" of the membranes.

343

344 **Principal component analysis**

345 With the aim of grouping the 5 selected strains based on their lipid composition and their ethanol
346 tolerance, the data obtained in the previous sections was used to perform a PCA (Fig. 10). The data
347 from the variables NIC, MIC, and the drop test growth value at 14% and 16% of ethanol in the plates,
348 related to the ethanol tolerance were used. For the lipid composition, the data of the

349 carboxyfluorescein release at the last time point; the data from the Laurdan experiments of the
350 differential GP value at 10% of ethanol and when no ethanol is present in the last time point, and the
351 PE abundance at 0% and 6% of ethanol in the media was used. The two commercial wine strains
352 MY3 and MY14 group together, and MY26 (the most sensitive to ethanol) and AJ4 (the most tolerant)
353 are the two strains that demonstrate the most differences among them. It is interesting to note that
354 MY26 is associated in the PCA with an accumulation of PE in the membrane at low ethanol
355 concentration and a higher membrane rigidity, and the most tolerant stain, AJ4, associated with a high
356 membrane fluidity in the presence of ethanol.

357

358 **Discussion**

359 In this study, we investigated the membrane properties of the selected yeast strains to try to understand
360 their different levels of ethanol tolerance. The mass spectrometry analysis of the lipid composition of
361 each strain in the absence of ethanol highlighted differences, in particular between MY29 and the
362 other strains, not only in the variety of species observed for the lipid classes but also in their saturation.
363 MY29 is a flor yeast. These yeasts constitute a separate phylogenetic group within *S. cerevisiae*
364 species. They are characterized by forming a layer on top of wine known as flor, which allows them
365 to access the oxygen during the fermentation of sherry wines, so they show different behavior and
366 thus physiological characteristics to wine yeast. Moreover, they have been reported to survive under
367 extreme conditions (ethanol content over 15%) (48, 49), which could relate to their membrane
368 structure.

369 Upon treatment with 6% ethanol, the lipid composition of MY29 underwent significant changes; the
370 composition was then found to be more similar to that of the other strains, suggesting that the
371 membrane of MY29 underwent more drastic changes than the other strains in response to ethanol.
372 The lack of significant differences at 6% ethanol suggests that each of the strains move towards a
373 more common lipid composition in response to ethanol. However, despite the fewer differences to
374 lipid composition at 6% ethanol between the strains, MY29 dominated the fermentation at this
375 concentration. In addition, the lipid composition of AJ4 was not significantly different from the other
376 strains at 6% ethanol, although it is the most tolerant to ethanol. It is possible that there may be further
377 adaptation of the membrane at higher ethanol concentrations than were investigated in this study, but
378 it is likely that other factors contribute to the ethanol tolerance of these strains.

379 Indeed, this has been suggested by other studies, where the relationship between H⁺-ATPase activity
380 and ergosterol content as well as the sterol to phospholipid and protein to phospholipid ratios are
381 important (45, 50, 51). Ethanol tolerance is a complex phenotype, and different mechanisms may lead
382 to improved tolerance. Fluidisation of the yeast membranes by ethanol is also known to activate the
383 unfolded protein response (UPR), and it is speculated that a better response could lead to greater

384 tolerance (52). Moreover, yeast cells can increase their tolerance to ethanol by other mechanisms,
385 such as the increase the biosynthesis of some amino acids, as tryptophan (53)
386 and trehalose accumulation (54).

387 Nevertheless, it is striking that yeast strains with different membrane compositions in the absence of
388 ethanol become more similar upon exposure, suggesting a common, or limited number, of membrane
389 compositions that maximize tolerance to ethanol.

390 Incorporation of longer acyl chains and a decrease in shorter chains has previously been shown to
391 occur in yeast in response to ethanol (46, 55); however, we did not observe any significant changes
392 in chain length. Our study does suggest that there were significant differences in saturation between
393 the species upon ethanol treatment. These changes occurred in GPGro and GPEth in MY29, and
394 occurred predominantly in DG and TG for the other strains, with shifts towards increased saturation
395 for AJ4 and increased unsaturation for MY3 and MY14. These changes appear to be complex and
396 specific to each strain. Documented changes to the membrane of yeast upon ethanol challenge are
397 conflicting (56); while some studies have shown that increased levels of unsaturated fatty acids are
398 linked to improved ethanol tolerance (46), changes to the unsaturation index may not necessarily be
399 associated with improved tolerance, or lead to the expected changes in membrane fluidity, and it is
400 rather the potential of the cell to alter its composition (45, 57). The lipid membrane is a highly
401 complex environment and multiple factors can influence membrane fluidity and permeability. Further
402 study of these strains is required to determine if their different compositions have similar biophysical
403 properties.

404 We investigated the fluidity of the membranes and the Laurdan assay demonstrated that the fluidity
405 of the membranes for each strain decreased over the duration of the fermentation, which has been
406 observed previously (58), and may be linked to nutrient depletion and changes in the growth rate of
407 the cells. In our study, the most tolerant strain, AJ4, underwent the largest changes in fluidity, where
408 the membranes were significantly more fluid at 10% ethanol than in the other conditions. AJ4 lipid-
409 containing liposomes were also the “leakiest” when compared to the other strains. This strain may
410 therefore be better able to tolerate the fluidizing effects of ethanol upon the membrane or to modulate
411 its membrane composition to lead to an increase in fluidity; this more fluid composition may allow
412 more efficient movement of ethanol across the membrane. The membranes of one of the least tolerant
413 strains, MY26, did not alter in fluidity in any of the conditions and liposomes comprised of MY26
414 lipids were less leaky when challenged with ethanol. In addition, our analysis of PE abundance shows
415 that MY26 contained significantly more PE than the other strains in both 0% and 6% ethanol, while
416 the most tolerant strain, AJ4, contained less PE in general than other strains. PE has a small headgroup
417 and can form hydrogen bonds with adjacent PE molecules (59). It influences lipid packing and
418 therefore membrane fluidity, where increased PE content results in less fluid membranes (60, 61),

419 consistent with our hypothesis. Lower PE content in relation to PC has been correlated with more
420 tolerant strains (46, 62). These findings suggest that more tolerant strains are more fluid and
421 permeable, while less tolerant strains are more rigid and less permeable. Several studies have
422 correlated membrane fluidity and ethanol tolerance, and many of these point to increased fluidity
423 being associated with more tolerant strains (45, 57), although another study suggests that less fluid
424 membranes are associated with more tolerant strains (58). In this study, we provide further support
425 for the concept that low PE content is beneficial for ethanol tolerance. This result can guide
426 engineering to improve ethanol tolerance towards the reduction of PE synthesis. This compound is
427 produced by four separate pathways, but the Psd pathway, which utilizes PS as a substrate is
428 predominant in *S. cerevisiae* (63,64), so future works can be addressed in this direction.
429 In summary, the lipid composition of most of the yeast strains in this study were comparable but there
430 were significant differences between these and the MY29 strain. Upon ethanol treatment, this
431 composition changed significantly and a more similar composition was reached, suggesting an
432 adaptation mechanism in common with the other strains. Changes in saturation were observed for
433 each of the strains upon ethanol treatment, but it is not clear if these changes have a direct impact
434 upon fluidity and tolerance, and it is likely that other factors beyond the scope of this study play a
435 critical role and further investigation is needed. The PE abundance of the least tolerant strain, MY26,
436 was significantly higher than in the other strains. Our investigation therefore suggests that the
437 membranes of more tolerant strains are more fluid and contain less PE. Overall, our results point to a
438 reduced set of desirable membrane compositions and features that promote ethanol tolerance with
439 increased fluidity and permeability appearing to be key.

440
441

442 **Material and methods**

443 **Strains and media conditions.**

444 The *Saccharomyces cerevisiae* yeast strains used in this study are listed in Table 1. A total number of
445 61 strains from different isolation sources were selected. These strains were maintained in GPY-agar
446 medium (%w/v: yeast extract 0.5, peptone, 0.5, glucose 2, agar 2). Yeast identity was confirmed by
447 sequencing the D1/D2 domain of the 26S rRNA gene (65).

448

449 **Drop test experiments. Assay in ethanol plates**

450 To assess yeast strains' ethanol tolerance, drop test experiments were carried out. Rectangular GPY
451 plates supplemented with different ethanol percentages (0, 6, 10, 14, 16 and 18%) were prepared.
452 Yeast cells were grown overnight at 28°C on GPY media and diluted to an OD₆₀₀= 0.1 in sterile water.
453 Then, serial dilutions of cells (10⁻¹ to 10⁻³) were transferred on the plates with replicates and incubated

454 at 28°C for ten days with the plates wrapped in parafilm to avoid ethanol evaporation. Each strain
455 was inoculated twice on the same plate but at different positions, and an exact replicate of the plate
456 was made. With this method, four biological replicates of each strain were performed. Growth values
457 were assigned to each of the replicates: 0 no growth, 1 weak growth, 2 intermediate growth and 3
458 remarked growth. Median growth values were assigned for each ethanol concentration. Hierarchical
459 clustering used in heatmap plot was elaborated using www.heatmapper.ca tool, (66) with Euclidean
460 distance measurement method and group clustering was based on growth in different ethanol media
461 averages (average linkage).

462

463 **Assay in liquid media.**

464 **Optical density measurements.**

465 GPY precultures of each strain were prepared and incubated at 28°C overnight. These cultures were
466 washed with sterile water and adjusted to an OD₆₀₀= 0.1 in each one of the culture media (YNB liquid
467 media supplemented with different ethanol percentages (0, 1, 6, 8, 10, 13, 16 and 18 %)). YNB is
468 composed of 6.7 g/L of aminoacids and ammonium sulfate (YNB, Difco) and supplemented with 20
469 g/L of D-glucose as carbon source. Growth was monitored in a SPECTROstar Omega instrument
470 (BMG Labtech, Offenburg, Germany) at 28°C. Nunc™ MicroWell™ 96 well plates (ThermoFisher
471 Scientific) wrapped in parafilm and with water in each of its 4 repositories were employed.
472 Measurements were taken at 600 nm every 30min, with 10 seconds of preshaking before each
473 measurement until 64 hours of growth monitoring. All the experiments were carried out in triplicate.

474

475 **Estimation of the NIC and MIC parameters.**

476 The basis of the technique, used as in (40), is the comparison of the area under the OD–time curve of
477 positive control (absence of ethanol, optimal conditions) with the areas of the tested condition
478 (presence of ethanol, increasing inhibitory conditions). As the amount of inhibitor in the well
479 increases, the effect on the growth of the organism also increases. This effect on the growth is
480 manifested by a reduction in the area under the OD–time curve relative to the positive control at any
481 specified time.

482 Briefly, the areas under the OD–time curves were calculated by integration using GCAT software
483 (<http://gcat-pub.glbrc.org/>). Then, for each ethanol condition and strain replicate, the fractional area
484 (f_a) was obtained by dividing the tested area between the positive control area ($f_a = (\text{test area}) /$
485 (positive control area). The plot of the f_a vs log₁₀ ethanol concentration produced a sigmoid-shape
486 curve that could be well fitted with the modified Gompertz function for decay (67) $f_a =$
487 $A+C \times \exp[-\exp(B(x - M))]$. After this modelling, the NIC (non-*inhibitory* concentration) and MIC
488 (Minimum Inhibitory Concentration) parameters could be estimated as in (66).

489 $NIC = 10[M - (1.718/B)]$ $MIC = 10[M + (1/B)]$

490 To check for significant differences among yeast species for NIC and MIC parameters, an analysis of
491 variance was performed using the one-way ANOVA module of Statistica 7.0 software. Tukey's test
492 was employed for mean comparison. ggplot2 package (68) implemented in R software, version 3.2.2
493 (RDevelopment Core Team 2011) was employed for graphic representation of these NIC and MIC
494 values.

495 **Strains selection and competition fermentation**

496 Competition fermentations were carried out in 30 mL GPY, GPY+6% ethanol and GPY+10% ethanol
497 in triplicate. 0.1 OD of each of the 5 strains (AJ4, MY3, MY14, MY26, MY29) were inoculated in
498 every initial culture. Every 3/5 days 1 mL of the culture was transferred into the corresponding fresh
499 media. After 5 and 10 rounds, culture plates of samples from every tube were obtained. 20 colonies
500 from every plate were randomly picked for their identification. This was carried out by means of
501 mitochondrial digestion profile identification (69), which allowed differentiation of all the strains
502 except for MY14 and MY29, which shared the same exact profile. As an alternative, as we had
503 available the genome sequences of MY14 and MY29 (70), we identified a divergent region among
504 these two strains which encompasses gene *MMS1*. We amplified a region of gene *MMS1* with primers
505 *fl* (AACGGATCCTTTTCCCAAC) and *rl* (CGGTCGCAAAAATTAACG) and used *RsaI*
506 digestion to differentiate specially these two strains. Theoretical results for digestion bands sizes in
507 an agarose gel were calculated based on Sanger sequencing of the amplicon for the strains of interest
508 (Figure S3).

509

510 **Lipid composition and membrane studies.**

511 **Lipid extraction and quantification by ammonium ferrothiocyanate assay**

512 Yeast precultures of each one of the five selected strains (AJ4, MY3, MY12, MY26 and MY29) were
513 first propagated in 25 mL of GPY media at 200 rpm and 28°C. The cultures were harvested after 24
514 h and total lipids were extracted using a modified Bligh and Dyer protocol (71). To quantify the lipids,
515 10 µL sample was taken from the above 100 µL reconstituted lipids in chloroform and added to 2 mL
516 chloroform with 1 mL of assay reagent (0.1M FeCl₃.6H₂O, 0.4 M ammonium thiocyanate) in a 15
517 mL glass tube. Samples were vortexed for 1 min and centrifuged at 14,500 g for 5 mins. The lower
518 layer was collected into quartz cuvettes. The absorbance was measured at 488 nm, and the
519 concentration of lipid was determined by comparison with a standard curve of a mixture of
520 phospholipid standards (POPC, POPE and POPG) (Sigma).

521

522 **Mass spectrometry of lipids present in the strains**

523 The lipids from each of the five yeast strains extracted as previously described were reconstituted in
524 100 μ L chloroform to contain 5 μ g/ μ L lipid as determined by ammonium ferrothiocyanate assay, and
525 then diluted 1 in 50 in solvent A (50:50 acetonitrile:H₂O, 5 mM ammonium formate and 0.1% v/v
526 formic acid). Analysis of 10 μ L samples was performed by LCMS. LC was performed on a U3000
527 UPLC system (Thermo scientific, Hemel Hempstead) using a Kinetex C18 reversed phase column
528 (Phenomenex, 2.6 μ m particle size, 2.1 mm x 150 mm), at a flow rate of 200 μ L/min with a gradient
529 from 10% solvent B to 100% solvent B (85:10:5 isopropanol: acetonitrile: H₂O, 5 mM ammonium
530 formate and 0.1% v/v formic acid) with the following profile: t=0 10% A, t=20 86%A, t=22 96%A,
531 t=26 95%A. MS analysis was carried out in positive and negative ionization mode on a Sciex 5600
532 Triple TOF. Source parameters were optimized on infused standards. Survey scans were collected in
533 the mass range 250-1250 Da for 250 ms. MSMS data was collected using top 5 information dependent
534 acquisition and dynamic exclusion for 5 s, using a fixed collision energy of 35V and a collision energy
535 spread of 10V for 200 ms per scan. ProgenesisQI[®] was used for quantification and LipidBlast
536 (<https://fiehnlab.ucdavis.edu/projects/LipidBlast>) for identification. All data were manually verified
537 and curated. Data were analysed by two-way ANOVA and Tukey's multiple comparisons test, where
538 n = 5. Data sets were uploaded to: <https://doi.org/10.17036/researchdata.aston.ac.uk.00000495>

539

540 **TLC analysis.**

541 Yeast lipids extracted as above after 24 h growth were analysed by TLC. Briefly, 20 μ g of lipid sample
542 and 10 μ g phospholipid lipid standards (POPE and POPS) (Sigma) were loaded onto silica gel TLC
543 plates (Sigma) and separated using chloroform/methanol/acetic acid/water 25:15:4:2. The plates were
544 air dried and sprayed with ninhydrin reagent (0.2% ninhydrin in ethanol) (Sigma) and charred at
545 100°C for 5 mins. Images of plates were captured with a digital camera and spot intensity was
546 determined using ImageJ software.

547

548 **Laurdan membrane fluidity assay.**

549 Yeast cultures were set up in GPY and incubated at 200 rpm and 28°C overnight. Then, 25 mL of
550 GPY media containing 0% ethanol, 6% ethanol or 10% ethanol was inoculated to an OD₅₉₅ of 0.5.
551 Samples were taken at different time points during the fermentation, and live yeast were diluted to an
552 OD₅₉₅ of 0.4 in GPY and incubated with 5 μ M Laurdan (6-dodecanoyl-2-dimethylaminonaphthalene)
553 for 1 h. Fluorescence emission of these cells stained with Laurdan was taken using a microplate reader
554 (Mithras, Berthold) with the following filters; λ_{ex} =460 λ_{em} =535. Generalized Polarization (GP),
555 derived from fluorescence intensities at critical wavelengths, can be considered as an index of
556 membrane fluidity and is calculated as $GP = (I_{460} - I_{535}) / (I_{460} + I_{535})$. Data were analyzed by one-way
557 ANOVA and Tukey's multiple comparisons test, where n = 3.

558

559 **Carboxyfluorescein dye leakage assay**

560 Lipids for each of the five selected yeast strains extracted as described previously were used to
561 generate 400 nm liposomes loaded with 100 mM Carboxyfluorescein (CF) in protein buffer (50 mM
562 tris, 50 mM NaCl, pH 7.4). Dye leakage assays were performed with at 0.125 mg/mL liposomes and
563 increasing concentrations of ethanol in protein buffer at room temperature, and the fluorescence
564 emission measured (λ_{ex} = 492 nm, λ_{em} =512 nm). Liposomes were treated with 5% Triton X-100 to
565 fully disrupt them, and fluorescence measurements were normalized to the maximum reading for each
566 liposome composition. Data were analyzed by one-way ANOVA and Tukey's multiple comparisons
567 test, where n = 3.

568

569 **PCA analysis**

570 To visualize the relationships among different ethanol tolerance parameters and lipid composition of
571 the selected *S. cerevisiae* strains, a principal component analysis (PCA) was performed using the
572 `prcomp` function and `ggbiplot` (0.55 version) and `ggplot` (3.2.1 version) implemented in R.

573

574 **Data availability**

575 The sequencing of the D1/D2 26S rRNA gene of the strains was deposited in GenBank with the
576 accession numbers MW559910-MW559970 (73).

577

578 **Acknowledgments**

579 ML-P was supported by a FPU contract from Ministerio de Ciencia, Innovación y Universidades
580 (ref. FPU15/01775). This work was supported by projects ERACoBioTech MeMBrane project
581 (UE) to AQ and AG, PCI2018-093190 (AEI/FEDER, UE) to AQ and BBSRC (BB/R02152X/1) to
582 AG.

583

584 **References**

585

- 586 1. Smith MG, Snyder M. 2006. Yeast as a model for human disease. *Curr Protoc Hum Genet*
587 15.6.1-15.6.8.
- 588 2. Legras JL, Merdinoglu D, Cornuet JM, Karst F. 2007. Bread, beer and wine: *Saccharomyces*
589 *cerevisiae* diversity reflects human history. *Mol Ecol* 16:2091–2102.
- 590 3. Arias García JA. 2008. Diversidad genética en las especies del complejo *Saccharomyces*
591 *sensu stricto* de fermentaciones tradicionales.

- 592 4. Badotti F, Vilaça ST, Arias A, Rosa CA, Barrio E. 2014. Two interbreeding populations of
593 *Saccharomyces cerevisiae* strains coexist in cachaça fermentations from Brazil. *FEMS Yeast*
594 *Res* 14:289–301.
- 595 5. Kodama K. 1993. Sake-brewing yeasts, p. 129–68. *In* Rose A.H., Harrison J.S., E (ed.), *The*
596 *Yeasts*, 2nd ed. London: Academic Press.
- 597 6. Stringini M, Comitini F, Taccari M, Ciani M. 2009. Yeast diversity during tapping and
598 fermentation of palm wine from Cameroon. *Food Microbiol* 26:415–420.
- 599 7. Suárez Valles B, Pando Bedriñana R, Fernández Tascón N, González Garcia A, Rodríguez
600 Madrera R. 2005. Analytical differentiation of cider inoculated with yeast (*Saccharomyces*
601 *cerevisiae*) isolated from Asturian (Spain) apple juice. *LWT - Food Sci Technol* 38:455–461.
- 602 8. van Zyl WH, Lynd LR, den Haan R, McBride JE. 2007. Consolidated bioprocessing for
603 bioethanol production using *Saccharomyces cerevisiae*, p. 205–235. *In* Olsson, L (ed.),
604 *Biofuels*. Springer Berlin Heidelberg, Berlin, Heidelberg.
- 605 9. Eberlein C, Leducq JB, Landry CR. 2015. The genomics of wild yeast populations sheds
606 light on the domestication of man’s best (micro) friend. *Mol Ecol* 24:5309–5311.
- 607 10. Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, Davey RP, Roberts IN, Burt
608 A, Koufopanou V, Tsai IJ, Bergman CM, Bensasson D, O’Kelly MJT, van Oudenaarden A,
609 Barton DBH, Bailes E, Nguyen AN, Jones M, Quail MA, Goodhead I, Sims S, Smith F,
610 Blomberg A, Durbin R, Louis EJ. 2009. Population genomics of domestic and wild yeasts.
611 *Nature* 458:337–41.
- 612 11. Alba-Lois, L. & Segal-Kischinevzky C. 2010. Beer & wine makers. *Nat Educ* 3:17.
- 613 12. Camarasa C, Sanchez I, Brial P, Bigey F, Dequin S. 2011. Phenotypic landscape of
614 *Saccharomyces cerevisiae* during wine fermentation: Evidence for origin-dependent
615 metabolic traits. *PLoS One* 6.
- 616 13. Franco-Duarte R, Mendes I, Umek L, Drumonde-Neves J, Zupan B, Schuller D. 2014.
617 Computational models reveal genotype–phenotype associations in *Saccharomyces cerevisiae*.
618 *Yeast* 31:265–277.
- 619 14. Suárez Valles B, Pando Bedriñana R, Fernández Tascón N, Querol Simón A, Rodríguez
620 Madrera R. 2007. Yeast species associated with the spontaneous fermentation of cider. *Food*
621 *Microbiol* 24:25–31.
- 622 15. Querol A, Fernández-Espinar MT, Del Olmo M, Barrio E. 2003. Adaptive evolution of wine
623 yeast. *Int J Food Microbiol* 86:3–10.
- 624 16. Querol A, Barrio E, Ramón D. 1994. Population dynamics of natural *Saccharomyces* strains
625 during wine fermentation. *Int J Food Microbiol* 21:315–323.
- 626 17. Schuller D, Cardoso F, Sousa S, Gomes P, Gomes AC, Santos MAS, Casal M. 2012. Genetic
627 diversity and population structure of *Saccharomyces cerevisiae* strains isolated from different
628 grape varieties and winemaking regions. *PLoS One* 7.

- 629 18. Pando Bedriñana R, Querol Simón A, Suárez Valles B. 2010. Genetic and phenotypic
630 diversity of autochthonous cider yeasts in a cellar from Asturias. *Food Microbiol* 27:503–
631 508.
- 632 19. Ganucci D, Guerrini S, Mangani S, Vincenzini M, Granchi L. 2018. Quantifying the effects
633 of ethanol and temperature on the fitness advantage of predominant *Saccharomyces*
634 *cerevisiae* strains occurring in spontaneous wine fermentations. *Front Microbiol* 9:1563.
- 635 20. Salvadó Z, Arroyo-López FN, Guillamón JM, Salazar G, Querol A, Barrio E. 2011.
636 Temperature adaptation markedly determines evolution within the genus *Saccharomyces*.
637 *Appl Environ Microbiol* 77:2292–302.
- 638 21. Salvadó Z, Arroyo-López FN, Barrio E, Querol A, Guillamón JM. 2011. Quantifying the
639 individual effects of ethanol and temperature on the fitness advantage of *Saccharomyces*
640 *cerevisiae*. *Food Microbiol* 28:1155–1161.
- 641 22. Jones RP, Greenfield PF, Wang L, Cai Y, Zhao X, Jia X, Zhang J, Liu J, Zhen H, Wang T,
642 Tang X LY, J. W. 1987. Ethanol and the fluidity of the yeast plasma membrane. *Yeast* 3:223–
643 232.
- 644 23. Lloyd D, Morrell S, Carlsen HN, Degn H, James PE, Rowlands CC. 1993. Effects of growth
645 with ethanol on fermentation and membrane fluidity of *Saccharomyces cerevisiae*. *Yeast*
646 9:825–833.
- 647 24. Marza E, Camougrand N, Manon S. 2002. Bax expression protects yeast plasma membrane
648 against ethanol-induced permeabilization. *FEBS Lett* 521:47–52.
- 649 25. Alexandre H, Ansanay-Galeote, Dequin S, Blondin B. 2001. Global gene expression during
650 short-term ethanol stress in *Saccharomyces cerevisiae*. *FEBS Lett* 498:98–103.
- 651 26. Yang K-M, Lee N-R, Woo J-M, Choi W, Zimmermann M, Blank LM, Park J-B. 2012.
652 Ethanol reduces mitochondrial membrane integrity and thereby impacts carbon metabolism
653 of *Saccharomyces cerevisiae*. *FEMS Yeast Res* 12:675–684.
- 654 27. Zinser E, Sperka-Gottlieb CDM, Fasch E V., Kohlwein SD, Paltauf F, Daum G. 1991.
655 Phospholipid synthesis and lipid composition of subcellular membranes in the unicellular
656 eukaryote *Saccharomyces cerevisiae*. *J Bacteriol* 173:2026–2034.
- 657 28. Peña C, Arango R. 2009. Evaluación de la producción de etanol utilizando cepas
658 recombinantes de *Saccharomyces cerevisiae* a partir de melaza de caña de azúcar. *Dyna*
659 159:153–161.
- 660 29. Ingram LO, Buttke TM. 1985. Effects of Alcohols on Micro-Organisms. *Adv Microb Physiol*
661 25:253–300.
- 662 30. Sánchez-Gallego JI. 2009. Efecto de la quercetina y la rutina frente al daño oxidativo
663 inducido en eritrocitos con distintos contenidos de colesterol. Universidad de Salamanca.
- 664 31. Simonin H, Beney L, Gervais P. 2008. Controlling the membrane fluidity of yeasts during
665 coupled thermal and osmotic treatments. *Biotechnol Bioeng* 100:325–333.

- 666 32. Chiou J-S, Krishna PR, Kamaya H, Ueda I. 1992. Alcohols dehydrate lipid membranes: an
667 infrared study on hydrogen bonding. *Biochim Biophys Acta - Biomembr* 1110:225–233.
- 668 33. Kranenburg M, Vlaar M, Smit B. 2004. Simulating induced interdigitation in membranes.
669 *Biophys J* 87:1596–1605.
- 670 34. Vanegas JM, Faller R, Longo ML. 2010. Influence of ethanol on lipid/sterol membranes:
671 Phase diagram construction from AFM imaging. *Langmuir* 26:10415–10418.
- 672 35. Lee AG. 2004. How lipids affect the activities of integral membrane proteins. *Biochim*
673 *Biophys Acta - Biomembr* 1666:62–87.
- 674 36. Montecucco C, Smith G a, Dabbeni-sala F, Johannsson A, Galante YM, Bisson R. 1982.
675 Bilayer thickness and enzymatic activity in the mitochondrial cytochrome c oxidase and
676 ATPase complex. *FEBS Lett* 144:145–148.
- 677 37. Yuan C, O’Connell RJ, Feinberg-Zadek PL, Johnston LJ, Treistman SN. 2004. Bilayer
678 thickness modulates the conductance of the BK channel in model membranes. *Biophys J*
679 86:3620–3633.
- 680 38. Leão C VUN. 1984. Effects of ethanol and other alkanols on passive proton influx in the
681 yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 11:774:43–8.
- 682 39. Tierney KJ, Block DE, Longo ML. 2005. Elasticity and phase behavior of DPPC membrane
683 modulated by cholesterol, ergosterol, and ethanol. *Biophys J* 89:2481–2493.
- 684 40. Arroyo-López FN, Salvadó Z, Tronchoni J, Guillamón JM, Barrio E, Querol A. 2010.
685 Susceptibility and resistance to ethanol in *Saccharomyces* strains isolated from wild and
686 fermentative environments. *Yeast* 27:1005–1015.
- 687 41. Lairón-Peris M, Pérez-Través L, Muñiz-Calvo S, Guillamón JM, Heras JM, Barrio E, Querol
688 A. 2020. Differential contribution of the parental genomes to a *S. cerevisiae* × *S. uvarum*
689 hybrid, inferred by phenomic , genomic , and transcriptomic analyses , at different industrial
690 stress conditions. *Front Bioeng Biotechnol* 8:1–20.
- 691 42. Rossouw D, Bagheri B, Setati ME, Bauer FF. 2015. Co-flocculation of yeast species, a new
692 mechanism to govern population dynamics in microbial ecosystems. *PLoS One* 10:1–17.
- 693 43. Alonso-del-Real J, Lairón-Peris M, Barrio E, Querol A. 2017. Effect of temperature on the
694 prevalence of *Saccharomyces non cerevisiae* species against a *S. cerevisiae* wine strain in
695 wine fermentation: Competition, physiological fitness, and influence in final wine
696 composition. *Front Microbiol* 8.
- 697 44. Beaven MJ, Charpentier C, Rose AH. 1982. Production and tolerance of ethanol in relation to
698 phospholipid fatty-acyl composition in *Saccharomyces cerevisiae* NCYC 431. *J Gen*
699 *Microbiol* 128:1447–1455.
- 700 45. Alexandre H, Rousseaux I, Charpentier C. 1994. Relationship between ethanol tolerance,
701 lipid composition and plasma membrane fluidity in *Saccharomyces cerevisiae* and *Kloeckera*
702 *apiculata*. *FEMS Microbiol Lett* 124:17–22.

- 703 46. Chi Z, Arneborg N. 1999. Relationship between lipid composition, frequency of ethanol-
704 induced respiratory deficient mutants, and ethanol tolerance in *Saccharomyces cerevisiae*. J
705 Appl Microbiol 86:1047–1052.
- 706 47. Learmonth RP, Gratton E. 2002. Assessment of membrane fluidity in individual yeast cells
707 by laurdan generalised polarisation and multi-photon scanning fluorescence microscopy, p.
708 241–252. In *Fluorescence Spectroscopy*. Springer.
- 709 48. Martínez P, Pérez Rodríguez L, Benítez T. 1997. Velum formation by flor yeasts isolated
710 from sherry wine. Am J Enol Vitic 48:55 LP – 62.
- 711 49. Naumov GI. 2017. Genetic polymorphism of sherry *Saccharomyces cerevisiae* yeasts.
712 Microbiol Russian Fed 86:19–31.
- 713 50. Aguilera F, Peinado RA, Millán C, Ortega JM, Mauricio JC. 2006. Relationship between
714 ethanol tolerance, H⁺-ATPase activity and the lipid composition of the plasma membrane in
715 different wine yeast strains. Int J Food Microbiol 110:34–42.
- 716 51. Shobayashi M, Mitsueda SI, Ago M, Fujii T, Iwashita K, Iefuji H. 2005. Effects of culture
717 conditions on ergosterol biosynthesis by *Saccharomyces cerevisiae*. Biosci Biotechnol
718 Biochem 69:2381–2388.
- 719 52. Navarro-Tapia E, Querol A, Pérez-Torrado R. 2018. Membrane fluidification by ethanol
720 stress activates unfolded protein response in yeasts. Microb Biotechnol 11:465–475.
- 721 53. Yoshikawa K, Tanaka T, Furusawa C, Nagahisa K, Hirasawa T, Shimizu H. 2009.
722 Comprehensive phenotypic analysis for identification of genes affecting growth under
723 ethanol stress in *Saccharomyces cerevisiae*. FEMS Yeast Res 9:32–44.
- 724 54. Bandara A, Fraser S, Chambers PJ, Stanley GA. 2009. Trehalose promotes the survival of
725 *Saccharomyces cerevisiae* during lethal ethanol stress, but does not influence growth under
726 sublethal ethanol stress. FEMS Yeast Res 9:1208–1216.
- 727 55. You KM, Rosenfield C-L, Knipple DC. 2003. Ethanol tolerance in the yeast *Saccharomyces*
728 *cerevisiae* is dependent on cellular oleic acid content. Appl Env Microbiol 69:1499–1503.
- 729 56. Henderson CM, Block DE. 2014. Examining the role of membrane lipid composition in
730 determining the ethanol tolerance of *Saccharomyces cerevisiae*. Appl Environ Microbiol
731 80:2966–2972.
- 732 57. Huffer S, Clark ME, Ning JC, Blanch HW, Clark DS. 2011. Role of alcohols in growth, lipid
733 composition, and membrane fluidity of yeasts, bacteria, and archaea. Appl Environ Microbiol
734 77:6400–6408.
- 735 58. Ishmayana S, Kennedy UJ, Learmonth RP. 2017. Further investigation of relationships
736 between membrane fluidity and ethanol tolerance in *Saccharomyces cerevisiae*. World J
737 Microbiol Biotechnol 33:1–10.
- 738 59. Murzyn K, Róg T, Pasenkiewicz-Gierula M. 2005. Phosphatidylethanolamine-
739 phosphatidylglycerol bilayer as a model of the inner bacterial membrane. Biophys J
740 88:1091–1103.60. Dawaliby R, Trubbia C, Delporte C, Noyon C, Ruyschaert JM, Van

- 741 Antwerpen P, Govaerts C. 2016. Phosphatidylethanolamine is a key regulator of membrane
742 fluidity in eukaryotic cells. *J Biol Chem* 291:3658–3667.
- 743 61. Ballweg S, Sezgin E, Doktorova M, Covino R, Reinhard J, Wunnicke D, Hänel I, Levental I,
744 Hummer G, Ernst R. 2020. Regulation of lipid saturation without sensing membrane fluidity.
745 *Nat Commun* 11:1–13.
- 746 62. Jurešić GC, Blagović B, Rupčić J. 2009. Alterations in phosphatidylcholine and
747 phosphatidylethanolamine content during fermentative metabolism in *Saccharomyces*
748 *cerevisiae* brewer's yeast. *Food Technol Biotechnol* 47:246–252.
- 749 63. Birner R, Bürgermeister M, Schneiter R, Daum G. 2001. Roles of phosphatidylethanolamine
750 and of its several biosynthetic pathways in *Saccharomyces cerevisiae*. *Mol Biol Cell* 12:997–
751 1007.
- 752 64. Bürgermeister M, Birner-Grünberger R, Nebauer R, Daum G. 2004. Contribution of different
753 pathways to the supply of phosphatidylethanolamine and phosphatidylcholine to
754 mitochondrial membranes of the yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta -*
755 *Mol Cell Biol Lipids* 1686:161–168.
- 756 65. Kurtzman CP, Robnett CJ. 1998. Identification and phylogeny of ascomycetous yeasts from
757 analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie Van*
758 *Leeuwenhoek* 98:331–371.
- 759 66. Babicki S, Arndt D, Marcu A, Liang Y, Grant JR, Maciejewski A, Wishart DS. 2016.
760 Heatmapper: web-enabled heat mapping for all. *Nucleic Acids Res* 44:W147–W153.
- 761 67. Lambert, R. J. W. and Pearson J. 2000. Susceptibility testing: Accurate and reproducible
762 minimum inhibitory concentration (MIC) and non-inhibitory concentration (NIC) values. *J*
763 *Appl Microbiol* 88:784–790.
- 764 68. Wickham H. 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer- Verlag, New York.
- 765 69. Querol A, Barrio E, Ramón D. 1992. A comparative study of different methods of yeast strain
766 characterization. *Syst Appl Microbiol* 15:439–446.
- 767 70. Morard M, Macías LG, Adam AC, Lairón-Peris M, Pérez-Torrado R, Toft C, Barrio E. 2019.
768 Aneuploidy and ethanol tolerance in *Saccharomyces cerevisiae*. *Front Genet* 10:82.
- 769 71. Spickett CM, Reis A, Pitt AR. 2011. Identification of oxidized phospholipids by electrospray
770 ionization mass spectrometry and LC-MS using a QQLIT instrument. *Free Radic Biol Med*
771 51:2133–2149.
- 772 72. Erny C, Raoult P, Alais A, Butterlin G, Delobel P, Matei-Radoi F, Casaregola S, Legras JL.
773 2012. Ecological success of a group of *Saccharomyces cerevisiae*/*Saccharomyces*
774 *kudriavzevii* hybrids in the Northern European wine-making environment. *Appl Environ*
775 *Microbiol* 78:3256–3265.
- 776 73. Lairon-Peris M, Querol A. Data from “Composition of membrane lipids reveals mechanisms
777 of ethanol tolerance in the model yeast *Saccharomyces cerevisiae*.” GenBank
778 <https://www.ncbi.nlm.nih.gov/nuccore> (accession no. MW559910-MW55997).

779

780 **Figure legends.**

781

782 **FIG 1.** Heatmap representation of growth values (from 0 to 3) of the analysed strains at plates with
783 increasing ethanol concentrations. Each line corresponds to a strain (AJ4, MY1-MY63) and each
784 column to a particular ethanol concentration (0%, 6%, 10%, 14%, 16% and 18%). The color key bar
785 at the top indicates growth values, from yellow (low growth value) to pink (high growth value).
786 Hierarchical clustering is showed on the left. Color dots on the right of the Figure indicate the
787 source/origin of each one of the strains and shapes their classification. In Fig. S1 can be seen one of
788 the four replicates from which these heatmap was constructed.

789

790 **FIG 2.** Representation of each strain NIC (yellow) and MIC (red) parameters of the selected strains
791 in relation with its ethanol tolerance (%). Values are averages from triplicate experiments and
792 standard deviation is represented too. Color dots on the right of the Figure indicate the source/origin
793 of each one of the strains and shapes their classification. Strains are ordered by MIC value.

794

795 **FIG 3.** Photograph of the drop tests in ethanol plates (A) and the NIC and MIC parameters (B) for
796 each one of the 5 selected strains.

797

798 **FIG 4.** Percentage of strains present in GPY+ethanol media determined by molecular identification
799 after 10 rounds of fermentations. Every biological replicate is indicated by letters *A*, *B* and *C* and
800 the ethanol concentration present in the media in the *X* axis.

801

802 **FIG 5.** Number of species identified by lipid class for AJ4, MY3, MY14, MY26 and MY29 strains
803 in the presence of 0% ethanol and 6% of ethanol. Lipids were extracted and analysed by LC-MS in
804 positive and negative ion mode (n = 5).

805

806 **FIG 6.** Percentage of saturated, monounsaturated and polyunsaturated chains by lipid class showing
807 significant changes for A) AJ4, MY3, MY14, MY26 and MY29 strains in the presence of 0%
808 ethanol, and B) AJ4, MY3, MY14, MY26 and MY29 strains in the presence of 6% ethanol. Lipids
809 were extracted and analysed by LC-MS in positive and negative ion mode (n = 5).

810

811 **FIG 7.** TLC analysis of phosphatidylethanolamine (PE) and phosphatidylserine (PS) abundance for
812 AJ4, MY3, MY14, MY26 and MY29 strains in the presence of 0%, 6% and 10% of ethanol.

813 Samples were loaded in triplicate and spot intensity was analyzed using ImageJ. Spot intensity is
814 plotted relative to phospholipid standards loaded onto each plate.

815

816 **FIG 8.** The effects of ethanol upon the fluidity of live yeast throughout the fermentation, measured
817 by changes to Laurdan generalized polarization (GP).

818

819 **FIG 9.** The effects of ethanol upon liposomes composed of lipids extracted from AJ4, MY3, MY14,
820 MY26 and MY29 strains normalized to the maximum amount of dye released upon treatment with
821 5% Triton X-100.

822

823 **FIG 10.** Plot of the first two factors of a Principal Component Analysis (PCA) of the five *S.*
824 *cerevisiae* strains regarding their lipid composition and their ethanol tolerance.

825 **TABLE 1.** List of the 61 *Saccharomyces* strains selected used in this work.

Strain name	Strain repository / Collection	Isolation source and origin	Strain properties / Description
Wine comercial fermentation strains			
MY1	Lallemand	Wine	White and rosé wines
MY2	Lallemand	Wine	White wines
MY3	Lallemand	Wine	Rosé and red wines
MY4	Lallemand	Wine	White and rosé wines
MY6	Lallemand	Wine	White, rosé and red wines
MY7	Lallemand	Wine	Red wines
MY8	Lallemand	Wine	Red wines
MY11	Lallemand	Wine	White wines
MY12	Lallemand	Wine	Red wines
MY13	Lallemand	Wine	White, rosé and red wines
MY14	Lallemand	Wine	Sparkling wines, fruit wines and ciders
MY15	Lallemand	Wine	White wines
MY16	Lallemand	Wine	White, rosé and red wines
MY17	Lallemand	Wine	White wines
MY18	Lallemand	Wine	Stuck fermentations
MY19	Lallemand	Wine	Red wines
MY20	Lallemand	Wine	Red wines
MY21	Lallemand	Wine	Red wines
MY51	Lallemand / AQ29	Wine	Red wines
MY62	Lallemand	Wine	Red wines ^a
MY63	Lallemand	Wine	White and rosé wines

826
827
828
829
830
831

Strain name	Strain repository / Collection	Isolation source and origin	Strain properties / Description
Wine non-commercial fermentation strains			
MY52	AQ1336	Wine, South Africa	-
MY53	AQ923	Wine, Spain	-
MY54	AQ924	Wine, Spain	-
MY55	AQ2371	Bili wine, West Africa	-
MY56	AQ2375	Bili wine, West Africa	-
MY61	I.CF 14 ^b	Wine, Hungary	High Temperature
MY28	AQ2492	Flor wine, Spain	-
MY29	AQ2356	Flor wine, Spain	-
MY30	AQ94	Flor wine, Spain	-
MY31	AQ636	Flor wine, Spain	-
Other commercial fermentation strains			
AJ4	Lallemand	Fermentations	
MY50	Lallemand	Fermenting cacao	-
MY60	Fermentis	Bioethanol	Ethanol Red
Other non-commercial fermentation strains			
MY25	AQ2579	<i>Agave salmiana</i> , Peru	-
MY26	AQ2493	<i>Agave salmiana</i> , México	-
MY27	AQ2591	Chicha de jora, Perú	-
MY32	AQ594	Sake, Japan	-
MY33	AQ1312	Sakeye, Japan	-
MY34	AQ1314	Sakeye, Japan	-
MY35	AQ2332	Chicha de jora, Perú	-
MY36	AQ2469	Chicha de jora, Perú	-
MY37	AQ2363	Masato, Perú	-
MY38	AQ2473	Masato, Perú	-
MY43	AQ1180	Cider, Ireland	-
MY44	AQ1182	Cider, Ireland	-
MY45	AQ1184	Cider, Ireland	-
MY46	AQ2851	Sugar cane, Brazil	-
MY47	AQ2543	Sugar cane, Brazil	-
MY48	AQ2506	Sugar cane, Brazil	-
MY57	AQ843	Beer, Belgium	-
MY58	AQ1323	Sorghum beer, Burkina Faso	-
MY49	AQ1085	Fermenting cacao, Indonesia	-
MY59	UFLA	Bioethanol	-

834
835
836
837
838

TABLE 1. Continuation

Strain name	Strain repository / Collection	Isolation source and origin	Strain properties / Description
Natural Environmental strains			
MY22	AQ2458	<i>Agelaia vicina</i> , Peru	-
MY23	AQ2163	<i>Quercus faginea</i> , Spain	-
MY24	AQ997	<i>Prunus armeniaca</i> , Hungary	-
Clinical strains			
MY39	AQ2587	Dietetic product, Spain	-
MY40	AQ2654	Faeces, Spain	-
MY41	AQ435	Vagina, Spain	-
MY42	AQ2717	Lung, Spain	-

839

840 AQ (Amparo Querol collection),



841 ^aMY62 is a *S. cerevisiae* strain containing a limited amount of *S. kudriavzevii* genome (72, 73)


842 ^b kindly provided by Professor Sipiczki

843 **Table 2.** Unsaturation index (UI) for lipids identified in each strain was calculated using the percentage of lipids with each number of unsaturated bonds: one unsaturation + (2 x
844 two unsaturations) + (3 x three unsaturations) + (4 x four unsaturations). Statistically significant differences between strains and ethanol conditions are highlighted in bold (two-
845 way anova and Tukey's multiple comparisons test). Errors (SD) are shown in brackets, n = 5.
846
847



Lipid species	0% ethanol					6% ethanol				
	AJ4	MY3	MY14	MY26	MY29	AJ4	MY3	MY14	MY26	MY29
CerP	41.38 (± 1.09)	42.50 (± 1.01)	41.86 (± 0.54)	41.90 (± 0.50)	40.23 (± 4.34)	83.25 (± 0.58)	82.93 (± 0.41)	84.39 (± 1.17)	83.93 (± 0.43)	83.25 (± 0.58)
CL	33.33 (± 13.92)	70.00 (± 8.15)	62.22 (± 23.36)	90.00 (± 31.83)	88.00 (± 35.91)	107.24 (± 34.56)	60.00 (± 4.08)	111.79 (± 32.31)	30.00 (± 19.96)	84.33 (± 32.53)
DG	86.76 (± 0.64)	89.02 (± 0.29)	88.99 (± 0.62)	89.25 (± 0.48)	90.03 (± 0.61)	89.10 (± 4.08)	89.91 (± 0.31)	88.31 (± 0.63)	87.65 (± 0.64)	89.71 (± 0.69)
GPA	104.16 (± 0.98)	103.65 (± 0.34)	107.60 (± 1.63)	104.22 (± 1.01)	103.12 (± 0.87)	103.96 (± 23.31)	104.26 (± 1.00)	105.53 (± 0.34)	104.08 (± 1.02)	105.09 (± 1.13)
GPCho	50.20 (± 0.57)	50.29 (± 0.17)	50.03 (± 0.43)	50.72 (± 0.32)	50.36 (± 0.35)	96.64 (± 19.96)	96.66 (± 0.49)	96.47 (± 0.59)	96.35 (± 0.50)	95.77 (± 0.71)
GPEth	108.72 (± 0.35)	108.20 (± 0.35)	108.72 (± 0.28)	109.36 (± 0.60)	97.36 (± 7.13)	109.69 (± 32.53)	110.43 (± 0.50)	110.04 (± 1.06)	109.81 (± 0.36)	108.73 (± 0.92)
GPGro	124.20 (± 0.97)	123.76 (± 1.02)	119.56 (± 1.01)	124.69 (± 2.03)	125.84 (± 5.31)	121.53 (± 3.14)	120.00 (± 0.31)	120.03 (± 2.85)	120.87 (± 2.67)	127.20 (± 2.24)
GPIIns	82.24 (± 1.21)	88.80 (± 3.42)	80.55 (± 2.65)	88.05 (± 3.25)	92.06 (± 2.02)	81.06 (± 2.40)	84.26 (± 1.81)	85.24 (± 4.14)	90.56 (± 2.16)	86.07 (± 2.39)
GPSer	96.68 (± 0.59)	95.85 (± 0.33)	95.92 (± 0.62)	97.30 (± 0.43)	95.06 (± 1.70)	97.83 (± 22.32)	96.80 (± 22.09)	96.52 (± 22.01)	97.00 (± 22.13)	96.68 (± 22.05)
MG	77.42 (± 17.68)	73.30 (± 16.58)	78.09 (± 17.71)	78.43 (± 17.77)	83.27 (± 18.95)	79.90 (± 1.88)	83.21 (± 1.26)	80.59 (± 2.78)	78.74 (± 0.90)	78.74 (± 1.52)
TG	116.88 (± 1.35)	118.43 (± 2.36)	118.00 (1.16)	114.93 (± 0.62)	118.19 (± 0.54)	118.75 (± 1.18)	120.76 (± 1.65)	125.55 (± 2.90)	118.22 (± 0.91)	122.29 (± 1.91)


848
849









- Origin**
- Natural
 - Agave
 - Flor
 - ★ Commercial strain
 - ▲ Clinical
 - Sake
 - Beer
 - ★ Bioethanol
 - Commercial wine
 - Chicha
 - Cider
 - ★ Cocoa
 - Cachaça
 - Wine
 - Masato





















PC2 (24.1% explained var.)

