Supporting Information

Disulfide Based Protecting Groups for the Cysteine Side Chain

Amit Chakraborty¹, Anamika Sharma^{1,2}, Fernando Albericio^{1,3,4*}, Beatriz G. de la Torre^{2*}

¹Peptide Science Laboratory, School of Chemistry and Physics, University of KwaZulu-Natal, Durban 4001, South Africa

²KwaZulu-Natal Research Innovation and Sequencing Platform (KRISP), School of Laboratory Medicine and Medical Sciences, College of Health Sciences, University of KwaZulu-Natal, Durban 4041, South Africa

³Institute for Advanced Chemistry of Catalonia (IQAC-CSIC), 08034 Barcelona, Spain

⁴CIBER-BBN, Networking Centre on Bioengineering, Biomaterials and Nanomedicine, and Department of Organic Chemistry, University of Barcelona, 08028 Barcelona, Spain

1. General

All solvents and reagents used in the experiments were bought from commercial suppliers and being used further without any purification unless otherwise mentioned. ¹H NMR and ¹³C NMR analysis were carried out using Bruker AVANCE III 400 MHz and 600 MHz spectrometer. The chemical shifts were showed in ppm (parts per million). For Analytical HPLC, Agilent 1100 system was used with Phenomenex AerisTMC18 column (3 μ m, 4.6 $\times$ 150 mm) (flow rate 1.0 mL/min and UV detection at 220 nm) and the data obtained thus were processed by Chemstation software over a 5-95 % gradient of MeCN (0.1% TFA)/H₂O (0.1% TFA) for 15 min unless otherwise mentioned. All mass spectrometry data were obtained from a Thermo Fisher Scientific UltiMate 3000 UHPLC-ISQTM EC single quadrupole mass spectrometer in positive ion mode over a 5-95 % gradient of MeCN (0.1% HCOOH)/H₂O (0.1% HCOOH) for 15 min if not otherwise specified. High Resolution Mass Spectrometry (HRMS) data were recorded using a Bruker ESI-QTOF mass spectrometer in positive ion mode. For High Efficiency Solid Phase Peptide Synthesis (HE-SPPS), Liberty Blue, CEM peptide synthesizer was used.

1.1 Solid-Phase Peptide Synthesis

All peptides were synthesized following standard Fmoc/*t*Bu based solid phase synthesis protocol (SPPS). Fmoc Rink Amide AM resin (0.74 mmol/g) was used as solid support for the peptides. Initially, the resin was washed using DMF (3×1 min), DCM (3×1 min) and DMF (3×1 min). Deprotection of Fmoc group was achieved by treatment of the resin with 20% piperidine/DMF (1×1 min and 1×6 min) followed by washing with DMF. The protected Fmoc-amino acids (3 eq.) were incorporated using DIC (3 eq.) and OxymaPure (3 eq.) in DMF, as coupling reagents, for 30-40 min at rt. This was repeated until the final peptide was achieved. Fmoc from the last coupled amino acid was removed as explained above. In case of acetylation in the *N*-terminal of the peptide, this was achieved by treating the peptidyl resin with acetic anhydride (10 eq.) and DIEA (20 eq.) in DMF for 30 min at rt followed by washing with DMF (3×1 min), DCM (2×1 min). Cleavage was performed after drying the peptidyl resin by treating with TFA/TIS/H₂O (95:2.5:2.5) for 1 h at rt. The cleavage mixture was then precipitated with Et₂O, centrifuged and the pellet was re-dissolved in H₂O/MeCN (1:1) for analysis by HPLC and LCMS.

2. LCMS characterization of all synthesized peptides

Serial number	Peptide	LCMS: m/z calcd.	LCMS: m/z obtained
1	H-Cys(Npys)-Gly-Phe-Leu-NH ₂	592.20 [M+H] ⁺	592.22 [M+H] ⁺
2	H-Cys(SIT)-Gly-Phe-Leu-NH ₂	540.27 [M+H] ⁺	540.31 [M+H] ⁺
3	H-Cys(MOT)-Gly-Phe-Leu-NH ₂	554.25 [M+H] ⁺	554.27 [M+H] ⁺
4	Ac-Ala-Cys(Stbu)-Leu-NH ₂	435.21 [M+H] ⁺	435.26 [M+H] ⁺
5	Ac-Asn-Cys(Stbu)-Asn-NH ₂	479.17 [M+H] ⁺	479.31 [M+H] ⁺
6	Ac-Ala-Cys(SIT)-Leu-NH ₂	449.23 [M+H] ⁺	449.45 [M+H] ⁺
7	Ac-Asn-Cys(SIT)-Asn-NH ₂	493.19 [M+H] ⁺	493.38 [M+H] ⁺
8	Ac-Ala-Cys(MOT)-Leu-NH ₂	463.20 [M+H] ⁺	463.27 [M+H] ⁺
9	Ac-Asn-Cys(MOT)-Asn-NH ₂	507.17 [M+H] ⁺	507.20 [M+H] ⁺
10	H-Cys(Stbu)-Tyr-Phe-Gln-Asn-Cys(Stbu)-Pro-Arg-Gly-NH ₂	632.26 [M+2H] ⁺ / 2 & 421.51 [M+3H] ⁺ / 3	632.48 [M+2H] ⁺ / 2 & 421.79 [M+3H] ⁺ / 3
11	H-Cys(SIT)-Tyr-Phe-Gln-Asn-Cys(SIT)-Pro-Arg-Gly-NH ₂	645.79 [M+2H] ⁺ / 2 & 430.86 [M+3H] ⁺ / 3	645.97 [M+2H] ⁺ / 2 & 431.25 [M+3H] ⁺ / 3
12	H-Cys(MOT)-Tyr-Phe-Gln-Asn-Cys(MOT)-Pro-Arg-Gly-NH ₂	659.77 [M+2H] ⁺ / 2 & 440.18 [M+3H] ⁺ / 3	660.30 [M+2H] ⁺ / 2 & 440.50 [M+3H] ⁺ / 3

3. Synthesis of Boc-Cys(PG)-Gly-Phe-Leu-NH₂

Peptide **Boc-Cys(Npys)-Gly-Phe-Leu-NH₂** was synthesized following the SPPS strategy as explained above. The synthesized peptide was confirmed by HPLC (for purity) and LCMS (for mass).

Then, portions of peptidyl resin (50 mg in each case) were taken in two reaction vessels separately and treated with 10 eq of *sec*-isoamyl mercaptan (**SIT**) in DMF and 10 eq of 2-methyloxolane-3-thiol (**MOT**) in DMF respectively for 5 min followed by washing with DMF (3 × 1 min) and DCM (2 × 1 min) to afford peptidyl resins of **Boc-Cys(SPG)-Gly-Phe-Leu-NH₂** (SPG = SIT and MOT). The resins upon drying were cleaved and processed as described above to obtain the crude peptides which were analysed and characterized by HPLC and LCMS. Later, both peptidyl resins were subjected to dithiol reduction study with β-mercaptoethanol (BME) and base stability study.

For the reduction by BME study, the peptidyl resins were treated three times for 5 min with solution consisting of BME in DMF (1:4) and 0.1 M DIEA. For the base stability study, the

peptidyl resins were treated with 20% piperidine in DMF (1 mL) for 2 h. The resins were then washed, dried, and cleaved. The HPLC analysis are shown in Fig. SI 1A for SIT and Fig. SI 1B for MOT

Figure SI 1: HPLC chromatograms of H-C(SIT)GFL-NH₂ (A), H-C(MOT)GFL-NH₂ (B) and subsequent thiol lability and base stability study.

4. Synthesis of Fmoc-Cys(SPG)-OH

Synthesis of Fmoc-Cys-OH (Scheme S1). The synthesis was carried out firstly by Fmoc protection of L-cystine as reported in literature with slight modification¹. Briefly, to an aqueous solution of L-cystine (1.0 g, 4.1 mmol), Fmoc-Amox (3.4 g, 10.3 mmol), dissolved in 50 mL of acetone was added dropwise and stirred at rt for 7 h while maintaining the pH 9-9.5 by addition of aqueous K₂CO₃ (0.2 M). After completion of reaction, the mixture was concentrated under reduced pressure to remove acetone followed by extraction with DCM (3 x 100 mL) until the unreacted Fmoc-Amox has been removed (as confirmed by TLC). The aqueous layer was then acidified with 6M HCl which precipitates bis-Fmoc-Cystine. The precipitate was then filtered and dried to be used for next step directly. Bis Fmoc-cystine was dissolved in a mixture of H₂O (105 mL) and MeCN (255 mL) followed by addition of 2M HCl and Zn powder (3 g, 46 mmol) for reduction at rt². The reaction was completed instantaneously. Zn was removed by filtration using celite. 6M HCl was added to the filtrate until precipitation was observed which upon filtration and drying afforded Fmoc Cysteine (overall yield, 69 %).

Scheme SI-1: Synthesis of Fmoc-Cysteine

General protocol for synthesis of Fmoc-Cys(SPG)-OH

To a solution of NCS (N-chlorosuccinimide) (0.8 g, 6.0 mmol) in DCM (22 mL) at -50 °C, was added thiol (5.8 mmol). The reaction was stirred at same temperature for about 15 min (SIT) or 30 min (MOT). The mixture was quickly poured into a stirring solution of Fmoc-Cys-OH (1.0 g, 2.9 mmol) in THF (25 mL) at -50 °C. The whole mixture was stirred for next 2 h maintaining the temperature at -50 °C. The reaction was then allowed to come to rt. The reaction mixture was then washed with acidified water (3 times). The organic layer was collected over MgSO₄, filtered and evaporated under reduced pressure. The crude was then purified using silica gel column chromatography, using n-Hexane/EtOAc as mobile phase to afford pure product.

Scheme SI-2: Synthesis of Fmoc-Cys(SPG)-OH where SPG = SIT or MOT

Fmoc-Cys(SIT)-OH

Yield = 0.68 g (52.5 %); White solid; **HPLC** [5-95 % of MeCN (0.1% TFA/ H₂O (0.1 %TFA) over 15 mins] t_R = 12.8 min; **¹H NMR** (600 MHz, CDCl₃): 7.31-7.76 (m, Ar-H, 8H), 5.72 (m, 1H), 4.73 (m, CH, 1H), 4.40 (d, J = 7.2 Hz, CH₂, 2H), 4.24 (t, J = 7.2 Hz, CH, 1H), 3.22 (m, CH₂, 1H), 3.15 (m, CH₂, 1H), 2.82

(m, CH, 1H), 1.95 (m, CH, 1H), 1.25 (m, CH₃, 3H), 0.95 (m, 2CH₃, 6H),; ¹³C NMR (150 MHz, CDCl₃): 175, 156, 143.7, 141.3, 127.8, 127.1, 125.2, 120, 67.5, 53.5, 47.1, 40.6, 32, 20.3, 18.1, 16.4; **HRMS**: m/z calcd. for C₂₃H₂₈NO₄S₂: [M+H]⁺ 446.1454; found 446.1545.

Fig SI 2: HPLC Chromatogram of **Fmoc-Cys(SIT)-OH** (inset: HRMS spectra)

Fig SI 3: ^1H NMR of Fmoc-Cys(SIT)-OH (400MHz, in CDCl_3)

Fig SI 4: ^{13}C NMR of Fmoc-Cys(SIT)-OH (100MHz in CDCl_3)

Fig SI 5: ^{13}C NMR (DEPT 135) of Fmoc-Cys(SIT)-OH (100MHz in CDCl_3)

Fmoc-Cys(MOT)-OH

Yield = 0.8 g (60.1 %); Off-white semi-solid; HPLC [5-95 % of MeCN (0.1% TFA/ H_2O (0.1 %TFA) over 15 mins] t_R = 10.7 and 10.8 min; ^1H NMR (400 MHz, CDCl_3): 7.31-7.76 (m, Ar-H, 8H), 5.88 (m, 1H), 4.70 (s, CH, 1H), 4.41 (m, CH_2 , 2H), 4.23 (m, CH, 1H), 4.11 (m, CH, 1H), 3.98 (m, CH_2 , 1H), 3.73 (m, CH_2 , 1H), 3.51 (m, CH_2 , 1H), 3.30 (m, CH_2 , 1H), 3.16 (m, CH, 1H), 2.35 (m, CH_2 , 1H), 2.10 (m, CH_2 , 1H), 1.28 (m, CH_3 , 3H); ^{13}C NMR (100 MHz, CDCl_3): 178.3, 156, 143.7, 141.3, 127.8, 127.1, 125.2, 120, 77.3, 67.4, 66, 54.3, 53.4, 47.1, 40.6, 33, 16.6; **HRMS:** m/z calcd. for $\text{C}_{23}\text{H}_{26}\text{NO}_5\text{S}_2$: $[\text{M}+\text{H}]^+$ 460.1247; found 460.1339.

Fig SI 6: HPLC Chromatogram of Fmoc-Cys(MOT)-OH (inset: HRMS spectra)

Fig SI 7: ¹H NMR of Fmoc-Cys(MOT)-OH (400MHz, in CDCl₃)

Fig SI 8: ^{13}C NMR of Fmoc-Cys(MOT)-OH (100MHz in CDCl_3)

Fig SI 9: ^{13}C NMR (DEPT 135) of Fmoc-Cys(MOT)-OH (100MHz in CDCl_3)

5. Peptide synthesis and Thiol lability study

5(A). Synthesis of Ac-Ala-Cys(SP_G)-Leu-NH₂ and Ac-Asn(Trt)-Cys(SP_G)-Asn(Trt)-NH₂

Fig SI 10: HPLC chromatograms of Ac-AC(SP_G)L-NH₂ (panel A) and Ac-NC(SP_G)N-NH₂ (panel B); SP_G = StBu (top row); SIT (middle row) and MOT (bottom row)

Reduction of disulfide by DTT:

The peptidyl resins Ac-Ala-Cys(SP_G)-Leu-NH-Rink amide resin and Ac-Asn(Trt)-Cys(SP_G)-Asn(Trt)-NH-Rink amide resin (SP_G = StBu, SIT and MOT) were treated with 5 eq. of DTT (1,4-dithiothreitol) in a solution of DMF /DIEA/H₂O (95:2.5:2.5) three times for 10 min each. Then, the resins were washed, dried and the peptides were cleaved. The crudes obtained were analysed by HPLC and LCMS (Fig. SI 11-16)

Figure SI 11: HPLC Chromatogram after deprotection of **SIT** from **Ac-AC(SIT)L-NH₂**
(LCMS: m/z calcd. [M+H]⁺ 347.17 found [M+H]⁺ 347.27)

Figure SI 12: HPLC Chromatogram after deprotection of **MOT** from **Ac-AC(MOT)L-NH₂**
(LCMS: m/z calcd. [M+H]⁺ 347.17 found [M+H]⁺ 347.25)

Figure SI 13: HPLC Chromatogram after deprotection of **StBu** from **Ac-AC(StBu)L-NH₂**

Figure SI 14: HPLC Chromatogram after deprotection of **SIT** from **Ac-NC(SIT)N-NH₂** (LCMS: m/z calcd. [M+H]⁺ 391.14 found [M+H]⁺ 391.20); [HPLC using 0-30 % gradient of MeCN (0.1% TFA)/H₂O (0.1% TFA) for 15 min]

Figure SI 15: HPLC Chromatogram after deprotection of **MOT** from **Ac-NC(MOT)N-NH₂** (LCMS: m/z calcd. [M+H]⁺ 391.14 found [M+H]⁺ 391.38); [HPLC using 0-20 % gradient of MeCN (0.1% TFA)/H₂O (0.1% TFA) for 15 min]

Figure SI 16: HPLC Chromatogram after deprotection of **StBu** from **Ac-NC(StBu)N-NH₂** [HPLC using 0-20 % gradient of MeCN (0.1% TFA)/H₂O (0.1% TFA) for 15 min]

5(B). Linear Vasopressin synthesis by Peptide synthesizer

Linear vasopressin using Fmoc-Cys(SP_G)-OH (SP_G = *St*Bu, SIT) were synthesized following HE-SPPS microwave assisted protocol taking Fmoc-Rink-Amide AM resin as solid support (0.05 mmol, 0.74 mmol/g). The protocol involved 2 min of coupling (Fmoc amino acid, DIC and Oxyma pure, all 5 eq.) and 1 min of deprotection (20% piperidine), both at 90 °C with no washing between coupling and deprotections steps. After the synthesis, the peptidyl resin was dried and cleaved using general method described above and analysed by HPLC and LCMS (Fig SI 17-18).

Figure SI 17: HPLC Chromatogram of linear Vasopressin synthesized using Fmoc-Cys(*St*Bu)-OH (LCMS: *m/z* calcd. $[M+2H]^+/2$ 632.26 and $[M+3H]^+/3$ 421.84; found $[M+2H]^+/2$ 631.94 and $[M+3H]^+/3$ 422.04)

Figure SI 18: HPLC Chromatogram of linear Vasopressin synthesized using Fmoc-Cys(SIT)-OH (LCMS: m/z calcd. $[M+2H]^+/2$ 645.79 and $[M+3H]^+/3$ 430.86; found $[M+2H]^+/2$ 646.12 and $[M+3H]^+/3$ 431.26)

References

- 1) Kumar, A.; Sharma, A.; Haimov, E.; El-Faham, A.; de la Torre, B. G.; Albericio, F. *Org. Process Res. Dev.* **2017**, *21*, 1533-1541.
- 2) Erlandsson, M.; Hällbrink, M. *Int. J. Pept. Res. Ther.* **2005**, *11*, 261-265.