



# Mercury Methylating Microbial Community Structure in Boreal Wetlands Explained by Local Physicochemical Conditions

Jingying Xu<sup>1</sup>, Van Liem-Nguyen<sup>2</sup>, Moritz Buck<sup>1,3</sup>, Stefan Bertilsson<sup>1,3</sup>, Erik Björn<sup>2</sup> and Andrea G. Bravo<sup>1,4\*</sup>

<sup>1</sup>Department of Ecology and Genetics, Limnology, Uppsala University, Uppsala, Sweden, <sup>2</sup>Department of Chemistry, Umeå University, Umeå, Sweden, <sup>3</sup>Swedish University of Agricultural Sciences, Uppsala, Sweden, <sup>4</sup>Institute of Marine Sciences, Spanish National Research Council, Spain

## OPEN ACCESS

### Edited by:

Vera I. Slaveykova,  
Université de Genève, Switzerland

### Reviewed by:

Tamar Barkay,  
Rutgers, The State University of New  
Jersey, United States  
Caitlin Marissa Gionfriddo,  
Oak Ridge National Laboratory,  
United States  
Cynthia C. Gilmour,  
Smithsonian Environmental  
Research Center, United States

### \*Correspondence:

Andrea G. Bravo  
jandriugarcia@gmail.com

### Specialty section:

This article was submitted to  
Freshwater Science,  
a section of the journal  
Frontiers in Environmental Science

**Received:** 09 December 2019

**Accepted:** 21 December 2020

**Published:** 18 February 2021

### Citation:

Xu J, Liem-Nguyen V, Buck M,  
Bertilsson S, Björn E and Bravo AG  
(2021) Mercury Methylating Microbial  
Community Structure in Boreal  
Wetlands Explained by Local  
Physicochemical Conditions.  
*Front. Environ. Sci.* 8:518662.  
doi: 10.3389/fenvs.2020.518662

The potent neurotoxin methylmercury (MeHg) is a major concern due to its negative effects on wildlife and human health. Boreal wetlands play a crucial role in Hg cycling on a global scale, and therefore, it is crucial to understand the biogeochemical processes involved in MeHg formation in this landscape element. By combining high-throughput *hgcA* amplicon sequencing with molecular barcoding, we reveal diverse clades of potential Hg<sup>II</sup> methylators in a wide range of wetland soils. Among Bacteria, Desulfuromonadota (14% of total reads), Desulfurobacterota\_A, and Desulfurobacterota (up to 6% of total reads), previously classified as Deltaproteobacteria, were important members of the *hgcA*+ microbial community in the studied wetlands. We also identified Actinobacteriota (9.4% of total reads), Bacteroidota (2% of total reads), and Firmicutes (1.2% of total reads) as members of the *hgcA*+ microbial community. Within Archaea, Methanosarcinales represented up to 2.5% of the total reads. However, up to half of the *hgcA*+ community could not be resolved beyond domain *Bacteria*. Our survey also shows that local physicochemical conditions, such as pH, nutrient concentrations, water content, and prevailing redox states, are important for shaping the *hgcA*+ microbial community structure across the four studied wetlands. Furthermore, we observed a significant correlation between Hg<sup>II</sup> methylation rate constants and the structure of the *hgcA*+ microbial community. Our findings expand the current knowledge on the *hgcA*+ microbial community composition in wetlands and the physicochemical factors underpinning spatial heterogeneity in such communities.

**Keywords:** wetlands, methylmercury, mercury methylation, *hgcA*, community composition, bacteria, mercury, drivers

## 1 INTRODUCTION

Wetlands cover 6% of the world's land surface and contain about 12% of the global carbon pool (Erwin, 2009). Hence, these systems play an important role in different global biogeochemical cycles, such as that of carbon and mercury (Hg) (Mitra et al., 2003; Tjerngren et al., 2012b). For example, wetlands take up and store carbon and release carbon dioxide and methane (Mitra et al., 2003). Also, wetlands have been identified as important sources or even sinks of the toxic methylmercury (MeHg)

(St. Louis et al., 1994; Hall et al., 2008; Tjerngren et al., 2012b). St. Louis et al. (1994) found 26–79 times higher yields of MeHg from wetland portions of catchments than from purely upland areas. Mass-balance estimates have indicated that purely upland catchments and lakes are sites of MeHg retention or demethylation (St. Louis et al., 1994), and catchments with wetland areas are sites of net MeHg production (St. Louis et al., 1994; Mitchell et al., 2008b; Hall et al., 2008). It is also shown that the creation of wetlands by the flooding of boreal soils led to increased MeHg production (Gilmour et al., 1998; Hall et al., 2008; Mitchell et al., 2008a; Mitchell et al., 2008b). The wetland type, percentage wetland area, or annual water yield appear to have a consistent effect on the magnitude of MeHg production (Louis et al., 1996). Furthermore, because wetlands are parts of large watersheds and thereby are hydrologically connected to adjacent systems, there is a high risk for MeHg that is produced in wetlands to be transported to downstream waters. This is critical as MeHg bioaccumulates and biomagnifies along the aquatic food webs, and this poses severe risks to the environment and humans (Boening, 2000). Therefore, an understanding of the potential MeHg formation in boreal wetlands is crucial for robust risk assessment and rational landscape management.

The methylation of  $\text{Hg}^{\text{II}}$  to MeHg in the environment is governed by two factors: the activities of microbial communities (Bravo et al., 2017) and the availability of  $\text{Hg}^{\text{II}}$  for use by these methylators (Schaefer and Morel, 2009; Jonsson et al., 2014; Mazrui et al., 2016). Some members of sulfate-reducing bacteria (SRB) (Gilmour et al., 1992; King et al., 2001; Gilmour et al., 2013), iron-reducing bacteria (FeRB) (Fleming et al., 2006; Kerin et al., 2006; Bravo et al., 2018b), methanogens (Hamelin et al., 2011; Yu et al., 2013), *Firmicutes* (Gilmour et al., 2013), acetogens, and recently also obligate fermenting lineages (McDaniel et al., 2020b; Peterson et al., 2020) have so far been implicated in  $\text{Hg}^{\text{II}}$  methylation. The identification of two genes (*hgcAB*) essential for  $\text{Hg}^{\text{II}}$  methylation (Parks et al., 2013) provides the means to more directly characterize the diversity of potential  $\text{Hg}^{\text{II}}$  methylators in contrasting environmental niches, such as soils (Liu et al., 2014; Liu et al., 2018; Vishnivetskaya et al., 2018; Xu et al., 2019), lakes, reservoirs (Bravo et al., 2018a; Bravo et al., 2018b; Jones et al., 2019; Jones et al., 2020), coastal sediments, animal guts, extremes of pH, salinity environments (Gilmour et al., 2013), the global ocean (Podar et al., 2015; Villar et al., 2020), Baltic Sea (Capo et al., 2020), and even in the Southern Ocean (Gionfriddo et al., 2016). However, for most of the recently discovered microorganisms carrying the *hgcA* gene (*hgcA+*), their role in the environment and their capacity to actually methylate  $\text{Hg}^{\text{II}}$  has yet to be demonstrated. Progress in that direction has been made by testing the capacity to methylate  $\text{Hg}^{\text{II}}$  of *hgcA+* isolates (Gilmour et al., 2018; Yu et al., 2018) or by amendment experiments with specific substrates or inhibitors (Schaefer et al., 2014; Christensen et al., 2017; Bravo et al., 2018a; Schaefer et al., 2020). However, although numerous studies show that *hgcA+* microorganisms are present and potentially active in a wide range of environments, the ecological underpinnings of the

*hgcA+* microbial community remains to be elucidated across systems. In lakes, previous studies reveal the composition of organic matter and the structure on the whole microbial community (Bravo et al., 2018a) as important for the *hgcA+* microbial community structure. However, still little is known on the factors influencing the structure and function of *hgcA+* microbial communities in the environment or the effect of the *hgcA+* microbial community composition on  $\text{Hg}^{\text{II}}$  methylation rate constants. Jones et al. (2020) amended freshwater lake sediments with different sulfate concentrations and observed that the experimental plots with the lowest sulfate concentrations presented the highest *hgcA+* diversity and the highest  $\text{Hg}^{\text{II}}$  methylation rate, highlighting that diverse *hgcA+* communities that do not reduce sulfate can methylate  $\text{Hg}^{\text{II}}$  as effectively as communities dominated by sulfate-reducing populations (Jones et al., 2020). Although this study emphasizes that the *hgcA+* microbial community structure is relevant for the  $\text{Hg}^{\text{II}}$  methylation rate, further evidence is needed to confirm the effect of compositional changes of *hgcA+* communities in  $\text{Hg}^{\text{II}}$  methylation rates.

In Swedish boreal wetlands, some studies have described the complex geochemical parameters determining  $\text{Hg}^{\text{II}}$  chemical speciation (Liem-Nguyen et al., 2017; Skyllberg, 2008),  $\text{Hg}^{\text{II}}$  methylation (Liem-Nguyen et al., 2021), and the organisms potentially involved in the process (Tjerngren et al., 2012b; Schaefer et al., 2014; Schaefer et al., 2020). A recent study performed a comprehensive thermodynamic speciation model, which provided new insights into the speciation of  $\text{Hg}^{\text{II}}$  in boreal wetland porewaters but did not demonstrate any clear relationship between  $\text{Hg}^{\text{II}}$  methylation rates and the calculated chemical speciation (Liem-Nguyen et al., 2021). In contrast, significant positive relationships were observed between  $\text{Hg}^{\text{II}}$  methylation rates and the sum of low molecular mass thiol compounds of biological origin. Although the authors could not conclude whether low molecular mass thiols might kinetically control the size and composition of the  $\text{Hg}^{\text{II}}$  pool available for microbial uptake and/or whether low molecular mass thiols were produced by  $\text{Hg}^{\text{II}}$  methylating microorganisms, their findings underline the role of microbial communities in determining  $\text{Hg}^{\text{II}}$  methylation rates. Knowledge about  $\text{Hg}^{\text{II}}$  methylating microbial composition, the factors controlling their activity, and their effects on  $\text{Hg}^{\text{II}}$  methylation rates will, thus, help us obtain a mechanistic understanding of these important ecosystem-level processes. The primary goal of the present paper is to expand on this and, more specifically, 1) map  $\text{Hg}^{\text{II}}$  methylating microbial communities in four boreal wetlands in Sweden, 2) identify geochemical characteristics important for shaping these communities, and 3) determine the interplay between *hgcA+* microbial communities and  $\text{Hg}^{\text{II}}$  methylation rates. For this purpose, high-throughput Illumina sequencing of amplified *hgcA* genes combined with molecular barcoding and detailed geochemical characteristics were carried out to study *hgcA+* microbial communities in 81 samples from boreal wetlands and, in this way, shed light on the organisms responsible for MeHg formation in the boreal landscape.

## 2 MATERIALS AND METHODS

### 2.1 Site Description

Four boreal wetlands situated within predominantly forested watersheds in Sweden were selected for this study. These systems cover variable biogeochemical features with regards to climate, acidity, organic carbon, sulfide concentration, and nutrient status as well as ancillary chemistry. The selected wetlands were categorized as 1) a northern group of poor-fen type of wetlands Storkälsmyran (SKM) and Kroksjön (KSN), the latter with some open water, and 2) a southern group of bog-fen sites, including Långedalen (LDN) and the more nutrient-rich southern wetland Gästern (GTN). A brief map of sampling sites within the wetlands and more detailed wetland descriptions can be found in Tjerngren et al. (2012a), Liem-Nguyen et al. (2017), and Liem-Nguyen et al. (2021).

### 2.2 Soil and Porewater Sampling

In total, 81 soil samples were collected from 17 sites across four boreal wetlands at 5 cm intervals starting from the top of the water table to 25 cm below this interface. The top few centimeters of the dry topsoil above the water table was discarded. On average, five soil profiles (0–25 cm) were collected at each site to probe the spatial variability within the sites. Soil samples were preserved in an icebox in the field prior to being transferred to the lab within 1 h. Once in the laboratory, an aliquot was immediately frozen at  $-80^{\circ}\text{C}$  for DNA extraction. Porewater samples were collected from the same depths as the soil samples using a custom device consisting of a vacuum pump and a perforated Teflon probe. During sampling,  $\text{N}_2$  gas was flushed through the tubing and bottle to keep the system anoxic.

Ziploc plastic bags were used to contain soil samples and were squeezed to minimize the remaining air in the bags and, hence, avoid further oxidation. Soil and porewater samples were handled inside a glovebox filled with  $\text{N}_2$  within 1 h of sampling at a field laboratory station located walking distance from the site. Within 1 h, porewater samples were vacuum filtrated through a  $0.22\ \mu\text{m}$  membrane filter (500 ml funnel filter, Sarstedt). More details on the sampling strategy can be found in Liem-Nguyen et al. (2017) and Liem-Nguyen et al. (2021).

### 2.3 Chemical Analyses

All the soil samples were analyzed for pH, water content, total Hg (THg), and MeHg. THg in soils was determined by solid combustion atomic absorption spectrometry using an AMA-254 Advanced Mercury Analyzer (LECO), and MeHg was acid extracted, ethylation derivatized, and determined by isotope dilution analysis (IDA) using gas chromatography inductively coupled plasma mass spectrometry (ICPMS) (Lambertsson et al., 2001). A detailed geochemical characterization of the studied wetland soils and porewater is presented in **Supplementary Tables S1, S2** and in Liem-Nguyen et al. (2021).

The  $\text{Hg}^{\text{II}}$  methylation potential ( $k_m$ ,  $\text{d}^{-1}$ ) was determined using two different enriched Hg isotope tracers, referred to as  $^{198}\text{Hg}^{\text{II}}(\text{aq})$  and  $^{200}\text{Hg}^{\text{II}}\text{-NOM}(\text{ads})$ . Detailed information on the preparation of the enriched Hg stable isotopes can be found in Liem-Nguyen et al. (2021). THg in porewater was measured by

IDA using Hg cold vapor generation coupled to ICPMS. MeHg in porewater was analyzed by IDA using purge-and-trap thermal desorption coupled to gas chromatography ICPMS (Lambertsson and Björn, 2004). Dissolved organic carbon was measured using a TOC-VCPH (Shimadzu), and  $\text{Fe}^{\text{II}}$  and  $\text{Fe}^{\text{III}}$  were determined by UV absorption spectrophotometry (Viollier et al., 2000). The concentrations of  $\text{SO}_4^{2-}$ ,  $\text{Cl}^-$ ,  $\text{PO}_4^{3-}$ ,  $\text{NH}_4^+$ , and  $\text{NO}_3^-$  in porewater were measured by ion chromatography. The  $\text{Hg}^{\text{II}}$  concentrations in porewater were calculated by subtracting MeHg from THg. Total thiol in freeze-dried porewater were determined using Sulfur K-edge X-ray absorption near edge structure spectroscopy (S K-edge XANES) and Hg LIII-edge extended X-ray absorption fine structure (EXAFS) spectroscopy as described by Liem-Nguyen et al. (2017). Concentrations of 16 low molecular mass (LMM) thiols were determined in porewater using solid-phase extraction online preconcentration hyphenated to liquid chromatography tandem mass spectrometry (SPE/LC-MS/MS) (Liem-Nguyen et al., 2015).

The certified reference materials MESS-3 ( $0.091 \pm 0.009\ \text{mg Hg/kg}$ ), ORMS-5 ( $26.2 \pm 1.3\ \text{pg Hg/g}$ ), and ERM-CC580 ( $0.075 \pm 0.004\ \text{mg/kg}$ ) were used for quality control of the total Hg and MeHg measurements in soil and porewater samples. The measured concentrations were in good agreement (95%–105%) with the reference materials. The chemical characteristics of the wetland sites can be also found elsewhere (Liem-Nguyen et al., 2017; Liem-Nguyen et al., 2021).

### 2.4 Microbiological Analyses

DNA was isolated from 0.2 g soil (wet weight) using the PowerSoil DNA isolation kit (Mo Bio Laboratories, Carlsbad, CA, United States). To target the functional gene *hgcA*, which plays an essential role in  $\text{Hg}^{\text{II}}$  methylation, previously published *hgcA* primers (*hgcA*<sub>261F</sub> = CGGCATCAAYGTCTGGTGYGC, *hgcA*<sub>912R</sub> = GGTGTAGGGGGTGCAGCCSGTRWARKT) (Schaefer et al., 2014) were adapted for high-throughput Illumina sequencing. Accordingly, HPLC-purified primers linked with generic adaptors were used for a first stage PCR. In a second stage PCR, primers binding to the generic adaptors but also featuring standard Illumina adaptors and sample-specific barcodes were used (*Multiplex\_fwd* = AATGATACGGCGACCACCGAGA {TCTACAC}-[i5 barcode] AACTCTTTCCCTACACGACG, *Multiplex\_rev* = CAAGCAG AAGACGGCATACGAGAT-[i7 barcode] GTGACTGGAGTTCAG ACGTGTGCTCTCCGATCT), enabling us to include all the first stage PCR products on a single lane for parallelized Illumina sequencing.

The *hgcA* was first amplified in 50  $\mu\text{L}$  volume with 1x Phusion GC Buffer, 0.2 mM dNTP mix, 5% DMSO, 0.1  $\mu\text{M}$  of each adaptor-linked primer, 7  $\mu\text{g}/\mu\text{L}$  BSA, 4  $\mu\text{L}$  extracted DNA template, and 1.0 U Phusion high fidelity DNA polymerase (NEB, United Kingdom) for an initial denaturation of 2 min at  $98^{\circ}\text{C}$  followed by 35 amplification cycles (10 sat  $96^{\circ}\text{C}$ , 30 sat  $56.5^{\circ}\text{C}$ , and 45 sat  $72^{\circ}\text{C}$ ) and a final extension at  $72^{\circ}\text{C}$  for 7 min. Following this initial step, a second PCR was conducted to add sample-specific molecular barcodes. Reactions were carried out in 20- $\mu\text{L}$  volumes using 1x Q5 reaction buffer, 0.2 mM dNTP mix, 0.1  $\mu\text{M}$  barcode primers, AMPure XP-

purified first stage PCR products, and 1.0 U Q5 high-fidelity DNA polymerase (NEB, United Kingdom) for an initial denaturation of 30 s at 98°C followed by 18 cycles (10 s at 98°C, 30 s at 66°C, and 30 s at 72°C) and a final extension at 72°C for 2 min. Amplicons of *hgcA* genes were assessed by gel electrophoresis and GelRed visualization on a 1% agarose gel (Invitrogen, United States) prior to a final purification by Agencourt AMPure XP (Beckman Coulter, United States) after both steps of PCR. Quantification of purified amplicons from the second stage PCR was performed using the PicoGreen kit (Invitrogen).

Subsequent amplicon sequencing and sequence processing was carried out, largely following the protocol described in Sinclair et al. (2015) using the MiSeq instrument using illumina MiSeq sequencing performed by the SNP/SEQ SciLifeLab facility hosted by Uppsala University using 2x300bp chemistry. Chimera identification and operational taxonomic unit (OTU) clustering at 3% sequence dissimilarity was done using UPARSE. Only the forward-read sequences were used in data analysis as PCR products were too short to obtain robustly overlapping forward and reserve reads. Low-quality sequences were filtered and trimmed using SICKLE (Joshi and Fass, 2011), and adaptors were removed by using CUTADAPT (Martin, 2011). Subsequent processing of reads was performed by USEARCH version 8.0, and reads were clustered at 60% identity cutoff using cd-hit-est. We further performed a manual check of the protein sequences using the knowledge from the original paper of Parks et al. (2013) that described the unique motif from *hgcA* [NVWCA(A/G/S)GK] and *hgcB* protein sequences [C(M/I)EC(G/S)(A/G)C]. Last, pplacer (Matsen et al., 2010), and a likelihood cutoff of 0.8 search was used for taxonomical annotation with a manually curated a GTDB-derived database release 89 (McDaniel et al., 2020a).

## 2.5 Statistical Analyses

Microbial community compositions between samples were compared at the OTU level using nonmetric multidimensional scaling (nMDS) based on Bray-Curtis similarities using R 3.3.2 (<https://www.r-project.org/>). Relative dissimilarities (or distances) among the samples were computed according to the resemblance matrix using the vegan package (Oksanen et al., 2017). Information on the common set of samples from community composition based on Bray-Curtis similarities and that from geochemical variables based on Euclidean distance were presented in a single ordination. A combined nMDS plot with bubble and vector plots of geochemical factors projected on the same ordination of community composition was constructed to reveal the relationship between community compositions and correlated geochemical variables (Clarke and Gorley, 2015). Pearson's correlation coefficients (R) between variables were tested using a significance level of alpha <0.05. The false discovery rate of significant correlations was corrected following Benjamini and Hochberg, (1995), and using the *p.adjust* function in R. ANOSIM, tests implemented in the vegan package were used to test for differences in microbial community structure across sites and depths. The function bioenv was used to find the best subset of environmental variables so that the Euclidean distances of scaled environmental variables have the

maximum (rank) correlation with community dissimilarities (Bray-Curtis). Standard Mantel tests were used to investigate correlations between the *hgcA*+ microbial community and the most influential environmental variables. Euclidean distance metrics were used for the most influential environmental variables (squared-root transformed), and Bray-Curtis dissimilarities were used for *hgcA*+ microbial community composition data (square-root transformed). Both distance matrices were calculated with the *vegdist* function. Redundancy analysis (RDA) was performed using the *rda* function in the R package vegan to further explore the relationships between Hg<sup>II</sup>-methylating microbial community composition and the studied environmental factors. RDA seeks a series of linear combinations of the explanatory variables that best explain the variation in the response matrix (i.e., Hg<sup>II</sup>-methylating microorganisms) but requires the number of explanatory variables to be below or equal to the number of observations (i.e., number of wetland soil samples) and the explanatory variables to not be intercorrelated. Intercorrelations between physicochemical variables were inspected by computing the variance inflation factor (VIF). All the variables with a VIF of >10 were excluded from RDA analysis. Prior to this analysis, the variables had been scaled and centered.

## 3 RESULTS

### 3.1 Distribution of *hgcA*+ Microbial Communities Across all Four Wetlands

A total of 4,535,861 high-quality *hgcA* sequences remained after quality control and chimera removal (2664–106,256 reads per sample). The data were rarefied to the remaining least covered sample (2664 sequences). The final combined sequence data set (215,784 reads) were clustered into 618 OTUs using a sequence identity threshold of 60%. In total, 24 taxa at the phyla level, 42 taxa at the class level, 45 taxa at the order level, and 53 taxa at the family level were detected from all the soil samples across the four wetlands. The combined richness detected for random subsets of analyzed samples approached saturation, suggesting that our subsampled data set covered the diverse *hgcA*+ microbial communities inhabiting the wetlands (**Supplementary Figure S1**).

Among Bacteria, Desulfuromonadota (14% of total reads), Desulfuobacterota\_A, and Desulfuobacterota (up to 6% of total reads) were abundant in all four studied sites (**Table 1**). These three phyla, previously classified as Deltaproteobacteria in the NCBI taxonomy, each contain confirmed Hg<sup>II</sup> methylators (Ranchou-Peyruse et al., 2009; Yu et al., 2012; **Table 1**). Within these three phyla, Desulfuromonadales is the most abundant order, representing up to 13.8% of the total reads (**Table 1**). Noteworthy is that, within the Desulfuromonadales order, the Geopsychrobacteraceae family reached 10% of the total reads. Actinobacteriota (Actinobacteria in the NCBI taxonomy) were also quite abundant, representing up to 9.5% of the total *hgcA*+ reads. About 27% of the total sequences could not be taxonomically assigned (Unclassified), and 35% of the reads could not be assigned beyond the rank of Bacteria (Unclassified Bacteria) (**Table 1**). Within the Firmicutes, the Syntrophomonadales and the Desulfitobacteriales orders contributed to 2% of the total reads (**Supplementary Table S3**). Archaea and, in particular, the order Methanosarcinales represented



**TABLE 1** | Relative *hgcA* sequence abundances (%) at levels of phyla for the combined 81-sample data set that includes all four wetlands.

	Northern		Southern		Total community percentage (%)
	SKM	KSN	LDN	GTN	
	Mean Std	Mean Std	Mean Std	Mean Std	
Archaea					
<i>Halobacterota</i>	0.28 ± 0.48	3.33 ± 3.47	3.24 ± 3.85	3.22 ± 3.05	2.6
<i>Thermoplasmatota</i>	0.02 ± 0.05	0.05 ± 0.17	0.03 ± 0.07	0.41 ± 1.59	0.1
Bacteria					
Unclassified Bacteria	51.78 ± 22.05	32.03 ± 18.51	34.11 ± 20.68	21.24 ± 17.86	34.7
<i>Acidobacteriota</i>	0.84 ± 1.51	0.32 ± 0.31	0.79 ± 1.17	0.19 ± 0.20	0.5
<i>Actinobacteriota</i>	5.29 ± 12.06	21.94 ± 20.29	1.99 ± 3.20	9.43 ± 12.32	9.4
<i>Bacteroidota</i>	1.12 ± 2.01	0.94 ± 0.94	3.57 ± 4.98	2.16 ± 2.09	2.0
<i>Chloroflexota</i>	0.01 ± 0.05	0.00 ± 0.01	0.00 ± 0.01	0.00 ± 0.00	0.0
<i>Desulfobacterota</i>	1.97 ± 2.30	0.98 ± 0.72	1.57 ± 2.36	0.23 ± 0.34	1.2
<i>Desulfobacterota_A</i>	8.91 ± 11.55	3.19 ± 5.65	0.99 ± 1.17	6.85 ± 12.46	4.8
<i>Desulfuromonadota</i>	3.86 ± 2.68	15.44 ± 14.10	12.90 ± 11.81	23.31 ± 18.74	13.8
<i>Elusimicrobiota</i>	0.58 ± 1.53	1.22 ± 2.03	0.03 ± 0.05	0.64 ± 1.61	0.6
<i>Fibrobacterota</i>	0.79 ± 1.42	0.19 ± 0.38	0.40 ± 0.48	0.17 ± 0.21	0.4
<i>Firmicutes_A</i>	0.02 ± 0.09	0.02 ± 0.04	0.01 ± 0.02	0.00 ± 0.01	0.0
<i>Firmicutes_B</i>	2.11 ± 3.61	1.09 ± 1.66	0.63 ± 1.52	1.16 ± 2.79	1.2
<i>Firmicutes_C</i>	0.00 ± 0.01	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.0
<i>Goldbacteria</i>	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.02 ± 0.04	0.0
<i>Margulisbacteria</i>	0.09 ± 0.19	0.00 ± 0.01	0.00 ± 0.00	0.00 ± 0.00	0.0
<i>Myxococcota</i>	0.06 ± 0.19	0.08 ± 0.19	0.01 ± 0.02	0.00 ± 0.00	0.0
<i>Nitrospirota</i>	0.68 ± 0.48	0.36 ± 0.45	0.31 ± 0.45	0.55 ± 0.85	0.5
<i>Spirochaetota</i>	1.50 ± 1.79	0.14 ± 0.18	0.18 ± 0.19	0.06 ± 0.13	0.4
<i>Synergistota</i>	0.00 ± 0.00	0.00 ± 0.02	0.00 ± 0.00	0.00 ± 0.00	0.0
<i>Verrucomicrobiota</i>	0.07 ± 0.12	0.00 ± 0.01	0.04 ± 0.12	0.00 ± 0.01	0.0
<i>WOR-3</i>	1.00 ± 2.88	0.42 ± 0.60	0.45 ± 1.64	0.11 ± 0.15	0.5
Unclassified	19.01 ± 13.71	18.27 ± 11.42	38.76 ± 15.51	30.25 ± 13.78	27.1

Mean ± SD: standard deviation.

about 3% of the total reads. At the OTU level, *hgcA*+ microbial communities were more similar within, as compared to between wetlands (Figure 1). Indeed, *hgcA*+ microbial community composition showed a significant clustering related to the studied site (SKM, KSN, LDN, and GTN; ANOSIM<sub>bySITE</sub>  $R = 0.38$ ,  $p < 0.001$ , Figure 1) with nMDS1 axis one reflecting a gradient from Northern (SKM and KSN) to Southern (LDN and GTN) wetlands. In contrast, no significant separation of communities was seen between different depth layers (ANOSIM<sub>byDEPTH</sub>  $R = 0.01$ ,  $p < 0.24$ , Figure 1A).



### 3.2 Potential Mercury Methylators Within Individual Wetlands


Among the classified *hgcA*+ prokaryotes, Syntrophales, OPB41, Desulfobacteriales, and Desulfuromonadales, Syntrophomonadales were the predominant orders in all four individual wetlands. SKM had the highest number of unclassified Bacteria (up to 52%, Figure 2). Spirochaetales, Syntrophomonadales, Desulfobacteriales, Syntrophales, and Syntrophorhabdadales featured higher abundances in SKM compared with the other wetlands (Supplementary Table S3 and Figure 2). In contrast, Desulfuromonadales reached their minimum relative abundance at SKM compared with the other wetlands. In SKM, *hgcA*+ microorganisms that grouped into SRB, including syntrophs, featured the highest relative abundances (Supplementary Table S3). In KSN, the *hgcA*+ community hosted

more Actinobacteriota (~22% of the total reads of KSN samples) compared with the other wetlands, and Bacteroidota were the least abundant among all the studied wetlands (Table 1). Compared to SKM, KSN presented a much higher relative abundance of Desulfuromonadales and Elusimicrobiales and also Methanosarcinales (Figure 2 and Supplementary Table S3). In the Southern wetlands, Desulfuromonadales, Desulfobacteriales, and Elusimicrobiales were more abundant in GTN compared to LDN (Figure 2 and Supplementary Table S3). In addition, the relative abundance of Methanosarcinales were similar in LDN and GTN and similar to that seen in KSN.

### 3.3 Geochemical Factors Affecting *HgcA*+ Microbial Communities Across Wetlands

We used RDA, a well-recognized multivariate analysis method in microbial ecology (Ramette, 2007), to link a matrix of multiple response variables (i.e., the abundance of the *hgcA*+ OTU) to a corresponding matrix of explanatory variables (i.e., the most influential geochemical parameters). However, as a prerequisite for RDA modeling, the geochemical variables selected should not be intercorrelated in order to optimize the precision in the prediction of the *hgcA*+ OTUs (Ramette, 2007). Hence, some variables were excluded, and in the final RDA model, the explanatory variables included phosphate ( $\mu\text{g/L}$ ),  $\text{Fe}^{\text{II}}/\text{Fe}^{\text{III}}$ , total\_LMM thiols (nM),  $\text{H}_2\text{S}$  ( $\mu\text{M}$ ), pH,  $\text{Hg}^{\text{II}}$  (pM),  $\text{Cl}^-$  (mg/L),  $\text{NH}_4^+$  ( $\mu\text{g/L}$ ),  $\text{SO}_4^{2-}$  (mg/L), water (%), DOC (mg/L), Si (mg/L)





**FIGURE 3 |** RDA biplot (2 first components) of the Hg<sup>II</sup> methylating microbial community (*hgcA* gene, OTU level) (response variable) constrained by physicochemical variables (predicting variable) and the studied sites (SKM, KSN, LDN, and GTN). The color-coding in the legend indicates the different sites. The shapes correspond to wetland core depth. The arrows and text summarize explanatory variables.

L), percentage of MeHg as THg in wetland soil, ionic strength, and K<sup>+</sup> (mg/L). The RDA model (**Figure 3**,  $F = 3$ ,  $p$ -value < 0.001) shows that the mentioned physicochemical variables explained 47% of the variation in the composition of *hgcA*+ microbial communities (RDA1 = 14%,  $p < 0.001$ ; RDA2 = 9.5%,  $p < 0.001$ ; RDA3 = 6.5%  $p < 0.001$ ; RDA4 = 4.6%,  $p = 0.003$ ) with an associated adjusted  $R^2$  of 31%. Hence, the RDA analysis emphasizes the importance of electron donors, such as SO<sub>4</sub><sup>2-</sup>; nutrients (phosphate); and organic matter (DOC and total LMM thiols) in shaping *hgcA*+ microbial communities and also points to a role of redox conditions (i.e., Fe<sup>II</sup>/Fe<sup>III</sup>, water saturation, and H<sub>2</sub>S). These results suggest that the environmental variables had a strong effect on the potential Hg<sup>II</sup> methylating microbial communities that is not merely due to geographical structuring. RDA and Mantel tests indicate that this large-scale spatial structuring of Hg<sup>II</sup> methylating microbial community composition is highly influenced by pH (Mantel  $R = 0.23$ ,  $p$ -value < 0.001, see **Figure 1B**), a physicochemical parameter that might act as an integrative factor of multiple chemical and landscape properties.

Pearson correlation analyses between the different orders and the measured physicochemical variables also identify pH as one of the main factors (**Supplementary Figure S4**). Positive significant correlations were observed between Syntrophomonadales and phosphate, DOC, and MeHg concentration and LMM thiols from biological origin. Similar correlations were observed for Spirochaetales. Interestingly, unclassified Desulfobacterota (Deltaproteobacteria) correlated positively with Fe<sup>III</sup> and the ionic strength, indicating that Fe<sup>III</sup>


might be an important geophysicochemical factor enhancing their metabolism.

### 3.4 Geochemical Factors Affecting *hgcA*+ Microbial Communities Within Each Wetland

At the SKM, GTN, and LDN sites, we observed a significant difference in *hgcA*+ microbial community composition as a function of the location within individual wetlands (ANOSIM<sub>byLOCATIONatSKM</sub>  $R = 0.6$ ,  $p < 0.001$ ; ANOSIM<sub>byLOCATIONatGTN</sub>  $R = 0.41$ ,  $p < 0.001$ ; ANOSIM<sub>byLOCATIONatLDN</sub>  $R = 0.64$ ,  $p < 0.001$ ), reflecting high spatial heterogeneity of *hgcA*+ microbial community composition at these sites. The variability across the locations where the different cores were collected was higher than across depths as none of the ANOSIM analyses performed with depth as structuring factor was significant (i.e., ANOSIM<sub>byDEPTHatSKM</sub>, ANOSIM<sub>byDEPTHatGTN</sub>, ANOSIM<sub>byDEPTHatLDN</sub>). In contrast, as illustrated in the NMDS (**Supplementary Figure S2**), *hgcA*+ microbial community composition was more similar across core replicates at KSN (ANOSIM<sub>byLOCATIONatKSN</sub>  $R = 0.02$ ,  $p = 0.35$ ). Interestingly, at KSN, depth appeared to be an important factor shaping the *hgcA*+ microbial community composition (ANOSIM<sub>byDEPTHatKSN</sub>  $R = 0.24$ ,  $p = 0.003$ ). NMDS of individual wetlands are presented in SI to visualize these differences (**Supplementary Figure S2**).

We used RDA analyses to investigate the geochemical factors influencing spatial and vertical variability in the *hgcA*+ microbial community composition across different locations within each individual wetland (**Figure 4**). Even if the explanatory variables shaping the *hgcA*+ microbial community composition differed across sites (**Figure 4**), RDA models identified nutrients (i.e., total-P, NO<sub>3</sub><sup>-</sup>, or PO<sub>4</sub><sup>3-</sup>), electron acceptors for well-described Hg<sup>II</sup> methylators (i.e., SO<sub>4</sub><sup>2-</sup> and Fe<sup>III</sup>), and local chemistry (e.g., pH, Al, K, etc.) as important factors predicting the *hgcA*+ community composition. Also, the ratio of Fe<sup>II</sup> to Fe<sup>III</sup>, water saturation, and/or H<sub>2</sub>S, which can be indicators of contrasting redox conditions and/or iron-reduction and sulfate-reduction activity, emerged as important predictors. Moreover, the concentration of the sum of LMM thiols of biological origin also came up as an important factor structuring *hgcA*+ microbial community composition (**Figure 4**).

More precisely, the explanatory variables of RDA<sub>SKM</sub> were total thiols (RSH), the Fe<sup>II</sup> to Fe<sup>III</sup> ratio, total LMM thiols from a biological origin, SO<sub>4</sub><sup>2-</sup> (mg/L), pH, water content, ionic strength, and PO<sub>4</sub><sup>3-</sup> (μg/L). The RDA<sub>SKM</sub> model (**Figure 4A**,  $F = 2.4$ ,  $p$ -value < 0.001) shows that the mentioned physicochemical variables explained 68% of *hgcA*+ microbial community composition (RDA1 = 26.7%  $p < 0.001$ ; RDA2 = 18.1%  $p < 0.001$ ) and an associated adjusted  $R^2$  of 40%. The RDA<sub>SKM</sub> model indicates that the *hgcA*+ microbial community composition of wetland soils at the SKM4 location (located close to the inlet) was likely more influenced by pH (i.e., landscape/catchment properties) and also by low H<sub>2</sub>S and high Fe<sup>III</sup> concentrations (**Supplementary Table S2**). In contrast, at SKM1 (located close to a small dam), *hgcA*+ microbial community composition was



more affected by nutrients and by local ferruginous geochemical conditions ( $Fe^{II}$  to  $Fe^{III}$  ratio). The  $RDA_{KSN}$  model (Figure 4B,  $F = 2$ ,  $p$ -value = 0.003) shows that pH, water content (%),  $PO_4^{3-}$  ( $\mu g/L$ ), the  $Fe^{II}$  to  $Fe^{III}$  ratio,  $H_2S$  ( $\mu M$ ), and  $K^+$  ( $mg/L$ ) concentrations explained 71% of the variation in the composition of *hgcA+* microbial communities ( $RDA_1 = 26.7\%$ ,  $p < 0.001$ ;  $RDA_2 = 18.1\%$ ,  $p < 0.001$ ) and an associated adjusted  $R^2$  of 36%. The  $RDA_{LDN}$  model (Figure 3C,  $F = 2.7$ ,  $p$ -value < 0.001) shows total P,  $SO_4^{2-}$  ( $mg/L$ ), water content (%),  $Fe^{III}$  ( $\mu M$ ), pH, Al ( $mg/L$ ), and LMM thiol of biological origin (*BiO\_LMM\_thiols* in nM) concentrations explained 63% (adjusted  $R^2$  of 39%) of the variation in the *hgcA+* microbial community composition. Finally, the  $RDA_{GTN}$  (Figure 4D,  $F = 2.3$ ,  $p$ -value < 0.001) shows  $H_2S$  ( $\mu M$ ), water content (%), pH,  $NO_3^-$  ( $\mu g/L$ ),  $Fe^{III}$  ( $\mu M$ ), LMM thiols of biological origin, and  $PO_4^{3-}$  ( $\mu g/L$ ) concentrations explained 70% (adjusted  $R^2$  of 39%) of the variation in the *hgcA+* microbial community composition.

### 3.5 Interplays Between *hgcA+* Microbial Community Composition and $Hg^{II}$ Methylation Rate Constants

$Hg^{II}$  methylation rate constants at these sites have been reported elsewhere (Liem-Nguyen et al., 2021) and are

presented in **Supplementary Table S1**. Liem-Nguyen et al. identified the concentration of  $Hg^{II}$  in porewater and the LMM thiol compounds of biological origin as the key factors controlling  $Hg^{II}$  methylation rate constants in SKM, KSN, LDN, and GTN wetlands. Here, we show a weak but still significant correlation between the *hgcA+* community composition and  $Hg^{II}$  methylation rate constants when considering the combined data set that included the four wetland sites (Mantel  $R = 0.13$ ,  $p$ -value = 0.003, **Supplementary Figure S5**). For visualization purposes, we present a correlation plot between *hgcA+* microbial community (Bray-Curtis) dissimilarity and the Euclidean distance of  $Hg^{II}$  methylation rate constants ( $k_m$  of spiked inorganic- $^{200}Hg$  bound to organic matter, **Supplementary Figure S5**). Also, we performed principal coordinate analysis of the *hgcA+* microbial community, and we show a positive significant correlation between the second component of principal coordinate analysis and  $Hg^{II}$  methylation rate constants, which supports the effect of the *hgcA+* community structure on the resulting  $k_m$  (**Supplementary Figure S5**). In contrast, when considering each wetland separately, we did not observe any significant correlation between  $Hg^{II}$  methylating microbial community structure and  $Hg^{II}$  methylation rate constants.



## 4 DISCUSSION

### 4.1 Community Composition of *hgcA*+ Microorganisms in Boreal Wetlands

The relationship between microbial community structure and function has emerged as an important yet elusive target in microbial ecology (Fuhrman, 2009). This is the first study to report a significant correlation between Hg<sup>II</sup> methylation rate constants and the composition of *hgcA*+ microorganisms in wetland ecosystems. In particular, we show that more diverse *hgcA*+ communities are related to higher Hg<sup>II</sup> methylation rates in wetlands. This has also been suggested by a recent study performed with freshwater lake sediments amended with different sulfate concentrations (Jones et al., 2020). Our study, therefore, emphasizes the relevance of describing the diversity of *hgcA*+ microbial communities in order to better understand the processes controlling MeHg formation in the environment.

Among the diverse *hgcA*+ microbial communities described at the studied wetlands, Deltaproteobacteria (Desulfuromonadota and Desulfobacterota\_A) was the predominant class (Table 1). Within this large class of bacteria, Desulfuromonadales, Desulfovibrionales, and Syntrophales orders were the most abundantly classified *hgcA*+ bacteria (Supplementary Table S3). Similar dominance of Deltaproteobacteria is seen in wetland and paddy soils in previous studies (Bae et al., 2014; Liu et al., 2014; Schaefer et al., 2014). Both SRB and FeRB can be found within these taxa. SRB were the first bacteria identified as responsible for Hg<sup>II</sup> methylation in aquatic systems and have been highlighted as the dominant Hg<sup>II</sup> methylators in various environments (Compeau and Bartha, 1985; Gilmour et al., 1992; King et al., 2000; Ullrich et al., 2001; Yu et al., 2010; Achá et al., 2012; Bravo et al., 2014; Gascón Díez et al., 2016; Ma et al., 2017; Liu et al., 2018; Jones et al., 2019). SRB are also capable of syntrophic fermentation of simple organic acids in the absence of SO<sub>4</sub><sup>2-</sup> as the terminal electron acceptors (McInerney et al., 2008; Plugge et al., 2011). Syntrophales were abundant members of the *hgcA*+ microbial community, reaching the highest abundance in SKM among all the wetlands. This could have resulted from the low concentrations of SO<sub>4</sub><sup>2-</sup> observed at this site (Tables 1 and Supplementary Table S1). Moreover, a recent study performing laboratory studies pinpointed syntrophy as an important metabolic pathway involved in Hg<sup>II</sup> methylation in nutrient-limited boreal peatlands (Hu et al., 2020), and we, therefore, suggest that this could be an important metabolic pathway involved in Hg<sup>II</sup> methylation in the studied wetlands. All these findings combined hint at the importance of sulfate-reducing syntrophs as potential important Hg<sup>II</sup> methylators in boreal wetlands.

FeRB were first implicated in Hg<sup>II</sup> methylation in a study by Fleming et al. (2006) and have since then been confirmed as important Hg methylators in various ecosystems, e.g., lake sediments, paddy soils, and wetland soils (Bae et al., 2014; Liu et al., 2014b; Schaefer et al., 2014). Several *Geobacter* strains (as

well as other studied FeRB) have been tested for Hg<sup>II</sup> methylation in laboratory conditions and displayed high MeHg production rates (Fleming et al., 2006; Kerin et al., 2006; Schaefer and Morel, 2009). Moreover, this metabolic guild has been identified as important Hg<sup>II</sup> methylators in both wetlands and sediments (Schaefer et al., 2014; Bravo et al., 2018b) and in forest soils (Xu et al., 2019). In this study, up to 10% of the total reads were associated to the Geopsychrobacteraceae family, which contains many *Geobacter* sp. already described as important methylators in isolates but also in the environment. The highest abundance of Geopsychrobacteraceae was observed at the more nutrient-rich site GTN and the lowest at the poor-fen SKM wetland (Supplementary Figure S6).

Firmicutes are relatively recently confirmed representatives of *hgcA*+ communities and have expanded the niches and metabolisms associated with MeHg production (Gilmour et al., 2013; Bae et al., 2014; Liu et al., 2014b). Methylation rates vary among the Firmicutes, and only a few species have been found to have methylation rates equal to some *Deltaproteobacteria* species living under similar conditions (Gilmour et al., 2013; Gilmour et al., 2018). Firmicutes represented only 1% of the total reads, although this exceeded 2% in one of the northern wetlands (Table 1). Methanogens were early on implicated in Hg<sup>II</sup> methylation (Wood, 1975). More recent work demonstrates that several methanogens have Hg<sup>II</sup> methylating properties (Gilmour et al., 2013; Yu et al., 2013) and that they may even be the primary methylators in some environments (Hamelin et al., 2011; Schaefer et al., 2014; Bravo et al., 2018b). *HgcA* genes from methanogens are important members of the *hgcA*+ microbial community in KSN, GTN, and LND (Table 1).

It is noteworthy that DNA extraction and PCR amplification are known to be prone to artifacts due to any combination of high primer mismatch, low abundance, and/or low DNA extraction efficiency that can significantly affect results. The primers inherently have preferences and also limitations in covering all species equally among different environmental samples. The primers used in this study (Schaefer et al. 2014) are predicted to cover less diversity than a more recently published primer set (Christensen et al., 2016; Gilmour et al., 2018). The high proportion of OTU attributed to unidentified species could be explained by the lack of identified organisms in the databases, primer mismatch, and/or the low conservation of the *hgcA* gene that is not ideal for biodiversity studies (Bravo and Cosio, 2020). Nevertheless, from our results, it is clear that *hgcA*+ microbial communities in wetlands are more diverse than previously thought. Furthermore, it is important to stress that the largest part of the microbial community could not be taxonomically identified, and we believe that efforts to develop techniques and increasing databases to enable a more extensive identification of the microorganisms involved in the formation of MeHg should be a research priority in the years to come.

## 4.2 Geochemical Factors Affecting Distribution of Hg<sup>II</sup> Methylating Communities

The comparison between gene assemblages with rates of biogeochemical processes, such as Hg<sup>II</sup> methylation rate constants, stems from the most fundamental level, where genes are prerequisites for any enzyme-catalyzed reaction and should, therefore, feature a correlation (Rocca et al., 2015). However, the relationship between gene and transcript abundances and the processes they facilitate can be obscured by various factors, such as habitat characteristics, complexity of biogeochemical pathways, and differences in the turnover time of nucleic acids and reaction products (Rocca et al., 2015). This study points to different physicochemical parameters as important factors determining the composition of the *hgcA*+ microbial community, which, in turn, also fundamentally controls Hg<sup>II</sup> methylation rates. Across all investigated wetlands, nMDS plots and RDA analyses show that there was more spatial variation in the diversity of *hgcA*+ microorganisms between the different wetlands than between different depths (Figures 1, 3). The combined *hgcA*+ microbial community showed changes in composition mainly related to pH (Figure 1, Mantel  $R = 0.23$ ,  $p$ -value  $< 0.001$ ) and PO<sub>4</sub><sup>3-</sup> concentration (Mantel  $R = 0.17$ ,  $p$ -value = 0.003). Previous work suggests that pH may act as an integrative factor of multiple landscape properties and could, thus, affect the *hgcA*+ microbial community in ways analogous to what has been shown for total bacterial communities (Niño-García et al., 2016). Indeed, in our study, we observe that changes in pH were strongly correlated to shifts in Fe<sup>III</sup>, Cl<sup>-</sup>, NH<sub>4</sub><sup>+</sup>, ionic strength, reduced thiols, total LMM thiols, Al, Mg, Mn, Na, and S (Supplementary Figure S3).

Concerning PO<sub>4</sub><sup>3-</sup>, it is not surprising that there is a significant correlation between this typically limiting nutrient and the composition of the *hgcA*+ microbial community in boreal wetlands. PO<sub>4</sub><sup>3-</sup> was identified as being strongly correlated to the composition of the *hgcA*+ microbial communities (Mantel  $R = 0.17$ ,  $p$ -value = 0.003) (Figure 3) and also when looking at variation within each individual wetland (Figure 4). Numerous studies have investigated the impact of nutrients, in particular P, for soil microbial community structure. The community composition of sulfate-reducing syntrophs and methanogens have, for example, been shown to depend on the position along nutrient gradients (Prechtel et al., 2001; Chauhan et al., 2004; Bae et al., 2015). Previous research has also repeatedly pointed out nutrient levels as strong regulators of bacterial growth and other metabolic activities, including MeHg production (Bigham et al., 2017; Bravo et al., 2017). Liem-Nguyen et al. (2016) also assessed the effects of variable nutrient N and P loadings to surface water and showed that increased nutrient loads caused increased phytoplankton biomass productivity and stimulated microbial activity and mercury methylation in sediments in which Hg<sup>II</sup> was presumably highly bioavailable. From all these studies, it can be concluded that nutrient

concentrations are an important factor shaping the *hgcA*+ microbial community.

Iron is also a key factor affecting Hg<sup>II</sup> methylation (Bravo et al., 2014). Fe<sup>III</sup> can act as a direct electron acceptor for FeRB and has been found to significantly influence Hg cycling and methylation (Fleming et al., 2006; Si et al., 2015). In addition, Fe<sup>III</sup>, or the Fe<sup>II</sup> to Fe<sup>III</sup> ratio, was significantly correlated to the composition of *hgcA*+ communities (Mantel  $R = 0.12$ ,  $p$ -value = 0.02) with a particularly strong correlation in wetlands SKM and GTN (Figures 3, 4). The availability of inorganic Hg<sup>II</sup> for uptake by methylating organisms has been found to limit the methylation process in laboratory conditions (Schaefer et al., 2011) and also in the studied sites (Liem-Nguyen et al., 2021). Fe<sup>III</sup> could directly affect *hgcA*+ communities by determining the availability of electron acceptors for FeRB and also indirectly via the relationship with Fe<sup>II</sup> that may influence Hg<sup>II</sup> availability, which can, in turn, be strongly affected by the interaction/complexation of Fe with organic material (Faganeli et al., 2012) and sulfur (Hsu-Kim et al., 2013). The relative abundance of the Geopsychrobacteraceae family correlated negatively with the Fe<sup>II</sup>/Fe<sup>III</sup> in porewater, likely indicating that more strongly reducing conditions limit their growth. Furthermore, the fact that the abundant unclassified Desulfobacterota (Deltaproteobacteria) correlated positively with Fe<sup>III</sup> and its ionic strength suggests an important role of *hgcA*+ FeRB in the studied wetlands.

Water content seems to be another important factor driving the composition of Hg<sup>II</sup> methylating communities (Figures 3, 4) and Hg<sup>II</sup> methylation rate constants (Supplementary Figure S3). Increasing moisture could result in elevated MeHg production from stimulation of microbial methylators, both by means of the organic material supplied via decaying vegetation and soils and from a concomitant oxygen depletion and lowering of the redox potential. It has previously been suggested that Hg<sup>II</sup> methylating microbes are stimulated by enhanced water saturation in organic forest soils as this create conditions that provide readily available electron donors to drive the formation and buildup of MeHg (Kronberg et al., 2016). Eklöf et al. (2018) also found that the highest % MeHg was associated with water-filled cavities and water-logged soil environments, conditions that would favor methylating microorganisms and enhance MeHg formation. In wetland KSN, where water content was generally very high (Supplementary Table S1), depth appeared to play a more important role in shaping the local *hgcA*+ community (Supplementary Figure S2, ANOSIM<sub>byDEPTHatKSN</sub>  $R = 0.24$ ,  $p = 0.003$ ). One explanation could be that the KSN soils provide vertically stratified heterogeneous habitats for microorganisms with varying fresh organic matter availability and nutrient levels (Fierer et al., 2003; Lorenz and Lal, 2005; Kramer and Gleixner, 2008). Furthermore, it is possible that other factors not considered in this study, such as the composition of the organic matter, may play a role in controlling the composition of the *hgcA*+ communities, and the strong influence of biologically derived LMM thiols support this.

## 5 CONCLUSION

Understanding the responses of bacterial communities to environmental factors is a major focus for research in microbial ecology, and this also applies to Hg<sup>II</sup> methylating microbial communities. Combining high-throughput *hgcA* amplicon sequencing with molecular barcoding reveals diverse clades of *hgcA*+ microorganisms in a wide range of wetland soils. This study confirms the predominant role of Deltaproteobacteria as *hgcA*+ microorganisms in boreal wetland soils and also unveils the presence of Firmicutes and Archaea as abundant members of methylating communities. By uncovering relationships between the composition of such communities, Hg<sup>II</sup> methylation rate constants, and the physicochemical factors, we propose an important role of pH, water saturation, nutrient status, Fe, and LMM thiols of biological origin for controlling the *hgcA*+ microbial community composition and their activities across the four wetlands. In the future, more attention should be paid to how specific groups of Hg<sup>II</sup> methylating microbes respond to various levels of limiting nutrients. Furthermore, it would be worthwhile to carry out more in-depth studies on the interactions between various Hg<sup>II</sup> methylating groups in contributing to the overall MeHg production. Our findings provide a better understanding of Hg<sup>II</sup> methylating microbial communities in boreal wetland soils. As more than 50% of wetlands are located in the high northern latitudes, the results reported here are likely broadly relevant for a wide range of wetlands.

## DATA AVAILABILITY STATEMENT

The datasets generated for this study can be found in the EBI Archive under accession number PRJEB20882.

## REFERENCES

- Achá, D., Hintelmann, H., and Pabón, C. A. (2012). Sulfate-reducing bacteria and mercury methylation in the water column of the lake 658 of the experimental lake area. *Geomicrobiol. J.* 29, 667–674. doi:10.1080/01490451.2011.606289
- Bae, H. S., Dierberg, F. E., and Ogram, A. (2014). Syntrophs dominate sequences associated with the mercury methylation-related gene *hgcA* in the water conservation areas of the Florida Everglades. *Appl. Environ. Microbiol.* 80, 6517–6526. doi:10.1128/AEM.01666-14
- Bae, H. S., Holmes, M. E., Chanton, J. P., Reddy, K. R., and Ogram, A. (2015). Distribution, activities, and interactions of methanogens and sulfate-reducing prokaryotes in the Florida everglades. *Appl. Environ. Microbiol.* 81, 7431–7442. doi:10.1128/AEM.01583-15
- Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc.* 57, 289. doi:10.1111/j.2517-6161.1995.tb02031.x
- Bigham, G. N., Murray, K. J., Masue-Slowey, Y., and Henry, E. A. (2017). Biogeochemical controls on methylmercury in soils and sediments: implications for site management. *Integr. Environ. Assess. Manage.* 13, 249–263. doi:10.1002/ieam.1822
- Boening, D. W. (2000). Ecological effects, transport, and fate of mercury: a general review. *Chemosphere* 40, 1335–1351. doi:10.1016/S0045-6535(99)00283-0

## AUTHOR CONTRIBUTIONS

EB, VL, JX, AB, SB, and JX. conceived the study. VL and EB, conducted the sampling campaigns. JX and AB performed the laboratory analyses with guidance from SB. MB performed the bioinformatics analyses. AB conducted all statistical analyses and built the figures with comments and suggestions from the rest of the co-authors. JX, AB, and SB. wrote the manuscript with significant assistance and comments from EB and VL.

## FUNDING

This project was carried out within the Swedish-Sino SMaREF (2013-6978) funded by the Swedish Research Council. This study was also supported by the Swedish Research Council (Grants No. 2011-7192, 2012-3892 and 2017-04422), a Wennergren foundation project to LVN and the Kempe Foundations (SMK-2745, SMK-1243) and the Marie Curie Individual Fellowship (H2020-MSCA-IF-2016; project-749645).

## ACKNOWLEDGMENTS

Sequencing was carried out at the SciLifeLab SNP/SEQ facility hosted by Uppsala University, and we acknowledge the Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) for access to storage and computational resources.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fenvs.2020.518662/full#supplementary-material>.

- Bravo, A. G., Bouchet, S., Tolu, J., Björn, E., Mateos-Rivera, A., and Bertilsson, S. (2017). Molecular composition of organic matter controls methylmercury formation in boreal lakes. *Nat. Commun.* 8, 14255. doi:10.1038/ncomms14255
- Bravo, A. G., and Cosio, C. (2020). Biotic formation of methylmercury: a bio-physicochemical conundrum. *Limnol. Oceanogr.* 65, 1010–1027. doi:10.1002/lno.11366
- Bravo, A. G., Cosio, C., Amouroux, D., Zopfi, J., Chevalley, P. A., Spangenberg, J. E., et al. (2014). Extremely elevated methyl mercury levels in water, sediment and organisms in a Romanian reservoir affected by release of mercury from a chlor-alkali plant. *Water Res.* 49, 391–405. doi:10.1016/j.watres.2013.10.024
- Bravo, A. G., Peura, S., Buck, M., Ahmed, O., Mateos-Rivera, A., Herrero Ortega, S., et al. (2018a). Methanogens and iron-reducing bacteria: the overlooked members of mercury-methylating microbial communities in boreal lakes. *Appl. Environ. Microbiol.* 84, e01774–18. doi:10.1128/AEM.01774-18
- Bravo, A. G., Zopfi, J., Buck, M., Xu, J., Bertilsson, S., Schaefer, J. K., et al. (2018b). Geobacteraceae are important members of mercury-methylating microbial communities of sediments impacted by waste water releases. *ISME J.* 12, 802–812. doi:10.1038/s41396-017-0007-7
- Capo, E., Bravo, A. G., Soerensen, A. L., Bertilsson, S., Pinhassi, J., Feng, C., et al. (2020). Deltaproteobacteria and spirochaetes-like bacteria are abundant putative mercury methylators in oxygen-deficient water and marine particles in the Baltic Sea. *Front. Microbiol.* 11, 574080. doi:10.3389/fmicb.2020.574080
- Chauhan, A., Ogram, A., and Reddy, K. R. (2004). Syntrophic-methanogenic associations along a nutrient gradient in the Florida Everglades. *Appl. Environ. Microbiol.* 70, 3475–3484. doi:10.1128/AEM.70.6.3475-3484.2004



- Christensen, G. A., Somenahally, A. C., Moberly, J. G., Miller, C. M., King, A. J., Gilmour, C. C., et al. (2017). Carbon amendments alter microbial community structure and net mercury methylation potential in sediments. *Appl. Environ. Microbiol.* 84, 1–14. doi:10.1128/AEM.01049-17
- Christensen, G. A., Wymore, A. M., King, A. J., Podar, M., Hurt, R. A., Santillan, E. U., et al. (2016). Development and validation of broad-range qualitative and clade-specific quantitative molecular probes for assessing mercury methylation in the environment. *Appl. Environ. Microbiol.* 82, 6068–6078. doi:10.1128/AEM.01271-16
- Clarke, K. R., and Gorley, R. N. (2015). *PRIMER v7: user manual/tutorial*. Plymouth UK: Prim.
- Compeau, G. C., and Bartha, R. (1985). Sulfate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50, 498–502. doi:10.1128/AEM.50.2.498-502.1985
- Eklöf, K., Bishop, K., Bertilsson, S., Björn, E., Buck, M., Skyllberg, U., et al. (2018). Formation of mercury methylation hotspots as a consequence of forestry operations. *Sci. Total Environ.* 613, 1069–1078.
- Erwin, K. L. (2009). Wetlands and global climate change: the role of wetland restoration in a changing world. *Wetl. Ecol. Manag.* 17, 71–84. doi:10.1007/s11273-008-9119-1
- Faganeli, J., Hines, M. E., Covelli, S., Emili, A., and Giani, M. (2012). Mercury in lagoons: an overview of the importance of the link between geochemistry and biology. *Estuar. Coast Shelf Sci.* 113, 126–132. doi:10.1016/j.ecss.2012.08.021
- Fierer, N., Schimel, J. P., and Holden, P. A. (2003). Variations in microbial community composition through two soil depth profiles. *Soil Biol. Biochem.* 35, 167–176. doi:10.1016/S0038-0717(02)00251-1
- Fleming, E. J., Mack, E. E., Green, P. G., and Nelson, D. C. (2006). Mercury methylation from unexpected sources: molybdate-inhibited freshwater sediments and an iron-reducing bacterium. *Appl. Environ. Microbiol.* 72, 457–464. doi:10.1128/AEM.72.1.457-464.2006
- Fuhrman, J. A. (2009). Microbial community structure and its functional implications. *Nature*, 459, 193. doi:10.1038/nature08058
- Gascón Díez, E., Loizeau, J. L., Cosio, C., Bouchet, S., Adatte, T., Amouroux, D., et al. (2016). Role of settling particles on mercury methylation in theoxic water column of freshwater systems. *Environ. Sci. Technol.* 50, 11672–11679. doi:10.1021/acs.est.6b03260
- Gilmour, C. C., Bullock, A. L., Mcburney, A., and Podar, M. (2018). Robust mercury methylation across diverse methanogenic archaea. *mBio* 9, e02403–e02417. doi:10.1128/mbio.02403-17
- Gilmour, C. C., Henry, E. A., and Mitchell, R. (1992). Sulfate stimulation of mercury methylation in freshwater sediments. *Environ. Sci. Technol.* 26, 2281–2287. doi:10.1021/es00035a029
- Gilmour, C. C., Podar, M., Bullock, A. L., Graham, A. M., Brown, S. D., Somenahally, A. C., et al. (2013). Mercury methylation by novel microorganisms from new environments. *Environ. Sci. Technol.* 47, 11810–11820. doi:10.1021/es403075t
- Gilmour, C. C., Riedel, G. S., Ederington, M. C., Bell, J. T., Gill, G. A., Stordal, M. C., et al. (1998). Methylmercury concentrations and production rates across a trophic gradient in the northern Everglades. *Biogeochemistry* 40, 327–345. ST-Methylmercury concentrations and pro. doi:10.1023/a:1005972708616
- Gionfriddo, C. M., Tate, M. T., Wick, R. R., Schultz, M. B., Zemla, A., Thelen, M. P., et al. (2016). Microbial mercury methylation in Antarctic sea ice. *Nat. Microbiol.* 1, 16127. doi:10.1038/nmicrobiol.2016.127
- Hall, B. D., Aiken, G. R., Krabbenhoft, D. P., Marvin-DiPasquale, M., and Swarzenski, C. M. (2008). Wetlands as principal zones of methylmercury production in southern Louisiana and the Gulf of Mexico region. *Environ. Pollut.* 154, 124–134. doi:10.1016/j.envpol.2007.12.017
- Hamelin, S., Amyot, M., Barkay, T., Wang, Y., and Planas, D. (2011). Methanogens: principal methylators of mercury in lake periphyton. *Environ. Sci. Technol.* 45, 7693–7700. doi:10.1021/es2010072
- Hsu-Kim, H., Kucharzyk, K. H., Zhang, T., and Deshusses, M. A. (2013). Mechanisms regulating mercury bioavailability for methylating microorganisms in the aquatic environment: a critical review. *Environ. Sci. Technol.* 47, 2441–2456. doi:10.1021/es304370g
- Hu, H., Wang, B., Bravo, A. G., Björn, E., Skyllberg, U., Amouroux, D., et al. (2020). Shifts in mercury methylation across a peatland chronosequence: from sulfate reduction to methanogenesis and syntrophy. *J. Hazard. Mater.* 387, 121967. doi:10.1016/j.jhazmat.2019.121967
- Jones, D. S., Johnson, N. W., Mitchell, C. P. J., Walker, G. M., Bailey, J. V., Pastor, J., et al. (2020). Diverse communities of hgcAB+ microorganisms methylate mercury in freshwater sediments subjected to experimental sulfate loading. *Environ. Sci. Technol.* 54, 14265–14274. doi:10.1021/acs.est.0c02513
- Jones, D. S., Walker, G. M., Johnson, N. W., Mitchell, C. P. J., Coleman Wasik, J. K., and Bailey, J. V. (2019). Molecular evidence for novel mercury methylating microorganisms in sulfate-impacted lakes. *ISME J.* 13, 1659. doi:10.1038/s41396-019-0376-1
- Jonsson, S., Skyllberg, U., Nilsson, M. B., Lundberg, E., Andersson, A., and Björn, E. (2014). Differentiated availability of geochemical mercury pools controls methylmercury levels in estuarine sediment and biota. *Nat. Commun.* 5, 4624. doi:10.1038/ncomms5624
- Joshi, N. A., and Fass, J. N. (2011). *Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files*. Available at: <https://github.com/najoshi/sickle>
- Kerin, E. J., Gilmour, C. C., Roden, E., Suzuki, M. T., Coates, J. D., and Mason, R. P. (2006). Mercury methylation by dissimilatory Iron-reducing bacteria. *Appl. Environ. Microbiol.* 72, 7919–7921. doi:10.1128/AEM.01602-06
- King, J. K., Kostka, J. E., Frischer, M. E., Saunders, F. M., and Jahnke, R. A. (2001). A quantitative relationship that demonstrates mercury methylation rates in marine sediments are based on the community composition and activity of sulfate-reducing bacteria. *Environ. Sci. Technol.* 35, 2491–2496. doi:10.1021/es001813q
- King, J. K., Kostka, J. E., Frischer, M. E., and Saunders, F. M. (2000). Sulfate-reducing bacteria methylate mercury at variable rates in pure culture and in marine sediments. *Appl. Environ. Microbiol.* 66, 2430–2437. doi:10.1128/AEM.66.6.2430-2437.2000
- Kramer, C., and Gleixner, G. (2008). Soil organic matter in soil depth profiles: distinct carbon preferences of microbial groups during carbon transformation. *Soil Biol. Biochem.* 40, 425–433. doi:10.1016/j.soilbio.2007.09.016
- Kronberg, R. M., Jiskra, M., Wiederhold, J. G., Björn, E., and Skyllberg, U. (2016). Methyl mercury formation in hillslope soils of boreal forests: the role of forest harvest and anaerobic microbes. *Environ. Sci. Technol.* 50, 9177–9186. doi:10.1021/acs.est.6b00762
- Lambertsson, L., and Björn, E. (2004). Validation of a simplified field-adapted procedure for routine determinations of methyl mercury at trace levels in natural water samples using species-specific isotope dilution mass spectrometry. *Anal. Bioanal. Chem.* 380, 871–875. doi:10.1007/s00216-004-2863-z
- Lambertsson, L., Lundberg, E., Nilsson, M., and Frech, W. (2001). Applications of enriched stable isotope tracers in combination with isotope dilution GC-ICP-MS to study mercury species transformation in sea sediments during *in situ*. *J. Anal. At. Spectrom.* 16, 1296–1301. doi:10.1039/b106878b
- Liem-Nguyen, V., Bouchet, S., and Björn, E. (2015). Determination of sub-nanomolar levels of low molecular mass thiols in natural waters by liquid chromatography tandem mass spectrometry after derivatization with p-(hydroxymercuri) benzoate and online preconcentration. *Anal. Chem.* 87, 1089–1096. doi:10.1021/ac503679y
- Liem-Nguyen, V., Jonsson, S., Skyllberg, U., Nilsson, M. B., Andersson, A., Lundberg, E., et al. (2016). Effects of nutrient loading and mercury chemical speciation on the formation and degradation of methylmercury in estuarine sediment. *Environ. Sci. Technol.* 50, 6983–6990. doi:10.1021/acs.est.6b01567
- Liem-Nguyen, V., Skyllberg, U., and Björn, E. (2021). Methylmercury formation in boreal wetlands in relation to chemical speciation of mercury(II) and concentration of low molecular mass thiols. *Sci. Total Environ.* 755, 142666. doi:10.1016/j.scitotenv.2020.142666
- Liem-Nguyen, V., Skyllberg, U., and Björn, E. (2017). Thermodynamic modeling of the solubility and chemical speciation of mercury and methylmercury driven by organic thiols and micromolar sulfide concentrations in boreal wetland soils. *Environ. Sci. Technol.* 51, 3678–3686. doi:10.1021/acs.est.6b04622
- Liu, X., Ma, A., Zhuang, G., and Zhuang, X. (2018). Diversity of microbial communities potentially involved in mercury methylation in rice paddies surrounding typical mercury mining areas in China. *Microbiologyopen* 7, e00577. doi:10.1002/mbo3.577
- Liu, Y. R., Yu, R. Q., Zheng, Y. M., and He, J. Z. (2014). Analysis of the microbial community structure by monitoring an Hg methylation gene (*hgcA*) in paddy soils along an Hg gradient. *Appl. Environ. Microbiol.* 80, 2874–2879. doi:10.1128/AEM.04225-13



- Lorenz, K., and Lal, R. (2005). The depth distribution of soil organic carbon in relation to land use and management and the potential of carbon sequestration in subsoil horizons. *Adv. Agron.* 88, 35–66. doi:10.1016/S0065-2113(05)88002-2
- Louis, V. L. S. T., Rudd, J. W. M., Kelly, C. A., Beaty, K. G., Flett, R. J., and Roulet, N. T. (1996). Production and loss of methylmercury and loss of total mercury from boreal forest catchments containing different types of wetlands. *Environ. Sci. Technol.* 30, 2719–2729. doi:10.1021/es950856h
- Ma, M., Du, H., Wang, D., Kang, S., and Sun, T. (2017). Biotically mediated mercury methylation in the soils and sediments of Nam Co Lake, Tibetan Plateau. *Environ. Pollut.* 227, 243–251. doi:10.1016/j.envpol.2017.04.037
- Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* 17, 10–12. doi:10.14806/ej.17.1.200
- Matsen, F. A., Kodner, R. B., and Armbrust, E. V. (2010). pplacer: linear time maximum-likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC Bioinformatics.* 11, 538. doi:10.1186/1471-2105-11-538
- Mazrui, N. M., Jonsson, S., Thota, S., Zhao, J., and Mason, R. P. (2016). Enhanced availability of mercury bound to dissolved organic matter for methylation in marine sediments. *Geochem. Cosmochim. Acta.* 194, 153–162. doi:10.1016/j.gca.2016.08.019
- McDaniel, E. A., Peterson, B., Stevens, S. L. R., Tran, P. Q., Anantharaman, K., and McMahon, K. D. (2020a). Expanded phylogenetic diversity and metabolic flexibility of microbial mercury methylation. *bioRxiv.* doi:10.1101/2020.01.16.909358
- McDaniel, E. A., Peterson, B. D., Stevens, S. L. R., Tran, P. Q., Anantharaman, K., and McMahon, K. D. (2020b). Expanded phylogenetic diversity and metabolic flexibility of mercury-methylating microorganisms. *mSystems.* 5, e00299–20. doi:10.1128/mSystems.00299-20
- McInerney, M. J., Struchtemeyer, C. G., Sieber, J., Mouttaki, H., Stams, A. J., Schink, B., et al. (2008). Physiology, ecology, phylogeny, and genomics of microorganisms capable of syntrophic metabolism. *Ann. N. Y. Acad. Sci.* 1125, 58–72. doi:10.1196/annals.1419.005
- Mitchell, C. P. J., Branfireun, B. A., and Kolka, R. K. (2008a). Assessing sulfate and carbon controls on net methylmercury production in peatlands: an *in situ* mesocosm approach. *Appl. Geochem.* 23, 503–518. doi:10.1016/j.apgeochem.2007.12.020
- Mitchell, C. P., Branfireun, B. A., and Kolka, R. K. (2008b). Spatial characteristics of net methylmercury production hot spots in peatlands. *Environ. Sci. Technol.* 42, 1010–1016. doi:10.1021/es0704986
- Mitra, S., Wassmann, R., and Vlek, P. L. G. (2003). *Global inventory of wetlands and their role in the carbon cycle.* Bonn: ZEF Discussion PapDev. Policy.
- Niño-García, J. P., Ruiz-González, C., and Del Giorgio, P. A. (2016). Interactions between hydrology and water chemistry shape bacterioplankton biogeography across boreal freshwater networks. *ISME J.* 10, 1755–1766. doi:10.1038/ismej.2015.226
- Oksanen, A. J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., Mcginn, D., et al. (2017). Package ‘vegan’ Software. Vienna, Austria: R Foundation for Statistical Computing.
- Parks, J. M., Johs, A., Podar, M., Bridou, R., Hurt, R. A., Smith, S. D., et al. (2013). The genetic basis for bacterial mercury methylation. *Science* 339, 1332–1335. doi:10.1126/science.1230667
- Peterson, B. D., McDaniel, E. A., Schmidt, A. G., Lepak, R. F., Tran, P. Q., Marick, R. A., et al. (2020). Microbial mercury methylation genes identified across diverse anaerobic microbial guilds in a eutrophic sulfate-enriched lake. *Environ. Sci. Technol.* 54, 15840–15851.
- Plugge, C. M., Zhang, W., Scholten, J. C., and Stams, A. J. (2011). Metabolic flexibility of sulfate-reducing bacteria. *Front. Microbiol.* 2, 81–88. doi:10.3389/fmicb.2011.00081
- Podar, M., Gilmour, C. C., Brandt, C. C., Soren, A., Brown, S. D., Crable, B. R., et al. (2015). Global prevalence and distribution of genes and microorganisms involved in mercury methylation. *Sci. Adv.* 1, e1500675. doi:10.1126/sciadv.1500675
- Prechtel, A., Alewell, C., Armbruster, M., Bittersohl, J., Cullen, J. M., Evans, C. D., et al. (2001). Response of sulphur dynamics in European catchments to decreasing sulphate deposition. *Hydrol. Earth Syst. Sci.* 5, 311–326. doi:10.5194/hess-5-311-2001
- Ramette, A. (2007). Multivariate analyses in microbial ecology. *FEMS Microbiol. Ecol.* 62, 142–160. doi:10.1111/j.1574-6941.2007.00375.x
- Ranchou-Peyruse, M., Monperrus, M., Bridou, R., Duran, R., Amouroux, D., Salvado, J. C., et al. (2009). Overview of mercury methylation capacities among anaerobic bacteria including representatives of the sulphate-reducers: implications for environmental studies. *Geomicrobiol. J.* 26, 1–8. doi:10.1080/01490450802599227
- Rocca, J. D., Hall, E. K., Lennon, J. T., Evans, S. E., Waldrop, M. P., Cotner, J. B., et al. (2015). Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed. *ISME J.* 9, 1693–1699. doi:10.1038/ismej.2014.252
- Schaefer, J. K., Kronberg, R. M., Björn, E., and Skjellberg, U. (2020). Anaerobic guilds responsible for mercury methylation in boreal wetlands of varied trophic status serving as either a methylmercury source or sink. *Environ. Microbiol.* 22, 3685. doi:10.1111/1462-2920.15134
- Schaefer, J. K., Kronberg, R. M., Morel, F. M., and Skjellberg, U. (2014). Detection of a key Hg methylation gene, hgcA, in wetland soils. *Environ. Microbiol. Rep.* 6, 441–447. doi:10.1111/1758-2229.12136
- Schaefer, J. K., and Morel, F. M. M. (2009). High methylation rates of mercury bound to cysteine by *Geobacter sulfurreducens*. *Nat. Geosci.* 2, 123–126. doi:10.1038/ngeo412
- Schaefer, J. K., Rocks, S. S., Zheng, W., Liang, L., Gu, B., and Morel, F. M. (2011). Active transport, substrate specificity, and methylation of Hg(II) in anaerobic bacteria. *Proc. Natl. Acad. Sci. U.S.A.* 108, 8714–8719. doi:10.1073/pnas.1105781108
- Si, Y., Zou, Y., Liu, X., Si, X., and Mao, J. (2015). Mercury methylation coupled to iron reduction by dissimilatory iron-reducing bacteria. *Chemosphere* 122, 206–212. doi:10.1016/j.chemosphere.2014.11.054
- Sinclair, L., Osman, O. A., Bertilsson, S., and Eiler, A. (2015). Microbial community composition and diversity via 16S rRNA gene amplicons: evaluating the Illumina platform. *PLoS One.* 10, e0116955. doi:10.1371/journal.pone.0116955
- Skjellberg, U. (2008). Competition among thiols and inorganic sulfides and polysulfides for Hg and MeHg in wetland soils and sediments under suboxic conditions: illumination of controversies and implications for MeHg net production. *J. Geophys. Res.* 113. doi:10.1029/2008JG000745
- St. Louis, V. L., Rudd, J. W. M., Kelly, C. A., Beaty, K. G., Bloom, N. S., and Flett, R. J. (1994). Importance of wetlands as sources of methyl mercury to boreal forest ecosystems. *Can. J. Fish. Aquat. Sci.* 51, 1065–1076. doi:10.1139/f94-106
- Tjerngren, I., Karlsson, T., Björn, E., and Skjellberg, U. (2012a). Potential Hg methylation and MeHg demethylation rates related to the nutrient status of different boreal wetlands. *Biogeochemistry* 108, 335–350. doi:10.1007/s10533-011-9603-1
- Tjerngren, I., Meili, M., Björn, E., and Skjellberg, U. (2012b). Eight boreal wetlands as sources and sinks for methyl mercury in relation to soil acidity, C/N ratio, and small-scale flooding. *Environ. Sci. Technol.* 46, 8052–8060. doi:10.1021/es300845x
- Ullrich, S. M., Tanton, T. W., Abdrashitova, S. A., and Svetlana, A. (2001). Mercury in the aquatic environment: a review of factors affecting methylation. *Crit. Rev. Environ. Sci. Technol.* 31, 241–293. doi:10.1080/20016491089226
- Villar, E., Cabrol, L., and Heimbürger-Boavida, L. E. (2020). Widespread microbial mercury methylation genes in the global ocean. *Environ. Microbiol. Rep.* 12, 277. doi:10.1111/1758-2229.12829
- Viollier, E., Inglett, P. W., Hunter, K., Roychoudhury, A. N., and Van Cappellen, P. (2000). The ferrozine method revisited: Fe(II)/Fe(III) determination in natural waters. *Appl. Geochem.* 15, 785–790. doi:10.1016/S0883-2927(99)00097-9
- Vishnivetskaya, T. A., Hu, H., Van Nostrand, J. D., Wymore, A. M., Xu, X., Qiu, G., et al. (2018). Microbial community structure with trends in methylation gene diversity and abundance in mercury-contaminated rice paddy soils in Guizhou, China. *Environ. Sci. Process. Impacts.* 20, 673–685. doi:10.1039/C7EM00558J
- Wood, J. M. (1975). Biological cycles for elements in the environment. *Naturwissenschaften* 62, 357–364. doi:10.1007/BF00625342
- Xu, J., Buck, M., Eklöf, K., Ahmed, O. O., Schaefer, J. K., Bishop, K., et al. (2019). Mercury methylating microbial communities of boreal forest soils. *Sci. Rep.* 9, 518. doi:10.1038/s41598-018-37383-z
- Yu, R. Q., Adatto, L., Montesdeoca, M. R., Driscoll, C. T., Hines, M. E., and Barkay, T. (2010). Mercury methylation in Sphagnum moss mats and its association with sulfate-reducing bacteria in an acidic Adirondack forest lake wetland. *FEMS Microbiol. Ecol.* 74, 655–668. doi:10.1111/j.1574-6941.2010.00978.x

- Yu, R. Q., Flanders, J. R., MacK, E. E., Turner, R., Mirza, M. B., and Barkay, T. (2012). Contribution of coexisting sulfate and iron reducing bacteria to methylmercury production in freshwater river sediments. *Environ. Sci. Technol.* 46, 2684–2691. doi:10.1021/es2033718
- Yu, R.-Q., Reinfelder, J. R., Hines, M. E., and Barkay, T. (2018). Syntrophic pathways for microbial mercury methylation. *ISME J.* 12, 1826. doi:10.1038/s41396-018-0106-0
- Yu, R., Reinfelder, J. R., Hines, M. E., and Barkay, T. (2013). Mercury methylation by the methanogen *Methanospirillum hungatei*. *Appl. Environ. Microbiol.* 79, 6325–6330. doi:10.1128/AEM.01556-13

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Xu, Liem-Nguyen, Buck, Bertilsson, Björn and Bravo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.