Chromosome-Level Genome Assembly of the Common Chaffinch (Aves: *Fringilla coelebs*): A Valuable Resource for Evolutionary Biology

María Recuerda ^{1,*†}, Joel Vizueta ^{1,2,†}, Cristian Cuevas-Caballé², Guillermo Blanco ¹, Julio Rozas ², and Borja Milá ¹

¹National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid, Spain

²Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia and Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Barcelona, Spain

†These authors contributed equally to this work.

*Corresponding author: E-mail: mariarecuerdacarrasco@gmail.com.

Accepted: 16 February 2021

Abstract

The common chaffinch, *Fringilla coelebs*, is one of the most common, widespread, and well-studied passerines in Europe, with a broad distribution encompassing Western Europe and parts of Asia, North Africa, and the Macaronesian archipelagos. We present a high-quality genome assembly of the common chaffinch generated using Illumina shotgun sequencing in combination with Chicago and Hi-C libraries. The final genome is a 994.87-Mb chromosome-level assembly, with 98% of the sequence data located in chromosome scaffolds and a N50 statistic of 69.73 Mb. Our genome assembly shows high completeness, with a complete BUSCO score of 93.9% using the avian data set. Around 7.8% of the genome contains interspersed repetitive elements. The structural annotation yielded 17,703 genes, 86.5% of which have a functional annotation, including 7,827 complete universal single-copy orthologs out of 8,338 genes represented in the BUSCO avian data set. This new annotated genome assembly will be a valuable resource as a reference for comparative and population genomic analyses of passerine, avian, and vertebrate evolution.

Key words: common chaffinch, *Fringilla coelebs*, reference genome, whole genome assembly.

Significance

High-quality reference genomes of wild, nonmodel species are very useful tools to understand how organisms evolve. If genomes are annotated, so that the specific genes are identified, we can make progress toward associating specific physical or behavioral traits with the genes that code for them, and thus further understand the evolutionary process. Here, we provide a high-quality, annotated genome of the common chaffinch, a common and widespread Eurasian finch, that will be a useful resource in studies related to evolution, phylogenomics, biogeography, and adaptation genomics, among others.

Introduction

The decreasing costs of DNA sequencing, along with advances in computational genomics, are promoting a rapid increase in the availability of high-quality reference genomes of nonmodel species, which greatly improves our capacity

to address a range of biological questions from a genomic perspective. Among them, the correct annotation of protein-coding genes in whole genomes allows to identify new genes involved in the process of evolutionary adaptation and provides a better understanding of the evolutionary mechanisms involved in the speciation process. Avian genomes are

© The Author(s) 2021. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

particularly suited for studying the molecular basis of speciation as they have a relatively simple architecture and are among the smallest within amniotes, ranging from 0.91 to 1.3 Gb (Gregory 2002). In the last decade, the number of bird reference genomes has increased dramatically (e.g. Dalloul et al. 2010; Warren et al. 2010; Zhang et al. 2012; Jarvis et al. 2014; Poelstra et al. 2014; Frankl-Vilches et al. 2015; Friis et al. 2018; Louha et al. 2020; Peñalba et al. 2020; Ducrest et al. 2020; Wang et al. 2020), providing major scientific breakthroughs in phylogenetics (Alström et al. 2018; Braun et al. 2019; Jarvis et al. 2015), comparative genomics (Zhang et al. 2014; Feng et al. 2020), adaptation genomics (Wirthlin et al. 2014; Lawson and Petren 2017), and genomic architecture (Poelstra et al. 2014; Vijay et al. 2016), among others. Moreover, the Ten-Thousand Bird Genomes (B10K) consortium has generated and analyzed over 300 avian genomes from 92.4% of bird families, providing an unprecedented genomic resource for avian comparative studies (Zhang et al. 2015, Feng et al. 2020).

The common chaffinch (Aves, Passeriformes, Fringillidae, *Fringilla coelebs*) is a widely distributed species, ranging from across Eurasia to the north of Africa, and has colonized three Macaronesian archipelagos in the Atlantic Ocean (Azores, Madeira, and the Canary Islands) (Collar et al. 2020). With about 15 currently recognized subspecies, the common chaffinch is an ideal system for testing hypotheses on the evolutionary process given its distribution across the continent and the colonization of several oceanic islands, recognized as excellent natural laboratories for studying evolution (Brown et al. 2013). Island systems have inspired the development of biogeographical theories (MacArthur and Wilson 1967) and are of central importance for understanding the role of area and isolation in colonization, extinction, and speciation rates (Valente et al. 2020), which are processes influencing global patterns of species richness (Losos and Schluter 2000). Species that have colonized insular environments, like the common chaffinch, are also excellent systems for the study of demographic events, such as bottlenecks leading to small effective population size (N_e) (Leroy et al. 2021), or the roles of drift and selection in the divergence process (Barton 1996). The common chaffinch has been intensively studied using molecular tools, so that the availability of a reference genome represents a valuable resource to improve our understanding of avian evolution, biogeography, and demography (Illera et al. 2018).

Results and Discussion

Assembly and Quality Control

The total length obtained by the HiRise software for the common chaffinch assembly was 994.87 Mb. Nevertheless, the estimate from k -mer metrics is 1.2 Gb. The discrepancy between these estimates could be caused by the presence of

repetitive elements given the assembly strategy used, which could have been improved including long-read sequencing technologies. This final assembly consists of 3,255 scaffolds, 3,239 over 1 kb, and an N50 of 69.73 Mb (see [supplementary table S1, Supplementary Material](#) online) with a sequence coverage of 249 $\times$. The use of Chicago and Hi-C libraries provided a clear improvement in quality by increasing 917 times the scaffold N50, reducing the number of scaffolds from 38,666 to 3,255 ([Supplementary table S1](#), see [supplementary methods](#) for details, [Supplementary Material](#) online). In fact, 98% of the total genome sequence maps in the 30 described chromosomes.

The chaffinch genome showed high synteny with the zebra finch genome (fig. 1), evidencing the completeness of the assembly, with all micro-chromosomes and the Z chromosome present in the assembly. In addition, the alignment between these genomes suggests the presence of several inversions in chromosomes 1, 1A, 2, 3, 5, 7, 8, and 9. Several studies have documented that inversions are very common in birds (Aslam et al. 2010; Völker et al. 2010; Skinner and Griffin 2012; Zhang et al. 2014). For instance, Hooper and Price (2017) identified 319 inversions on the 9 largest autosomes combined in 81 independent clades. No putative contaminations were detected and 89.6% of the reads were mapped in the genome assembly (see [Supplementary fig. S1, Supplementary Material](#) online). The mean GC content of the assembly was 41.86% (± 11 SD). The common chaffinch genome assembly included 7,832 complete copies (93.9%) out of the 8,338 BUSCO data set from avian genomes, among which 7,816 were single-copy orthologs and 16 were duplicated. Only 1.8% of the gene models were fragmented, and 4.3% were missing in the genome. These few missing gene models could represent divergent or lost genes in our species, but also could be related with putative errors during the assembly process or missing data.

Repetitive Regions

Overall, 7.82% of the genome assembly are repeats (~ 78 Mb), of which 85.4% are transposable elements (TEs). The most abundant TEs are LINES (53.5%) followed by LTR (29.4%), DNA elements (4.1%), and SINES (1.4%), with the remaining 11.6% unclassified. The rest of repeats (14.6%) contained simple repeats (75.4%), low complexity repeats (18.5%), satellites (4.2%) and small RNA (1.9%) (see [supplementary table S2, Supplementary Material](#) online). The number of repetitive regions is within the expected range in birds, which is at 4–10% of the genome (Zhang et al. 2014).

A total of 111,076 microsatellites, with motif length ranging between 2 and 20 bp, were identified in the common chaffinch genome (see [supplementary fig. S3, Supplementary Material](#) online; their genomic locations are shown in [supplementary file S1](#) in the Figshare repository). The most common k -mer sizes conforming the microsatellites were 2 (68.2%), 3 (15.9%) and 4 (8.2%) (see [supplementary](#)

FIG. 1.—(a) Circos plot comparing the zebra finch (right hemisphere) and the common chaffinch (left hemisphere) genome assemblies. The common chaffinch chromosomes marked with an asterisk (*) show inversions with respect to the zebra finch assembly. (b) Linear synteny plots of the common chaffinch chromosomes showing inversions relative to the zebra finch generated with the R package *genoPlotR* (Guy et al. 2010). The zebra finch assembly (top) is compared with the common chaffinch assembly (bottom), and numbers designate specific chromosomes.

file S1). The most common length of the microsatellites was 10 bp (40.4%), followed by 12 bp (13%) and 15 bp (8.8%) (see [supplementary file S1](#) for the length distribution of microsatellites). In addition, the number of microsatellites was positively correlated with the sequence length ([Supplementary fig. S3](#), [Supplementary Material](#) online; see [supplementary file S1](#) for the frequency of occurrence in every scaffold).

Gene Annotation and Function Prediction

Our annotation pipeline combining both *de novo* and homology-based predictions inferred 21,831 proteins encoded by 17,703 genes in the common chaffinch genome

with a mean length of 15,818 bp ([Table 1](#)). The common chaffinch genome annotation (see [supplementary file S2](#) in Figshare) included 7,850 complete copies (94.2%) out of the 8,338 of BUSCO avian data set used, retrieving all expected copies with a slight increase from that estimated in the unannotated genome (see above). Among the complete BUSCO genes, 7,827 were single-copy orthologs (99.7%) and 23 were duplicated (0.3%). Around 1.9% (162) of the gene models were fragmented and 3.9% showed no significant matches (326).

Over all predicted proteins, 19,458 (89.1%) provided positive BLASTP hits against the Uniprot SwissProt database, and

Table 1

Genome Statistics and Predicted ncRNAs of the *Fringilla coelebs* Genome Compared with Other Similarly Sized Avian Species (*Melospiza melodia*, *Taeniopygia guttata*, *Ficedula albicollis*, *Manacus vitellinus*, and *Geospiza fortis*), Modified from Louha et al. (2020).

	<i>F. coelebs</i>	<i>M. melodia</i>	<i>T. guttata</i>	<i>F. albicollis</i>	<i>M. vitellinus</i>	<i>G. fortis</i>
Number of genes	17,703	15,086	17,561	16,763	18,976	14,399
Mean gene length (bp)	15,818	14,457	26,458	31,394	27,847	30,164
Number of CDSs	17,703	15,086	17,561	16,763	18,976	14,399
Mean CDs length (bp)	1,679	1,325	1,677	1,942	1,929	1,766
Number of exons	221,872	131,940	171,767	189,043	190,390	164,721
Mean exon length (bp)	165	153	255	253	264	195
Mean number of exons/gene	10.16	8.67	10.25	12.22	11.51	11.41
Number of introns	200,041	116,724	153,909	171,236	171,089	149,563
Mean intron length (bp)	1,902	1,695	2,930	3,257	3,294	2,813
Total proteins	21,831					
ncRNA						
tRNA	325	267	184	179		
miRNA	140	166	302	510		
snRNA	18	16	44	32		
snoRNA	126	154	241	199		
rRNA	5	8	100	22		
lncRNA	17	20	908	1473		

19,617 (89.9%) against the annotated proteins from the zebra finch genome. In addition, InterProScan identified 18,551 (85%) specific protein-domain signatures in the predicted peptides. The combination of the annotation from these databases allowed assigning a functional annotation with GO terms to 19,425 proteins (89%) assigned to 15,309 genes (86.5%; [supplementary file S3](#) in Figshare).

tRNAs and Other Noncoding RNA Prediction

The search by tRNAscan-SE ([supplementary file S4](#) in Figshare) identified 325 tRNAs in the common chaffinch genome, of which 167 decode for the standard twenty amino acids. Among all the tRNAs detected, 131 presented low scores and therefore were categorized as pseudogenes (i.e. lacking tRNA-like secondary structures). There were no suppressor tRNAs, 1 had undetermined isotopes, 25 were chimeric and 15 included introns within their sequences. One of the tRNAs was predicted to code for selenocysteine (sequences and structures of the predicted tRNAs are available in File S5 in Figshare). In addition, the search against both tRNA databases (GtRNAdb and tRNAdb) yielded positive results in many other species, suggesting that tRNA prediction in our assembly was correct. Moreover, our searches using Infernal identified 354 ncRNAs, which were classified as follows: 39 CREs, 2 Ribozymes, 7 Gene, 140 miRNAs, 126 snoRNAs, 18 snRNAs, 5 rRNAs, and 17 lncRNAs (File S6 in Figshare). The number of tRNAs predicted in the common chaffinch genome is the highest when compared with other passerine species (i.e., *M. melodia*, *T. guttata*, and *F. albicollis*), but the other types of ncRNAs present similar values to the *M. melodia* genome and lower than the other two species

([Table 1](#)), probably because we applied a strict threshold to avoid an excess of false positives.

Conclusions

We provide here a high-quality assembly for the common chaffinch, a valuable resource as a reference genome to address a range of biological questions from a genomic perspective. Moreover, our annotation provides useful information to detect candidate genes involved in adaptation and divergence processes. The combination of the Chicago and shotgun sequencing with the HiRise assembly approach lead to a highly contiguous chromosome-level genome assembly. The genome assembly size was 994.87 Mb, with the 30 chromosomes accounting for 98% of it. Although the expected length of the genome was 1.2 Gb, closer to those obtained in other avian species by flow cytometry (Gregory 2002), the BUSCO analyses showed that both the assembly and structural annotation encode 93.9% and 94.2% complete copies out of the 8,338 orthologous conserved genes in avian species, respectively. This discrepancy of the genome size could be caused by the absence of large repetitive elements in the assembly. The structural annotation predicted 17,703 coding genes, with most of them (86.5%) assigned to functional annotation and GO terms.

Materials and Methods

Sample Collection and Genome Assembly

A blood sample was extracted from a common chaffinch female captured in Torreiglesias, Segovia, Spain, in 2017 and frozen immediately in liquid nitrogen. The sample was

handled by Dovetail Genomics for DNA extraction, sequencing and genome assembly using the HiRise pipeline (Putnam et al. 2016). The absence of a Z chromosome in our first assembly (GenBank assembly accession: GCA_015532645.1) led us to conduct a second assembly presented here, which includes sex-linked scaffolds used to reconstruct the chaffinch Z chromosome (see [supplementary methods, Supplementary Material](#) online, for details). Gene completeness in the chaffinch genome assembly (and in the annotated gene set) was assessed through BUSCO (Benchmarking Universal Single-Copy Orthologs) v4.0.5 (Seppey et al. 2019) by using the 8,338 single-copy orthologous genes in the Aves lineage group odb10, using chicken as the Augustus reference species.

Identification of Repetitive Regions and Gene Annotation

Repetitive regions were identified and masked prior to gene prediction. First, repeats were modelled ab initio using Repeat Modeler 1.0.11 (Smit and Hubley 2019) in scaffolds longer than 100 Kb with default options. The repeats obtained were merged with known bird repeat libraries from the RepBase database (RepBase-20181026) (Bao et al. 2015), Dfam_Consensus-20181026 and repeats from the zebra finch (obtained from B10K). The resulting repeat library was compared against the complete assembly with Repeat Masker 4.0.7 (Smit et al. 2015) and the identified regions were soft-masked. For the identification and description of microsatellites in the common chaffinch genome assembly we used GMATA v.2.01 (Wang and Wang 2016), with sequence motif length between 2 and 20 bp.

Gene prediction was conducted with BRAKER v2.1.5 (Hoff et al. 2016) and GeMoMa v1.7.1 (Keilwagen et al. 2016, 2018). First, the conserved orthologous genes from BUSCO Aves_odb10 were used as proteins from short evolutionary distance to train Augustus (Gremme et al. 2005; Stanke et al. 2006; see figure 3B in Hoff et al. 2019). The predicted proteins were combined with homology-based annotations using the zebra finch (GCF_008822105.2; Warren et al. 2010) and chicken (GCF_000002315.6; Hillier et al. 2014) annotated genes with GeMoMa pipeline, obtaining the final reported gene models. We applied a similarity-based search approach to assist the functional annotation of the chaffinch predicted proteins, using the UniProt SwissProt database, the annotated proteins from the zebra finch genome (Warren et al. 2010; UniProt Consortium 2014) and InterProScan v5.31 (Jones et al. 2014). The functional annotation, including Gene Ontology terms, was integrated from all searches providing a curated set of chaffinch coding genes (see [supplementary methods, Supplementary Material](#) online, for details).

Noncoding RNA Prediction and Identification

For the prediction and functional classification of Transfer RNAs (tRNAs) in the common chaffinch genome we used

tRNAscan-SE v2.0 (Lowe and Chan 2016). The tRNA search across the genome and the identification of ncRNA (noncoding RNA) homologues was conducted using the software package Infernal v1.1.1 (Nawrocki 2014) (see [supplementary methods, Supplementary Material](#) online, for details). For comparative purposes, we added our results to those from Louha et al. (2020), which compared different genome assemblies of avian species.

Supplementary Material

[Supplementary data](#) are available at *Genome Biology and Evolution* online.

Acknowledgments

This research was supported by the Spanish Ministry of Economy and Competitiveness (CGL2015-66381P to B.M. and G.B.) and the Spanish Ministry of Science and Innovation (PGC2018-098897-B-I00 from to B.M.). M.R. was supported by a doctoral fellowship from the Spanish Ministry of Education, Culture, and Sport (FPU16/05724).

Data Availability

The chaffinch genome assembly has been deposited at NCBI under BioProject PRJNA674347 with accession number JADKPM000000000, the raw data are available at SRA NCBI database with accession numbers SRR12998620–SRR12998622, and the annotation and all described data sets are publicly accessible in Figshare (<https://doi.org/10.6084/m9.figshare.13296122.v3>).

Literature Cited

- Alström P, et al. 2018. Complete species-level phylogeny of the leaf warbler (Aves: Phylloscopidae) radiation. *Mol Phylogenet Evol.* 126:141–152.
- Aslam ML, et al. 2010. A SNP based linkage map of the turkey genome reveals multiple intrachromosomal rearrangements between the turkey and chicken genomes. *BMC Genomics* 11(1):647–611.
- Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob DNA.* 6:11.
- Barton NH. 1996. Natural selection and random genetic drift as causes of evolution on islands. *Philos Trans R Soc Lond B Biol Sci.* 351(1341):785–795.
- Braun EL, Cracraft J, Houde P. 2019. Resolving the avian tree of life from top to bottom: the promise and potential boundaries of the phylogenomic era. In: Kraus R, editor. *Avian genomics in ecology and evolution*. Cham (Switzerland): Springer. p. 151–210.
- Brown RM, et al. 2013. Evolutionary processes of diversification in a model island archipelago. *Annu Rev Ecol Evol Syst.* 44(1):411–435.
- Collar N, Newton I, Bonan A. 2020. Finches (*Fringillidae*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E, editors. *Handbook of the birds of the world alive*. Barcelona (Spain): Lynx Edicions. Available from: <https://www.hbw.com/node/52376> [accessed 2020 Jan 15].

- Dalloul RA, et al. 2010. Multi-platform next-generation sequencing of the domestic turkey (*Meleagris gallopavo*): genome assembly and analysis. *PLoS Biol.* 8(9):e1000475.
- Ducressat AL, et al. 2020. New genome assembly of the barn owl (*Tyto alba alba*). *Ecol Evol.* 10(5):2284–2298.
- Feng S, et al. 2020. Dense sampling of bird diversity increases power of comparative genomics. *Nature* 587(7833):252–257.
- Frankl-Vilches C, et al. 2015. Using the canary genome to decipher the evolution of hormone-sensitive gene regulation in seasonal singing birds. *Genome Biol.* 16(1):19.
- Friis G, et al. 2018. Genome-wide signals of drift and local adaptation during rapid lineage divergence in a songbird. *Mol Ecol.* 27(24):5137–5153.
- Gregory TR. 2002. Animal genome size database. Available from: <http://www.genomesize.com>.
- Gremme G, Brendel V, Sparks ME, Kurtz S. 2005. Engineering a software tool for gene structure prediction in higher organisms. *Inf Softw Technol.* 47(15):965–978.
- Guy L, Roat Kultima J, Andersson SG. 2010. genoPlotR: comparative gene and genome visualization in R. *Bioinformatics* 26(18):2334–2335.
- Hillier LW, et al. 2014. Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature* 423:695–777.
- Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1: unsupervised RNA-Seq-based genome annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics* 32(5):767–769.
- Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-genome annotation with BRAKER. In: Kollmar M, editor. *Gene prediction*. New York: Humana. p. 65–95.
- Hooper DM, Price TD. 2017. Chromosomal inversion differences correlate with range overlap in passerine birds. *Nat Ecol Evol.* 1(10):1526–1534.
- Illera JC, et al. 2018. Acoustic, genetic, and morphological analyses of the Canarian common chaffinch complex *Fringilla coelebs* ssp. reveals cryptic diversification. *J Avian Biol.* 49(12):1–12.
- Jarvis ED, et al. 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science* 346(6215):1320–1331.
- Jarvis ED, et al.; The Avian Phylogenomics Consortium. 2015. Phylogenomic analyses data of the avian phylogenomics project. *GigaScience* 4(1):s13742-014.
- Jones P, et al. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* 30(9):1236–1240.
- Keilwagen J, et al. 2016. Using intron position conservation for homology-based gene prediction. *Nucleic Acids Res.* 44(9):e89.
- Keilwagen J, Hartung F, Paulini M, Twardziok SO, Grau J. 2018. Combining RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC Bioinformatics* 19(1):189.
- Lawson LP, Petren K. 2017. The adaptive genomic landscape of beak morphology in Darwin's finches. *Mol Ecol.* 26(19):4978–4989.
- Leroy T, et al. 2021. Forthcoming. Endemic island songbirds as windows into evolution in small effective population sizes. *Curr Biol.* Available from: 10.1016/j.cub.2020.12.040.
- Losos JB, Schluter D. 2000. Analysis of an evolutionary species–area relationship. *Nature* 408(6814):847–850.
- Louha S, Ray DA, Winker K, Glenn TC. 2020. A high-quality genome assembly of the North American Song Sparrow, *Melospiza melodia*. *G3 (Bethesda)* 10(4):1159–1166.
- Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res.* 44(W1):W54–W57.
- MacArthur RH, Wilson EO. 1967. *The theory of island biogeography*. Princeton (NJ): Princeton University Press. 203 p.
- Nawrocki EP. 2014. Annotating functional RNAs in genomes using Infernal. In: Gorodkin J, Ruzzo W, editors. *RNA sequence, structure, and function: computational and bioinformatic methods*. Totowa (NJ): Humana Press. p. 163–197.
- Peñalba JV, et al. 2020. Genome of an iconic Australian bird: High-quality assembly and linkage map of the superb fairy-wren (*Malurus cyaneus*). *Mol Ecol Resour.* 20(2):560–578.
- Poelstra JW, et al. 2014. The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. *Science* 344(6190):1410–1414.
- Putnam NH, et al. 2016. Chromosome-scale shotgun assembly using an in vitro method for long-range linkage. *Genome Res.* 26(3):342–350.
- Seppy M, Manni M, Zdobnov EM. 2019. BUSCO: assessing genome assembly and annotation completeness. In: Kollmar M, editor. *Gene prediction*. New York: Humana. p. 227–245.
- Skinner BM, Griffin DK. 2012. Intrachromosomal rearrangements in avian genome evolution: evidence for regions prone to breakpoints. *Heredity (Edinb.)* 108(1):37–41.
- Smit A, Hubley R. 2019. RepeatModeler-1.0. 11. Institute for Systems Biology. Available from: <http://www.repeatmasker.org/RepeatModeler/>
- Smit AFA, Hubley R, Green P. 2015. RepeatMasker Open-4.0. 2013–2015. Available from: <http://www.repeatmasker.org>.
- Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics.* 7:62.
- UniProt Consortium. 2014. UniProt: a hub for protein information. *Nucleic Acids Res.* 43:D204–D212.
- Valente L, et al. 2020. A simple dynamic model explains the diversity of island birds worldwide. *Nature* 579(7797):92–96.
- Vijay N, et al. 2016. Evolution of heterogeneous genome differentiation across multiple contact zones in a crow species complex. *Nat Commun.* 7(1):13195.
- Völker M, et al. 2010. Copy number variation, chromosome rearrangement, and their association with recombination during avian evolution. *Genome Res.* 20(4):503–511.
- Wang X, Wang L. 2016. GMATA: an integrated software package for genome-scale SSR mining, marker development and viewing. *Front Plant Sci.* 7:1350.
- Wang W, et al. 2020. First de novo whole genome sequencing and assembly of the bar-headed goose. *PeerJ.* 8:e8914.
- Warren WC, et al. 2010. The genome of a songbird. *Nature* 464(7289):757–762.
- Wirthlin M, Lovell PV, Jarvis ED, Mello CV. 2014. Comparative genomics reveals molecular features unique to the songbird lineage. *BMC Genomics* 15:1082.
- Zhang G, et al. 2015. Genomics: bird sequencing project takes off. *Nature* 522(7554):34.
- Zhang G, et al.; Avian Genome Consortium. 2014. Comparative genomics reveals insights into avian genome evolution and adaptation. *Science* 346(6215):1311–1320.
- Zhang G, Parker P, Li B, Li H, Wang J. 2012. The genome of Darwin's finch (*Geospiza fortis*). *GigaScience. Database.* Available from: 10.5524/100040.

Associate editor: Bonnie Fraser