

Unique spicules may confound species differentiation: taxonomy and biogeography of *Melonanchora* Carter, 1874 and two new related genera (Myxillidae: Poecilosclerida) from the Okhotsk Sea

Andreu Santín¹, María-Jesús Uriz², Javier Cristobo^{3,4}, Joana R. Xavier^{5,6} and Pilar Ríos^{3,7}

¹ Institut de Ciències del Mar (ICM-CSIC), Barcelona, Catalonia, Spain

² Centre d'Estudis Avançats de Blanes (CEAB-CSIC), Blanes, Catalonia, Spain

³ Instituto Español de Oceanografía. Centro Oceanográfico de Gijón., Gijón, Asturias, Spain

⁴ Departamento de Ciencias de la Vida, EU-US Marine Biodiversity Group, Universidad de Alcalá, Alcalá de Henares, Madrid, Spain

⁵ CIIMAR—Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Porto, Portugal

⁶ Department of Biological Sciences, University of Bergen, Bergen, Norway

⁷ Departamento de Biología Animal, Universidad de Málaga, Málaga, Spain

ABSTRACT

Sponges are amongst the most difficult benthic taxa to properly identify, which has led to a prevalence of cryptic species in several sponge genera, especially in those with simple skeletons. This is particularly true for sponges living in remote or hardly accessible environments, such as the deep-sea, as the inaccessibility of their habitat and the lack of accurate descriptions usually leads to misclassifications. However, species can also remain hidden even when they belong to genera that have particularly characteristic features. In these cases, researchers inevitably pay attention to these peculiar features, sometimes disregarding small differences in the other “typical” spicules. The genus *Melonanchora* Carter, 1874, is among those well suited for a revision, as their representatives possess a unique type of spicule (spherancorae). After a thorough review of the material available for this genus from several institutions, four new species of *Melonanchora*, *M. tumultuosa* sp. nov., *M. insulsa* sp. nov., *M. intermedia* sp. nov. and *M. maeli* sp. nov. are formally described from different localities across the Atlanto-Mediterranean region. Additionally, all *Melonanchora* from the Okhotsk Sea and nearby areas are reassigned to other genera; *Melonanchora kobjakovae* is transferred to *Myxilla* (*Burtonanchora*) while two new genera, *Hanstoreia* gen. nov. and *Arhythmata* gen. nov. are created to accommodate *Melonanchora globogilva* and *Melonanchora tetrademitifera*, respectively. *Hanstoreia* gen. nov. is closest to *Melonanchora*, whereas *Arhythmata* gen. nov., is closer to *Stelodoryx*, which is most likely polyphyletic and in need of revision.

Submitted 5 August 2021

Accepted 27 October 2021

Published 22 December 2021

Corresponding author

Andreu Santín, santin@icm.csic.es

Academic editor

Alexander Ereskovsky

Additional Information and
Declarations can be found on
page 50

DOI 10.7717/peerj.12515

© Copyright

2021 Santín et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biodiversity, Marine Biology, Taxonomy, Zoology

Keywords Porifera, *Melonanchora*, Deep-Sea, Spicule evolution, New species, Atlanto-Mediterranean, Okhotsk Sea, Aleutian Islands, Poecilosclerida, Biogeography

INTRODUCTION

Accurate species-level taxonomy is a fundamental keystone for conservation assessment, planning, and management (Myers *et al.*, 2000; Groves *et al.*, 2017). The differentiation between cryptic species (as in Knowlton, 1993), is of paramount importance for effective conservation policies (Lohman *et al.*, 2010). While cryptic species are a widespread phenomenon among both terrestrial and marine phyla (e.g., Baker, 1984; Mayer & Helversen, 2001; Concepción *et al.*, 2008; Crespo & Pérez-Ortega, 2009; Dennis & Hellberg, 2010; Lohman *et al.*, 2010; Payo *et al.*, 2013; Golestani *et al.*, 2019), the assumed lack of barriers to gene flow in marine habitats (Hellberg, 2009) contributed to the assumption that benthic organisms have greater distribution ranges and phenotypic plasticity than their terrestrial counterparts (Knowlton, 1993). As a result of this assumption, many benthic species were considered to be geographically widespread or even cosmopolitan (Klautau *et al.*, 1999). Recent studies have generally demoted this idea (e.g., Klautau *et al.*, 1999; van Soest, Hooper & Hiemstra, 1991; van Soest & Hooper, 1993). The dispersal capabilities vary greatly among benthic species even within the same phyla (Uriz *et al.*, 1998) and they can be differentially reduced by natural barriers (Allcock *et al.*, 1997; Waters & Roy, 2004). In this sense, some invertebrate Phyla, such as sponges and corals, produce short-lived, free larvae that are seemingly incapable of countering apparently weak marine barriers such as littoral currents or substrate discontinuity, often resulting in extremely low dispersal capabilities (Hellberg, 2009). In sponges, for instance, genetically structured populations, even at short spatial scales, have been repeatedly reported (Duran *et al.*, 2004; Duran, Pascual & Turon, 2004; Calderon *et al.*, 2007; Blanquer, Uriz & Caujapé-Castells, 2009; Blanquer & Uriz, 2010; Guardiola, Frotscher & Uriz, 2016), which favours speciation and makes the existence of widely distributed or cosmopolitan species unlikely.

Species complexes and cryptic species are particularly prevalent among sponges with few diagnostic characters (Klautau *et al.*, 1999; Uriz, Garate & Agell, 2017a, 2017b), especially when these characters exhibit environmental plasticity (Maldonado *et al.*, 1999; Xavier *et al.*, 2010; De Paula *et al.*, 2012). For example, the sponge complex *Chondrilla nucula* Schmidt, 1862, was once considered as having a circumtropical distribution (Klautau *et al.*, 1999), *Stylocordyla borealis* (Lovén, 1868) was reported as occurring at both poles (Uriz *et al.*, 2010), the Atlanto-Mediterranean *Scopalina lophyropoda* Schmidt, 1862 and *Hemimycale columella* (Bowerbank, 1874) both contained several morphologically cryptic species revealed by molecular markers (Blanquer & Uriz, 2008; Uriz, Garate & Agell, 2017a, 2017b) and the excavating sponges *Cliona celata* Grant, 1826 and *Cliona viridis* (Schmidt, 1862), which are known to be “species complexes” which remain partially unresolved (Xavier *et al.*, 2010; De Paula *et al.*, 2012; Escobar, Zea & Sánchez, 2012; Leal *et al.*, 2016; Gastaldi *et al.*, 2018). Cryptic species complexes are also prevalent in sponge genera without mineral (spicules) or organic skeletons (spongin fibres), such as

Hexadella Topsent, 1896, where species are almost indistinguishable based solely on morphological or histological characteristics (Reveillaud et al., 2010, 2012). However, species can also remain hidden even when they belong to genera that have particularly characteristic spicules. In these cases, researchers inevitably pay attention to these peculiar spicules, sometimes disregarding small differences in the other “typical” spicules.

Some genera of Poecilosclerida, one of the most diverse orders in terms of spicule diversity (Hooper & Van Soest, 2002), possess unique spicular types that greatly facilitate their identification. Examples include dianciastras in *Hamacantha Gray, 1867* (Hajdu, 1994; Hajdu & Castello-Branco, 2014), clavidiscs in *Merlia Kirkpatrick, 1908* (Vacelet & Uriz, 1991), discorhabds in *Latrunculia du Bocage, 1869* (Samaai, Gibbons & Kelly, 2006) or thraustoxeas in *Rhabderemia Topsent, 1890* (van Soest & Hooper, 1993). Nevertheless, because taxonomists historically have focused on these particular spicules (van Soest, Hooper & Hiemstra, 1991), differences in other apparently banal spicules have been disregarded. As a consequence, some of these genera (e.g., *Rhabderemia van Soest & Hooper (1993)*, *Acarnus, Gray, 1867*, *van Soest, Hooper & Hiemstra (1991)*, *Merlia, Vacelet & Uriz (1991)* or *Trachytedania Ridley, 1881* (Cristobo & Ugorri (2001)) contain or contained, until recently, few formally described species that were considered as having a widespread geographic distribution. Moreover, only the well-described species are usually recognised and reported in the literature (van Soest, Hooper & Hiemstra, 1991), while those with poor or imprecise descriptions remain forgotten, a trend which is aggravated for sponges living in remote or hardly accessible environments, such as the deep-sea (Reveillaud et al., 2010). Despite the challenges involved, comprehensive reviews of such genera are considered extremely useful for the discovery of cryptic species (Reveillaud et al., 2012) and to test biogeographical and evolutionary hypotheses (van Soest & Hooper, 1993; Cárdenas et al., 2007).

The genus *Melonanchora Carter, 1874*, is among those well suited for such revisions, as (i) it possesses a unique spicule type (spherancorae); (ii) currently contains only five formally accepted species (van Soest et al., 2021) (iii) only two out of the five species are commonly recorded over large geographical areas (Baker et al., 2018) and (iv) the three remaining species seem to be endemic to the Okhotsk Sea and nearby Pacific Islands (Koltun, 1958, 1970; Lehnert, Stone & Heimler, 2006a) and present clear differences with their Atlantic counterparts (Lehnert, Stone & Heimler, 2006a). Finally, *Melonanchora* representatives occur within Vulnerable Marine Ecosystems (VMEs) across the Atlanto-Mediterranean region, thus being in need of accurate identifications for the evaluation of the conservation status of the sponge grounds where they occur (Best et al., 2010; ICES, 2012).

In this context, this paper: (1) reviews the status of all the species currently allocated to *Melonanchora* with particular emphasis in the Pacific species, apparently endemic to the Okhotsk Sea, and their relationships with other Myxillidae; (2) provides a reliable guide for their identification; (3) describes new species of the genus; (4) and discusses the biogeographical implications of the circumpolar distribution of this genus.

MATERIALS AND METHODS

Museum material and sample treatment

The materials for this study consisted of samples from natural history museums and other scientific institutions and unregistered individuals from surveys across the North Atlantic (Life+ INDEMARES, NEREIDA and ABIDES) as well as specimens from authors' own collections. The institutions are abbreviated in the text as follow:

Canadian Museum of Nature, Canada (CMNI), using the prefix CMNI; Gothenburg Natural History Museum, Sweden (GNM), using the prefix GNM; Museo Civico di Storia Naturale di Genova, Italy (MSNG), using the prefix MSNG; Museum of Biology of Lund, Sweden (MZLU), using the prefix MZLU; Naturalis Biodiversity Center, The Netherlands (NBC, previously ZMA), using the prefix ZMA.POR. and ZMA.POR.P; National Museum of Natural History, Smithsonian Institution, Unites States (NMNH, previously USNM) using the prefix NMNH-USNM; Musée Zoologique de la Ville de Strasbourg, France (MZS) using the prefix MZS; Museu de Ciències Naturals (Zoologia) de Barcelona, Spain (MZB), using the prefix MZB; National History Museum, United Kingdom (NHMUK, previously BMNH), using the prefix NHMUK; Swedish Museum of Natural History, Sweden (NRM), using the prefix NRM; Yale Peabody Museum of Natural History, Unites States (YPM), using the prefix YPM IZ; Museum für Naturkunde, Germany (ZMB) using the prefix ZMB; Jean Vacelet's personal collections (JV) and Manuel Solórzano's personal collections (MS).

DNA was extracted from small pieces of tissue of four samples (MSNG Vis4.7, CMNI 2018-0107, GNM Por624, NMNH-USNM 1082996) using QIAGEN's DNeasy Blood and Tissue kit, following the instructions of the manufacturer. Amplification and sequencing of the mitochondrial cytochrome c oxidase subunit I (COI) were attempted but proved unsuccessful, with only two samples yielding an amplicon but resulting in sequencing of non-target DNA (bacteria). This was likely due to the low quantity and integrity of the DNA in the samples, as assessed by spectrophotometry using a DeNovix DS-11 FX.

All known species of *Melonanchora* were represented in the studied material, with the holotypes for all species but *Melonanchora tetradedritifera* [Koltun, 1970](#) being examined. Spicule preparations for both optical and scanning electron microscopy (SEM) were performed according to [Cristobo et al. \(1993\)](#) and [Uriz, Garate & Agell \(2017a\)](#). Optical observations were performed using a Leica DM IRB inverted microscope from the Instituto de Ciencias del Mar (ICM-CSIC), whereas SEM observation were conducted using an ITACHI TM3000 TableTop Scanning Electron Microscope from the Center for Advanced Studies of Blanes (CEAB-CSIC), Spain, a JEOL-6100 SEM from the University of Oviedo (UO), Spain, and a HITACHI S-3500 N scanning electron microscope from the Institut de Ciències del Mar (ICM-CSIC), Spain. Spicule sizes are given as ranges with average values (in italics) \pm Standard Deviation (e.g., MIN.–MEAN \pm SD–MAX.). Unless otherwise stated, spicule measurements were performed on 40 spicules per spicule type. The species classification adopted in the study follows that currently proposed by [Morrow & Cárdenas \(2015\)](#) and the World Porifera Database ([van Soest et al., 2021](#)). A key to *Melonanchora* can be found in [Supplemental Material 1](#).

Finally, the electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is: [urn:lsid:zoobank.org:pub:F1A22CAA-DE1F-434D-9A6B-F00853C40FF5]. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central SCIE and CLOCKSS.

RESULTS

Systematic Description

Phylum PORIFERA *Grant, 1836*

Class DEMOSPONGIAE *Sollas, 1885*

Subclass Heteroscleromorpha *Cárdenas, Pérez & Boury-Esnault, 2012*

Order POECILOSCLERIDA *Topsent, 1928*

Family MYXILLIDAE *Dendy, 1922*

Genus *Melonanchora* *Carter, 1874*

Type species:

Melonanchora elliptica *Carter, 1874*: 212 (by monotypy).

Diagnosis:

From encrusting to massive-globular growth form, with paper-like, easily detachable thin ectosome, bearing fistular processes. Ectosomal skeleton composed of smooth strongyles to tylotes with somewhat asymmetrical ends, whereas the choanosome is mainly composed of smooth strongyles or styles. Microscleres include typically two categories of anchorate isochelae, rarely three, and spherancorae (amended from *van Soest, 2002*).

Remarks:

The genus *Melonanchora* was erected by *Carter (1874)* for *Melonanchora elliptica* on the account of this species singular anchorate-derived chelae (spherancorae), placing it tentatively with the “*Halichondria*” family concept built around *H. (= Myxilla) incrustans* (*Johnston, 1842*). The genus was later included in Desmacidonidae *Schmidt (1880)* until *Lundbeck (1910)*, and later *Topsent (1928)*, transferred it to Myxillidae. Simultaneously, Hentschel had it assigned it to Dendoricellidae¹ (*Hentschel, 1929*), but this assignment was not widely accepted (*Alander, 1935*) and was quickly disregarded.

The family Myxillidae has been redefined over the years (*Hajdu, van Soest & Hooper, 1994*; *Desqueyroux-Faúndez & van Soest, 1996*; *van Soest, 2002*) and the genus *Melonanchora* fits well within the current definition of Myxillidae established in the Systema Porifera (*Hooper & Van Soest, 2002*), which is restricted to “those genera which combine the possession of anchorate chelae with diactinal ectosomal tornotes

¹ While Hentschel assigned it to Dendoricellidae, he later wrongfully referred *Melonanchora* as part of Tedanidae within the text.

[oxeotes and tylotes] and choanosomal styles in a reticulate arrangement”. Yet, after re-examination of all the available *Melonanchora* material, the current definition of the genus ([van Soest, 2002](#)) needs to be amended to better allocate the new species here described or re-described, including: presence of smooth strongyles (*Melonanchora emphysema* ([Schmidt, 1875](#)), *M. tumultuosa* sp. nov., *Melonanchora intermedia* sp. nov.) as choanosomal megascleres and the possession of two to three chelae categories (*M. intermedia* sp. nov., *M. maeli* sp. nov.).

Nevertheless, the main diagnostic character of the genus, the spherancorae, remains unaltered since Carter’s original description (See “The Origin of Spherancorae”). Aside from spherancorae, Carter also added the presence of a papillated paper-thin like ectosome (Figs. 1A, 1C, and 1F) as an additional diagnostic character ([Carter, 1874](#)). Although this feature is shared with other deep-sea genera such as *Cornulum* [Carter, 1876](#) or *Coelosphaera* [Thomson, 1873](#) ([Lehnert & Stone, 2015](#); [Schejter, Cristobo & Ríos, 2019](#)), *Melonanchora* differs from the later in its white-translucent coloration, brittle and loose appearance and its characteristic wart-shaped papillae, which may make external identification feasible at the genus level ([Stone, Lehnert & Reiswig, 2011](#)).

Melonanchora elliptica [Carter, 1874](#)
(Figs. 1A, 2, 3)

Synonymy:

Melonanchora elliptica [Carter, 1874](#): 212, pl. XIII figs 6–12, pl. XV figs. 35a–35b; [Vosmaer, 1885](#): 31, pl. I fig. 14, pl. V figs. 69–70 (*partim*); [Topsent, 1892](#): 101–102; [Fristedt, 1887](#): 454, pl. 25 fig. 5, 55 (*partim*); [Arnesen, 1903](#): 15–16, pl. II fig. 4, pl. V fig. 4; [Topsent, 1904](#): 144, pl. IV fig. 10; [Lundbeck, 1905](#): 213–216, pl. VII figs. 4–6, pl. XX figs. 1a–1o; [Lundbeck, 1909](#): 402–403; [Arndt, 1913](#): 116; [Topsent, 1913](#): 44; [Topsent, 1928](#): 246; [Hentschel, 1929](#): 966; [Burton, 1931](#): 4; [Alander, 1935](#): 5; [Arndt, 1935](#): 71–73, Fig. 141; [Koltun, 1959](#): 122–123, fig. 76; [Ríos & Cristobo, 2017](#): 169; [Baker et al., 2018](#): 20–25, figs. 5–7; [Dinn & Leys, 2018](#): 63.

Not: *M. elliptica*; [Schmidt, 1880](#): 85, pl. IX fig. 8.

Material examined.

Holotype: NHMUK 1882.7.28.54a, between the north coast of Scotland and the Faroe Islands; *HMS Porcupine* expedition (1869), ca. 800 m depth, 1869. (two slides); NHMUK-Norman Coll. N°50 10.1.1.1417, *HMS Porcupine* expedition (1869); NHMUK 1954.3.9.301 N°50; NHMUK - Norman Coll. -H. J. Carter Slide Coll. 1954.3.9.301; ZMB Por 3042, between the North coast of Scotland and the Faroe Islands, North Atlantic Ocean (59.85166, –6.03333).

Additional specimens examined:

CMNI 2018-0107, Saglek Bank, Labrador Sea, North Atlantic Ocean (60.45213, –61.26894), 427 m depth, 2016-07-21, collected by Dinn, Curtis ([Dinn & Leys, 2018](#)); MZLU L936/3483, Trondheim Fjord, Norway (63.494092, 10.31647), 1936; NRM 113070, off Lindenows Fjord, Greenland, North Atlantic Ocean (60.06666, –34.25), 237.9 m depth,




Figure 1 External appearance of various *Melonanchora* species. (A) External view of *Melonanchora elliptica* (MZLU L935/3858), *p* indicates some ectosomal papillae; (B) Individual of *Melonanchora emphysema* (*Me*) attached to coral rubble (GNM Porifera 416); (C) Holotype of *Melonanchora tumultuosa* sp. nov. (GNM Porifera 624), *Ect* indicates the ectosome, *Ch* indicates the choanosome, *Cha* indicates the choanosomal cavities, *Os* indicates the oscules; (D) Individual of *Arythmata tetradentifera* (NMNH-USNM 148959); (E) Holotype of *Melonanchora insulsa* sp. nov. (MZS Po165); (F) Holotype of *Hanstoreia globogilva* (NMNH-USNM 1082996), *p* indicates some ectosomal papillae and *Os* indicates the oscules; (G) Holotype of *Melonanchora maeli* sp. nov. (ZMA.POR.7269), *p* indicates some ectosomal papillae and *Ch* the choanosome.

Full-size DOI: [10.7717/peerj.12515/fig-1](https://doi.org/10.7717/peerj.12515/fig-1)

1885 (*Fristedt, 1887*); YPM IZ 006552.PR, Laurentian Channel, Nova Scotia, North Atlantic Ocean (44.5667, -56.6958), *USFC Albatross*, 218 m depth, 1885; NHMUK-Norman Collection 1910.1.1.588, Hardanger Fjord, ca. 180 m depth, 1882; NHMUK-Sott-Ryen Coll., 1931.6.1.19, Folden Fjord, Norway (*Burton, 1931*); NHMUK Norman Coll. 1910.1.1.1418, Norway, 1882; NHMUK-Norman Coll. 1910.1.1.1419, Norway, 1882; NHMUK-Norman Coll. 1910.1.1.1420, Norway, 1882; NHMUK-Norman Coll. 1910.1.1.1421 (*Fristedt, 1887*); NHMUK-Norwegian Coll. 1982.9.6.14.a., Norway, 1885; ZMA.POR.P.10797, North of Hammerfest, Norway, Arctic Ocean (72.15003, 22.71246),

R/V Willem Barents expedition (1880–84), 265 m depth, 1881 ([Vosmaer, 1885](#)); ZMA. POR.1548, North of Hammerfest, Norway, Arctic Ocean (72.15, 22.68333), *R/V Willem Barents* expedition (1880–84), 265 m depth, 1881 ([Vosmaer, 1885](#)).

Unregistered material:

NR0509_43, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,554 m depth (NEREIDA Coll.); NR0509_49, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,137 m depth (NEREIDA Coll.); NR0509_52, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 870 m depth (NEREIDA Coll.); NR0509_73, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,122 m depth (NEREIDA Coll.); NR0509_82a, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,127 m depth (NEREIDA Coll.); NR0610_21, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,055 m depth (NEREIDA Coll.); NR0709_5, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,248 m depth (NEREIDA Coll.).

Description:

Usually massive-globular sponge ([Fig. 1A](#)), more rarely encrusting (CMN 2018-0107), with an easily detachable paper-like thin ectosome bearing abundant fistular processes. The choanosome shows several scattered pores and channels. Colour whitish translucent outside, cream-orange in the choanosome.

Skeleton:

Ectosomal skeleton consists of tangential tylostrongyles with a criss-cross arrangement ([Fig. 2C](#)). Choanosomal skeleton with scattered poorly defined tracts ([Fig. 2B](#)) of styles to substyles and abundant organic content. Microscleres are distributed thorough the choanosome without any clear discernible pattern, yet, in some individuals (including the holotype), spherancorae form a dense palisade between the ectosome and the choanosome and might also cover the choanosomal tracts ([Fig. 2D](#)).

Spicular complement:

Styles, tylostrongyles, two categories of chelae, and spherancorae ([Figs. 3A–3G](#)).

Ectosomal tylostrongyles ([Fig. 3B](#)): Unevenly, slightly flexuous unequally thinning towards both ends, with a more or less central swelling and, differentially inflated ends (strongyle to tylote appearance).

Size range: $560.3\text{--}624.3 \pm 32.2\text{--}666.5 \mu\text{m} \times 7.8\text{--}11.8 \pm 3\text{--}17.3 \mu\text{m}$

Choanosomal styles ([Fig. 3A](#)): Entirely smooth, slightly curved towards its distal end. In general, they have the point markedly acerate, but points can also be blunt to various degrees in some spicules (stylostrongyles) ([Fig. 3F](#)).

Size range: $782.5\text{--}830.8 \pm 50\text{--}908.1 \mu\text{m} \times 17.2\text{--}19.3 \pm 1.1\text{--}20.5 \mu\text{m}$

Isochelae I ([Fig. 3E, c'](#)): Small anchorate isochelae, with a straight shaft, well-developed fimbriae and spatulated alae. The distal alae slightly point outwards, giving a “V” lateral appearance to both ends.




Figure 2 Optical microscope imaging of *Melonanchora* spicules. (A) General view of the spicules of *Melonanchora* (NHMUK 1882.7.28.54a) un light microscopy. C. I indicates the largest chelae category, C. II indicates the smallest chelae category, and Sph indicates spherancorae; (B) View of the loose choanosomal tracts off *Melonanchora elliptica* (NHMUK 1882.7.28.54a) (C) View of the characteristic criss-cross like pattern of the ectosome of *Melonanchora* (NHMUK–Norman Coll. 1910.1.1.1421); (D) Spherancorae covering the choanosomal tracts in *Melonanchora elliptica* (NHMUK 1882.7.28.54a). [Full-size](#) DOI: 10.7717/peerj.12515/fig-2

Size range: $24.2\text{--}26.6 \pm 3.4\text{--}29 \mu\text{m}$

Isochelae II (Fig. 3D, b’): large isochelae with a straight shaft, well-developed fimbriae and spatulated alae. The distal alae slightly point outwards, giving a “V” lateral appearance to both ends.

Size range: $48.3\text{--}51.1 \pm 3.8\text{--}58 \mu\text{m}$

Spherancorae (Fig. 3C, a’): Unique to the genus, with an oval shape and slightly pointed ends, which might resemble a rugby ball. It possesses fimbriae on its internal face, which may be free or fused to various degrees.

Size range: $48.3\text{--}51.2 \pm 2.7\text{--}53.1 \times 23.1\text{--}28.3 \pm 1.6\text{--}29.2 \mu\text{m}$

Geographic distribution and ecological remarks:

Melonanchora elliptica is a common amphi-Atlantic species (Fig. 4) also occurring in Arctic waters (Carter, 1877), as far as the Barents Sea (Koltun, 1959; Katckova et al., 2018). It has been recorded from the coasts of Norway (Vosmaer, 1885; Topsent, 1913), Faroe Plateau (Carter, 1874; Lundbeck, 1905), Porcupine Seamount (Könnecker & Freiwald, 2005; van Soest & De Voogd, 2015) and Rockall Bank (van Soest & Lavaleye, 2005),




Figure 3 *Melonanchora elliptica* spicule plate. Spicular set for *Melonanchora elliptica* (sample NHMUK 1882.7.28.54a., holotype). (A) Choanosomal style; (B) Ectosomal tylostrongyle; (C) Spherancorae; (D) Large chelae category (Chelae II); (E) small chelae category (Chelae I); (F) Detail of the styles' acerate end; (G) General view of *M. elliptica*'s spicules by SEM imaging. (a') Spherancora

Figure 3 (continued)

(b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b'), (c') 300 μm ; (C)–(F) 30 μm and (G) 500 μm . Images (A) to (E) and (G) were taken from sample NHMUK 1882.7.28.54a (holotype). Images for F were taken from both NHMUK 1882.7.28.54a (holotype) and CMNI 2018-0107. [Full-size !\[\]\(2bdfe261b986065ee0ac76460d6528c9_img.jpg\) DOI: 10.7717/peerj.12515/fig-3](https://doi.org/10.7717/peerj.12515/fig-3)

Greenland and Iceland ([Lundbeck, 1905](#); [Burton, 1959](#)), the Galician coast ([Ríos & Cristobo, 2017](#)), the Azores archipelago ([Topsent, 1892, 1904, 1928](#)) and the area within the Labrador Peninsula and the Newfoundland Seas ([Topsent, 1913](#); [Michaud & Pelletier, 2006](#); [Baker et al., 2018](#)), from 80 to 1,554 m depth. In the Canadian coasts and the Gulf of Maine, the species is commonly found on sponge grounds on trawlable areas ([Maciolek et al., 2008, 2011](#)) and it has been observed to be an occasional nursery ground for the octopus *Rossia palpebrosa* [Owen, 1935](#) ([Wareham Hayes, Fuller & Shea, 2017](#)). Nevertheless, its role and ecological significance in Vulnerable Marine Ecosystems (VMEs) are still poorly understood and in need of further research.

Remarks:

Melonanchora elliptica is the type species of the genus, first described from a specimen collected during the *HMS Porcupine* expedition (1869) in the Northeast Atlantic ([Carter, 1874](#)). The holotype description referred to a soft roundish sponge with a thin paper like ectosome with papillate projections that lodge pores and oscula. However, while the pore areas are indeed located at the wart-like papillae, the oscula are not at their tip ([Figs. 1C; 1F](#)), as initially claimed ([Carter, 1874](#); [Vosmaer, 1885](#)) but on the ectosome ([Lundbeck, 1905](#)), yet they are visible only after a careful examination. The conspicuous ectosome is loosely attached to the choanosome here and there, which, together with its fragility, might contribute to its rip off during trawl sampling ([Vosmaer, 1885](#); [Topsent, 1892](#)). Collected individuals without ectosome, appear smooth, porous, and lack the characteristic papillae. However, the presence of spherancorae facilitates the species identification, even after the ectosome's detachment ([Baker et al., 2018](#)).

While Carter's original description was precise, the illustrations were not sufficiently accurate. Thus, subsequent authors ([Vosmaer, 1885](#); [Topsent, 1892, 1904](#)) referred to Schmidt's redescription based on specimens from the Caribbean ([Schmidt, 1880](#)) rather than Carter's description of the type specimen for their species identification. However, Schmidt's material (MZS Po165) was in fact another species (described below as *Melonanchora insulsa* sp. nov.) clearly differing from *M. elliptica* in the shape of chelae and spherancorae. Finally, Topsent's individuals from the Azores are insufficiently described ([Table 1](#)) and were not available. While it is clear that they belong to *Melonanchora*, it is impossible to ascertain based on Topsent's descriptions if they unequivocally belong to *M. elliptica* or to any other North Atlantic *Melonanchora* species.

Melonanchora emphysema ([Schmidt, 1875](#))
([Figs. 1B; 5; 6](#))




Figure 4 Distribution map for north Atlantic *Melonanchora* species. Distribution map for the North Atlantic *Melonanchora* species: *Melonanchora elliptica* (green circle), *Melonanchora emphysema* (orange square), *Melonanchora tumultuosa* sp. nov. (red triangle); *Melonanchora maeli* sp. nov. (dark green square); *Melonanchora intermedia* sp. nov. (purple square); *Melonanchora insulsa* sp. nov. (dark blue square). Projected view (UTM Zone 31N (WGS84)) with geographic (WGS84) coordinates indicated for reference. The 1,000 m depth isobaths is represented by a grey line. Geographic and bathymetric data used was obtained from <http://www.natureallearthdata.com>.

Full-size DOI: 10.7717/peerj.12515/fig-4

Synonymy:

Desmacidon emphysema Schmidt, 1875: 118.

Melonanchora elliptica; Alander, 1935: 5 (*partim*).

Melonanchora emphysema; Vosmaer, 1885: 31, pl. I fig. 14, pl. V figs. 69–70 (*partim*); Thiele, 1903: 393; Lundbeck, 1905: 213–216, pl. XX fig. 2a–2d; Lundbeck, 1909: 402–403; Arndt, 1913: 116; Hentschel, 1929: 966–967; Arndt, 1935: 73, Fig. 142; Alander, 1942: 57 (*partim*); Vacelet, 1969: 200–201, fig. 38; Solórzano & Durán, 1982: 105–106, fig. 5c; Solórzano, 1990: 755–777, L. 92; Solórzano, 1991: 34; Ríos & Cristobo, 2017: 169; Santín *et al.*, 2021: Tab. 1.

Not *Melonanchora emphysema*; van Soest, 1993: 210, Tab. 2; Pulitzer-Finali, 1983: 561.

Material examined.

Holotype:

ZMB Por 2680, North Sea, from a Fjord of the southern coasts of Norway; ZMB Por 6571, North Sea, from a Fjord of the southern coasts of Norway.

Additional specimens examined:

GNM Porifera 416, Skagerrak, Sweden, 80–100 m depth, 1934, (Alander, 1935, 1942); GNM Porifera 290, Norra Kosterområdet Säcken, Baltic Sea (59.01441, 11.11977), 80 m depth, 1934, (Alander, 1935, 1942); GNM Porifera 390, Norra Kosterområdet Säcken, Baltic Sea (59.01441, 11.11977), 80 m depth, 1927, (Alander, 1935, 1942); MZB 2019–1740–Blanes Canyon, north-western Mediterranean Sea (41.50722, 2.93388),

Table 1 Comparative table between all known records of *Melonanchora elliptica* Carter, 1874, including the locality (Loc.) and depth of the sample, as well as the measurement of their spicular complement.

Author	Loc./Depth	Ectosomal megascleres	Choanosomal megascleres	Isochelae	Spherancorae
<i>Melonanchora elliptica</i> Carter, 1874					
Carter (1874)	Faroe Plateau*/ 'deep-sea'	(St) ca. 750 μm	(S) ca. 495 μm	Present	Present
Reexamination <i>van Soest</i> (2002)	Faroe Plateau*/ 'deep-sea'	(St) 450–650 \times 13–15 μm	(S) 650–860 μm	(I) 22–44 μm (II) 60 μm	48–68 μm
Reexamination <i>This study</i> (ZMB Por 3042)	Faroe Plateau */'deep-sea'	(St) 500–561.9 \pm 34.4–611.2 \times 14.7–15.9 \pm 1.1–19.6 μm	(S) 730–804.3 \pm 78.9–1176 \times 14.7–19.2 \pm 2.1–22.2 μm	(I) 22.8–25 \pm 1.5–27.6 μm (II) 48.9– 61 \pm 2.4–66.3 μm	58.8–62.4 \pm 2.2–68.3 \times 27.6–29.7 \pm 1.8–31.3 μm
<i>Vosmaer</i> (1885)	Barents Sea	Present	Present	Present	Present
Reexamination <i>This study</i> (ZMA.POR. P.10797)	Barents Sea	(St) 584–678 \pm 55.9–762 \times 13.8–16.8 \pm 1.7–18.6 μm	(S) 738–994.3 \pm 89.9–1146 \times 15–19.1 \pm 2.7–23.7 μm	(I) 24–27.8 \pm 1.5–31 μm (II) 63–71.8 \pm 2.3–81 μm	63–67.5 \pm 2.2–72 \times 26–28.9 \pm 1.7–30.5 μm
<i>Fristedt</i> (1887)	East Greenland/ 580 m	(St) 500 μm	<i>nm</i>	(I) 15 μm (II) 60 μm	70 μm
<i>Arnesen</i> (1903)	Between Bergen and Trondheim/ 100–180 m	<i>nm</i>	(S) ca. 1000 μm	(I) <i>nm</i> (II) 68 μm	60 μm
<i>Lundbeck</i> (1905)	North Atlantic/ 105–1,460 m	(St) 410–620 \times 8–17 μm	(S) 680–860 \times 14–21 μm	(I) 21–28 μm (II) 47–61 μm	54–68 \times 24–38 μm
<i>Arndt</i> (1935)	North Atlantic/ 'deep-sea'	(St) 410–620 μm	(S) 680–860 μm	(I) 21–28 μm (II) 47–75 μm	54–68 μm
<i>Koltun</i> (1959)	Barents Sea/ 106–385 m	(St) 410–620 \times 8–17 μm	(S) 680–904 \times 14–27 μm	<i>nm</i>	<i>nm</i>
<i>Baker et al.</i> (2018)	Davis Strait/ 537–1,132 m	(St) 528.1–594.7–655.5 \times 14.2–19.3–23.9 μm (St) 575.9–618.6–661.5 \times 18.3–21.6– 24.8 μm (St) 497.4–613.1–725.5 \times 15.7–19.5–22.2 μm (St) 504.4–568–629.1 \times 16–19.2–22.7 μm (St) 498.4–553–603 \times 15.7–18.6–22.3 μm	(S) 689.7–842.8–902.8 \times 11.1–15.1–21.1 μm (S) 730.2–778.4–822.4 \times 13.3–15.5–17.9 μm (S) 701.8–759.8–827.4 \times 12–14.5–19 μm (S) 743.5–814.3–879.1 \times 11.3–14.4–18.8 μm (S) 682.2–758.4–835.4 \times 13.5–17.4–20.5 μm	(I) 23.1–25.4–28.8 μm (II) 40.4–57.4–67.6 μm (I) 22.7–24.9–27 μm (II) 44.7–54.8–61.6 μm (I) 21.4–25.1–29.1 μm (II) 50.9–56.9–60.8 μm (I) 23.2–26–27.2 μm (II) 48.2–52.5–57.7 μm (I) 21.5–24.4–26.3 μm (II) 42.1–59–82.8 μm	48–57.2–65.7 \times 24–29.7–35.9 μm 54.1–62.8–68 \times 26.9–31–36.9 μm 51.2–57.9–63.4 \times 23.7–30.1–37.5 μm 46.3–55.8–61.7 \times 25.6–29–33.2 μm 41.5–49.5–57.5 \times 27.8–31.8–37.9 μm
<i>Dinn & Leys</i> (2018)	Saglek Bank, Northern Labrador Sea/427 m	(T) 554–623–693 \times 12.6–15.5–18.6 μm .	(S) 749–833–923 \times 18.5–23–26 μm	(I) 18–22–27.6 μm (II) 35–55–64 μm	43–50–53 μm
Reexamination <i>This study</i> (CMNI 2018- 0107)	Saglek Bank, Northern Labrador Sea/427 m	(St) 560.3–624.3 \pm 32.2–667.6 \times 7.8–11.8 \pm 3–17.3 μm .	(S) 782.5–830.7 \pm 50–908 \times 19.3–21.5 \pm 1.2–23.1 μm	(I) 24.1–24.9 \pm 1.2–29 μm (II) 48.3–51 \pm 3.8–59 μm	48.3–51.2 \pm 2.6–53.1 \times 26.5–29 \pm 0.7–29.8 μm
<i>This study</i> (NR0509_43)	Flemish Cap, Tail Grand Bank/ 1,554 m	(St) 533–645 \times 6–13 μm	(S) 619–803 \times 14–18 μm	(I) 21–26 μm (II) 46–66 μm	48–64 \times 20–33 μm

(Continued)

Table 1 (continued)

Author	Loc./Depth	Ectosomal megascleres	Choanosomal megascleres	Isochelae	Spherancorae
<i>This study</i> (NR0509_49)	Flemish Cap, Tail Grand Bank/ 1,137 m	(St) 488–610 × 8–17 μm	(S) 601–1000 × 15–27 μm	(I) 20–30 μm (II) 50–67 μm	52–61 × 19–28 μm
<i>This study</i> (NR0509_52)	Flemish Cap, Tail Grand Bank/ 1,122 m	(St) 504–598 × 12–16 μm	(S) 751–1086 × 16–24 μm	(I) 21–35 μm (II) 55–77 μm	55–66 × 26–39 μm
<i>This study</i> (NR0509_73)	Flemish Cap, Tail Grand Bank/870 m	(St) 555–625 × 11–17 μm	(S) 767–910 × 15–24 μm	(I) 25–29 μm (II) 39–70 μm	51–63 × 23–34 μm
<i>This study</i> (NR0509_82a)	Flemish Cap, Tail Grand Bank/ 1,127 m	(St) 538–676 × 12–20 μm	(S) 637–867 × 17–20 μm	(I) 22–28 μm (II) 51–71 μm	58–68 × 27–39 μm
<i>This study</i> (NR0620_21)	Flemish Cap, Tail Grand Bank/ 1,248 m	(St) 532–842 × 10–19 μm	(S) 722–902 × 10–22 μm	(I) 19–27 μm (II) 38–52 μm	46–59 × 25–35 μm
<i>This study</i> (NR0709_5)	Flemish Cap, Tail Grand Bank/ 1,055 m	(St) 518–845 × 11–20 μm	(S) 705–833 × 13–22 μm	(I) 23–33 μm (II) 37–63 μm	50–62 × 26–35 μm
<i>This study</i> (NHMUK Norman Coll. 1910.1.1.1418)	Norway/unknown	(St) 479.5–602.8 ± 24.1–673 x 14.3–16.4 ± 2.2–19.1 μm	(S) 765–863.8 ± 59.5–925.7 x 15.3–19.8 ± 1.5–21.7 μm	(I) 24.3–27.1 ± 2.4–33.3 μm (II) 61–72.6 ± 8–82 μm	67–75.6 ± 5.4–82.6 × 27.1–31.7 ± 4.3–35.4 μm
<i>This study</i> (NHMUK Norman Coll. 1910.1.1.1419)	Norway/unknown	(St) 548–570.3 ± 10.3–628 x 13.7–15.8 ± 1.8–18.7 μm	(S) 745.6–880.1 ± 34.9–936 x 14.9–18.5 ± 1.3–23.5 μm	(I) 26–27.2 ± 0.8–28.5 μm (II) 67.3–75.5 ± 1.4–78 μm	67–75.2 ± 6.5– 83 × 23.7–33.1 ± 6.5–36 μm
<i>Melonanchora cf. elliptica</i> Carter, 1874					
<i>Topsent</i> (1892)	Azores/736–1,267 m	(St) Present	(S) Present	(I) <i>nm</i> (II) 55 μm	70 μm
<i>Topsent</i> (1904)	Azores/523–1,360 m	<i>nm</i>	<i>nm</i>	(I) 18–21 μm (II) <i>nm</i>	<i>nm</i>
<i>Topsent</i> (1913)	Norwegian coast/ 440 m	<i>nm</i>	<i>nm</i>	<i>nm</i>	<i>nm</i>
<i>Topsent</i> (1928)	Azores/650–950 m	<i>nm</i>	<i>nm</i>	(I) 19–23 μm (II) 40–41 μm	43 × 26 μm
	Azores/1,378 m	<i>nm</i>	<i>nm</i>	(I) 20–23 μm (II) 72 μm	72 × 35 μm

Notes:

(S) indicates styles; (St) indicates strongyles; (T); indicates tylostyles.

* indicates this is the holotype of the species; *nm* indicates a spicular type that was not mentioned on a description, yet it is assumed was present on the samples.

‘ABIDES’ survey, 684 m depth, 2018 (*Santín et al., 2021*); ZMA.POR.P.10800 Outer Hebrides, Scotland, North-East Atlantic (56.80588, -7.42903), 2006; ZMA.POR.20192 Outer Hebrides, Scotland, North-East Atlantic (56.80588, -7.42903), 2006; ZMA.POR.P.10799 West of Hvasser, Norway, Baltic Sea (59° 04′ 42.06″N 10° 43′ 55.379″E), 2006; ZMA.POR.20559.b West of Hvasser, Norway, Baltic Sea (59.07835, -10.73204), 2006; ZMA.POR.20473.b West of Hvasser, Norway, Baltic Sea (59.07835, -10.73204), 2006; ZMA.POR.20551 West of Hvasser, Norway, Baltic Sea (59.07835, -10.73204), 2006; ZMA.

POR.P.10798 Outer Hebrides, Scotland, North-East Atlantic (56.8071, -7.43025), 2006; ZMA.POR.20353.a Outer Hebrides, Scotland, North-East Atlantic (56.8071, -7.43025), 2006; ZMA.POR.P.10795 West of Ireland, North-East Atlantic (55.50093, -15.78839), attached to *Madrepora* debris, 2005; ZMA.POR.P.20020 West of Ireland, North-East Atlantic (55.50093, -15.78839), attached to *Madrepora* debris, 2005; ZMA.POR.20020 West of Ireland, North-East Atlantic (55.50093, -15.78839), attached to *Madrepora* debris, 2005; ZMA.POR.P.10829 West of Hvasser, Norway, Baltic Sea (59.07577, 10.73552), 2007; ZMA.POR.20467 West of Hvasser, Norway, Baltic Sea (59.07577, 10.73552), 2007; ZMA.POR.P.10828 Outer Hebrides, Scotland, North-East Atlantic (56.8059, -7.44183), 2006; ZMA.POR.20175.b Outer Hebrides, Scotland, North-East Atlantic (56.8059, -7.44183), 2006; ZMA.POR.P.10827 Outer Hebrides, Scotland, North-East Atlantic (56.80563, -7.426029), 2006; ZMA.POR.20335 Outer Hebrides, Scotland, North-East Atlantic (56.80563, -7.426029), 2006.

Unregistered material:

AVILES_0710-48DR5, Avilés Canyon System, Cantabrian Sea (43.80333, -6.15583), 128 m depth (INTEMARES AVILES Coll.); MS, off Bares (44.055, -7.64638), Spanish coasts, 500 m depth; JV, Cassidaigne Canyon (42.95, 5.38333), 360 m depth ([Vacelet, 1969](#)); Galician Bank, west of Galician coast, Spain (42.58305, -11.58305) ca. 700 m depth; Baixo do Placer do Cabezo de Laxe (43, -9.03333), Galicia Coast, Spain, Fishermen's by-catch, 58 m depth, 1981 ([Duran & Solórzano, 1982](#); [Solórzano, 1990, 1991](#)).

Description:

Mostly encrusting, rarely massive-encrusting (GNM Porifera 416), with an easily detachable paper-like ectosome bearing fistular processes. Fistulae might be absent in small encrusting individuals. Colour whitish translucent in the ectosome, cream-orange in the choanosome while in alcohol.

Skeleton:

Ectosomal skeleton formed by intertwined tangential tylostrongyles. The choanosomal skeleton is ill defined, with scattered tracts of tylostrongyles identical to those conforming the ectosome. Microscleres mostly scattered thorough the choanosome without any clear discernible pattern.

Spicule complement:

Tylostrongyles, two categories of chelae, and spherancorae ([Figs. 5A-5E](#) and [Figs. 6A-6F](#)).

Ectosomal and choanosomal tylostrongyles ([Figs. 5A; 6A](#)): of similar shape to those of *M. elliptica*: they are unevenly and slightly flexuous, enlarged at the central zone and narrowing toward unequal tyloid ([Fig. 6F](#)), giving them the appearance from strongyles to tylostrongyles.

Size range: $492.7-508.1 \pm 13-521.6 \mu\text{m} \times 9.7-10.6 \pm 2.8-14.5 \mu\text{m}$

Isochelae I ([Figs. 5D, c'; 6E, c'](#)): Small isochelae with a straight shaft, gently bending to its ends, with three spatulated alae and well-formed fimbriae.




Figure 5 *Melonanchora emphysema* spicule plate. Spicular set for *Melonanchora emphysema* (sample ZMB Por 2680, holotype). (A) Ectosomal and chonasomoal tylostrongyle; (B) Spherancorae; (C) Large chelae category (Chelae II); (D) small chelae category (Chelae I); (E) General view of *M. emphysema*'s spicules by SEM imaging. (a') Spherancora (b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (a'), (b'), (c') 200 µm; (B), (C), (D) 30 µm and (E) 500 µm. [Full-size !\[\]\(5f471a71b78d7676bc356df190b88ab4_img.jpg\) DOI: 10.7717/peerj.12515/fig-5](https://doi.org/10.7717/peerj.12515/fig-5)

Size range: $24.1\text{--}26.6 \pm 2.8\text{--}28.9$ µm

Isochelae II (Figs. 5C, b'; 6C, b'): very similar to isochelae I, but bigger in size.

Size range: $48.3\text{--}51.5 \pm 5.5\text{--}58$ µm




Figure 6 *Melonanchora cf. emphysema* spicule plate. Spicular set for *Melonanchora cf. emphysema* from Laxe, Galicia coast, Spain (unregistered sample). (A) Ectosomal and chonasomoal tylostrongyle; (B) Spherancorae; (C) Large chelae category (Chelae II); (D) Chelae II with reduced alae; (E) small chelae category (Chelae I); (F) Detail of the tyles. (a') Spherancora (b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (a'), (b'), (c') 175 µm; (B), (C), (D) 20 µm and (F) 12 µm. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac_img.jpg\) DOI: 10.7717/peerj.12515/fig-6](https://doi.org/10.7717/peerj.12515/fig-6)

Spherancorae (Figs. 5B, a'; 6B, a'): Elongated-ovoid (Fig. 5B) to stadium shaped (Fig. 6B) with teeth-like fimbriae on its internal surface, which may be fused at various degrees. Size range: $37.6\text{--}38.8 \pm 1.1\text{--}40.5 \times 25.1\text{--}27.6 \pm 1.6\text{--}28.9$ µm

Geographic distribution:

Originally described from the coasts of Norway ([Schmidt, 1875](#)), the species is known from deep Atlantic and Arctic waters ([Fig. 4](#)), including Greenland and Iceland, ([Lundbeck, 1905, 1909, 1910](#)), Faroe Islands ([Hentschel, 1929](#)), Porcupine Bank ([van Soest & De Voogd, 2015](#)), Baltic Sea ([Alander, 1935, 1942](#)), the Spanish coasts ([Solórzano, 1990; Ríos & Cristobo, 2017](#); this paper), and the coasts of Norway ([Vosmaer, 1885; Arndt, 1913](#)) including the Svalbard archipelago ([Gulliksen et al., 1999](#)). The species had also been tentatively recorded from the Atlantic Canadian coast ([Baker et al., 2018; Murillo et al., 2018](#)), yet these records correspond to *Melonanchora tumultuosa* sp. nov., thus its presence in the west Atlantic area remaining unconfirmed. Additionally, the species has also been sparsely recorded from the Mediterranean Sea and nearby areas: the Gulf of Lyon ([Vacelet, 1969; Santín et al., 2021](#)) and the northern coasts of Spain ([Solórzano & Durán, 1982; Solórzano, 1990, 1991; Ríos & Cristobo, 2017](#); this study). The species appears to be a frequent inhabitant of cold-water corals communities ([Könnecker & Freiwald, 2005; van Soest & De Voogd, 2015](#)), yet it might also occur attached to rocky substrata or debris.

Remarks:

[Schmidt \(1875\)](#) poorly described *Desmacidon emphysema* from the coast of Norway, a species characterized by the presence of a papillate ectosome and smooth megascleres enlarged at the middle, with unequally swelled ends. While Schmidt accurately reported spherancorae in his *M. emphysema* samples from the Caribbean ([Schmidt, 1880](#)), he missed these spicules in the Northern Sea samples, mistaking them with diatoms ([Schmidt, 1875](#)), which led to his misclassification of *M. emphysema* in the genus *Desmacidon*, until amended by [Thiele \(1903\)](#). Furthermore, Schmidt's incomplete description ([Table 2](#)) led several authors to consider the species a synonym of *M. elliptica* ([Vosmaer, 1885; Arnesen, 1903](#)) while others claimed that a clear distinction existed ([Thiele, 1903; Lundbeck, 1905](#)). The problem mainly arose as the main distinguishing feature between both species relies on its choanosomal megascleres, with *M. elliptica* possessing styles and *M. emphysema* possessing strongyles ([Lundbeck, 1905](#)), yet several authors had described samples with blunt-ended styles as choanosomal megascleres ([Vosmaer, 1885; Baker et al., 2018](#)).

The re-examination of Schmidt holotype (ZMB Por 2680) however leaves no doubt about the validity of the species. As previously pointed out ([Thiele, 1903; Lundbeck, 1905](#)), *M. emphysema*'s choanosomal megascleres are exclusively tylostrongyles identical to its ectosomal ones, while its spherancorae are smaller or equal in size to the large isochelae ([Table 2](#)). Conversely, in *M. elliptica* there is a clear distinction between the choanosomal (styles) and ectosomal (tylostrongyles) megascleres and, additionally, the spherancorae are within the size range of the large isochelae ([Table 1](#)). Thus, individuals identified as *M. emphysema* with blunt-ended diactines in two clear categories do not correspond to this species but to a new one, *Melonanchora tumultuosa* sp. nov. (described below). Finally, in the Mediterranean and nearby areas, *M. emphysema* tylostrongyles are almost half in size than those in the North Atlantic specimens (average length ca. 400 vs. 600 µm; [Table 2](#)), and it has been suggested that they might correspond to a yet

Table 2 Comparative table between all known records of *Melonanchora emphysema* (Schmidt, 1875), including the locality (Loc.) and depth of the sample, as well as the measurement of their spicular complement.

Author	Loc./Depth	Ectosomal megascleres	Choanosomal megascleres	Isochelae	Spherancorae
<i>Melonanchora emphysema</i> (Schmidt, 1875)					
Schmidt (1875)	Haugesund, Norway*/193 m	(St) Present	nm	Present	nm
Reexamination <i>This study</i> (ZMB Por 2680)	Haugesund, Norway*/193 m	(St) 500–570 ± 15.9–627 × 10.9–15.8 ± 3.1–18.5 µm	Same as in ectosome	(I) 19.6–24.7 ± 2.7–29.4 µm (II) 55.3–60.2 ± 3.9–68.6 µm	40.4–44.3 ± 1.8–58 × 23.1–25.6 ± 1.3–28 µm
Thiele (1903)	North Atlantic	(St) ca. 650 µm	Same as in ectosome	(I) 21 µm (II) 60 µm	50 µm
Lundbeck (1905)	North Atlantic/375–1,460 m	(St) 440–610 × 10–14 µm	Same as in ectosome	(I) 24–30 µm (II) 57–71 µm	50–56 × 28 µm
Alander (1942)	Skandia, Sweden/85 m	Present	Present	Present	Present
Reexamination <i>This study</i> (GNM Porifera 390)	Skandia, Sweden/85 m	(St) 492.7–508.1 ± 13–521.7 × 9.7–10.6 ± 2.8–14.5 µm	Same as in ectosome	(I) 24.2–26.6 ± 2.7–29 µm (II) 48.3–51.5 ± 5.5–58 µm	37.6–38.9 ± 1–42.6 × 21.6–24.3 ± 1.6–29 µm
Vacelet (1969)	Mediterranean/360–370 m	(St) 330–490 × 8.5–18 µm	Same as in ectosome	(I) 22 µm (II) 40–53 µm	40–45 × 20 µm
Reexamination <i>This study</i> (unregistered)	Mediterranean/360–370 m	(T) 389.3–418.6 ± 11.7–477 × 12.2–14.6 ± 1.3–17.6 µm	Same as in ectosome	(I) 21.4–22.9 ± 0.9–25.3 µm (II) 41.2–45 ± 1.2–55.1 µm	38.4–41.3 ± 1.5–44.5 × 17.1–19.7 ± 2.3–22.7 µm
<i>This study</i> (ZMA.POR.P.10800)	Scotland/-	(St) 342–472.8 ± 61.8–540 × 5.4–6.9 ± 0.8–7.8 µm	Same as in ectosome	(I) 22.8–24.3 ± 1–25.8 µm (II) 48–52.5 ± 5.6–63 µm	37.8–41.7 ± 2.8–44.4 × 18–19.5 ± 1.3–21 µm
<i>This study</i> (unregistered)	Galicia Bank/500 m	(T) 439.2–479.9 ± 30.4–537.6 × 12.2–15.5 ± 1.8–18.7 µm	Same as in ectosome	(I) 20.7–23.4 ± 1.5–25.4 µm (II) 42–51.2 ± 4.3–57.2 µm	37.2–41.2 ± 2–44.6 × 17.3–20.6 ± 1.2–23.4 µm
<i>This study</i> (unregistered)	Galicia Bank/500 m	(T) 429.2–482.2 ± 29.7–538.9 × 11.8–15 ± 1.7–18.7 µm	Same as in ectosome	(I) 20.2–22.8 ± 1.9–27.3 µm (II) 40.6–54 ± 4.8–62.7 µm	34.7–41.2 ± 4–54.5 × 17.2–20.2 ± 2–23.5 µm
<i>This study</i> (MZB 2019–1740)	Gulf of Lyon/684 m	(T) 253.6–375.6 ± 48.7–426.1 µm × 8.8–10.1 ± 1.7–13.7 µm	Same as in ectosome	(I) 20.5–24.1 ± 3.7–30.4 µm (II) 44.3–53 ± 4.2–60 µm	41.2–43.7 ± 2.1–46.6 × 18.3–20.5 ± 2.7–26.3 µm
<i>Melonanchora cf. emphysema</i> (Schmidt, 1875)					
Solórzano & Duran (1981)	Galicia Coast, Spain*/58 m	(St) 316–345 × 9 µm	Same as in ectosome	(I) 22–26 µm (II) 44–51 µm	27–40 µm
Reexamination <i>Solórzano</i> (1990)	Galicia Coast, Spain*/58 m	(St) 316–345 × 8–9 µm	Same as in ectosome	(I) 22–26 µm (II) 44–51 µm	27–40 × 18–20 µm

(Continued)

Table 2 (continued)

Author	Loc./Depth	Ectosomal megascleres	Choanosomal megascleres	Isochelae	Spherancorae
Reexamination <i>This study</i> (unregistered)	Galicia Coast, Spain*/58 m	(T) 302.6–345.8 ± 24–384.5 × 4.9–6.83 ± 0.8–8 µm	Same as in ectosome	(I) 16.5–20 ± 1.4–22.2 µm (II) 35–44 ± 3.9–50 µm	31.9–36.2 ± 2.3–40.5 × 14.2–17.2 ± 2.1–20.5 µm
<i>This study</i> (AVILES_0710–48DR5)	Cantabrian Sea/ 128 m	(T) 274–329.6 ± 30.6–387.6 × 4.6–6.1 ± 0.8–7.6 µm	Same as in ectosome	(I) 15.4–18 ± 1.3–20.7 µm (II) 33.6–44 ± 3.8–48.9 µm	34.7–37.2 ± 1.2–39.3 × 12.6–16 ± 2–19.9 µm

Notes:
 (S) indicates styles; (St) indicates strongyles; (T); indicates tylostyles.
 * indicates this is the holotype of the species; *nm* indicates a spicular type that was not mentioned on a description, yet it is assumed was present on the sample/s.

undescribed species (*Vacelet, 1969*). In this sense, reexamination of all known Mediterranean material did in fact reveal a new species, *Melonanchora intermedia* sp. nov. (described below), occurring within western Mediterranean mesohypoxic environments (*Pulitzer-Finali, 1983; Díaz, Ramírez-Amaro & Ordines, 2021*). However, no major differences could be observed with *M. emphysema* specimens from other deep-sea Mediterranean and nearby areas other than the aforementioned size of their tylostongyles (Table 2). Additionally, the Mediterranean and Iberian specimens' spherancorae (Fig. 6B) closely match a stadium-shaped appearance, which is characteristic of *M. emphysema*. However, it must be noted that one specimen from the Galician coast and another one from the Cantabrian Sea possess relatively smaller and thinner tylostongyles (ca. 330 µm length vs. ca. 6 µm width) when compared with all other *M. emphysema* records (Table 2), and, in the Galician sample, an additional category of chelae with reduced alae could be observed in very low numbers (Fig. 6D). Nevertheless, said chelae are absent from all other Iberian or Mediterranean *M. emphysema* material. Given the high variability in megasclere size observed within all *Melonanchora* species (Tables 1–3), as well as the poor conservation status of these deviant samples, it would be unwise to erect a new species based solely on the megascleres size. Yet, the possibility that those specimens correspond in fact to a cryptic species cannot be entirely ruled out, and its identity should be further clarified if more individuals with said characteristics were to be discovered.

Melonanchora tumultuosa sp. nov.
(Figs. 1C; 7)

Synonymy:

Melonanchora elliptica; *Vosmaer, 1885*: 31, pl. I fig. 14, pl. V figs. 69–70 (*partim*); *Lundbeck, 1905*: 213–216, pl. VII figs. 4–6, pl. XX figs. 1a–1o (*partim*); *Lundbeck, 1909*: 402–403 (*partim*); *Alander, 1935*: 5 (*partim*).
Melonanchora emphysema; *Alander, 1942*: 57 (*partim*); *Baker et al., 2018*: 26–30, figs. 8–10.
 Not *Melonanchora elliptica* *Carter, 1874*: 212.

Material examined.

Table 3 Comparative table for all new species of *Melonanchora*, as well as the closely related genus *Hanstoreia* gen. nov., described on this work, including: the locality (Loc.) and depth of the sample, as well as the measurement of their spicular complement.

Author	Loc./Depth	Ectosomal megascleres	Choanosomal megascleres	Isochelae	Spherancorae
<i>Melonanchora tumultuosa</i> sp. nov.					
<i>Vosmaer (1885)</i>	–	Present	Present	Present	Present
Reexamination <i>This study</i> (ZMA.POR. P.10796)	Norway/256 m	(St) 483–542.6 ± 38.3–600 µm × 10.6–12.9 ± 3.2–19.3 µm	(St) 627.9–802.3 ± 42.2–924.5 µm × 11.6–18.3 ± 1.5–24.4 µm	(I) 21.2–26.5 ± 3.8–28.9 µm (II) 48.6–68.6 ± 8.1–72.9 µm	48.3–67.5 ± 6.8–78.62 × 18.9–22.3 ± 1.6–25.2 µm
<i>Baker et al. (2018)</i>	Davis Strait/ 537–1132 m	(St) 485.1–599.8–673.3 × 12.7–15.6–20 µm (St) 537.5–582.6–670.8 × 12.0–14.4–17.4 µm (St) 509.9–569.8–611.6 × 11.3–14.7–17.9 µm	(St) 831.1–913.6–981.6 × 15.7–19.5–22.7 µm (St) 823.5–884.6–957.8 × 13.5–19.2–24 µm (St) 672.6–770.9–860.1 × 17.4–20–23.9 µm	(I) 22.6–25.8–32.2 µm (II) 43.3–59–66.4 µm (I) 22.2–24.3–27.1 µm (II) 44–49.5–56.8 µm (I) 20.5–22.7–25.4 µm (II) 49.5–52.3–56.3 µm	53.2–57.5–63.7 × 23.1–27.7–35.3 µm 52.8–54.9–59.3 × 24.9–30.4–36.0 µm 57.5–61.7–65.1 × 23.9–26.9–28.8 µm
<i>This study</i> (NR0509_82b)	Flemish Cap, Tail Grand Bank/ 1,027 m	(St) 548–657 × 11–17 µm	(St) 716–873 × 14–22 µm	(I) 22–26 µm (II) 49–68 µm	56–67 × 25–38 µm
<i>This study</i> (NR0610_30)	Flemish Cap, Tail Grand Bank/613 m	(St) 544–657 × 8–18 µm	(St) 483–823 × 8–13 µm	(I) 24–32 µm (II) 38–67 µm	47–65 × 22–34 µm
<i>This study</i> (GNM Porifera 624)	Sydkoster Island, Sweeden*/100 m.	(St) 483–542.6 ± 38.3–600 × 10.6–12.9 ± 3.2–19.3 µm	(St) 627.9–802.3 ± 42.2–924.5 × 11.6–18.3 ± 1.5–24.4 µm	(I) 21.2–26.5 ± 3.8–28.9 µm (II) 48.6–68.6 ± 8.1–72.9 µm	48.3–67.5 ± 6.8–78.6 × 18.9–22.3 ± 1.6–25.2 µm
<i>This study</i> (NHMUK, 83.12.13.70.89)	Unknown	(St) 483–542.6 ± 38.3–600 × 10.6–12.9 ± 3.2–19.3 µm	(St) 768–895.7 ± 38.3–993 × 15.7–19.8 ± 1.6–24 µm	(I) 18.5–21 ± 2.6–25 µm (II) 55.7–76.1 ± 2.9–79 µm	62.8–70 ± 4.9–78 × 22.1–24.5 ± 1.9–29.3 µm
<i>This study</i> (NHMUK Norman Coll. 1898.5.7.38)	Norway	(St) 490–550.4 ± 38.9–607.6 × 10.8–13.1 ± 3.3–19.6 µm	(St) 637–712.7 ± 31.3–813.5 × 11.8–14.7 ± 1.5–21.1 µm	(I) 21.3–26.5 ± 2.5–29 µm (II) 40.2–57.7 ± 8.2–69.6 µm	48.3–60 ± 4.2–67.6 × 25.1–27 ± 1.5–29 µm
<i>This study</i> (ZMA.POR. P.10825)	Norway/130–150 m	(St) 528–617 ± 52.2–667 × 12.8–15 ± 2–18 µm	(St) 642–696 ± 58.8–804.3 × 14.7–18.6 ± 2.7–21.9 µm	(I) 24–28.9 ± 4.4–32 µm (II) 54–72.3 ± 8.7–81 µm	56.6–64.3 ± 6.4–72.3 × 18–23.8 ± 2.8–27.4 µm
<i>This study</i> (ZMA.POR. P.10822)	Norway/130–150 m	(St) 402–499.5 ± 60.5–540 × 12–13.7 ± 1.8–16.1 µm	(St) 645–756 ± 88–1026 × 12.5–19.3 ± 1.9–21 µm	(I) 23–27.6 ± 4.1–30 µm (II) 51–70.1 ± 9.2–78 µm	52.2–58.8 ± 7.9–74 × 23.4–25.9 ± 2.8–30 µm
<i>This study</i> (ZMA.POR.4977)	Norway/130–150 m	(St) 462–515.5 ± 54.8–582 × 11.9–14.2 ± 1.6–16.5 µm	(St) 601.3–719.5 ± 79.3–1002 × 13.3–18.2 ± 2.7–22.7 µm	(I) 24–29 ± 2.6–33 µm (II) 60–71.5 ± 7.1–84 µm	48–55.6 ± 6.2–72 × 24–25.9 ± 2.4–30 µm
<i>Melonanchora intermedia</i> sp. nov.					
<i>Pulitzer-Finali (1983)</i>	Corsica, Mediterranean Sea*/128 m	(St) 380–490 × 6–11 µm	Same as in ectosome	(I) 19–21 µm (II) 32–49 µm	37–43 µm

(Continued)

Table 3 (continued)

Author	Loc./Depth	Ectosomal megascleres	Choanosomal megascleres	Isochelae	Spherancorae
Reexamination <i>This study</i> (MSNG R.N. N IS.4.7)	Corsica, Mediterranean Sea*/128 m	(St) 369–411.8 ± 14.5–475.3 × 7.2–9.7 ± 1.5–11 µm	Same as in ectosome	(I) 19–21.5 ± 0.7–22.7 µm (II) 30.1–35.2 ± 2.9–38.6 µm (III) 33.2–39.5 ± 5.1–47.8 µm	38.9–44.4 ± 6.7–51.2 × 20–21.8 ± 1.9–24.2 µm
<i>Díaz, Ramírez- Amaro & Ordines</i> (2021)	Mallorca Channel, Mediterranean Sea/ 104–138 m	(T) 359–446–556 × 5– 8 –11 µm	Same as in ectosome	(I) 14–18–21 µm (II) <i>nm</i> (III) 29–42 –47 µm	36–40–46 × 14–19–23 µm
<i>Melonanchora insulsa</i> sp. nov.					
<i>Schmidt</i> (1880)	Gulf of Mexico*/ 'deep-sea'	-	-	(I) 23 µm (II) 68 µm	60 µm
Reexamination <i>This study</i> (MZS Po165)	Gulf of Mexico*/ 'deep-sea'	(St) 593.6–656.7 ± 36.2–701 × 16.1–17.1 ± 1.2–19.5 µm	(S) 813.4–989 ± 41.2–1121.7 × 19.3–20.7 ± 1.4–22.5 µm	(I) 27.2–30.9 ± 3.4–35.8 µm (II) 48.6–52.3 ± 5.1–68 µm	52.9–56.5 ± 4.2–62.1 × 22–24.3 ± 1.7–26.6 µm
<i>Melonanchora maeli</i> sp. nov.					
<i>This study</i> (ZMA.POR.7269)	Cape Verde*/'deep- sea'	(T) 531.6–590.9 ± 37.9–627.9 × 9.7–10.3 ± 0.5–10.6 µm	(S) 637.6–918.5 ± 75.6–1062.6 × 17.3–19.2 ± 1.3–21.3 µm	(I) 17.4–19.8 ± 1.7–23.2 µm (II) 27–29.3 ± 1.2–31.9 µm (III) 45.4–49.6 ± 2–53.1 µm	48.3–50.2 ± 1.7–53.2 × 17.4–19.2 ± 1.5–21.3 µm
<i>Hanstoreia globogilva</i> (Lehnert, Stone & Heimler, 2006a)					
Lehnert et al. (2006a)	Aleutian Islands*/ 190 m	(T) 640–680 × 10–12 µm	(Ac) 660–670 × 20–30 µm	(I) 23–25 µm (II) <i>nm</i>	(I) 65–93 µm (II) 65–93 µm
Reexamination <i>This study</i> (NMNH-USNM 1082996)	Aleutian Islands*/ 190 m	(T) 598.9–675 ± 22.5–724.5 × 9.7–10.9 ± 2.2–14.5 µm	(Ac) 589.3–638.3 ± 30–677.3 × 27–28 ± 1.1–29 µm	(I) 23.1–25.2 ± 1.1–27 µm (II) 48–64.4 ± 6.8–67.6 µm	(I) 77.3–86.9 ± 2.8–91.8 × 27–30 ± 2.3–33.8 µm (II) <i>nm</i>

Notes:

(S) indicates styles; (St) indicates strongyles; (T); indicates tylostyles.

* indicates this is the holotype of the species; *nm* indicates a spicular type that was not mentioned on a description, yet it is assumed was present on the samples.

Holotype (here designated): GNM Porifera 624, Kostergrundet, Sydkoster Island, Sweden, 100 m depth.

Additional specimens examined:

NHMUK–Icelandic Coll. 1958.1.1.633, Iceland, North Atlantic Ocean (63.55, –11.41666), 1936; NHMUK Norman Coll. 1898.5.7.38, Norway, 1893; NHMUK, 83.12.13.70.89; MZLU L935/3858, Koster, Säcken, Sweden, Baltic Sea (59.00971, 11.11471), 1934, (*Alander, 1935; 1942*); ZMA.POR.P.10796, Northwest of Tromsø, Norway, Arctic Ocean (72.60138, 24.95), *R/V Willem Barents* expedition (1880–84), 256 m depth, 1881 (*Vosmaer, 1885*); ZMA.POR.P.10825, Marsteinsboen, Norway, North East Atlantic (60.12583, 4.98944), 130–150 m depth, on stone, 1982; ZMA.POR.P.10822, Marsteinsboen, Norway, North East Atlantic (60.12583, 4.98944), 130–150 m depth, on stone, 1982; ZMA.POR.P.10824, Marsteinsboen, Norway, North East Atlantic (60.12583, 4.98944), 130–150 m depth, on stone, 1982; ZMA.POR.4977, Marsteinsboen, Norway, North East Atlantic

(60.12583, 4.98944, 130–150 m depth, on stone, 1982; ZMA.POR.P.10823, off Saengsbokt, Bergen, Norway, North East Atlantic (60.36666, 4.81666), 350–600 m depth, 1982; ZMA.POR.4976, off Saengsbokt, Bergen, Norway, North East Atlantic (60.36666, 4.81666), 350–600 m depth, 1982.

Unregistered material:

NR0509_82b, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 1,127 m depth (NEREIDA Coll.); NR0610_30a, Flemish Cap, Tail Grand Bank, North Atlantic Ocean, 613 m depth (NEREIDA Coll.).

Description:

Massive-globular sponge, with an easily detachable paper-like thin ectosome bearing abundant fistular processes (typical of the genus). The choanosome is orange-cream in colour and the ectosome results whitish, yet translucent, in alcohol.

Skeleton:

Spicule arrangement as in the other species of the genus (*viz. M. elliptica*), with its main distinguishing feature being the presence of strongyles as choanosomal megascleres.

Spicule complement:

Tylostrongyles, strongyles, two categories of isochelae, and spherancorae (Figs. 7A–7F)

Ectosomal tylostrongyles (Fig. 7B): As in other *Melonanchora*, they are slightly flexuous, with a more or less central swelling. The tips can be strongyloid or slightly tylote often vaguely unequal.

Size range: $483\text{--}542.6 \pm 38.3\text{--}600 \mu\text{m} \times 10.6\text{--}12.9 \pm 3.2\text{--}19.3 \mu\text{m}$

Choanosomal strongyles (Fig. 7A): Entirely smooth, with asymmetrical ends (one clearly rounded and the other blunt but somewhat narrower. More or less curved throughout its entire length.

Size range: $627.9\text{--}802.3 \pm 42.2\text{--}924.5 \mu\text{m} \times 11.6\text{--}18.3 \pm 1.5\text{--}24.4 \mu\text{m}$

Isochelae I (Fig. 7E, c'): Anchorate, with a straight shaft, gently bending to its ends, with three-spatulated alae.

Size range: $21.2\text{--}26.5 \pm 3.8\text{--}28.9 \mu\text{m}$

Isochelae I (Fig. 7D, b'): Similar to isochelae I, but smaller in size.

Size range: $48.6\text{--}68.6 \pm 8.1\text{--}72.9 \mu\text{m}$

Spherancorae (Fig. 7C, a'): With a prolate-oval shape, and dentate fimbriae on its internal face, which might be free or fused at various degrees. The junction points of each couple of opposite alae can be observed in most spicules, with the resulting fused shaft being slightly asymmetrical.

Size range: $48.3\text{--}67.5 \pm 6.8\text{--}78.62 \times 18.9\text{--}22.3 \pm 1.6\text{--}25.2 \mu\text{m}$




Figure 7 *Melonanchora tumultuosa* spicule plate. Spicular set for *Melonanchora tumultuosa* sp. nov. (sample GNM Por 624, holotype). (A) Choanosomal strongyle; (B) Ectosomal tylostrongyle; (C) Spherancorae; (D) Large chelae category (Chelae II); (E) small chelae category (Chelae I), (F) General view of *M. tumultuosa* sp. nov. spicules by SEM imaging. (a') Spherancora (b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b'), (c') 300 μm ; (C), (D), (E) 30 μm and (F) 500 μm . [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.12515/fig-7](https://doi.org/10.7717/peerj.12515/fig-7)

Geographic distribution and type locality:

The species presents an amphi-Atlantic distribution (Fig. 4), being sympatric with *M. elliptica*. Its type locality is the Sydkoster Island, Sweden, yet, known records for the species also include Iceland (NHMUK–1958.1.1.633) the Davis Strait (Baker et al., 2018) and Norwegian coasts (Vosmaer, 1885; this paper).

Etymology:

From the latin *tumultuosus*, meaning full of commotion. It refers to the confusion that samples of this species have caused between *M. elliptica* and *M. emphysema* during the past century.

Remarks:

Specimens of *M. tumultuosa* sp. nov. had been considered by several authors to be *M. emphysema* because of their possession of both ectosomal and choanosomal strongyles (Baker et al., 2018). Close re-examination of the *M. emphysema* type revealed only one type of megascleres, which is present in both ectosome and choanosome (Fig. 5A), whereas in *M. tumultuosa* sp. nov., two different types of strongyles characterise either the ectosome (Fig. 7B) or the choanosome (Fig. 7A).

Additionally, it had been suggested that those *Melonanchora* with two strongyle categories could in fact be *M. elliptica* individuals with styles modified into strongyles (Baker et al., 2018). In this regard, sponge spicules might vary in shape due to environmental conditions (Bell, Barnes & Turner, 2002) and/or silica abundance (Uriz et al., 2003) even to the point not expressing one or more spicule types (Maldonado & Uriz, 1996; Maldonado et al., 1999). However, *M. elliptica* and *M. tumultuosa* sp. nov. co-occur in their areas of distribution, even at local scales (Baker et al., 2018), weakening such an idea. Finally, *M. tumultuosa* sp. nov., spherancorae shape is mostly prolate (Fig. 7C), commonly with asymmetrical shafts and rounded ends, whereas they are clearly spheroidal in *M. elliptica*, with slightly pointed ends (Fig. 3C), which is translated in an overall slender spherancorae for *M. tumultuosa* sp. nov, compared to *M. elliptica* (average width 29.7 vs. 22.3 µm respectively; Tables 1 & 3).

Melonanchora intermedia sp. nov.

(Fig. 8)

Synonymy:

Melonanchora emphysema; Pulitzer-Finali, 1983: 561; Díaz, Ramírez-Amaro & Ordines, 2021: 42–43, fig. 16.

Not *Melonanchora emphysema* (Schmidt, 1875: 118).

Material examined.

Holotype (here designated): MSNG Vis4.7–off Calvi, Corsica (42.53333, 8.6), depth 128 m, detrital, dredge, 18 July 1975. R.N. N IS.4.7 (Pulitzer-Finali, 1983).




Figure 8 *Melonanchora intermedia* spicule plate. Spicular set for *Melonanchora intermedia* sp. nov. (sample MSNG Vis4.7, holotype). (A) Ectosomal and chonasomoal tylostrongyle; (B) Spherancorae; (C) small chelae category (Chelae I); (D) Large chelae category (Chelae II); (E) Anisochelae; (F) Detail of the tylostrongyle's ends. (a') Spherancora (b') Chelae II (c') Anisochelae and (d') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (a'), (b'), (c'), (d') 200 µm; (B), (C), (D), (E), (F) 20 µm. [Full-size !\[\]\(5f471a71b78d7676bc356df190b88ab4_img.jpg\) DOI: 10.7717/peerj.12515/fig-8](https://doi.org/10.7717/peerj.12515/fig-8)

Description:

Small subglobular individual attached to rocky debris. It possesses a paper-like ectosome with the warty-like papillae typical of the genus, yet with just a few papillae.

Skeleton:

Ill-defined paucispiculate tracts in the choanosomal area, and a clear crisscross pattern can be observed in the ectosome. Microscleres are abundantly scattered throughout the choanosome.

Spicule complement:

Tylostrongyles, three categories of chelae and spherancorae (Figs. 8A–8F).

Ectosomal and choanosomal tylostrongyles (Fig. 8A): from more or less straight to entirely bent on its length. They show a wider central zone, narrowing asymmetrically toward differently marked tylotoid ends (Fig. 8F), giving the spicule a variable shape between strongyles to tylostrongyles.

Size range: $369.6\text{--}411.8 \pm 14.5\text{--}475.3 \mu\text{m} \times 7.2\text{--}9.7 \pm 1.5\text{--}11 \mu\text{m}$

Isochelae I (Fig. 8C, d'): anchorate, with a gently curved shaft and irregularly spatulated rounded alae, often with a malformed tooth in one or both of the extremes.

Size range: $19\text{--}21.5 \pm 0.7\text{--}22.7 \mu\text{m}$

Isochelae II (Fig. 8D, b'): With an almost straight shaft and three alae, presenting a prominent fusion between the lateral alae and the shaft.

Size range: $30.1\text{--}35.2 \pm 2.9\text{--}38.6 \mu\text{m}$

Isochelae III (Fig. 8E, c'): With a long, gently curved shaft and slightly asymmetrical ends, e.g., the alae of one extreme are ca. 1.5 longer than those of the opposite extreme (anisochelae appearance). Alae are usually flat and with a straight end (occasionally with a bifid appearance), occupying ca. $\frac{1}{4}$ of the spicule size.

Size range: $33.2\text{--}39.5 \pm 5.1\text{--}47.8 \mu\text{m}$

Spherancorae (Fig. 8B, a'): with an elongated shape, and fimbriae on its internal face, which can be free or fused to varying degrees. Spherancorae with incompletely fused alae are present.

Size range: $38.9\text{--}44.4 \pm 6.7\text{--}51.2 \times 20\text{--}21.8 \pm 1.9\text{--}24.2 \mu\text{m}$

Geographic distribution and type locality:

The species seems so far to be endemic to the Mediterranean Sea (Fig. 4), having only been recorded from its type locality off Calvi, on the Corsica island (Pulitzer-Finali, 1983) and, more recently, from the Mallorca Channel (Díaz, Ramírez-Amaro & Ordines, 2021).

Regarding its ecology, while records are still scarce it appears to occur at rhodolith beds and rocky environments close to the limit of the continental shelf, between 104 to 134 m depth.

Etymology:

From the Latin *intermedia* (“in between”). The name refers to its unique possession of a third intermediate category of isochelae, contrary to almost all other *Melonanchora* species, which only possess two.

Remarks:

The closest species to *M. intermedia* sp. nov. would be *M. emphysema*, a typical deep-sea species also recorded from the Mediterranean Sea. Both species share the presence of tylostrongyles as their only megascleres, yet their microscleres present clear divergences, with isochelae being smaller in size in *M. intermedia* sp. nov. compared to *M. emphysema* (average length (I) 21.5 and (II) 35.2 vs. (I) 24.7 and (II) 60.2 μm , respectively; Tables 2 & 3), as well as the presence of a third category of chelae with flat, slightly asymmetrical ends in *M. intermedia* sp. nov. In this sense, in their description of ‘*Melonanchora emphysema*’ Díaz, Ramírez-Amaro & Ordines (2021) only mentions two chelae categories with no apparent aberrant morphologies, which could cast doubts about its placement between *M. emphysema* or *M. intermedia* sp. nov. Nevertheless, all spicular categories mentioned in Díaz, Ramírez-Amaro & Ordines (2021) fall within the size range of *M. intermedia* sp. nov. (Table 3), and its biggest isochelae category possess flat ends, which is one of the defining characteristics of *M. intermedia* sp. nov. Regarding the fact that only two chelae categories could be identified in his specimen, it is possible that isochelae II and III might have been confused in optical microscopy as, in fact, Fig. 16 of that same publication depicts a isochelae with rounded alae which matches in size (ca. 35 μm) the isochelae II category of the holotype. It is also interesting to note that the smallest isochelae category in the holotype of *M. intermedia* sp. nov. usually showed alae with aberrant morphologies (Fig. 8C), a feature that was not described for the Mallorca specimen. As so, this might point out that the presence and/or abundance of certain of chelae types within this species might be subjected to a certain degree of intraspecific variation. Finally, the Mallorca specimen shares with *M. intermedia* sp. nov. a subglobular appearance, as well as depth range and habitat (100–140 m depth) which further supports its inclusion as *M. intermedia* sp. nov. as opposed to *M. emphysema*, which appears to be an encrusting sponge mostly limited to the deep-sea and other cold-water environments.

Melonanchora insulsa sp. nov.

(Fig. 1E; 9)

Synonymy:

Melonanchora elliptica Schmidt, 1880: 85, pl. IX fig. 8.Not *Melonanchora elliptica* Carter, 1874: 212.

Material examined.

Holotype (here designated): MZS Po165, Gulf of Mexico, USCSS *Blake* expedition (1878–79) in the Gulf of Mexico, (24, –86), deep-sea dredging, 1879.

Description:

A small (less than 1 cm^2), thin fragment of choanosome, and some scrapped pieces of ectosome (Fig. 1E). Although we cannot report on the sponge’s original shape, Schmidt 1 (880) described the sample as a crust growing on an euptectellid glass sponge from the genus *Regadrella*.

Skeleton:

The ectosomal skeleton consists of tangential strongyles with a criss-cross arrangement, whereas the choanosomal skeleton is formed by ill-defined style-made tracts. Microscleres are widespread throughout the choanosome without a clear discernible pattern.

Spicule complement:

Styles, strongyles, two categories of chelae, spherancorae (Figs. 9A–9F).

Ectosomal strongyles (Fig. 9B): slightly flexuous, with more or less unequal ends.

Size range: $593.6\text{--}656.7 \pm 36.2\text{--}701 \times 16.1\text{--}17.1 \pm 1.2\text{--}19.5 \mu\text{m}$

Choanosomal styles (Fig. 9A): entirely smooth, mostly straight, with acerate points (Fig. 9F), sometimes slightly curved towards its distal end.

Size range: $813.4\text{--}989 \pm 41.2\text{--}1121.7 \times 19.3\text{--}20.7 \pm 1.4\text{--}22.5 \mu\text{m}$

Isochelae I (Fig. 9E): Smaller in size, and with a more prominent fusion between the lateral alae and the shaft.

Size range: $27.2\text{--}30.9 \pm 3.4\text{--}35.8 \mu\text{m}$

Isochelae II (Fig. 9D): With a gently curved shaft, and spatulated alae.

Size range: $48.6\text{--}52.3 \pm 5.1\text{--}68 \mu\text{m}$

Spherancorae (Fig. 9C): with an elliptical slightly asymmetrical shape, and teeth-like fimbriae on its internal face, which might be free or fused to different extent. Ridges of the spherancorae are unequally, gently bent, giving its ellipsoid shape a slightly asymmetrical appearance.

Size range: $52.9\text{--}56.5 \pm 4.2\text{--}62.1 \times 22\text{--}24.3 \pm 1.7\text{--}26.6 \mu\text{m}$

Geographic distribution and type locality:

The species is so far only known from the Gulf of Mexico (East of the Campache Escarpment, 24.0°N 86.0°W), and was collected from deep waters (Fig. 4).

Etymology:

From the latin *in-* (“not”) + *salsus* (“salted”), meaning insipid, tasteless. The name refers to the original description of the specimen made by Schmidt (1880), who regarded the sample as boring or “*uninteressanten*”.

Remarks:

Schmidt (1880) unambiguously stated that this individual from the Gulf of Mexico belonged to *M. elliptica*. However, the two types of chelae in *M. elliptica*’s have a straight shaft with free alae pointing outwards (Figs. 3D–3E), whereas in *M. insulsa* sp. nov. chelae show a slightly bent shaft and its alae are more parallel to the later (Figs. 9D–9E). Apart from their morphological differences, the smaller isochelae category appears to be bigger in *M. insulsa* sp. nov. (average length 30.9 μm ; Table 3) when compared with those from *M. elliptica* (average length 26.6 μm ; Table 1). Moreover, *M. elliptica*’s spherancorae are regularly oval (Fig. 3C), whereas *M. insulsa*’ spherancorae are irregular,




Figure 9 *Melonanchora insulsa* spicular plate. Spicular set for *Melonanchora insulsa* sp. nov. (sample MZS Po165, holotype). (A) Choanosomal style; (B) Ectosomal tylostrongyle; (C) Spherancorae; (D) Large chelae category (Chelae II); (E) small chelae category (Chelae I); (F) Detail of the styles' acerate end as seen in SEM imaging. (a') Spherancora (b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b'), (c') 300 μ m; (C), (D), (E) 30 μ m and (F) 100 μ m.

Full-size DOI: [10.7717/peerj.12515/fig-9](https://doi.org/10.7717/peerj.12515/fig-9)

somewhat asymmetrical ellipsoids (Fig. 9C), supporting the distinction of *M. insulsa* sp. nov. as a different species from *M. elliptica*.

Melonanchora maeli sp. nov.
(Fig. 1G; 10)

Synonymy:

Melonanchora emphysema; van Soest, 1993: 210, Tab. 2.

Not *Melonanchora emphysema* (Schmidt, 1875: 118).

Material examined.

Holotype (here designated): ZMA.POR.7269, Ponta Tremorosa, Ilha de Santiago, Cape Verde, (14.8833, -23.5333), 1986; ZMA.POR.P. 10826, Ponta Tremorosa, Ilha de Santiago, Cape Verde, (14.8833, -23.5333), 1986 (microscopic slide).

Description:

A small sub-globular sponge, covered with abundant, proportionally big, bulbous fistules which arise from a paper-thin like ectosome (Fig. 1G). The ectosome is only attached here and there to the cavernous choanosome, making the former easily detachable. The choanosome is beige-orange and the ectosome is somewhat whitish, yet translucent.

Skeleton:

The ectosomal skeleton consists of tangential tyloles with a more or less developed criss-cross arrangement, whereas the choanosomal skeleton is formed by ill-defined style-made tracts. Microscleres are widespread thorough the choanosome without a clear discernible pattern.

Spicule complement:

Styles, tyloles, three categories of chelae and spherancorae (Figs. 10A–10H). The sample was contaminated with tetractinellid spicules from an unidentified specimen stored altogether with the holotype.

Ectosomal tylostrongyles (Fig. 10B): slightly flexuous, with clearly marked tyloles at both ends. Very regular in size.

Size range: $531.3\text{--}590.9 \pm 37.9\text{--}627.9 \times 9.7\text{--}10.3 \pm 0.5\text{--}10.6 \mu\text{m}$

Choanosomal styles (Fig. 10A): entirely smooth and mostly straight to slightly bent, always with acerate endings. The heads vary between those of true styles to true tylostyles (Fig. 10G), albeit the later are rare.

Size range: $637.6\text{--}918.5 \pm 75.6\text{--}1062.6 \times 17.3\text{--}19.2 \pm 1.3\text{--}21.3 \mu\text{m}$

Isochelae I (Fig. 10F; d'): Small anchorate chelae, with a straight, short shaft, long fimbriae and spatulated alae.

Size range: $17.4\text{--}19.8 \pm 1.7\text{--}23.2 \mu\text{m}$




Figure 10 *Melonanchora maeli* spicular plate. Spicular set for *Melonanchora maeli* sp. nov. (sample ZMA.POR.7269, holotype). (A) Choanosomal style; (B) Ectosomal tylostongyle; (C) Spherancorae; (D) Large chelae category (Chelae III); (E) Intermediate chelae category (Chelae II); (F) Small chelae category (Chelae I); (G) Head of a style modified into a tylostyle; (H) Detail of a spherancora lateral view. (a') Spherancorae (b') Chelae III (c') Chelae II and (d') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b'), (c'), (d') 300 µm; (C), (D), (E), (F) 30 µm; (G) 400 µm and (H) 20 µm. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac_img.jpg\) DOI: 10.7717/peerj.12515/fig-10](https://doi.org/10.7717/peerj.12515/fig-10)

Isochelae II (Fig. 10E; c'): The least abundant of all three chelae categories, with a slightly bent shaft, in intermediate size between isochelae I and III, with short, slender alae. Only 29 spicules could be measured.

Size range: 27–29.3 ± 1.2–31.9 μm

Isochelae III (Fig. 10D; b’): The biggest of the three isochelae categories, it is strikingly similar to isochelae II, with a long, slightly bent shaft and reduced slim alae. Yet, the alae are more reduced in regards to the general size of the spicule, and they are widely opened in respect to each other, contrary to isochelae II, where the separation between alae isn’t obvious.

Size range: 45.4–49.6 ± 2–53.1 μm

Spherancorae (Figs. 10C, 10H; a’): with an elongated oval shape, almost straight with just a subtle curvature near the tips, and teeth-like fimbriae on its internal face. It usually shows a slightly asymmetrical appearance.

Size range: 48.3–50.2 ± 1.7–53.2 × 17.4–19.2 ± 1.5–21.3 μm

Geographic distribution and type locality:

This is the southernmost species of *Melonanchora* known to date, and, the only species of the genus to occur in Cape Verde archipelago (14° 52′ 59.88″N 23° 31′ 59.88″W) (Fig. 4).

Etymology:

The species is dedicated to *Mael*, the Elder God of the Seas in the world of Malaz, co-created by Steven Erikson and Ian C. Esslemont, in recognition of the vast and unique universe of their novels.

Remarks:

Originally identified as *M. emphysema* (van Soest, 1993), the specimen appears to be new to science. While its spicule complement would place it close to *M. elliptica* or *M. insulsa* sp. nov. due to the possession of styles as choanosomal megascleres, the presence of three chelae categories easily tells it apart from those. Additionally, the shape of the chelae is very different to that of the abovementioned species, with considerably reduced alae in two of the chelae categories (Figs. 10D and 10E), a feature which isn’t shared by any other *Melonanchora* species. Furthermore, its spherancorae are almost straight (Fig. 10C), whereas in most other *Melonanchora* species a clear oval morphology can be observed.

Genus *Hanstoreia* gen. nov.

Diagnosis:

Massive-globular growth form, with paper-like, easily detachable thin ectosome, bearing multiple fistular processes. Ectosomal skeleton going from no apparent organization to an ill-defined crisscross of smooth strongyles to tylotes with somewhat asymmetrical ends, whereas the choanosome is composed of ill-defined acanthostyles tracts. Microscleres include typically two categories of anchorate isochelae, rarely three, with at least one in the form of acanthose, incomplete ‘spherancorae’.

Type species:

Melonanchora globogilva Lehnert, Stone & Heimler, 2006a: 9–13 (here designated).




Figure 11 Distribution map for North Pacific species previously in *Melonanchora*. Distribution map for *Hanstoreia globogilva* (red diamond), *Myxilla (B.) kobjakovae* (green square) and *Arythmata tetraditifera* (purple circle). Projected view (UTM Zone 31N (WGS84)) with geographic (WGS84) coordinates indicated for reference. A grey line represents the 1,000 m depth isobaths. Geographic and bathymetric data used was obtained from <http://www.natureearthdata.com>.

Full-size DOI: 10.7717/peerj.12515/fig-11

Etymology:

The genus is dedicated to a much esteemed and dearly missed Nordic colleague, Hans Tore Rapp (University of Bergen), in recognition of his exceptional contributions on taxonomy and ecology of deep-sea sponges of the boreal and Arctic regions.

Remarks:

Hanstoreia globogilva was recently described from the Pacific Ocean (Fig. 11), and tentatively assigned to the genus *Melonanchora*, yet it presented some unique spicule types absent from their Atlantic counterparts (Lehnert, Stone & Heimler, 2006a). In this sense, the species clearly resembled *M. elliptica*, the type species of *Melonanchora*, yet it possessed acanthostyles (Fig. 12A) as choanosomal megascleres and particular isochelae with dentate fimbria (Fig. 12C) along the internal face of its alae and shaft, which were reminiscent of spherancorae, the main diagnostic feature for *Melonanchora*.

The placement of this species within *Melonanchora* was initially based on its external morphology (Fig. 1F) and, under the consideration that other *Melonanchora* species (viz. *M. tetraditifera* Koltun, 1970 and *M. kobjakovae* Koltun, 1958) had been previously described with incomplete 'spherancorae' (Koltun, 1958, 1970). However, SEM observation of Koltun's species (this study, Figs. 13–14) proved that those species did not bear true spherancorae but more or less complete cleistochelae, and therefore both *M. tetraditifera* and *M. kobjakovae* need to be reassigned to other genera (See below).

While the dissimilarities between *H. globogilva* and *Melonanchora* are quite clear (smooth vs. acanthose choanosomal megascleres, complete vs. incomplete ‘spherancorae’), they also share several traits (mainly two categories of smooth isochelae, ectosomal tylostrongyles to strongyles, a thin translucent paper-like ectosome and a more or less subspherical external morphology) thus, arguments both in favour and against erecting a new genus for *H. globogilva* could be made. Regarding *H. globogilva*’s external appearance, within Poecilosclerida there are other unrelated genera apart from *Melonanchora* (viz. *Cornulum*, family Acarnidae; *Coelosphaera*, family Coelosphaeraeidae) which might present a subglobular appearance and possess a warty, paper-like ectosome. Thus, external appearance alone does not represent a reliable character for genus assignment.

Regarding its spicular complement and skeletal arrangement, *H. globogilva* is indubitably closer to *Melonanchora* than to any other genera within Myxillidae, but it still presents major differences with the former. In this sense, *H. globogilva* possesses true acanthostyles, which are lacking from any other *Melonanchora* representative so far. Furthermore, all known *Melonanchora* possess an ectosomal arrangement of tangential tylostrongyles forming a dense, well-defined crisscross pattern (Fig. 2C), whereas in the choanosome megascleres are mostly arranged in spicule tracts, with some free spicules in between. On the contrary, in the case of *H. globogilva* tylostrongyles in the ectosome are mostly arranged in a confused manner, whereas their choanosomal tracts are ill-defined and with abundant free spicules in between. Nevertheless, the main difference between *H. globogilva* and *Melonanchora* would be that of its supposed incomplete ‘spherancorae’. In this sense, *H. globogilva* possesses unique, acanthose square-shaped chelae, which might be reminiscent of spherancorae while still in formation. Nevertheless, it has already been proven that unique microscleres, including chelae derivatives, might have evolved independently by phylogenetically distant species. In this sense, a similar case to that of *H. globogilva* and *Melonanchora* would be that of the proposed synonymy of *Abyssocladia Lévi, 1964* with *Phelloderma Ridley & Dendy, 1886* by *van Soest & Hajdu (2002)*. *Abyssocladia* was known from just three ill-known species while *Phelloderma* was monotypic, but both genera appeared to share the possession of a unique, apparently identical chelae type in the form of ‘abyssochelae’. Nevertheless, and as noted by the authors, both species greatly differed in all other aspects, including general shape, skeletal architecture and the rest of its spicular complement. The discovery of additional species of *Abyssocladia* casted additional doubts about the genus status, which was then revived and reassigned to the family Cladorizhidae based on its similar skeletal arrangement, presence of sigmancistras and shared carnivorous habit (*Vacelet, 2006*). Finally, the use of molecular markers demonstrated that *Abyssocladia* and *Phelloderma* were not closely related (*Vargas et al., 2013*), and thus that, despite their striking similarity, their unique chelae had developed independently. Lastly, it is also worth noticing that if included in *Melonanchora*, *H. globogilva* would be the sole representative of the genus in the Pacific, whereas all other species occur in the North Atlantic.

While *Melonanchora* and *H. globogilva* could be arguably closer to each other than to other Myxillidae, based on previous precedents we have decided to erect a new genus, *Hanstoreia* gen. nov. to allocate *H. globogilva*, rather than including it in *Melonanchora*.

Hanstoreia globogilva (Lehnert, Stone & Heimler, 2006a)
(Figs. 1F; 12)

Synonymy:

Melonanchora globogilva Lehnert, Stone & Heimler, 2006a: 9–13, fig. 4a–4f, fig. 5a–5d;
Stone, Lehnert & Reiswig, 2011: 88, Appendix IV. 168–169.

Melonanchora globoblanca Lehnert, Stone & Heimler, 2006a: 12 (misspelling of the former).

Material examined.

Holotype: NMNH-USNM 1082996, north of Amlia Island, Aleutian Islands (58.46902, –173.59802), 190 m depth, 2006.

Description:

Sub-spherical shape, with an easily detachable paper-like thin ectosome bearing abundant bulbous fistules (Fig. 1F). The choanosome is light-yellow and the ectosome is somewhat translucent-whitish, in life.

Skeleton:

The ectosomal skeleton consists on a loose crisscross of spicules arranged perpendicularly to the surface here and there, yet for most of it no clear arrangement can be discerned. The choanosome consists of ill-arranged tracts of tylotes and acanthostyles, without a clear discernible orientation, and with the tylotes being restricted to the upper areas of the choanosome. Microscleres are abundant and concentrate towards the choanosomal tracts.

Spicule complement:

Tylotes, acanthostyles, and three chelae categories, one of them in the form of incomplete ‘spherancorae’ (Figs. 12A–12F).

Ectosomal tylotes (Fig. 12B): Unevenly flexuous, with a central thickening, unequally thinning towards both ends, which show variable tyles with variable swellings.

Size range: $598.9\text{--}675 \pm 22.5\text{--}724.5 \times 9.7\text{--}10.9 \pm 2.2\text{--}14.5 \mu\text{m}$

Choanosomal acanthostyles (Fig. 12A): Slightly curved along its length, with an acerate point. Spines are short and stout, moderately abundant along the entire shaft but the tip. Size range: $589.3\text{--}638.3 \pm 30\text{--}677.3 \times 27\text{--}28 \pm 1.1\text{--}29 \mu\text{m}$

Isochelae I (Fig. 12E): with a straight shaft, well-developed fimbriae and spatulated alae, the lateral ones largely fused with the shaft.

Size range: $23.1\text{--}26.2 \pm 1.1\text{--}27 \mu\text{m}$

Isochelae II (Fig. 12D): Almost identical to isochelae I, but bigger in size.

Size range: $48.3\text{--}64.4 \pm 6.8\text{--}67.6 \mu\text{m}$

Spherancorae (Fig. 12C): Uncompleted, with free teeth, resembling chelae. As in all other *Melonanchora*, dentate fimbriae cover its internal face.

Size range: $77.3\text{--}86.9 \pm 2.8\text{--}91.8 \times 27\text{--}30 \pm 2.3\text{--}33.8 \mu\text{m}$




Figure 12 *Hanstoreia globogilva* spicule plate. Spicular set for *Hanstoreia globogilva* (sample NMNH-USNM 1082996, holotype). (A) Choanosomal acanthostyle; (B) Ectosomal tylostongyle; (C) Spherancorae; (D) Large chelae category (Chelae II); (E) small chelae category (Chelae I); (F) General view of *H. globogilva*'s spicules by SEM imaging. (a') Spherancora (b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b'), (c') 300 μm ; (C), (D), (E) 30 μm and (F) 500 μm . [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac_img.jpg\) DOI: 10.7717/peerj.12515/fig-12](https://doi.org/10.7717/peerj.12515/fig-12)

Geographic distribution:

The species appears to be rare, as it has only been seldomly recorded from deep bottoms around the Aleutian Archipelago (Lehnert, Stone & Heimler, 2006a; Stone, Lehnert & Reiswig, 2011) (Fig. 11).

Remarks:

The original description mentions a second category of spherancorae-isochelae with outer dented margins which could not be found again upon re-examination of the type material. As they are in the same size-range as the incomplete 'spherancorae', they are here regarded as likely to constitute aberrant modifications or developmental stages of *H. globogilva*'s unique chelae. Additionally, the re-examination of the type material made it clear the existence of a second, larger, isochelae category almost identical to its smallest one but much less abundant, which might explain its absence from the species' original description.

Genus *Myxilla* Schmidt, 1862Subgenus (*Burtonanchora*) Laubenfels, 1936

Type species:

Myxilla (*Burtonanchora*) *crucifera* Wilson, 1925 (by original designation).

Diagnosis:

Myxilla with smooth choanosomal styles. Chelae are three-teethed, with occasional polydentate modifications (amended from van Soest, 2002).

Myxilla (*Burtonanchora*) *kobjakovae* (Koltun, 1958)

(Fig. 13)

Synonymy:

Melonanchora kobjakovae Koltun, 1958: 58, fig. 13; Koltun, 1959: 122, fig. 75; pl. XVII, fig. 4; pl. XVIII, fig. 2; Javnov, 2012 (*partim*): 65–66.

Material examined:

Syntype (here designated): NHMUK 1963.7.29.23, Southern Kuril Islands, Pacific coast, *R/V Toporok* Kuril-Sakhalin expedition (1946–49) (Stns 127, 128), Deep-sea dredging, 1949. Exchanged with V. M. Koltun in July 1963.

Description:

The sponge is tubular, digitate or funnel shaped, with a long stem. Its surface is smooth, with the oscules being located on the top of the finger-like processes in the digitate forms. Colour bright orange in life, and from ochre to dark-brown, in alcohol.

Skeleton:

Choanosomal skeleton consisting of a dense isodictial reticulation of multispicular tracts embedded in spongin fibres without echinating spicules. Ectosomal skeleton formed by a tangential layer of more or less disarranged spicules.

Spicule complement:

Styles, strongyles, and two categories of chelae (Figs. 13A–13E).




Figure 13 *Myxilla kobjakovae* spicule set Spicular set for *Myxilla (B.) kobjakovae* (sample NHMUK 1963.7.29.23, holotype). (A) Choanosomal style; (B) Ectosomal strongyle; (C) Large chelae category (Chelae I); (D) Small chelae category (Chelae II); (E) Style's aberrant end; (F) close up view of the strongyles microspinned end; (F') close up view of the strongyles' microspinned other end. (a') Chelae I (b') Chelae II relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b') 150 μm ; (C), (D), (E) 30 μm and (F), (F') 10 μm . [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.12515/fig-13](https://doi.org/10.7717/peerj.12515/fig-13)

Ectosomal strongyles (Fig. 13B): Straight, short and stout, with a subtle swelling at each end (Fig. 13f, f'), finished in a ring of weak spines, typical of *Myxilla*. They can also be found scattered through the choanosome.

Size range: 140.3–190.3–323.8 \pm 12.2 \times 7.1–9.8–12.5 \pm 2.1 μm

Choanosomal styles (Fig. 13A): slightly curved along its length, with an acerate distal end and a proximal end sometimes vaguely inflated.

Size range: 327.5–397.5–567.3 \pm 23.2 \times 17.8–20.3–22.6 \pm 1.9 μm

Isochelae I (Fig. 13D): Unusual small anchorate isochelae with three prominent alae ending in a double hook-like termination. The alae of both ends almost contact each other, somewhat resembling a cleistochelae. Fimbriae are well developed, and present an inner hook on its lower part which points towards the interior of the chelae.

Size range: 29.2–33.3–35.7 \pm 2.8 μm

Isochelae II (Fig. 13C): Anchored, three-teethed chelae, with spatulated alae. It has clear, well developed fimbriae, which expand from the shaft.

Size range: 60.1–79.7–87.6 ± 7.8 μm

Geographic distribution:

So far, the species has only been recorded from the Okhotsk Sea, at the Kuril, Iturup and Urup islands (Koltun, 1958, 1959; Javnov, 2012; Guzii et al., 2018) and the Kamchatka peninsula (Calkina, 1969) at depths ranging from 28 to 231 m (Fig. 11).

Remarks:

Myxilla (B.) *kobjakovae* was initially assigned to *Melonanchora* based on the presence of smooth choanosomal megascleres and spherancorae (Koltun, 1958). Yet, after re-examining the holotype, we verified that those supposed spherancorae were in fact cleistochelae derivatives (Fig. 13D). Additionally, *M. kobjakovae* clearly deviates from *Melonanchora* species in growth form, lack of a paper-like ectosome, and type of megascleres. Besides *Melonanchora*, just two other Myxillidae genera possess smooth megascleres: *Myxilla* (*Burtonanchora*) Laubenfels, 1936 and *Stelodoryx* Topsent, 1904. Both genera resemble each other in most aspects (Lehnert & Stone, 2015), yet *Stelodoryx* is defined as possessing polydentate anchorate isochelae whereas *Myxilla* (B.) has exclusively three-teethed anchorate isochelae (van Soest, 2002). However, *Myxilla* (B.) *asigmata* Topsent, 1901 has been observed to possess chelae with 3–5 alae (Ríos & Cristobo, 2007), implying that the definition of *Myxilla* should be modified to include the eventual possession of polydentate chelae. On the other hand, as a result of the inclusion of some other genera as synonyms of *Stelodoryx* by van Soest (2002), some of the current species of *Stelodoryx* possess three-teethed chelae (viz. *Stelodoryx lissostyla* (Koltun, 1959)). As to, whether *Stelodoryx* and *Myxilla* are synonymous or two different genera is unclear and in need of a taxonomic revision.

The presence of polydentate chelae, while not specific enough, is still used as the main classifying feature to distinguish *Myxilla* and *Stelodoryx* (Bertolino et al., 2007; Lehnert & Stone, 2015). Thus, the new species is here referred to *Myxilla* (*Burtonanchora*) due to the possession of three-teethed anchorate chelae, yet it differs from most other *Myxilla* (B.) in the absence of sigmas, possession of two chelae categories, one of them in the form of cleistochelae, and its stalked growth form. Further reclassification of the species should not be ruled out in light of a broader Myxillidae review.

Finally, the species description in the Russian Fauna of the East seas (Javnov, 2012) depicts varying morphologies for *M. kobjakovae*. While polymorphism is common in sponges, the huge variations depicted in the Russian individuals, which range from the typical digitate-branching orange sponge, to conical-shaped or tubular-rimmed, cream coloured individuals (Javnov, 2012) suggest they may represent a different related species.

Genus *Arhythmata* gen. nov.

Type species:

Melonanchora tetradedritifera Koltun, 1970 (here designated).

Diagnosis:

Lamellate sponge, apparently resulting from coalescent digitations, with the surface slightly uneven. Ectosome thin, coriaceous, easy to detach, with subectosomal cavities. Oscula are large and unevenly spread. Choanosome crossed by numerous canals. The ectosomal skeleton is a tangential layer of strongyles perpendicular to the choanosomal spicule tracts. The choanosomal skeleton consists of a loose isodictial reticulation of multispicular style tracts embedded in spongin. The spicule complement consists of smooth choanosomal styles, ectosomal tylotes with spiny heads and three categories of polydentate chelae, among which, at least one is asymmetrically modified. So far, monotypic genus restricted to the deep-sea areas around the Okhotsk Sea.

Etymology:

From the Latin *arhythmus*, meaning “inharmonious” or “of unequal measure”, referring to the asymmetry of the alae of *A. tetradedritifera*'s peculiar chelae.

Remarks:

Arhythmata tetradedritifera was originally described as *Melonanchora tetradedritifera* based on the possession of smooth choanosomal styles, two categories of chelae, and spherancorae (Koltun, 1970). However, Koltun misidentified unique, modified chelae as spherancorae (See “The Origin of Spherancorae”), and described styles and tylostrongyles that highly differed in shape from those of other *Melonanchora* species. This spicule combination draws the species closer to *Myxilla* (*Burtonanchora*) and *Stelodoryx* as they are the only Myxillidae genera with smooth styles. However, in contrast to *M. (B.) kobjakovae*, *A. tetradedritifera* possesses polydentate (4–5) chelae, which will place the species closer to *Stelodoryx* than to *Myxilla*. However, while *Myxilla* (*Burtonanchora*) (13 accepted species; van Soest et al., 2021) represents a narrowed, well-defined, portion of *Myxilla* (91 accepted species; van Soest et al., 2021), *Stelodoryx* (18 accepted species; van Soest et al., 2021), represents an amalgam of spicule types on a rather small genus (Lehnert & Stone, 2015). Indeed, the actual concept of *Stelodoryx* is only distinguished from *Myxilla* by the presence of polydentate chelae, yet little attention has been paid to the other spicule complements (Lévi, 1993). Megascleres in *Stelodoryx* include both smooth (viz. *Stelodoryx flabellata* Koltun, 1959) or spiny (viz. *Stelodoryx mucosa* Lehnert & Stone, 2015) ectosomal tylotes or tornotes, or even styles (viz. *Stelodoryx siphofuscus* Lehnert & Stone, 2015); with choanosomal acanthostyles (viz. *S. mucosa*), smooth styles (viz. *S. siphofuscus* or *S. mucosa*), microspined styles (viz. *Stelodoryx lissostyla* (Koltun, 1959)), oxeas (viz. *Stelodoryx oxeata* Lehnert, Stone & Heimler, 2006a, 2006b) or even strongyles (viz. *S. flabellata*). Additionally, chelae may be three-teethed (viz. *S. lissostyla*) or polydentate, with teeth varying from four to seven, having from one (viz. *S. flabellata*) to three (viz. *S. oxeata*) chelae categories, with occasional accompanying sigmas (viz. *S. oxeata* or *S. mucosa*). Thus *Stelodoryx*, with just 18 species, harbours a spicule variability that might equal those of all four subgenera of *Myxilla* together (van Soest, 2002). With a combination of strongyles with microspined head and smooth styles, the closest relative to *A. tetradedritifera* within *Stelodoryx* would be *Stelodoryx jamesorri*

Lehnert & Stone, 2020 which has already been signalled as of difficult allocation within the genus *Stelodoryx* (*Lehnert & Stone, 2020*). While both species share several common traits (stout choanosomal smooth styles, ectosomal tylotes to strongyles with microspined heads and the possession of two categories of peculiar polydentate chelae), both species differ in the possession of third, unique chelae category for *A. tetradeditifera* and in their skeletal organization, being plumoreticulate in *Stelodoryx jamesorri*, as opposed to the isodyctial reticulation observed in *A. tetradeditifera*. Finally, *Stelodoryx pluridentata* (*Lundbeck, 1905*) and *Stelodoryx strongyloxeata* *Lehnert & Stone, 2020*, would also be arguably close to *A. tetradeditifera*, but they possess ectosomal styles instead of strongyles (*Lévi, 1993; Lehnert & Stone, 2020*), with additional sigmas in the former (*Lévi, 1993*) and choanosomal strongyloxeas in the later (*Lehnert & Stone, 2020*).

Consequently, a new genus, *Arhythmata* gen. nov., is here erected to properly accommodate *Melanonchora tetradeditifera*, with a diagnosis based on the combination of ectosomal microspined strongyles, smooth choanosomal styles in an isodyctial arrangement, and three polydentate chelae categories and, from which at least one is modified into an asymmetrical chelae, a rare feature within Poecilosclerida, which has been considered of taxonomic value for other genera (e.g., *Echinostylnos* spp.; *Lévi, 1993*), and which are here termed *retortochelae* (*Fig. 14C*) and defined as “*asymmetrical stout chelae in which alae are not facing their direct opposite, but the space in-between opposing alae*”. Interestingly enough, *retortochelae* appear to be very rare within Porifera, with *Echinostylnos* *Topsent, 1927* being the only other genera with asymmetrically twisted chelae, albeit not all its accepted species possess such (*Carvalho et al., 2016*). On the other hand, their stout, somewhat clesitocheliferous morphology is also relatively unusual within chelae, just being common in two other genera: *Abyssocladia* (known as abyssochelae) and *Phelloderma*. Despite their rarity, molecular analyses have shown that said chelae have been independently acquired (*Vargas et al., 2013; Göcke, Hajdu & Janussen, 2016*), thus being safe to assume that this is also the case for *Arhythmata* gen. nov. Finally, while currently the genus remains monotypic, this might change in the future upon a proper re-examination of the genus *Stylodoryx*, which is much in need of revision.

Arhythmata tetradeditifera (*Koltun, 1970*)
(*Figs. 1D, 14*)

Synonymy:

Melanonchora tetradeditifera *Koltun, 1970*: 209, fig. 22.

Material examined.

NMNH-USNM 148959, AB120069, South of Amlia Island, Central Aleutian Islands, Pacific coast, (51.8392, -173.906), 337 m depth, July 2012; NMNH-USNM 1478958, AB120046, South of Kanaga Island, Central Aleutian Islands, Pacific coast, (51.5587, 177.622), 358 m depth, July 2012.




Figure 14 *Arhythmata tetradentifera* spicule plate. Spicular set for *Arythmata tetradentifera* (sample NMNH-USNM 148959). (A) Choanosomal style; (B) Ectosomal strongyle; (C) and (C') Retortochelae; (D) Large chelae category (Chelae II); (E) Style's aberrant end; (f) close up view of the strongyles microspinned end; (f') close up view of the strongyles' microspinned other end. (a') Retortochelae (b') Chelae II and (c') Chelae I relative sizes when compared with that of the megascleres. Scale bars for (A), (B), (a'), (b'), (c') 300 µm; (C), (D), (E) 50 µm and (f), (f') 10 µm.

Full-size DOI: [10.7717/peerj.12515/fig-14](https://doi.org/10.7717/peerj.12515/fig-14)

Description:

As described in the genus definition (Fig. 1D). All the examined samples contained sand grains through the choanosome. Additionally, the colour when dry is dark brown, close to kobicha (kelp like) or taupe (brown-greyish), whereas the ectosome is whitish with wheat-like shadings.

Skeleton:

The ectosomal skeleton consists of a somewhat confused tangential layer of strongyles perpendicular to the choanosomal spicule tracts, which consists of a loose isodictial reticulation of multispicular style tracts embedded in spongin.

Spicule complement:

Styles, strongyles, three categories of chelae (Figs. 14A–14D).

Ectosomal strongyles (Fig. 14B): Short, straight, with both ends slightly spinose and slight inflated somewhat unequally (Fig. 14f, f'); a distal thorn is present, which gives them the appearance of tornote-like strongyles.

Size range: 270.5–307.8–357.4 ± 24.3 × 9.6–10.3–14.5 ± 1 μm

Choanosomal styles (Fig. 14A): Entirely smooth, slightly curved along its length, almost doubling in width the tylostrongyles.

Size range: 521–608–685 ± 54.3 × 24.1–29.3–33.8 ± 2.3 μm

Isochelae I (Fig. 14E): Small ancorate pentadentate, with a short shaft.

Size range: 48.3–60.4–67.7 ± 7.3 μm

Isochelae II (Fig. 14D): ancorate pentadentate isochelae, with a comparatively large, almost straight shaft.

Size range: 67.7–70.6–87.3 ± 3.4 μm

Retortochelae (Fig. 14C): Asymmetrical, almost ovoid, ancorate isochelae with a curved, somewhat twisted shaft and four five, long teeth. The upper and lower teeth are not facing each other but slightly displaced, in such a way that each tooth occupies the space between two opposite teeth and vice versa. This makes the chelae asymmetrical, with the alae looking as if they have been sculpted with notches and tips to accommodate the opposing alae.

Size range: 77.3–88.6–106 ± 2 × 48.3–49.1–53.1 ± 2 μm

Geographic distribution:

Currently, the species has only been located at the deep-sea waters (338 to 3,335 m depth) of the Okhotsk Sea, mostly around the Simushir Islands (Koltun, 1970; Downey, Fuchs & Janussen, 2018) and the Aleutian Islands (Fig. 11).

Remarks:

Although the holotype of this species could not be examined, the studied material fits well with Koltun's original description, in terms of spicule types and sizes (Koltun, 1970).

However, the species has been observed to possess two different chelae categories, mainly distinguished by its size and shaft lengths, which were not described by Koltun, while the spherancorae mentioned in the original description are, in fact, modified chelae with a twisted shaft, long teeth and an ovoid contour (retortochelae; Fig. 14C).

Arhythmata tetradedritifera represents a new addition to the already diverse Myxillidae fauna of the Okhotsk deep-sea and nearby areas. During the past years, several new species from the area have been included in Myxillidae (Lehnert, Stone & Heimler, 2006a, 2006b; Lehnert & Stone, 2015), which might partially respond to a high abundance of endemic benthic fauna in the area (Downey, Fuchs & Janussen, 2018). Although the genus remains monotypic for the time being, further exploration in the deep bottoms of the Okhotsk Sea and nearby areas might result in the discovery of additional species.

DISCUSSION

Diversity and biogeography of the genus *Melonanchora*

In contrast to most sponge genera, *Melonanchora* shows a quite narrow distribution, restricted to the circumpolar Arctic and some North Atlantic areas. Additionally, only one species, *M. elliptica* could be considered common across its distribution area ([Fristedt, 1887](#); [Lundbeck, 1905](#); [van Soest & De Voogd, 2015](#); [Baker et al., 2018](#)). Despite initial misidentification of fossil spherancorae ([Hinde & Holmes, 1892](#)), there are no known fossil records for the genus, thus making discussion about its origin and radiation, tentative.

Contrary to biogeographic distributions of other sponge genera, which suggest they may have a Tethyan or Gondawan origin (e.g., *Acarnus*, [van Soest, Hooper & Hiemstra, 1991](#); *Rhabderemia*, [van Soest & Hooper, 1993](#); [Hajdu & Desqueyroux-Faúndez, 2008](#); *Hamigera*, [Santín et al., 2020](#)), the current distribution of *Melonanchora* might be better explained by trans-Arctic exchanges. The opening of the Bering Strait during the late Pliocene (ca. 5.3 Ma; [Vermeij, 1991](#)), allowed a massive interchange of species among northern areas of the Atlantic and the Pacific ([Vermeij, 1991](#)), which is supported by both the fossil record ([Reid, 1990](#)) and molecular studies ([Dodson et al., 2007](#); [Coyer et al., 2011](#)). This exchange did not just occur among vagile fauna ([Dodson et al., 2007](#)), but also among benthic species ([Reid, 1990](#)), including sponges ([Ereskovsky, 1995](#)). Benthic species are known to have crossed the strait, in the several opening and closing events of the strait during the glacial and interglacial periods ([Coyer et al., 2011](#)). Additionally, during these glacial and interglacial periods, species expanded or contracted their distribution areas as a result of climate changes and their associated biotic and abiotic factors, which provided new suitable habitats ([Jansson & Dynesius, 2002](#)). The common ancestor for both *Melonanchora* (Atlantic) and *Torentendalia* gen. nov. (Pacific), might have expanded from Pacific to Atlantic waters during one of the several events that opened the Bering Strait, with the aforementioned genera resulting from the isolation of its Pacific and Atlantic populations. Once in the Atlantic, it could have expanded further south towards the tropical regions during the glacial periods ([Ereskovsky, 1995](#)). Thus, *M. maeli* sp. nov. and *M. insulsa* sp. nov., the only representatives of the genus close to the equator, might be a legacy of this latitudinal migration, being now confined to “deep-sea refugia” due to previous climatic changes ([Ereskovsky, 1995](#); [Convey et al., 2009](#)). Finally, the Mediterranean *M. intermedia* sp. nov. might represent a recent speciation from *M. emphysema*, which might have entered the Mediterranean after the Messinian Salinity Crisis, as hypothesized for other Mediterranean sponges ([Boury-Esnault, Pansini & Uriz, 1992](#); [Xavier & Van Soest, 2012](#)). However, the lack of fossil records in their current distribution area ([Ereskovsky, 1995](#)) and the lack of phylogenetic data, paired with the scarcity of material of most *Melonanchora* species, makes it difficult to properly assess the vicariant events that led to its diversification, leaving the field open for future research efforts.

The origin of spherancorae

The order Poecilosclerida [Topsent, 1928](#), build around the exclusive presence of chelae is, with over 2.500 formally described species ([van Soest et al., 2021](#)) possibly the most diverse group within Porifera ([Hooper & Van Soest, 2002](#)). The high taxon diversity parallels that of its chelae, with basic chelae morphotypes (palmate, anchorate, and arcuate) described for the first time by [Levinsen \(1893\)](#) and [Lundbeck \(1905, 1910\)](#), and several modifications of the formers ([Hajdu, van Soest & Hooper, 1994](#); [Hooper & Van Soest, 2002](#)).

In his initial description of *Melonanchora*, [Carter \(1874\)](#) assumed that the two chelae categories present his specimen where in fact early developmental stages of the unique, “melon-shaped” chelae, which characterized the genus or even, the last developmental stage of anchorate chelae ([Vosmaer, 1885](#)). While this view was soon refuted, and the “melon-shaped” chelae was recognized as a separate chelae type ([Schmidt, 1880](#)), it was not until 1885 that they were given a specific designation, “*mel*”, based on their unique shape ([Vosmaer, 1885](#)). However, the name would remain unsettled for the following years, with several authors following Vosmaer’s proposal as *melonanchoras* ([Fristedt, 1887](#); [Levinsen, 1893](#); [Arnesen, 1903](#)), while others followed Topsent’s proposed designation ([Topsent, 1892](#)) of *spherancisters* ([Thiele, 1903](#); [Topsent, 1904](#)). Topsent’s proposal however, was based on his perception that each shaft of the chelae resembled a diancistra ([Topsent, 1892](#)). However, diancistras are sigmoid derivatives ([Hajdu, van Soest & Hooper, 1994](#)) whereas spherancorae are true chelae derivatives ([Levinsen, 1893](#)). Nevertheless, the term “*melonanchora*” was identical to that of the genus, which could lead to confusion. As so, Lundbeck settled the dispute in 1905, when he designated these unique chelae as spherancorae, highlighting its chelae nature and unique oval morphology ([Lundbeck, 1905](#)).

Regarding the spherancora’s unique morphology, the common presence of developmental stages in several individuals has given a proper view of their chelae nature ([Levinsen, 1893](#)) as well of their developmental stages. Spherancorae start as slim anchorate chelae, with a thin shaft and three teeth (Fig. 15.1), of the same width. Later, those three teeth expand, until they coalesce (Fig. 15a), forming four indistinguishable shafts, all being at approximately right angles in respect to each other, and giving the spherancorae its characteristic oval shape (Fig. 15.2). While not usually visible as they occur on the internal shaft’s view, the junction points of the alae usually develop into a swelling in adult spherancorae (Fig. 15c). Right after the arcs are formed, the spherancorae begin the development of its internal “teeth-brims” (Fig. 15.3), as in other teathed chelae, (e.g., *Guitarra solorzanoi*; [Cristobo, 1998](#)). The internal dentate fimbriae are regularly arranged along the internal surface of the *Melonanchora*’s shaft (Fig. 15.4; 15.5; 15.5’), yet the teeth are not fused to the shafts, but are free and protrude from a small ridge formed at side of the shafts (Fig. 15c). The length and a degree of fusion vary between individuals of the same species, ranging from the most common free teeth forms (Fig. 15b), to partially joined teeth, or even almost coalescent teeth. This intraspecific variability

regarding the fusion degree of the alae might partially reflect silica availability at the time the spicules were formed (Uriz *et al.*, 2003), as it has been reported for other sponge taxa (e.g., *Bavestrello*, *Bonito* & *Sará*, 1993; *Cárdenas* & *Rapp*, 2013).

While the spherancora's morphology seems to be rather conservative between *Melonanchora* species, *H. globogilva* possesses unique acanthose chelae, which would resemble incomplete 'spherancorae' (Fig. 12C). These chelae present non-coalescenting alae and internal teeth-brims, which might loosely resemble those in placocheles (Cristobo, 1998), yet this is likely to be anecdotal, and of little to no taxonomical significance. Nevertheless, the architecture of this third chelae category could be consistent with that of the developmental stages of true spherancorae (Fig. 15), as its teeth-brims are not restricted to the alae, but are present all along the shaft's internal surface, as true spherancorae. As so, *H. globogilva*'s unique chelae might point towards a common ancestor between both *Melonanchora* and *Toretendalia* gen. nov., and represent, in fact, ancestral 'spherancorae' (Lehnert, Stone & Heimler, 2006a), further supporting its chelae ancestry.

Confusion between spherancorae and other spicular types is highly unlikely, yet there are a few spicular types that could, or have been, confused with spherancorae. Placocheles and derivatives (Fig. 16C) are a complex group of microscleres, synapomorphic for the family Guitarridae (Uriz & Carballo, 2001; Hajdu & Lerner, 2002), which share with spherancorae the possession of teeth-brims along the shafts and alae (Hajdu, van Soest & Hooper, 1994). While the possible affinity of Guitarridae with Myxillidae was eventually proposed (van Soest, 1988), this was poorly supported, among others, by the likely palmate origin of placocheles (Hajdu, van Soest & Hooper, 1994), which are absent in Myxillidae. The development of teeth-brims among chelae, while not a common trait, should be regarded a homoplastic character acquired independently by several taxa. Apart from placocheles, both cleistochelae (*viz.* *M. (B.) kobjakovae*) and clavidiscs (Hinde & Holmes, 1892; Ivanik, 2003) have been interpreted at some point as spherancorae due to their ovoidal morphology. Fossil *Merlia* species (*viz.* *Merlia morlandi* (Hinde & Holmes, 1892); *Merlia* sp. Ivanik, 2003; Lukowiak, Pisera & Stefanska, 2019) have been confused with *Melonanchora* due to the similarity between clavidiscs (Fig. 16D) and spherancorae (Fig. 16A) lateral view. Nevertheless, clavidiscs are synapomorphic for *Merlia* and believed to be sigmancistra derivatives (Hooper & Van Soest, 2002), contrary to the spherancora's chelae origin. Coincidentally, the lateral view of cleistochelae (Fig. 16B) has also been misinterpreted as spherancorae, with which they share their chelae origin and the presence of partially fused alae. However, cleistochelae lack the inner teeth-brims and present a single arc (2D byplan), resulting from the fusion of all free alae in a single piece, whereas spherancorae present two arcs (3D byplan), as they result from the fusion of each one of the free alae with its opposing counterpart.

Finally, and despite their unique morphology amongst sponge microscleres, the function of spherancorae, as that of many other microscleres, remains unclear. In this sense, while megascleres possess a clear architectural role in the sponge skeleton, microscleres are mostly believed to play a consolidating or defensive role, if any




Figure 15 Developmental stages of spherancorae. Formation process of a spherancorae. 1. Initial stages of formation; the chelae origin can still be observed, with a full formed shaft (*s*) and free alae (*al*) still visible; 2. Fusion phase; the alae coalesce forming the four shafts; alae's junction points (*jp*) are visible (a.); 3. Thickening phase; the shafts start to thicken, and start forming the ridges (*r*) from which the fimbriae will later develop; 4. Fimbriae development phase; fimbriae start developing on the ridges, while the shafts continue thickening; 5. Fully formed spherancorae, with complete, free fimbriae (*f*) clearly visible (b.); 5'. Internal view of a spherancorae, visible due to the braking of a shaft; the junction point (*jp*) of the alae is still visible on the internal side of the shafts as a swelling (c.), while it is observable that fimbriae (*f*) are mostly free, only attached to the shafts (*s*) by its base. Scale bar for Figures 1–5 is 20 µm, whereas for figures a., b., and c. is 10 µm. All images were taken from *Melonanchora tumultuosa* sp. nov. (NHMUK Norman Coll. 1898.5.7.38). [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90_img.jpg\) DOI: 10.7717/peerj.12515/fig-15](https://doi.org/10.7717/peerj.12515/fig-15)

(*Uriz et al., 2003*). In the *M. elliptica* holotype, spherancorae were observed to concentrate and form a dense palisade on the outer layer of the choanosome as well as surrounding the aquiferous canals, which could imply a defensive role, or a possible role in the architecture of the aquiferous system, yet this was not observed in any other of the samples analysed, and remains speculative.




Figure 16 Comparison between spherancorae and other fused chelae. (A) Spherancorae from *Melonanchora elliptica* (NHMUK 1882.7.28.54a); (B) cleistochelae from *Clathria* sp. (NHMUK 1910.10.12.18); (C) placochelae from *Guitarra dendyi* (Kirkpatrick, 1907) (Ríos pers. Coll.); (D) Clavidisc from *Merlia normani* Kirkpatrick, 1908 (Uriz pers. Coll.). [Full-size !\[\]\(fcc3264021d438d9732560e78099f674_img.jpg\) DOI: 10.7717/peerj.12515/fig-16](https://doi.org/10.7717/peerj.12515/fig-16)

ACKNOWLEDGEMENTS

The authors would like to thank Inés Fernández, Alejandra Calvo and Cristina Boza from the IEO Gijón for their help with samples from the Galician Bank, as well as Mar Sacau, from the IEO Vigo for allowing us to use the material from the NEREIDA surveys, Inês Gregório (CIIMAR) for her support with the molecular work performed and lastly, Christine Morrow for critically reviewing the manuscript's English use. Additionally, the first author would like to thank Maria Pascual for her invaluable hospitality in Gijón; Tetiana Stefanska for her help getting access to Russian literature; Patricia Baena and Marina Biel for their help with some samples; Carlota Ruiz, for listening to endless conjecturations and her aid with bibliography; Jordi Grinyó, for always being there with a helping hand; Alfredo Quintana (University of Oviedo), María García (CEAB-CSIC) and José Manuel Fortuño (ICM-CSIC) for their technical assistance during SEM image acquisition and finally, to all the personnel in the Centro Oceanográfico de Gijón (IEO) for their warmth during his stay there. The authors would also like to thank the inestimable help of all the museum's curators and staff: Dr. Tom White from the NHMUK, Maria Taviano, from the MSNG, Dr. Jean-Marc Gagnon from the CMNI, Dr. Maria Mostadius from the MZLU, Dr. Marie Meister from the MZS, Dr. Carsten Lüter from the ZMB, Allen Collins and Lisa Comer from the NMNH, Eric A. Lazo-Wasem from the YPM, Bram

van der Bijl from the NBC, Manuel R. Solórzano for providing samples from Galicia, and last, but not least, Prof. Jean Vacelet, whom allowed us access to his personal collections, for which the authors are very grateful of.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research has been performed in the scope of the SponGES project, which received funding from the European Union's Horizon 2020 Research and Innovation Programme under grant agreement no. 679849. This study was funded by the European Commission LIFE C "Nature and Biodiversity" call, and included in the INDEMARES (07/NAT/E/000732) and INTEMARES (LIFE15 IPE ES 012) projects. The Biodiversity Foundation, of the Ministry of Environment, was the institution responsible for coordination these projects. The present investigation was undertaken as part of the NAFO Potential Vulnerable Marine Ecosystems-Impacts of Deep-Sea Fisheries project (NEREIDA) (Grant Agreement S12.770786), which is supported by Spain's General Secretary of the Sea (SGM), Spain's Ministry for the Rural and Marine Environment, the Spanish Institute of Oceanography, the Geological Survey of Canada, the Canadian Hydrographic Service, Fisheries and Oceans Canada, the UK's Centre for the Environment Fisheries and Aquaculture Science (Cefas), the Russian Polar Research Institute of Marine Fisheries and Oceanography, and the Russian P.P. Shirshov Institute of Oceanology (RAS). Sample MZB 2019-1740 was collected under the ABRIC project (Ref. RTI2018-096434-B-I00) funded by the Spanish Ministry of Science and Innovation. Finally, AS was the recipient of the 2019 Young Scientist Best Paper Award of the Dept. of Marine Biology and Oceanography at the Institute of Marine Sciences, which provided funding for the SEM imaging, alongside with the grant Consolidate SGR378 Benthic Ecology from the Generalitat de Catalunya awarded to MJU with the institutional support of the 'Severo Ochoa Centre of Excellence' accreditation (CEX2019-000928-S). There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

European Union's Horizon 2020 Research and Innovation Programme: 679849.

European Commission LIFE C "Nature and Biodiversity" call.

INDEMARES: 07/NAT/E/000732.

INTEMARES: LIFE15 IPE ES 012.

NAFO Potential Vulnerable Marine Ecosystems-Impacts of Deep-Sea Fisheries project (NEREIDA): S12.770786.

Spain's General Secretary of the Sea (SGM).

Spain's Ministry for the Rural and Marine Environment.

Spanish Institute of Oceanography.

Geological Survey of Canada.

Canadian Hydrographic Service.
 Fisheries and Oceans Canada.
 UK's Centre for the Environment Fisheries and Aquaculture Science (Cefas).
 Russian Polar Research Institute of Marine Fisheries and Oceanography.
 Russian P. P. Shirshov Institute of Oceanology (RAS).
 ABRIC project: Ref. RTI2018-096434-B-I00.
 Spanish Ministry of Science and Innovation.
 Institute of Marine Sciences.
 Generalitat de Catalunya: SGR378.
 Severo Ochoa Centre of Excellence: CEX2019-000928-S.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Andreu Santín conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- María-Jesús Uriz conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Javier Cristobo performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Joana R Xavier performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Pilar Ríos performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The specimen are deposited in the following locations:

National History Museum: NHMUK 1882.7.28.54a; NHMUK - Norman Coll. N°50 10.1.1.1417; NHMUK 1954.3.9.301 N°50; NHMUK-Norman Coll. -H. J. Carter Slide Coll. 1954.3.9.301; NHMUK -Norman Collection 1910.1.1.588; NHMUK-Sott-Ryen Coll., 1931.6.1.19; NHMUK Norman Coll. 1910.1.1.1418; NHMUK-Norman Coll. 1910.1.1.1419; NHMUK-Norman Coll. 1910.1.1.1420; NHMUK-Norman Coll. 1910.1.1.1421; NHMUK-Norwegian Coll. 1982.9.6.14.a.; NHMUK-Icelandic Coll. 1958.1.1.633; NHMUK Norman Coll. 1898.5.7.38; NHMUK, 83.12.13.70.89; NHMUK 1963.7.29.23

Museum für Naturkunde: ZMB Por 3042; ZMB Por 2680; ZMB Por 6571.

Canadian Museum of Nature: CMNI 2018-0107.

Museum of Biology of Lund: MZLU L936/3483; MZLU L935/3858.

Swedish Museum of Natural History: NRM 113070.

Yale Peabody Museum of Natural History: YPM IZ 006552.PR.

Naturalis Biodiversity Center: ZMA.POR.P.10797; ZMA.POR.1548; ZMA.POR.P.10800; ZMA.POR.20192; ZMA.POR.P.10799; ZMA.POR.20559.b; ZMA.POR.20473.b; ZMA.POR.20551; ZMA.POR.P.10798; ZMA.POR.20353.a; ZMA.POR.P.10795; ZMA.POR.P.20020; ZMA.POR.20020; ZMA.POR.P.10829; ZMA.POR.20467; ZMA.POR.P.10828; ZMA.POR.20175.b; ZMA.POR.P.10827; ZMA.POR.20335; ZMA.POR.P.10796; ZMA.POR.P.10825; ZMA.POR.P.10822; ZMA.POR.P.10824; ZMA.POR.4977; ZMA.POR.P.10823; ZMA.POR.4976; ZMA.POR.7269; ZMA.POR.P. 10826

Gothenburg Natural History Museum: GNM Porifera 416; GNM Porifera 290; GNM Porifera 390; GNM Porifera 624.

Museo Civico di Storia Naturale di Genova: MSNG Vis4.7.

Museu de Ciències Naturals (Zoologia) de Barcelona: MZB 2019–1740.

Musée Zoologique de la Ville de Strasbourg: MZS Po165.

National Museum of Natural History, Smithsonian Institution: NMNH-USNM 1082996; NMNH-USNM 148959; NMNH-USNM 1478958.

Additional information on each of the analyzed specimens can be found on the species descriptions.

New Species Registration

The following information was supplied regarding the registration of a newly described species:

Publication LSID: lsid:zoobank.org:pub:F1A22CAA-DE1F-434D-9A6B-F00853C40FF5
Arhythmata gen. nov.: urn:lsid:zoobank.org:act:25BD6F3B-D818-432B-854C-71AB54DB72BA

Hanstoreia gen. nov.:

urn:lsid:zoobank.org:act:B89C3FC5-8EDA-4D53-9FFC-9417E75E97E7

Melonanchora insulsa sp. nov.: urn:lsid:zoobank.org:act:1082F7BF-4584-47A3-B3E5-E0E8E81CC9E7

Melonanchora intermedia sp. nov.: urn:lsid:zoobank.org:act:986651DC-2C16-4EC8-8D66-D28FC963EBB0

Melonanchora maeli sp. nov.: urn:lsid:zoobank.org:act:48A1B8FE-9C46-4D9C-986E-A8EDF6383537

Melonanchora tumultuosa sp. nov.: urn:lsid:zoobank.org:act:1E7784B2-9BBE-4854-9BDF-0616D8BA8F0A

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.12515#supplemental-information>.

REFERENCES

- Alander H. 1935.** Additions to the Swedish sponge fauna. *Arkiv för Zoologi* **28B**(5):1–6.
- Alander H. 1942.** Sponges from the Swedish west-coast and adjacent waters. Ph.D. Thesis. (University of Lund, H. Struves: Göteborg)1–95.

- Allcock AL, Brierley AS, Thorpe JP, Rodhouse PG. 1997.** Restricted gene flow and evolutionary divergence between geographically separated populations of the Antarctic octopus *Pareledone turqueti*. *Marine Biology* **129**(1):97–102 DOI [10.1007/s002270050150](https://doi.org/10.1007/s002270050150).
- Arndt W. 1913.** Zoologische Ergebnisse der ersten Lehr-Expedition der Dr. P. Schottländerschen Jubiläums-Stiftung. *Jahresbericht der Schlesischen Gesellschaft für Vaterländische Kultur* **90**(1):110–136.
- Arndt W. 1935.** Porifera. In: Grimpe G, Wagler E, eds. *Die Tierwelt der Nord- und Ostsee*. Leipzig: Akademische Verlagsgesellschaft m.b.H., 1–140.
- Arnesen E. 1903.** Spongien von der norwegischen Küste. II. Monaxonida: Halichondrina. *Bergens Museum Årbog* **1903**:1–30 pls I-VII.
- Baker RJ. 1984.** A sympatric cryptic species of mammal: a new species of *Rhogeessa* (Chiroptera: Vespertilionidae). *Systematic Biology* **33**(2):178–183 DOI [10.2307/2413018](https://doi.org/10.2307/2413018).
- Baker E, Odenthal B, Tompkins G, Walkusz W, Siferd T, Kenchington E. 2018.** Sponges from the 2010–2014 paamiut multispecies trawl surveys, Eastern Arctic and Subarctic: class demospongiae, subclass heteroscleromorpha, order poecilosclerida, families crellidae and myxillidae. *Canadian Technical Report of Fisheries and Aquatic Sciences* **3253**:1–52 DOI [10.13140/RG.2.2.35195.62249](https://doi.org/10.13140/RG.2.2.35195.62249).
- Bavestrello G, Bonito M, Sará M. 1993.** Silica content and spicular size variation during an annual cycle in *Chondrilla nucula* Schmidt (Porifera, Demospongiae) in the Ligurian Sea. *Scientia Marina* **57**(4):421–425.
- Bell J, Barnes D, Turner J. 2002.** The importance of micro and macro morphological variation in the adaptation of a sublittoral demosponge to current extremes. *Marine Biology* **140**(1):75–81 DOI [10.1007/s002270100665](https://doi.org/10.1007/s002270100665).
- Bertolino M, Schejter L, Calcinai B, Cerrano C, Bremec C. 2007.** Sponges from a submarine canyon of the Argentine Sea. In: Custódio MR, Lôbo-Hajdu G, Hajdu E, Muricy G, eds. *Porifera Research: Biodiversity, Innovation, and Sustainability*. Vol. 28. Rio de Janeiro: Museu Nacional, Série Livros, 189–201.
- Best M, Kenchington E, MacIsaac K, Wareham VE, Fuller SD, Thompson AB. 2010.** Sponge identification guide NAFO area. *Scientific Council Studies* **43**:1–50 DOI [10.2960/S.v43.m1](https://doi.org/10.2960/S.v43.m1).
- Blanquer A, Uriz MJ. 2008.** ‘A posteriori’ searching for phenotypic characters to describe new cryptic species of sponges revealed by molecular markers (Dictyonellidae: *Scopalina*). *Invertebrate Systematics* **22**(5):489–502 DOI [10.1071/IS07004](https://doi.org/10.1071/IS07004).
- Blanquer A, Uriz MJ. 2010.** Population genetics at three spatial scales of a rare sponge living in fragmented habitats. *BMC Evolutionary Biology* **10**(1):13 DOI [10.1186/1471-2148-10-13](https://doi.org/10.1186/1471-2148-10-13).
- Blanquer A, Uriz MJ, Caujapé-Castells J. 2009.** Small-scale spatial genetic structure in *Scopalina lophyropoda*, an encrusting sponge with philopatric larval dispersal and frequent fission and fusion events. *Marine Ecology Progress Series* **380**:95–102 DOI [10.3354/meps07931](https://doi.org/10.3354/meps07931).
- Boury-Esnault N, Pansini M, Uriz MJ. 1992.** A new *Discorhabdella* (Porifera, Demospongiae), a new Tethyan relict of pre-Messinian biota? *Journal of Natural History* **26**(1):1–7 DOI [10.1080/00222939200770011](https://doi.org/10.1080/00222939200770011).
- Bowerbank JS. 1874.** *A monograph of the British Spongiadae*. Vol. 3. London: Ray Society. i–xvii, 1–367, pls I–XCII.
- Burton M. 1931.** The Folden Fiord. Report on the sponges collected by Mr. Soot-Ryven in the Folden Fiord in the year 1923. *Tromsø Museum Skrifter* **1**(13):1–8.
- Burton M. 1959.** Spongia. In: Fridriksson A, Tuxen SL, eds. *The Zoology of Iceland*. 2(3–4). Copenhagen & Reykjavik: Ejnar Munksgaard, 1–71.

- Calderon I, Ortega N, Duran S, Becerro M, Pascual M, Turon X. 2007.** Finding the relevant scale: clonality and genetic structure in a marine invertebrate (*Crambe crambe*, Porifera). *Molecular Ecology* **16(9)**:1799–1810 DOI [10.1111/j.1365-294X.2007.03276.x](https://doi.org/10.1111/j.1365-294X.2007.03276.x).
- Calkina AV. 1969.** On the characteristics of epifauna, western Kamchatka shelf. In: *Proceedings of the All-Union Scientific and Research Institute of Marine Fisheries and Oceanography (VNIRO)*, Vol. 65. 248–257 [in Russian].
- Carter HJ. 1874.** Descriptions and Figures of Deep-sea Sponges and their Spicules from the Atlantic Ocean, dredged up on board H. M. S. 'Porcupine', chiefly in 1869; with Figures and Descriptions of some remarkable Spicules from the Agulhas Shoal and Colon, Panama. *Annals and Magazine of Natural History* (4) **14(79)**:207–221 245–257, pls XIII–XV.
- Carter HJ. 1876.** Descriptions and Figures of Deep-Sea Sponges and their Spicules, from the Atlantic Ocean, dredged up on board H. M. S. 'Porcupine', chiefly in 1869 (concluded). *Annals and Magazine of Natural History* (4) **18(105)**:226–240 (106): 307–324; (107): 388–410; (108): 458–479, pls XII–XVI DOI [10.1080/00222937608682035](https://doi.org/10.1080/00222937608682035).
- Carter HJ. 1877.** Arctic and Antarctic sponges & c. *Annals and Magazine of Natural History*. (4) **20(115)**:38–42.
- Carvalho MS de, Lopes DA, Cosme B, Hajdu E. 2016.** Seven new species of sponges (Porifera) from deep-sea coral mounds at Campos Basin (SW Atlantic). *Helgoland Marine Research* **70(10)**:1–33 DOI [10.1186/s10152-016-0461-z](https://doi.org/10.1186/s10152-016-0461-z).
- Concepción GT, Crepeau MW, Wagner D, Kahng SE, Toonen RJ. 2008.** An alternative to ITS, a hypervariable, single-copy nuclear intron in corals, and its use in detecting cryptic species within the octocoral genus *Carijoa*. *Coral Reefs* **27(2)**:323–336 DOI [10.1007/s00338-007-0323-x](https://doi.org/10.1007/s00338-007-0323-x).
- Convey P, Stevens MI, Hodgson DA, Smellie JL, Hillenbrand C-D, Barnes DKA, Clarke A, Pugh PJA, Linse K, Craig Cary S. 2009.** Exploring biological constraints on the glacial history of Antarctica. *Quaternary Science Reviews* **28**:3035–3048 DOI [10.1016/j.quascirev.2009.08.015](https://doi.org/10.1016/j.quascirev.2009.08.015).
- Coyer JA, Hoarau G, Van Schaik J, Luijckx P, Olsen JL. 2011.** Trans-Pacific and trans-Arctic pathways of the intertidal macroalga *Fucus distichus* L. reveal multiple glacial refugia and colonizations from the North Pacific to the North Atlantic. *Journal of Biogeography* **38(4)**:756–771 DOI [10.1111/j.1365-2699.2010.02437.x](https://doi.org/10.1111/j.1365-2699.2010.02437.x).
- Crespo A, Pérez-Ortega S. 2009.** Cryptic species and species pairs in lichens: a discussion on the relationship between molecular phylogenies and morphological characters. *Anales del Jardín Botánico de Madrid* **66(1)**:71–81 DOI [10.3989/ajbm.2225](https://doi.org/10.3989/ajbm.2225).
- Cristobo FJ. 1998.** *Guitarra solorzanoi* (Porifera, Demospongiae) a new species from the Galician coast (Northeast Atlantic). *Ophelia* **48(1)**:25–34 DOI [10.1080/00785236.1998.10428675](https://doi.org/10.1080/00785236.1998.10428675).
- Cristobo FJ, Urgorri V. 2001.** Revision of the genus *Trachytedania* (Porifera: Poecilosclerida) with a description of *Trachytedania ferrolensis* sp.nov. from the north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom* **81**:569–579 DOI [10.1017/S0025315401004258](https://doi.org/10.1017/S0025315401004258).
- Cristobo FJ, Urgorri V, Solórzano MR, Ríos P. 1993.** Métodos de recogida, estudio y conservación de las colecciones de poríferos. In: Thomas B, Palacios F, Martínez-López MC, eds. *Símpoio Internacional sobre Preservación y Conservación de Colecciones de Historia Natural (Comunicaciones sobre la Situación, Preservación y Conservación de Colecciones de Historia Natural)*. Vol. 2. 277–287 ISBN 84-7483-908-4.
- Cárdenas P, Pérez T, Boury-Esnault N. 2012.** Sponge systematics facing new challenges. In: Becerro MA, Uriz MJ, Maldonado M, Turon X, eds. *Advances in Sponge Science: Phylogeny,*

- Systematics, Ecology. Advances in Marine Biology*. Vol. 61. London, United Kingdom: Academic Press 2012, 79–209.
- Cárdenas P, Rapp HT. 2013.** Disrupted spiculogenesis in deep-water Geodiidae (Porifera, Demospongiae) growing in shallow waters. *Invertebrate Biology* **132**(3):173–194 DOI [10.1111/ivb.12027](https://doi.org/10.1111/ivb.12027).
- Cárdenas P, Xavier J, Tendal OS, Schander C, Rapp HT. 2007.** Redescription and resurrection of *Pachymatisma normani* (Demospongiae: Geodiidae), with remarks on the genus *Pachymatisma*. *Journal of the Marine Biological Association of the United Kingdom* **87**:1511–1525 DOI [10.1017/S0025315407058286](https://doi.org/10.1017/S0025315407058286).
- De Paula TS, Zilberberg C, Hajdu E, Lôbo-Hajdu G. 2012.** Morphology and molecules on opposite sides of the diversity gradient: four cryptic species of the *Cliona celata* (Porifera, Demospongiae) complex in South America revealed by mitochondrial and nuclear markers. *Molecular Phylogenetics and Evolution* **62**(1):529–541 DOI [10.1016/j.ympev.2011.11.001](https://doi.org/10.1016/j.ympev.2011.11.001).
- Dendy A. 1922.** Report on the Sigmatotetraxonida collected by H. M. S. 'Sealark' in the Indian Ocean. In: *Reports of the Percy Sladen Trust Expedition to the Indian Ocean in 1905*. Vol. 7. London: Transactions of the Linnean Society of London, 1–164 pls 1–18.
- Dennis AB, Hellberg ME. 2010.** Ecological partitioning among parapatric cryptic species. *Molecular Ecology* **19**(15):3206–3225 DOI [10.1111/j.1365-294X.2010.04689.x](https://doi.org/10.1111/j.1365-294X.2010.04689.x).
- Desqueyroux-Faúndez R, van Soest RWM. 1996.** A review of Iophonidae, Myxillidae and Tedaniidae occurring in the South East Pacific (Porifera: Poecilosclerida). *Revue suisse de Zoologie* **103**(1):3–79 DOI [10.5962/bhl.part.79938](https://doi.org/10.5962/bhl.part.79938).
- Dinn C, Leys SP. 2018.** *Field guide to sponges of eastern Canadian Arctic*. Edmonton: Department of Biological Sciences, University of Alberta, Edmonton AB, T6G 2E9, 1–102.
- Dodson JJ, Tremblay S, Colombani F, Carscadden JE, Lecomte F. 2007.** Trans-Arctic dispersals and the evolution of a circumpolar marine fish species complex, the capelin (*Mallotus villosus*). *Molecular Ecology* **16**(23):5030–5043 DOI [10.1111/j.1365-294X.2007.03559.x](https://doi.org/10.1111/j.1365-294X.2007.03559.x).
- Downey RV, Fuchs M, Janussen D. 2018.** Unusually diverse, abundant and endemic deep-sea sponge fauna revealed in the Sea of Okhotsk (NW Pacific Ocean). *Deep Sea Research Part II: Topical Studies in Oceanography* **154**:47–58 DOI [10.1016/j.dsr2.2018.02.005](https://doi.org/10.1016/j.dsr2.2018.02.005).
- du Bocage JV. 1869.** New siliceous sponges from Portugal and the island of Saint-Iago (Archipelago of Cape Verde). *Jornal de Sciencias Mathematicas, Physicas e Naturaes* **2**(6):159–162 pls X–XI.
- Duran S, Pascual M, Estoup A, Turon X. 2004.** Strong population structure in the marine sponge *Crambe crambe* (Poecilosclerida) as revealed by microsatellite markers. *Molecular Ecology* **13**(3):511–522 DOI [10.1046/j.1365-294x.2004.2080.x](https://doi.org/10.1046/j.1365-294x.2004.2080.x).
- Duran S, Pascual M, Turon X. 2004.** Low levels of genetic variation in mtDNA sequences over the western Mediterranean and Atlantic range of the sponge *Crambe crambe* (Poecilosclerida). *Marine Biology* **144**(1):31–35 DOI [10.1007/s00227-003-1178-5](https://doi.org/10.1007/s00227-003-1178-5).
- Duran C, Solórzano M. 1982.** Aportaciones al conocimiento del macrozoobentos de la zona infralitoral rocosa de Galicia mediante la utilizacion de la escafandra autonoma: I. Demosponjas. *Trabajos Compostelanos de Biología* **9**:49–67.
- Díaz JA, Ramírez-Amaro S, Ordines F. 2021.** Sponges of Western Mediterranean seamounts: new genera, new species and new records. *PeerJ* **9**(1–4):e11879 DOI [10.7717/peerj.11879](https://doi.org/10.7717/peerj.11879).
- Ereskovsky AV. 1995.** Materials to the Faunistic Study of the White and Barents seas sponges. 6. The origin of the White and Barents seas sponge faunas. *Berliner Geowissenschaftliche Abhandlungen* **16**:715–730.

- Escobar D, Zea S, Sánchez JA. 2012.** Phylogenetic relationships among the Caribbean members of the *Cliona viridis* complex (Porifera, Demospongiae, Hadromerida) using nuclear and mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* **64**(2):271–284 DOI [10.1016/j.ympev.2012.03.021](https://doi.org/10.1016/j.ympev.2012.03.021).
- Fristedt K. 1887.** Sponges from the Atlantic and Arctic Oceans and the Behring Sea. Vega-Expeditionens Vetenskap. *Iakttagelser (Nordenskiöld)* **4**:401–471 pls 22–31.
- Gastaldi M, De Paula TS, Narvarte MA, Lôbo-Hajdu G, Hajdu E. 2018.** Marine sponges (Porifera) from the Bahía San Antonio (North Patagonian Gulfs, Argentina), with additions to the phylogeography of the widely distributed *Cliona* aff. *celata* and *Hymeniacion perlevis*, and the description of two new species. *Marine Biology Research* **14**(7):682–716 DOI [10.1080/17451000.2018.1506136](https://doi.org/10.1080/17451000.2018.1506136).
- Golestani H, Crocetta F, Padula V, Camacho-García Y, Langeneck J, Poursanidis D, Pola M, yokeş MB, Cervera JL, Jung D-W, Gosliner TM, Araya JF, Hooker Y, Schrödl M, Valdés A. 2019.** The little *Aplysia* coming of age: from one species to a complex of species complexes in *Aplysia parvula* (Mollusca: Gastropoda: Heterobranchia). *Zoological Journal of the Linnean Society* **187**(2):279–330 DOI [10.1093/zoolinnean/zlz028](https://doi.org/10.1093/zoolinnean/zlz028).
- Grant RE. 1826.** Notice of a New Zoophyte (*Cliona celata* Gr.) from the Firth of Forth. *Edinburgh New Philosophical Journal* **1**:78–81.
- Grant RE. 1836.** Animal kingdom. In: Todd RB, ed. *The Cyclopaedia of Anatomy and Physiology*. Vol. 1. London: Sherwood, Gilbert, and Piper, 107–118, 1–813.
- Gray JE. 1867.** Notes on the arrangement of sponges, with the descriptions of some new genera. *Proceedings of the Zoological Society of London* **1867**(2):492–558 pls XXVII–XXVIII.
- Groves CP, Cotterill FPD, Gippoliti S, Robovský J, Roos C, Taylor PJ, Zinner D. 2017.** Species definitions and conservation: a review and case studies from African mammals. *Conservation Genetics* **18**(6):1247–1256 DOI [10.1007/s10592-017-0976-0](https://doi.org/10.1007/s10592-017-0976-0).
- Guardiola M, Frotscher J, Uriz MJ. 2016.** High genetic diversity, phenotypic plasticity, and invasive potential of a recently introduced calcareous sponge, fast spreading across the Atlanto-Mediterranean basin. *Marine Biology* **163**(5):123 DOI [10.1007/s00227-016-2862-6](https://doi.org/10.1007/s00227-016-2862-6).
- Gulliksen B, Palerud R, Brattegard T, Sneli J. 1999.** Distribution of marine benthic macro-organisms at Svalbard (including Bear Island) and Jan Mayen. -Research Report for DN 1999-4. Directorate for Nature Management.
- Guzii AG, Makarieva TN, Denisenko VA, Dmitrenok PS, Popov RS, Kuzmich AS, Fedorov SN, Krasokhin VB, Kim NY, Stonik VA. 2018.** Melonoside B and melonosins A and B, lipids containing multifunctionalized ω -hydroxy fatty acid amides from the far eastern marine sponge *Melonanchora kobjakovae*. *Journal of Natural Products* **81**(12):2763–2767 DOI [10.1021/acs.jnatprod.8b00785](https://doi.org/10.1021/acs.jnatprod.8b00785).
- Göcke C, Hajdu E, Janussen D. 2016.** *Phelloderma* (Porifera: Demospongiae) and its relation to other Poecilosclerida, including description of *P. oxychaetoides* sp. nov., and redescription of the type species, *P. radiatum* Ridley & Dendy, 1886. *Journal of the Marine Biological Association of the United Kingdom* **96**(3):597–604 DOI [10.1017/S0025315414000538](https://doi.org/10.1017/S0025315414000538).
- Hajdu E. 1994.** A phylogenetic interpretation of hamacanthids (Demospongiae, Porifera), with the redescription of *Hamacantha popana*. *Journal of Zoology* **232**(1):61–77 DOI [10.1111/J.1469-7998.1994.TB01559.X](https://doi.org/10.1111/J.1469-7998.1994.TB01559.X).
- Hajdu E, Castello-Branco C. 2014.** *Hamacantha* (*Hamacantha*) *boomerang* sp. nov. from deep-sea coral mounds at Campos Basin, SW Atlantic, and redescription of *H.(H.) schmidtii* (Carter, 1882) (Hamacanthidae, Poecilosclerida, Demospongiae). *Zootaxa* **3753**(4):384–390 DOI [10.11646/zootaxa.3753.4.7](https://doi.org/10.11646/zootaxa.3753.4.7).

- Hajdu E, Desqueyroux-Faúndez R. 2008.** A reassessment of the phylogeny and biogeography of *Rhabderemia* Topsent, 1890 (Rhabderemiidae, Poecilosclerida, Demospongiae). *Revue Suisse de Zoologie* **115**(2):377–395.
- Hajdu E, Lerner C. 2002.** Family guitarridae denty, 1924. In: Hooper JNA, Van Soest RWM, eds. *Systema Porifera. A Guide to the Classification of Sponges*. Vol. 1. New York, Boston, Dordrecht, London, Moscow: Kluwer Academic/Plenum Publishers, 651–655.
- Hajdu E, van Soest RWM, Hooper JNA. 1994.** Proposal for a phylogenetic subordinal classification for poecilosclerid sponges. In: Van Soest RWM, van Kempen TMG, Braekman JC, eds. *Sponges in Time and Space. Biology, Chemistry, Paleontology. Proceedings of the 4th International Porifera Congress. Amsterdam, Netherlands, 19–23 April 1993*. Netherlands: A. A. Balkema.
- Hellberg ME. 2009.** Gene flow and isolation among populations of marine animals. *Annual Review of Ecology, Evolution, and Systematics* **40**:291–310 DOI [10.1146/annurev.ecolsys.110308.120223](https://doi.org/10.1146/annurev.ecolsys.110308.120223).
- Hentschel E. 1929.** Die Kiesel- und Hornschwämme des Nördlichen Eismees. In: Römer F, Schaudinn F, Brauer A, Arndt W, eds. *Fauna Arctica. Eine Zusammenstellung der arktischen Tierformen mit besonderer Berücksichtigung des Spitzbergen-Gebietes auf Grund der Ergebnisse der Deutschen Expedition in das Nördliche Eismeer im Jahre 1898*. Vol. 5 (4). Jena: G.Fischer, 857–1042 pls XII–XIV.
- Hinde GJ, Holmes WM. 1892.** On the sponge-remains in the Lower Tertiary Strata near Oamaru, Otago, New Zealand. *Journal of the Linnean Society. Zoology* **24**(151):177–262 pls 7–15 DOI [10.1111/j.1096-3642.1892.tb02480.x](https://doi.org/10.1111/j.1096-3642.1892.tb02480.x).
- Hooper JNA, Van Soest RWM. 2002.** Order poecilosclerida topsent, 1928. In: Hooper JNA, van Soest RWM, eds. *Systema Porifera: A Guide to the Classification of Sponges*. Vol. 1. New York, Boston, Dordrecht, London, Moscow: Kluwer Academic/Plenum Publishers, 403–408.
- ICES. 2012.** Report of the ICES Advisory Committee 2012. ICES Advice, 2012. Books 1–11, 2184.
- Ivanik MM. 2003.** *Paleogene Spongiofauna of the East-European platform and adjacent regions*. Ukraine: Kiev: Institute of geological sciences NAS of Ukraine, 202.
- Jansson R, Dynesius M. 2002.** The fate of clades in a world of recurrent climate change: Milankovitch oscillations and evolution. *Annual Review of Ecology and Systematics* **33**:741–777 DOI [10.1146/annurev.ecolsys.33.010802.150520](https://doi.org/10.1146/annurev.ecolsys.33.010802.150520).
- Javnov SV. 2012.** *Invertebrates of the Far Eastern seas of Russia (polychaetes, sponges, bryozoans, etc.)*/S.V. Yavnov. Vladivostok: Russian Island. [in Russian].
- Johnston G. 1842.** *A history of British sponges and lithophytes*. Edinburgh: W.H. Lizars. i–xii, 1–264, pls I–XXV.
- Katckova ES, Morozov GS, Ljubina OS, Saburov RM. 2018.** Biogeographic composition of the sponges (porifera) fauna in the western part of the Barents Sea. Biosystems: organization, behavior, management. In: *71st All-Russian School-Conference of Young Scientists with International Participation, 17–20 April Nizhny Novgorod*. [in Russian].
- Klautau M, Russo CA, Lazoski C, Boury-Esnault N, Thorpe JP, Solé-Cava AM. 1999.** Does cosmopolitanism result from overconservative systematics? A case study using the marine sponge *Chondrilla nucula*. *Evolution* **53**(5):1414–1422 DOI [10.1111/j.1558-5646.1999.tb05406.x](https://doi.org/10.1111/j.1558-5646.1999.tb05406.x).
- Knowlton N. 1993.** Sibling species in the sea. *Annual Review of Ecology and Systematics* **24**(1):189–216 DOI [10.1146/annurev.es.24.110193.001201](https://doi.org/10.1146/annurev.es.24.110193.001201).
- Koltun BM. 1958.** Silicon sponges (Cornacuspongida) of the region of the southern Kuril Islands and the waters surrounding southern Sakhalin. *Issledovaniya dal'nevostochnykh Morei SSR* **5**:42–77 [in Russian].

- Koltun VM. 1959.** Cornosiliceous sponges of the northern and far eastern seas of the U.S.S.R. *Opredeliteli po faune SSR, izdavaemye. Zoologicheskim muzeem Akademii nauk* **67**:1–236 [Translated from Russian to English by the Fisheries Research Board of Canada, Translation Series, 1842, 1–442].
- Koltun VM. 1970.** Sponge fauna of the Northwestern Pacific from the shallows to the hadal depths, Fauna of the Kurile-Kamchatka Trench and its environment. In: Bogorov VG, ed. *Proceedings of the Shirshov Institute of Oceanology*. Vol. 86. 179–233 [Translated from Russian to English by the Israel Program for Scientific Translations, Jerusalem 1972].
- Könnecker G, Freiwald A. 2005.** *Plectroninia celtica* n. sp. (Calcarea, Minchinellidae), a new species of “pharetronid” sponge from bathyal depths in the northern Porcupine Seabight, NE Atlantic. *Facies* **51**(1–4):53–59 DOI [10.1007/s10347-005-0068-3](https://doi.org/10.1007/s10347-005-0068-3).
- Laubenfels MW de. 1936.** A discussion of the sponge fauna of the dry tortugas in particular and the West Indies in general, with material for a revision of the families and orders of the Porifera. Carnegie Institute of Washington Publication. 467 (Tortugas Laboratory Paper 30) 1–225, pls 1–22.
- Leal CV, De Paula TS, Lobo-Hajdu G, Schoenberg CH, Esteves EL. 2016.** Morphological and molecular systematics of the ‘*Cliona viridis* complex’ from south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* **96**(2):313–322 DOI [10.1017/S0025315415001642](https://doi.org/10.1017/S0025315415001642).
- Lehnert H, Stone RP. 2015.** New species of sponges (Porifera, Demospongiae) from the Aleutian Islands and Gulf of Alaska. *Zootaxa* **4033**(4):451–483 DOI [10.11646/zootaxa.4033.4.1](https://doi.org/10.11646/zootaxa.4033.4.1).
- Lehnert H, Stone RP. 2020.** Three new species of Poecilosclerida (Porifera, Demospongiae, Heteroscleromorpha) from the Aleutian Islands. *Alaska Zootaxa* **4851**(1):137–150 DOI [10.11646/zootaxa.4851.1.5](https://doi.org/10.11646/zootaxa.4851.1.5).
- Lehnert H, Stone R, Heimler W. 2006a.** New species of Poecilosclerida (Demospongiae, Porifera) from the Aleutian Islands, Alaska, USA. *Zootaxa* **1155**:1–23 DOI [10.11646/zootaxa.1155.1.1](https://doi.org/10.11646/zootaxa.1155.1.1).
- Lehnert H, Stone R, Heimler W. 2006b.** New species of deep-sea demosponges (Porifera) from the Aleutian Islands (Alaska, USA). *Zootaxa* **1250**:1–35 DOI [10.5281/zenodo.173010](https://doi.org/10.5281/zenodo.173010).
- Levinsen GMR. 1893.** *Studier over Svampe-Spicula: Cheler og Ankere. Videnskabelige Meddelelser fra Dansk naturhistorisk Forening i Kjøbenhavn 1893*. [In Danish], 1–20, pl. 1.
- Lohman DJ, Ingram KK, Prawiradilaga DM, Winker K, Sheldon FH, Moyle RG, Ng PKL, Ong PS, Wang LK, Braile TM, Astuti D, Meier R. 2010.** Cryptic genetic diversity in widespread Southeast Asian bird species suggests that Philippine avian endemism is gravely underestimated. *Biological Conservation* **143**(8):1885–1890 DOI [10.1016/j.biocon.2010.04.042](https://doi.org/10.1016/j.biocon.2010.04.042).
- Lovén S. 1868.** Om en märklig i Nordsjön lefvande art af Spongia. Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar **25**(2):105–121.
- Lukowiak M, Pisera A, Stefanska T. 2019.** Uncovering the hidden diversity of Paleogene sponge fauna of the East European Platform through reassessment of the record of isolated spicules. *Acta Palaeontologica Polonica* **64**(4):871–895 DOI [10.4202/app.00612.2019](https://doi.org/10.4202/app.00612.2019).
- Lundbeck W. 1905.** Porifera. (Part II.) Desmacidonidae (pars.). *The Danish Ingolf-Expedition* **6**(2):1–219 pls I–XX.
- Lundbeck W. 1909.** The Porifera of East Greenland. *Meddelelser om Grønland* **29**:423–464.
- Lundbeck W. 1910.** Porifera. (Part III.) Desmacidonidae. *The Danish Ingolf-Expedition* **6**(3):1–124.
- Lévi C. 1964.** Spongiaires des zones bathyale, abyssale et hadale. *Galathea Report. Scientific Results of The Danish Deep-Sea Expedition Round the World 7:1950–1952* 63–112.

- Lévi C. 1993. Porifera Demospongiae: Spongiaires bathyaux de Nouvelle-Calédonie, récoltés par le 'Jean Charcot' Campagne BIOCAL, 1985. In: Crosnier A, ed. *Résultats des Campagnes MUSORSTOM*. Vol. 11. Paris: Mémoires du Muséum national de l'Histoire naturelle, (A), 9–87.
- Maciolek NJ, Dahlen DT, Diaz RJ, Hecker B. 2011. Outfall benthic monitoring report: 2010 results. Boston: Massachusetts Water Resources Authority. Report 2011-14, 43.
- Maciolek NJ, Doner SA, Dahlen DT, Diaz RJ, Hecker B, Hunt C, Smith WK. 2008. Outfall benthic monitoring interpretive report: 1992–2007 results. Boston: Massachusetts Water Resources Authority. Report 2008-20, 149.
- Maldonado M, Carmona MC, Uriz MJ, Cruzado A. 1999. Decline in Mesozoic reef-building sponges explained by silicon limitation. *Nature* 401(6755):785–788 DOI 10.1038/44560.
- Maldonado M, Uriz MJ. 1996. Skeletal morphology of two controversial poecilosclerid genera (Porifera Demospongiae): *Discorhabdella* and *Crambe*. *Helgoländer Meeresuntersuchungen* 50:369–390 DOI 10.1007/BF02367110.
- Mayer F, Helversen OV. 2001. Cryptic diversity in European bats. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 268(1478):1825–1832 DOI 10.1098/rspb.2001.1744.
- Michaud MH, Pelletier E. 2006. Sources and fate of butyltins in the St. Lawrence Estuary ecosystem. *Chemosphere* 64(7):1074–1082 DOI 10.1016/j.chemosphere.2005.12.002.
- Morrow C, Cárdenas P. 2015. Proposal for a revised classification of the Demospongiae (Porifera). *Frontiers in Zoology* 12(1):7 DOI 10.1186/s12983-015-0099-8.
- Murillo FJ, Kenchington E, Tompkins G, Beazley L, Baker E, Knudby A, Walkusz W. 2018. Sponge assemblages and predicted archetypes in the eastern Canadian Arctic. *Marine Ecology Progress Series* 597:115–135 DOI 10.3354/meps12589.
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403(6772):853 DOI 10.1038/35002501.
- Owen R. 1835. Mollusca–Cephalopoda Nov. Gen.-Rossia. (Owen.). In: Ross J, ed. *Appendix to the narrative of a second voyage in search of a North-West Passage, and of a residence in the Arctic regions during the years 1829, 1830, 1831, 1832, 1833. (Volume II, Appendix, Natural History)*, 120 + cxliv. London: A.B. Webster, xcii–xcix.
- Payo DA, Leliaert F, Verbruggen H, D'hondt S, Calumpong HP, De Clerck O. 2013. Extensive cryptic species diversity and fine-scale endemism in the marine red alga *Portieria* in the Philippines. *Proceedings of the Royal Society B: Biological Sciences* 280(1753):20122660 DOI 10.1098/rspb.2012.2660.
- Pulitzer-Finali G. 1983. A collection of Mediterranean Demospongiae (Porifera) with, in appendix, a list of the Demospongiae hitherto recorded from the Mediterranean Sea. *Annali del Museo civico di storia naturale Giacomo Doria* 84:445–621.
- Reid DG. 1990. Trans-Arctic migration and speciation induced by climatic change: the biogeography of *Littorina* (Mollusca: Gastropoda). *Bulletin of Marine Science* 47(1):35–49.
- Reveillaud J, Allewaert C, Pérez T, Vacelet J, Banaigs B, Vanreusel A. 2012. Relevance of an integrative approach for taxonomic revision in sponge taxa: case study of the shallow-water Atlanto-Mediterranean *Hexadella* species (Porifera: Ianthellidae: Verongida). *Invertebrate Systematics* 26(3):230–248 DOI 10.1071/IS11044.
- Reveillaud J, Remerie T, van Soest R, Erpenbeck D, Cárdenas P, Derycke S, Xavier JR, Rigaux A, Vanreusel A. 2010. Species boundaries and phylogenetic relationships between Atlanto-Mediterranean shallow-water and deep-sea coral associated *Hexadella* species (Porifera, Ianthellidae). *Molecular Phylogenetics and Evolution* 56(1):104–114 DOI 10.1016/j.ympev.2010.03.034.

- Ridley SO. 1881.** XI. Spongida. Horny and Siliceous Sponges of Magellan Straits, S.W. Chili, and Atlantic off SW Brazil. In: Gunther A, ed. *Account of the Zoological Collections made during the Survey of H.M.S. 'Alert' in the Straits of Magellan and on the Coast of Patagonia*. London: Proceedings of the Zoological Society of London, 107–141 pls X–XI.
- Ridley SO, Dendy A. 1886.** Preliminary report on the Monaxonida collected by H. M. S. Challenger. Part I. *Annals and Magazine of Natural History* **18**:325–351, 470–493
DOI [10.1080/00222938609459982](https://doi.org/10.1080/00222938609459982).
- Ríos P, Cristobo J. 2007.** Sponges of the genus *Myxilla* Schmidt, 1862 collected in Antarctic waters by Spanish Antarctic expeditions. In: Custódio MR, Lôbo-Hajdu G, Hajdu E, Muricy G, eds. *Porifera Research: Biodiversity, Innovation, and Sustainability*. Vol. 28. Rio de Janeiro: Museu Nacional, Série Livros, 525–546.
- Ríos P, Cristobo P. 2017.** Porifera. In: Bañón R, ed. *Inventario de la biodiversidad marina de Galicia: Proyecto LEMGAL*. Santiago de Compostela: Consellería do Mar, Xunta de Galicia, 159–172.
- Samaai T, Gibbons MJ, Kelly M. 2006.** Revision of the genus *Latrunculia* du Bocage, 1869 (Porifera: Demospongiae: Latrunculiidae) with descriptions of new species from New Caledonia and the Northeastern Pacific. *Zootaxa* **1127**:1–71 DOI [10.11646/zootaxa.1127.1.1](https://doi.org/10.11646/zootaxa.1127.1.1).
- Santín A, Grinyó J, Uriz MJ, Gili JM, Puig P. 2020.** First deep-sea *Hamigera* (Demospongiae: Porifera) species associated with Cold-Water Corals (CWC) on antipodal latitudes of the world. *Deep Sea Research Part I: Oceanographic Research Papers* **164**:103325
DOI [10.1016/j.dsr.2020.103325](https://doi.org/10.1016/j.dsr.2020.103325).
- Santín A, Grinyó J, Uriz MJ, Lo Iacono C, Gili JM, Puig P. 2021.** Mediterranean coral provinces as a sponge diversity reservoir: is there a mediterranean cold-water coral sponge fauna? *Frontiers in Marine Science* **8**:671 DOI [10.3389/fmars.2021.662899](https://doi.org/10.3389/fmars.2021.662899).
- Schejter L, Cristobo J, Ríos P. 2019.** *Coelosphaera* (*Coelosphaera*) *koltuni* sp. nov. (Porifera: Demospongiae): a new species from South Orkney Islands, Antarctica. *Marine Biodiversity* **49**(4):1987–1996 DOI [10.1007/s12526-019-00959-7](https://doi.org/10.1007/s12526-019-00959-7).
- Schmidt O. 1862.** *Die Spongien des adriatischen Meeres*. Vol. i–viii. Leipzig: Wilhelm Engelmann, 1–88 pls 1–7.
- Schmidt O. 1875.** Spongien. Die Expedition zur physikalisch-chemischen und biologischen Untersuchung der Nordsee im Sommer 1872. *Jahresbericht der Commission zur Wissenschaftlichen Untersuchung der Deutschen Meere in Kiel* **2–3**:115–120 pl. I.
- Schmidt O. 1880.** Die Spongien des Meerbusen von Mexico (Und des caraibischen Meeres). In: Schmidt O, ed. *Reports on the dredging under the supervision of Alexander Agassiz, in the Gulf of Mexico, by the USS 'Blake'*. Jena: Gustav Fischer, 33–90 pls V–X.
- Sollas WJ. 1885.** A Classification of the Sponges. *Annals and Magazine of Natural History* **16**(95):395 DOI [10.1080/00222938509459901](https://doi.org/10.1080/00222938509459901).
- Solórzano MR. 1990.** Poríferos del litoral gallego: estudio faunístico, distribución e inventario. PhD thesis, Universidad de Santiago de Compostela, 1036.
- Solórzano MR. 1991.** Inventario dos Poríferos do litoral galego (Porifera). Cadernos da Area de Ciencias Biolóxicas (Inventarios). In: Coruña A, ed. *Seminario de Estudos Galegos*. Vol. VII. O Castro-Sada, 54.
- Solórzano M, Durán C. 1982.** Nota preliminar sobre la fauna de poríferos asociados a *Dendrophyllia cornigera* (Lamarck, 1816) frente a las costas de Galicia (NW de España). *Actas del IIº Simposio ibérico de Estudios del Bentos Marino, Barcelona (19–22 de marzo de 1981)* **III**:101–110.

- Stone RP, Lehnert H, Reiswig H. 2011.** A guide to the deep-water sponges of the Aleutian Island Archipelago. NOAA Professional Paper NMFS 12:187.
- Thiele J. 1903.** Beschreibung einiger unzureichend bekannten monaxonen Spongien. *Archiv für Naturgeschichte* **69**(1):375–398 pl. XXI.
- Thomson CW. 1873.** *The Depths of the Sea*. London: Macmillan and Co., 527.
- Topsent E. 1890.** Notice préliminaire sur les spongiaires recueillis durant les campagnes de l'Hirondelle. *Bulletin de la Société zoologique de France* **15**:26–32, 65–71
[DOI 10.5962/bhl.part.18721](https://doi.org/10.5962/bhl.part.18721).
- Topsent E. 1892.** Contribution à l'étude des Spongiaires de l'Atlantique Nord (Golfe de Gascogne, Terre-Neuve, Açores). *Résultats des Campagnes Scientifiques Accomplies par le Prince Albert I. Monaco* **2**:1–165 pls I–XI.
- Topsent E. 1896.** Matériaux pour servir à l'étude de la faune des spongiaires de France. *Mémoires de la Société zoologique de France* **9**:113–133.
- Topsent E. 1901.** Spongiaires. Résultats du voyage du S.Y. 'Belgica' en 1897–99 sous le commandement de A. de Gerlache de Gomery. Expédition antarctique belge. *Zoologie* **4**:1–54 pls I–VI.
- Topsent E. 1904.** Spongiaires des Açores. *Résultats des Campagnes Scientifiques Accomplies par le Prince Albert I. Monaco* **25**:1–280 pls 1–18.
- Topsent E. 1913.** Spongiaires provenant des campagnes scientifiques de la 'Princesse Alice' dans les Mers du Nord (1898–1899–1906–1907). *Résultats des Campagnes Scientifiques Accomplies par le Prince Albert I de Monaco* **45**:1–67 pls I–V.
- Topsent E. 1927.** Diagnoses d'Éponges nouvelles recueillies par le Prince Albert Ier de Monaco. *Bulletin de l'Institut Océanographique Monaco* **502**:1–19.
- Topsent E. 1928.** Spongiaires de l'Atlantique et de la Méditerranée provenant des croisières du Prince Albert Ier de Monaco. *Résultats des Campagnes Scientifiques Accomplies par le Prince Albert I. Monaco* **74**:1–376.
- Uriz MJ, Carballo JL. 2001.** Phylogenetic relationships of sponges with placocheleae or related spicules (Poecilosclerida, Guitarridae) with a systematic revision. *Zoological Journal of the Linnean Society* **132**:411–428 [DOI 10.1111/j.1096-3642.2001.tb02468.x](https://doi.org/10.1111/j.1096-3642.2001.tb02468.x).
- Uriz MJ, Garate L, Agell G. 2017a.** Molecular phylogenies confirm the presence of two cryptic Hemimycale species in the Mediterranean and reveal the polyphyly of the genera *Crella* and *Hemimycale* (Demospongiae: Poecilosclerida). *PeerJ* **5**:e2958 [DOI 10.7717/peerj.2958](https://doi.org/10.7717/peerj.2958).
- Uriz MJ, Garate L, Agell G. 2017b.** Redescription and establishment of a holotype and three paratypes for the species *Hemimycale mediterranea* sp. nov. *PeerJ* **5**:e3426
[DOI 10.7717/peerj.3426](https://doi.org/10.7717/peerj.3426).
- Uriz MJ, Gili JM, Orejas C, Pérez-Porro AR. 2010.** Do bipolar distributions exist in marine sponges? *Stylocordyla chupachups* sp. nv. (Porifera: Hadromerida) from the Weddell Sea (Antarctic), previously reported as *S. borealis* (Lovén, 1868). *Polar Biology* **34**(2):243–255
[DOI 10.1007/s00300-010-0876-y](https://doi.org/10.1007/s00300-010-0876-y).
- Uriz MJ, Maldonado M, Turon X, Martí R. 1998.** How do reproductive output, larval behaviour, and recruitment contribute to adult spatial patterns in Mediterranean encrusting sponges? *Marine Ecology Progress Series* **167**:137–148 [DOI 10.3354/meps167137](https://doi.org/10.3354/meps167137).
- Uriz MJ, Turon X, Becerro MA, Agell G. 2003.** Siliceous spicules and skeleton frameworks in sponges: origin, diversity, ultrastructural patterns, and biological functions. *Microscopy Research and Technique* **62**(4):279–299 [DOI 10.1002/jemt.10395](https://doi.org/10.1002/jemt.10395).

- Vacelet J. 1969.** Eponges de la Roche du Large et de l'étage bathyal de Méditerranée (Récoltes de la soucoupe plongeante Cousteau et dragages). *Mémoires du Muséum national d'Histoire naturelle*. *Mémoires du Muséum national d'Histoire naturelle (A, Zoologie)* **59(2)**:145–219.
- Vacelet J. 2006.** New carnivorous sponges (Porifera, Poecilosclerida) collected from manned submersibles in the deep Pacific. *Zoological Journal of the Linnean Society* **148(4)**:553–584 DOI [10.1111/j.1096-3642.2006.00234.x](https://doi.org/10.1111/j.1096-3642.2006.00234.x).
- Vacelet J, Uriz MJ. 1991.** Deficient spiculation in a new species of Merlia (Merliida, Demospongiae) from the balearic Islands. In: Reitner J, Keupp H, eds. *Fossil and Recent Sponges*. Vol. i–xviii. Berlin, Heidelberg & New York: Springer-Verlag, 170–178.
- van Soest RWM. 1988.** *Tetrapocillon atlanticus* n. sp. (Porifera, Poecilosclerida) from the Cape Verde islands. *Beaufortia* **38(2)**:37–46.
- van Soest RWM. 1993.** Affinities of the Marine Demospongiae Fauna of the Cape Verde Islands and Tropical West Africa. *Courier Forschungsinstitut Senckenberg* **159**:205–219.
- van Soest RWM, Hooper JN, Hiemstra F. 1991.** Taxonomy, phylogeny and biogeography of the marine sponge genus *Acarnus* (Porifera: Poecilosclerida). *Beaufortia* **42(3)**:49–88.
- van Soest RWM. 2002.** Family myxillidae denty, 1922. In: Hooper JNA, Van Soest RWM, eds. *Systema Porifera. A Guide to the Classification of Sponges*. Vol. 1. New York, Boston, Dordrecht, London, Moscow: Kluwer Academic/Plenum Publishers, 602–620.
- van Soest RWM, Boury-Esnault N, Hooper JNA, Rützler K, de Voogd NJ, Alvarez B, Hajdu E, Pisera AB, Manconi R, Schönberg C, Klautau M, Picton B, Kelly M, Vacelet J, Dohrmann M, Díaz MC, Cárdenas P, Carballo JL, Ríos P, Downey R. 2021.** World porifera database. Available at <http://www.marinespecies.org/porifera> (accessed 15 March 2021).
- van Soest RWM, De Voogd NJ. 2015.** Sponge species composition of north-east Atlantic cold-water coral reefs compared in a bathyal to inshore gradient. *Journal of the Marine Biological Association of the United Kingdom* **95(7)**:1461–1474 DOI [10.1017/S0025315413001410](https://doi.org/10.1017/S0025315413001410).
- van Soest RWM, Hajdu E. 2002.** Family myxillidae denty, 1922. In: Hooper JNA, Van Soest RWM, eds. *Systema Porifera. A Guide to the Classification of Sponges*. Vol. 1. New York, Boston, Dordrecht, London, Moscow: Kluwer Academic/Plenum Publishers, 621–624.
- van Soest RWM, Hooper JNA. 1993.** Taxonomy, phylogeny, and biogeography of the marine sponge genus *Rhabderemia* Topsent, 1890 (Demospongiae, Poecilosclerida). In: Uriz MJ, Rützler K, eds. *Recent Advances in Ecology and Systematics of Sponges*. Vol. 57. Spain: Scientia Marina, 319–351.
- van Soest RWM, Lavaleye MS. 2005.** Diversity and abundance of sponges in bathyal coral reefs of Rockall Bank, NE Atlantic, from boxcore samples. *Marine Biology Research* **1(5)**:338–349 DOI [10.1080/17451000500380322](https://doi.org/10.1080/17451000500380322).
- Vargas S, Erpenbeck D, Goecke C, Hall KA, Hooper JN, Janussen D, Woerheide G. 2013.** Molecular phylogeny of *Abyssocladia* (Cladorhizidae: Poecilosclerida) and *Phelloderma* (Phellodermidae: Poecilosclerida) suggests a diversification of chelae microscleres in cladorhizid sponges. *Zoologica Scripta* **42(1)**:106–116 DOI [10.1111/j.1463-6409.2012.00560.x](https://doi.org/10.1111/j.1463-6409.2012.00560.x).
- Vermeij GJ. 1991.** Anatomy of an invasion — the trans-Arctic interchange. *Paleobiology* **17**:281–307 DOI [10.1017/S0094837300010617](https://doi.org/10.1017/S0094837300010617).
- Vosmaer GCJ. 1885.** The sponges of the 'Willem Barents' expedition 1880 and 1881. *Bijdragen tot de Dierkunde* **12(3)**:1–47.
- Wareham Hayes VE, Fuller S, Shea E. 2017.** Egg deposition by *Rossia palpebrosa* (Cephalopoda: Rossiinae) in deep-sea sponges, in temperate Northwest Atlantic and fringes of polar Canadian Arctic. In: *Poster presentation, 10th World Sponge Conference, Galway. Book of abstracts*. 190.

- Waters JM, Roy MS. 2004.** Phylogeography of a high-dispersal New Zealand sea-star: does upwelling block gene-flow? *Molecular Ecology* **13(9)**:2797–2806
[DOI 10.1111/j.1365-294X.2004.02282.x](https://doi.org/10.1111/j.1365-294X.2004.02282.x).
- Wilson HV. 1925.** Silicious and horny sponges collected by the U.S. Fisheries Steamer ‘Albatross’ during the Philippine Expedition, 1907–10, pls 37–52. in: contributions to the biology of the Philippine Archipelago and adjacent regions. *Bulletin of the United States National Museum* **100(2)**:273–532.
- Xavier JR, Rachello-Dolmen PG, Parra-Velandia F, Schönberg CHL, Breeuwer JAJ, Van Soest RWM. 2010.** Molecular evidence of cryptic speciation in the cosmopolitan excavating sponge *Cliona celata* (Porifera, Clionidae). *Molecular Phylogenetics and Evolution* **56(1)**:13–20
[DOI 10.1016/j.ympev.2010.03.030](https://doi.org/10.1016/j.ympev.2010.03.030).
- Xavier JR, Van Soest RW. 2012.** Diversity patterns and zoogeography of the Northeast Atlantic and Mediterranean shallow-water sponge fauna. *Hydrobiologia* **687(1)**:107–125
[DOI 10.1007/s10750-011-0880-4](https://doi.org/10.1007/s10750-011-0880-4).