

Bridging the genotype–phenotype gap for a Mediterranean pine by semi-automatic crown identification and multispectral imagery

Filippo Santini^{1,2} , Shawn Carlisle Kefauver^{3,4} , José Luis Araus^{3,4} , Víctor Resco de Dios^{1,2,5} , Saray Martín García⁶, Delphine Grivet⁷  and Jordi Voltas^{1,2} 

¹Joint Research Unit CTFC – AGROTECNIO, Av. Alcalde Rovira Roure 191, Lleida E-25198, Spain; ²Department of Crop and Forest Sciences, University of Lleida, Av. Alcalde Rovira Roure 191, Lleida E-25198, Spain; ³AGROTECNIO (Center for Research in Agrotechnology), Av. Alcalde Rovira Roure 191, Lleida E-25198, Spain; ⁴Integrative Crop Ecophysiology Group, Plant Physiology Section, Faculty of Biology, University of Barcelona, Barcelona E-08028, Spain; ⁵School of Life Science and Engineering, Southwest University of Science and Technology, 99 Qinglong Ave., Mianyang 621010, China; ⁶Föra Forest Technologies, c/Eduardo Saavedra 38, Soria E-42004, Spain; ⁷Department of Forest Ecology and Genetics, Forest Research Centre, INIA, Carretera A Coruña km 7.5, Madrid E-28040, Spain

Summary

Author for correspondence:

Filippo Santini

Email: filippo.santini@pvcf.udl.cat

Received: 30 January 2020

Accepted: 31 July 2020

New Phytologist (2021) 229: 245–258

doi: 10.1111/nph.16862

Key words: genome-wide association study (GWAS), genomics, *Pinus halepensis*, remote sensing, SNPs, unmanned aerial vehicle (UAV).

- Progress in high-throughput phenotyping and genomics provides the potential to understand the genetic basis of plant functional differentiation. We developed a semi-automatic methodology based on unmanned aerial vehicle (UAV) imagery for deriving tree-level phenotypes followed by genome-wide association study (GWAS).
- An RGB-based point cloud was used for tree crown identification in a common garden of *Pinus halepensis* in Spain. Crowns were combined with multispectral and thermal orthomosaics to retrieve growth traits, vegetation indices and canopy temperature. Thereafter, GWAS was performed to analyse the association between phenotypes and genomic variation at 235 single nucleotide polymorphisms (SNPs).
- Growth traits were associated with 12 SNPs involved in cellulose and carbohydrate metabolism. Indices related to transpiration and leaf water content were associated with six SNPs involved in stomata dynamics. Indices related to leaf pigments and leaf area were associated with 11 SNPs involved in signalling and peroxisome metabolism. About 16–20% of trait variance was explained by combinations of several SNPs, indicating polygenic control of morpho-physiological traits.
- Despite a limited availability of markers and individuals, this study provides a successful proof-of-concept for the combination of high-throughput UAV-based phenotyping with cost-effective genotyping to disentangle the genetic architecture of phenotypic variation in a widespread conifer.

Introduction

The rapid development of genotyping and phenotyping technologies is narrowing the knowledge gap between genomics and phenomics (Houle *et al.*, 2010; Großkinsky *et al.*, 2015). Hundreds of individuals can be characterized with an ever-growing number of genetic markers, covering large parts of the genome. As a consequence, the amount of genome-wide association studies (GWAS) seeking to understand the genetic basis underlying phenotypic differentiation in plants has increased exponentially (Lobos *et al.*, 2017). Common garden experiments, in which individuals from contrasting geographical origins grow under the same environmental conditions, are often used in ecological studies to understand to what extent individual differences are controlled by genetic effects

(McKown *et al.*, 2014; Baison *et al.*, 2019). Combining accurate and rapid phenotyping with genomic scans in common gardens can therefore result in an effective method to study the genetic basis of adaptation in plant species (de Villemerueil *et al.*, 2016).

Efforts towards understanding the genetic basis of adaptation in forest tree species have been limited by a lack of labour-saving phenotyping techniques and by a restricted amount of molecular information. Genome-wide sets of molecular markers are needed to gain information on the loci contributing to the genetic architecture of complex traits (Grattapaglia & Resende, 2011). Because most traits are controlled by many loci with small effects, a high density of loci is required to achieve a solid accounting of phenotypic variation (Savolainen *et al.*, 2007; White *et al.*, 2007). Once available for only a few model

organisms, sets of molecular markers are now accessible for non-model species such as forest trees (Khan & Korban, 2012; Jaramillo-Correa *et al.*, 2015).

Furthermore, the development of suitable phenotyping approaches is limited by the costs associated with manual phenotypic measurements of anatomical and physiological traits. These costs derive from the long life cycles of forest trees and the considerable dimensions of adult individuals (Ludovisi *et al.*, 2017). As an alternative, unmanned aerial vehicle (UAV) imagery is emerging as an effective high-throughput phenotyping tool to indirectly infer variation in common garden experiments of forest trees (Santini *et al.*, 2019a). Based on specific wavelengths, UAV-based multispectral imagery allows the calculation of vegetation indices, which have been used as proxies to obtain information about traits such as crown characteristics, leaf anatomy and content of photosynthetic pigments in leaves (Santini *et al.*, 2019a,b). Additionally, water status and total transpiration in trees can be inferred through thermal images (Ludovisi *et al.*, 2017; Santini *et al.*, 2019a). Moreover, innovative techniques of photogrammetry can be applied to RGB (red, green, blue) images to obtain 3D reconstructions of trees. These techniques aim at the automatic identification of single trees in forests and at estimating growth parameters such as tree diameter or height (Nevalainen *et al.*, 2017). Therefore, high-resolution remote sensing data coupled with efficient algorithms for automatic crown identification represents a potentially powerful approach for fast and accurate individual tree phenotyping (Wallace *et al.*, 2016; Santini *et al.*, 2019a). In previous studies, we developed a UAV-derived phenotyping methodology that successfully detected adaptive variation in leaf area, crown architecture, transpiration rate and photosynthetic pigments in *Pinus halepensis* Mill. (Santini *et al.*, 2019a) and *Pinus nigra* Arnold (Santini *et al.*, 2019b). Here, we further developed and combined this methodology with genomic information in a common garden of *P. halepensis* to evaluate the potential of linking novel phenotyping technology with high-throughput genotyping to identify candidate genes involved in adaptive differentiation for a widespread forest species.

Aleppo pine (*P. halepensis*) is a drought-resistant species and the most common conifer across the Mediterranean basin. Common garden experiments have revealed a wide differentiation among populations in traits such as aerial growth (Schiller & Atzmon, 2009; Voltas *et al.*, 2018), phenology (Klein *et al.*, 2013), water uptake patterns (Voltas *et al.*, 2015), hydraulic conductivity (Tognetti *et al.*, 1997) and reproductive effort (Climent *et al.*, 2008). With the goal of unveiling the genetic basis underlying this differentiation, which remains largely unexplored, a set of single nucleotide polymorphism (SNP) markers was developed by Pinosio *et al.* (2014). Ruiz-Daniels *et al.* (2018, 2019) used these markers to genotype natural populations, and detected several SNPs putatively associated with adaptation to local conditions. A limitation of these approaches derives from the complex neutral genetic structure of this species (i.e. unrelated to local adaptation), which has been shaped by past population dynamics (Serra-Varela *et al.*, 2017; Ruiz Daniels *et al.*, 2018). The strong genetic structure separating Western and Central-Eastern Mediterranean populations, with further subdivisions within

each group (Ruiz Daniels *et al.*, 2018), is prone to generate false positive associations related to nonadaptive differentiation among populations with divergent phenotypes, making the identification of loci associated with phenotypic changes challenging (Yu *et al.*, 2006). Previous analyses performed in natural populations can be complemented with common-garden information for the purpose of identifying SNPs putatively involved in relevant adaptive traits for this pine species by controlling for the effects of environmental variation and neutral structure. In this regard, a preliminary GWAS based on the set of SNPs developed by Pinosio *et al.* (2014) and performed in a common garden has provided the first insights into the genetic basis of variation in traits including growth, reproduction, water use efficiency and wood anatomy in Aleppo pine (Rodríguez Quilón, 2017).

In this study, we used *P. halepensis* populations grown in a common garden as a model system to provide a novel approach for high-throughput phenotyping of a suite of traits related to tree morphology and physiology, which are seldom considered in genotype–phenotype association studies. We aimed at deepening the genetic control of these traits by combining phenotyping through semi-automatic crown identification and multispectral imagery with genotyping at the individual level. We hypothesized that, for a species encompassing a large functional variability in characteristics related to water and carbon economy, the genetic basis of physiological and morphological variability is identifiable, at least in part, through high-throughput UAV imagery. Specifically, the objectives of this study were to: develop a high-throughput methodology for deriving tree-level information of growth traits, together with vegetation indices and canopy temperature; identify putative genes underlying the variation in such traits, test for multigenic control of phenotypic variation and estimate the proportion of phenotypic variation explained by genotypes; and provide insight into the potential of combining UAV imagery with genotypic information in disentangling adaptive variation in forest species.

Materials and Methods

Study site and plant material

The study was carried out on adult individuals of Aleppo pine growing in a common garden experiment in Altura (39°49'29"N, 00°34'22"W, 640 m asl; Castellón province, eastern Spain) (Supporting Information Fig. S1). Seeds from 56 natural populations of *P. halepensis* were collected in 1995 and planted in a forest nursery in Spain (Table S1; Fig. S2). These populations cover a large part of the species' distribution range and they are representative of the wide variety of environmental conditions in which *P. halepensis* is currently found. For each population, open-pollinated seeds were collected from 20–30 trees, spaced at least 100 m apart, and subsequently bulked into population seedlots. This sampling strategy was implemented to minimize genetic relatedness among the half-sib families and to collect a representative amount of intrapopulation variability. In 1997, 1-yr-old seedlings were planted systematically (2.5 × 2.5 m spacing) at the study site in experimental units consisting of

linear plots of four individuals. Four replicates were established following a Latinized row–column design for a total of 896 individuals (16 per population) tested in the trial. Height and diameter at breast height (DBH) were ground-measured for all trees in 2013 (at age 16 yr).

UAV flights

Flights were performed at noon on 26 July 2016 (at age 19 yr) using an octocopter (Mikrokopter OktoXL; HiSystems GmbH, Moormerland, Germany) flying under remote control at *c.* 100 m a.g.l. (above ground level). Tree crown cover fraction at the trial was 65% (Santini *et al.*, 2019a). A significant crown overlapping was observed between neighbouring trees (Fig. S1). To ensure high-quality image registration and an overlapping between pictures of *c.* 80%, forward motion was kept to less than 5 m s⁻¹ during flights. The duration of the flights was 10 min and the total area covered was *c.* 0.6 ha, corresponding to the extension of the common garden.

The UAV was equipped with three different cameras that were mounted on a two-servo gimbal (MK HiSight SLR2; HiSystems) nadir-looking in consecutive flights. First, a multispectral camera (MCA12; Tetracam Inc., Chatsworth, CA, USA) was operated to capture 15.6-megapixel calibrated-reflectance images at 10 wavelengths (450 ± 40, 550 ± 10, 570 ± 10, 670 ± 10, 700 ± 10, 720 ± 10, 840 ± 10, 860 ± 10, 900 ± 20, 950 ± 40 nm) in the visible and near-infrared (NIR) regions of the spectrum. Second, RGB images were obtained using a Mirrorless Interchangeable Lens Camera (Lumix GX7; Panasonic, Osaka, Japan). Finally, an FLIR thermal camera (Tau2 640; FLIR Systems, Nashua, NH, USA) was used for the acquisition of thermal images. Further details on camera calibration, validation of reflectance data and thermal data retrieval are reported in Methods S1 and S2.

Image processing and crown identification

In previous studies, we used a manual approach to derive plot-level phenotypic data from UAV imagery in this same trial (Santini *et al.*, 2019a). Here, we developed a workflow, schematized in Fig. 1, for the semi-automatic acquisition of tree-level phenotypic data. Two orthomosaics (one multispectral and one thermal) were obtained from raw multispectral and thermal images using the Agisoft PHOTOSCAN PROFESSIONAL software (Agisoft LLC, St Petersburg, Russia). The spatial resolution of the multispectral and thermal orthomosaics were 7 cm² and 29 cm² per pixel respectively. The geographical coordinates of four ground control points were obtained with 1-m accuracy using a GPS device (Juno 5B; Trimble Inc., Sunnyvale, CA, USA) and were used for georeferencing the orthomosaics.

RGB images were analysed through structure-from-motion (SfM) photogrammetry in the software Agisoft PHOTOSCAN PROFESSIONAL to obtain a georeferenced 3D dense point cloud of the common garden, in which the height of each point was expressed in metres above sea level. The software FUSION (McGaughey, 2012) was used to classify vegetation and soil points of the dense point cloud, and a digital terrain model

(DTM) was obtained thereafter (Fig. 2a). Subsequently, the dense point cloud and the DTM were combined in FUSION to obtain a normalized point cloud, in which the height of each point is expressed in metres above the ground. Finally, we obtained the canopy height model (CHM) from the normalized point cloud using the function `grid_canopy` implemented in the package `LIDR` (Roussel & Auty, 2018) of the R environment (R Core Team, 2019) (Fig. 2b).

The CHM was used to identify treetops using an algorithm based on local maximum filter (Popescu & Wynne, 2004) implemented in the function `tree_detection` in the R package `LIDR` (Roussel & Auty, 2018). The height of each treetop was retrieved from the CHM as an imagery-derived estimation of tree height (H_{UAV}). Finally, single crowns were segmented (i.e. outlined) in the CHM using the function `mcws` in the `FORESTTOOLS` package (Plowright, 2018), which performs a watershed segmentation (Meyer & Beucher, 1990) guided by the locations of the treetops (Fig. 2b). Tree crown area was calculated as the projection of the outlined crown on the ground. Crown area was considered as a surrogate of DBH (Lockhart *et al.*, 2005) (Fig. S3).

The accuracy of tree identification was visually checked in the software QGIS (v.3.6; QGIS Development Team, 2019) and the crowns of the misidentified trees were manually corrected (Fig. 2c,d). The obtained georeferenced crown shapes were used to identify the individual crowns from multispectral and thermal orthomosaics; that is, by using the crown shapes as template images corresponding to single trees, crowns were cropped from the orthomosaics (Fig. S3). Therefore, two images of the crown (one multispectral and one thermal) were retrieved for each of the 806 living trees of the common garden (see Results section) and used to calculate tree-level vegetation indices and canopy temperature.

Vegetation indices and canopy temperature

We calculated 10 vegetation indices (Table 1) from multispectral images for each pixel of every image, which corresponded to a particular tree crown. An average value was obtained afterwards for each of the indices per image (tree), resulting in individual tree data. Indices related to leaf area were calculated including the normalized difference vegetation index (NDVI), the optimized soil adjusted vegetation index (OSAVI), the renormalized difference vegetation index (RDVI) and the enhanced vegetation index (EVI). Additionally, we also calculated the modified chlorophyll absorption reflectance index (MCARI) and the transformed chlorophyll absorption ratio index (TCARI). The latter indices are related to both leaf area and Chl content (Daughtry, 2000). The ratio between TCARI and OSAVI (TCARI/OSAVI index) was calculated as a better estimate of Chl content, that is free of leaf area effects (Zarco-Tejada *et al.*, 2004). Furthermore, we calculated the carotenoid reflectance index 2 (CRI2), the anthocyanin reflectance index 2 (ARI2) and the water band index (WBI) to investigate other photosynthetic pigments and water content in needles. Finally, we used thermal images to retrieve crown-level estimates of canopy temperature. A detailed description of vegetation indices is available in Methods S3.




Fig. 1 Flowchart showing the different steps followed to carry out the study of association genetics in *Pinus halepensis* using red, green, blue (RGB), multispectral and thermal imagery retrieved with an unmanned aerial vehicle (UAV) as phenotyping tools.




Fig. 2 (a) Digital terrain model and (b) canopy elevation model of the common garden of *Pinus halepensis* obtained through structure-from-motion analysis of red, green, blue (RGB) images. The correctly identified treetops (TTOPS, red dots), the false positive tree tops (i.e. no trees identified as trees; TTOPS_FP, light blue dots) and the false negative treetops (i.e. trees not identified; TTOPS_FN) are plotted on the canopy elevation model together with the segmented crown shapes (red polygons). An area of the trial (black square) has been enlarged to show the difference between crown shape before (c) and after (d) manual correction of the misidentified treetops.

Genotyping

A subset of 375 individuals belonging to 28 out of the 56 tested populations (Table S1; Fig. S2) were genotyped in a previous study (Pinosio *et al.*, 2014; Ruiz Daniels *et al.*, 2018). For this purpose, 1-yr-old needles were collected from the top third part of the crown in July 2013. After DNA extraction, the individuals were genotyped using two sets of molecular markers. The first set consisted of eight nuclear simple sequence repeat (SSR) loci, and was used to assess both population structure and relatedness

between individuals (i.e. kinship matrix). A detailed description of sample collection and SSR amplification is reported in Ruiz Daniels *et al.* (2018).

The second set comprised 294 SNPs included in loci derived from transcriptomes of *P. halepensis* evaluated in their responses to fire, as well as from resequenced loci first identified in loblolly pine (*Pinus taeda*) and identified from previous studies as candidates involved in adaptation of the latter species (Pinosio *et al.*, 2014). This panel of markers thus comprised candidate genes related to wood anatomy, growth, phenology and, also, to a wide

Table 1 Vegetation indices (VIs) considered in this study.

Index	Descriptor	Wavelengths	Formula	Reference
NDVI	Leaf area	Red, NIR	$(R_{840} - R_{670}) / (R_{840} + R_{670})$	Rouse <i>et al.</i> (1974)
OSAVI	Leaf area	Red, NIR	$(R_{840} - R_{670}) / (R_{840} + R_{670} + 0.16) \times 1.16$	Rondeaux <i>et al.</i> (1996)
RDVI	Leaf area	Red, NIR	$(R_{840} - R_{670}) / (R_{840} + R_{670})^{1/2}$	Roujean & Breon (1995)
EVI	Leaf area	Blue, red, NIR	$2.5 \times (R_{840} - R_{670}) / [(R_{840} + 6 \times R_{670} - 7.5 \times R_{450}) + 1]$	Huete <i>et al.</i> (2002)
MCARI	Leaf Chl content; leaf area	Green, red, NIR	$[(R_{700} - R_{670}) - 0.2 \times (R_{700} - R_{550})] \times (R_{700} / R_{670})$	Daughtry (2000)
TCARI	Leaf Chl content; leaf area	Green, red, NIR	$3 \times (R_{700} - R_{670}) - 0.2 \times (R_{700} - R_{550}) \times (R_{700} / R_{670})$	Haboudane <i>et al.</i> (2002)
TCARI/OSAVI	Leaf Chl content	Green, red, NIR	–	Haboudane <i>et al.</i> (2002)
ARI2	Anthocyanin content	Blue, NIR	$R_{840} \times (1/R_{550} - 1/R_{700})$	Gitelson <i>et al.</i> (2001)
CRI2	Carotenoid content	Blue, NIR	$1/R_{550} - 1/R_{700}$	Gitelson <i>et al.</i> (2002)
WBI	Water content	NIR	R_{900} / R_{950}	Peñuelas <i>et al.</i> (1993)

R indicates the reflectance in a single wavelength (in nm). NIR, near infra-red.

range of responses to abiotic stresses such as drought, cold and oxidative stress (Methods S4). This second set was used to identify genotype–phenotype associations.

Statistical analyses

Phenotypic data To validate the image-derived estimations of growth traits, H_{UAV} and crown area were compared by means of simple correlations to ground-based measurements of height and DBH obtained in 2013. To remove the effects of heterogeneous growing conditions on phenotypic records, tree-level estimates of H_{UAV} , crown area, vegetation indices and canopy temperature were subjected to mixed-effects linear models. ANOVAs consisted of column and replicate as fixed terms and column by replicate interaction, row nested to replicate, and row nested to replicate by column interaction as random terms. The tree-level residuals of this model were individuals' phenotypic data which retained only genotypic variation (i.e. they were largely free of environmental effects, which were accounted for by trial design). This dataset was subsequently used for genotype–phenotype association analyses (see the 'GWAS' section in Material and Methods), and was also used as input for a principal component analysis (PCA), to summarize the information retrieved by vegetation indices and to interpret tree-level relationships among traits. To this end, the loadings of H_{UAV} , crown area, vegetation indices and canopy temperature were plotted for the first two components of the PCA. Simple correlations were also calculated among phenotypic variables.

SSR data Individual genotypes at SSR loci were used to calculate relatedness between individuals. For this, a matrix of kinship between each pair of individuals was obtained from SSRs with the software SPAGEDi 1.3 (Hardy & Vekemans, 2015) using the kinship coefficient developed by Loiselle *et al.* (1995). In addition, the genetic structure of populations was inferred using the Bayesian clustering method implemented in STRUCTURE (Pritchard *et al.*, 2000). We ran STRUCTURE varying the numbers of possible genetic clusters (K) from one to 10, and each K was replicated 10 times. Each run consisted of 1×10^5 burn-in iterations and 1×10^6 data collection iterations. The different runs for the same K were then averaged using the software CLUMPP (Jakobsson & Rosenberg, 2007). The most likely K was selected calculating an empirical

statistic, ΔK , based on the second derivative of the likelihood of K (Evanno *et al.*, 2005). In addition, the genetic structure obtained through Bayesian clustering was evaluated with an independent approach. For this, we conducted a principal coordinates analysis (PCoA) on a matrix of pairwise genetic distances (G'_{ST}) between populations, and the first two components were plotted to summarize relationships between populations. Both the calculation of genetic distances and PCoA were carried out using the program GENALEX (Peakall & Smouse, 2006).

GWAS A GWAS was performed to test for the association between genotypes at single loci and each phenotypic trait (i.e. H_{UAV} , crown area, canopy temperature and vegetation indices). The residuals obtained through ANOVA (as phenotypes) were associated with genotypes at 235 SNPs out of the original 294. These 235 SNPs were those showing high-quality genotypes and a frequency of the minor allele > 0.05 . A total of 375 individuals, for which both genotypic and phenotypic data were available, were used for this analysis.

A mixed-effects linear model (MLM) was fitted independently for all pair combinations of SNPs and phenotypic traits following the procedure of Yu *et al.* (2006) implemented in TASSEL 5.0 (Bradbury *et al.*, 2007). The probability of membership of each individual to each genetic cluster detected from clustering analysis (see Results section) was included to avoid false positive associations related to large-scale genetic structure. Moreover, the kinship matrix was also included in the model to control for finest genetic structure (Yu *et al.*, 2006). After the MLMs were fitted, a correction for multiple testing was performed on the P -values using the false discovery rate method (Storey & Tibshirani, 2003) implemented in the R QVALUE package (Storey *et al.*, 2019).

SNP loci for which significant associations emerged with phenotypic traits were annotated from homology with other plant species as follows. First, the 200 bp sequence in which the SNPs were originally identified (Pinosio *et al.*, 2014) was used as query in the BLASTN tool. The best target sequence was subsequently used as query in BLASTX. The best match from BLASTX was retrieved to obtain the annotation of the SNPs. The result of BLASTN was also used to identify the effect of polymorphisms on proteins. For this, the 200 bp sequence was aligned with the sequence of the best match derived from BLASTN. The sequence

derived from this alignment was then translated to proteins based on the GenBank information. The software DNASP 6 (Rozas *et al.*, 2017) was used to check if the SNPs were located in a non-coding or coding region and, in the latter case, the resultant change in the amino acids sequence (synonymous/non-synonymous).

Multilocus association tests GWAS tests for significant contributions of a single locus to trait variation, but it does not provide an estimate for the total number of loci contributing to it. Alternatively, the number of SNPs explaining trait variation and the overall phenotypic variance explained by combinations of SNPs were estimated using the multilocus mixed model (MLMM) approach proposed by Segura *et al.* (2012). This approach consists of a stepwise mixed-model regression with forward inclusion of SNP markers. Before the stepwise analysis, a null model is calculated by incorporating all the possible SNPs, the kinship matrix and the genetic structure. The variance is then partitioned into genetic variance (i.e. explained by the kinship matrix, the population structure and the SNPs) and residual variance. At each step of the stepwise regression, the SNP with the lowest *P*-value is incorporated into the model, and the *P*-values of the other SNPs are recalculated, as well as the explained genetic and residual variance. The stepwise regression is terminated when the explained genetic variance reaches a value proximal to that of the null model (i.e. incorporating all the SNPs). We used the extended Bayesian information criterion (eBIC; Chen & Chen, 2008) to select the model that best fitted our data among the different steps of the MLMM. The SNPs included as cofactors and the percentage of variance explained (PVE) were recorded for the best-fitting model. An MLMM was performed for each phenotypic trait analysed in the GWAS.

Data availability

Individual genotypic data are available at the Dryad repository (<https://datadryad.org/stash/dataset/doi:10.5061/dryad.pt3b974>). Phenotypic data will be made available at a Mendeley repository.

Results

UAV-derived individual crown delineation

A total of 736 trees (92%) were successfully identified through automatic canopy segmentation, while 72 trees were not recognized by the segmentation algorithm and were manually identified. There were also 13 cases of false tree identification. Altogether, 806 georeferenced crown shapes (discarding dead trees) were obtained (Fig. 2).

Canopy segmentation and PCA analysis

Based on the detected treetops and tree crowns, individual records of H_{UAV} , crown area, vegetation indices and canopy temperature were obtained for the 806 individuals of the common

garden (Fig. S4). Mean H_{UAV} and crown area among trees were 5.80 ± 1.04 m and 6.52 ± 1.99 m² respectively (mean \pm SD). H_{UAV} correlated with ground-based height measurements at the tree level ($r=0.85$, $P<0.001$), indicating a reliable image-derived estimation of tree height. Crown area correlated with ground-measured DBH ($r=0.70$, $P<0.001$) (Fig. S5).

The PCA loadings summarized the relationships among vegetation indices, canopy temperature, H_{UAV} and crown area (Fig. 3). Simple correlations between traits are reported in Table S2. Indices related to leaf area (i.e. NDVI, RDVI and OSAVI) grouped together in the plot of loadings, and they were unrelated to TCARI/OSAVI (informative of Chl content). MCARI and TCARI (indicative of both leaf area and Chl content) and EVI (indicative of leaf area) were grouped in between the other leaf area indices (NDVI, OSAVI and RDVI) and TCARI/OSAVI (Fig. 3; Table S2). WBI (informative of water content), ARI2 (informative of anthocyanin content) and CRI2 (informative of carotenoid content) were less well explained by the first two PCA dimensions and were negatively associated with TCARI/OSAVI. Canopy temperature was negatively associated with H_{UAV} and crown area, which, in turn, were positively related to indices describing leaf area (Table S2). However, canopy temperature, H_{UAV} and crown area were poorly represented in the first two axes of the PCA (Fig. 3).




Fig. 3 Plot of the loadings of the first two PCA axes describing the relationships between unmanned aerial vehicle (UAV)-derived growth traits, vegetation indices and canopy temperature in the common garden of *Pinus halepensis*. NDVI, normalized difference vegetation index; OSAVI, optimized soil adjusted vegetation index; RDVI, re-normalized difference vegetation index; EVI, enhanced vegetation index; MCARI, modified chlorophyll absorption reflectance index; TCARI, transformed chlorophyll absorption ratio index; CRI2 carotenoid reflectance index 2; ARI2, anthocyanin reflectance index 2; WBI, water band index; T, canopy temperature; H_{UAV} , UAV-retrieved tree height.

Neutral genetic data

The Bayesian clustering performed on SSR genotypes indicated that the most probable number of clusters was two (Fig. S6). In this scenario, Western Mediterranean populations comprising individuals from the Iberian Peninsula and the Balearic islands were separated from Central–Eastern Mediterranean ones (i.e. Greek, Italian and Tunisian) (Fig. 4). One Tunisian population showed a high degree of admixture. Possible clustering also emerged at $K=4$ or $K=7$, but with a lower probability than the $K=2$ scenario (Fig. S6). In addition, considering four or seven genetic clusters did not result in clear geographical distinction between genetic groups. The scenario with two main genetic clusters was confirmed by PCoA performed on a matrix of population genetic distances (Fig. S7). The first two axes of the PCoA identified a clear clustering in the two above-mentioned groups. The scenario with two main genetic clusters also agreed with previously reported evidence (Ruiz Daniels *et al.*, 2018). Probabilities of individual assignment to the two clusters at $K=2$ were therefore used for correcting genotypic–phenotypic associations and to avoid false positives.

GWAS

GWAS revealed 12 significant associations between SNPs and growth traits (i.e. H_{UAV} and crown area) with $P < 10^{-3}$ (Table 2). These associations were significant also after correction for multiple testing ($q < 0.05$). H_{UAV} was associated with 10 SNPs, which individually explained between 5% and 10% of phenotypic variance. Two SNPs were associated with crown area, each explaining *c.* 5% of the phenotypic variance.

Sixteen significant associations emerged between SNPs and vegetation indices (Table 2). Among those, seven associations

occurred with indices related to leaf area, five with WBI (indicative of leaf water content), two with CRI2 (indicative of carotenoid content) and one with ARI2 (indicative of anthocyanin content). Most associations were also significant ($q < 0.05$) or marginally significant ($q < 0.1$) after correction for multiple testing, with the exception of the SNP associated with ARI2 ($q = 0.14$). These associations explained individually a small proportion of phenotypic variance (range 3.5–5.5%). No associations emerged with TCARI/OSAVI (indicative of Chl content). Finally, one SNP was associated with canopy temperature, explaining *c.* 4% of the phenotypic variance. However, this association was not significant after correction for multiple testing ($q = 0.11$).

Some SNPs were associated with more than one trait (Table 2), as follows: SNP 91 was associated with EVI (indicative of leaf area) and WBI (indicative of leaf water content), SNP 108 was associated with H_{UAV} , EVI and WBI, SNP 133 was associated with H_{UAV} and WBI, SNP 241 was associated with EVI and MCARI, and, finally, SNP 273 and SNP 350 were both associated with H_{UAV} and EVI.

The majority of SNPs for which significant associations emerged were annotated in known genes (Table 3; Table S3). For some of the annotated markers, it was possible to retrieve the molecular and biological functions of the homologous protein. In most cases, however, SNPs were located in noncoding regions or resulted in synonymous polymorphisms (Table S3).

Multilocus association test

The multilocus association test performed through MLM revealed that combinations of the 235 SNPs explained a relatively high proportion of the phenotypic variance of several traits (Table 4). Some, but not all, of the SNPs identified by the single-




Fig. 4 (a) Bar plot showing results of the assignment test with $K=2$ for the 375 individuals genotyped in the common garden of *Pinus halepensis*. Each individual is represented by a vertical line divided into two coloured segments representing the probabilities that the individual is assigned to the group comprising Western (in red) or Central-Eastern (in blue) Mediterranean populations. Population tags are reported in Supporting Information Table S1. (b) Pie charts showing the percentage of assignment to the two groups in each population. The charts are plotted on the actual geographical coordinates of populations.

Table 2 Results of the genome-wide association study (GWAS).

Trait	Indicator	Marker	<i>q</i>	PVE (%)
Leaf area	EVI	SNP108 ^A	0.01	4.26
	EVI	SNP241 ^A	0.01	3.75
	EVI	SNP273 ^B	0.01	4.64
	EVI	SNP350 ^A	0.01	4.43
	EVI	SNP91 ^A	0.01	4.60
	MCARI	SNP241 ^A	0.07	3.52
Photosynthetic pigments	TCARI	SNP151 ^A	0.07	4.54
	ARI2	SNP258 ^B	0.14	4.87
	CRI2	SNP67 ^A	0.05	5.00
	CRI2	SNP201 ^B	0.06	3.95
Water content	CRI2	SNP204 ^B	0.03	4.96
	WBI	SNP91 ^A	0.02	5.15
	WBI	SNP108 ^A	0.02	4.07
	WBI	SNP133 ^A	0.02	4.28
	WBI	SNP265 ^B	0.02	4.46
Canopy temperature	WBI	SNP350 ^A	0.02	4.82
	T	SNP140 ^B	0.11	4.34
Growth traits	Crown area	SNP2 ^B	0.05	4.94
	Crown area	SNP9 ^A	0.05	5.38
	<i>H</i> _{UAV}	SNP18 ^A	<0.001	7.39
	<i>H</i> _{UAV}	SNP108 ^A	<0.001	6.64
	<i>H</i> _{UAV}	SNP133 ^A	<0.001	6.43
	<i>H</i> _{UAV}	SNP159 ^A	<0.001	6.85
	<i>H</i> _{UAV}	SNP206 ^B	<0.001	5.70
	<i>H</i> _{UAV}	SNP217 ^A	<0.001	5.08
	<i>H</i> _{UAV}	SNP250 ^B	<0.001	5.50
	<i>H</i> _{UAV}	SNP273 ^B	<0.001	5.52
	<i>H</i> _{UAV}	SNP340 ^B	<0.001	5.70
	<i>H</i> _{UAV}	SNP350 ^A	<0.001	10.03

The single nucleotide polymorphisms (SNPs) associated with traits with $P < 10^{-3}$ are reported, together with the corrected *q*-value and the percentage of variance explained (PVE).

EVI, enhanced vegetation index; MCARI, modified chlorophyll absorption reflectance index; TCARI, transformed chlorophyll absorption reflectance index; ARI2, anthocyanin reflectance index 2; CRI2, carotenoid reflectance index 2; WBI, water band index; T, canopy temperature; *H*_{UAV}, tree height derived from drone imagery.

^AFrom resequenced genes of loblolly pine.

^BFrom *Pinus halepensis* transcriptome.

locus association test were present in the multilocus association models. Notably, a combination of 10 and nine SNPs explained 18% and 20% of the phenotypic variance of EVI (related to leaf area) and WBI (related to leaf water content) respectively. For *H*_{UAV}, 16% of the phenotypic variance was explained by a combination of only two SNPs, both identified also by GWAS. By contrast, the best MLM explained a low percentage of the phenotypic variance for several traits including NDVI, RDVI, OSAVI and those indices related to photosynthetic pigments. Finally, a combination of nine SNPs explained 16% of the phenotypic variance in canopy temperature.

Discussion

Canopy segmentation and recovery of phenotypic information

Our study is, to the best of our knowledge, the first to apply automatic crown identification in a common garden of a forest

species. This approach was highly effective in identifying single trees and providing tree-level estimations of growth-related traits. Although ground-based and UAV data were collected over different years, the intertree variability in height estimated through RGB-derived imagery agreed with ground-based measurements. Similarly, crown area was a good indicator of stem diameter, as reported elsewhere for other species (Lockhart *et al.*, 2005; Filipescu *et al.*, 2012), including *Pinus* spp. (Pretzsch *et al.*, 2015).

Developing cost- and time-effective phenotyping approaches is fundamental for characterizing the genetic basis underlying phenotypic differentiation (Lobos *et al.*, 2017). Our procedure retrieved growth information through a single flight and few hours of computation. Several studies have shown that the approach based on RGB-derived dense point clouds – obtained with inexpensive devices – can provide accurate estimations of growth traits in natural forests and fruit orchards, comparable to those obtained with more expensive technologies such as light detection and ranging (LiDAR) (Zarco-Tejada *et al.*, 2014; Wallace *et al.*, 2016; Weiss & Baret, 2017). In this regard, the continuous development and optimization of algorithms for automatic crown identification can facilitate the implementation of this methodology to forest genetic trials. Another advantage is that it allows the estimation of tree-level values of vegetation indices, which are surrogates of meaningful morphophysiological traits (Roberts *et al.*, 2016; Santini *et al.*, 2019b).

Traits such as leaf area or canopy transpiration can hardly be retrieved for a significant number of adult trees through ground-based phenotyping techniques. This issue prevented ground-based validation of vegetation indices and thermal data in this study. However, the indices that we considered have already been widely used and validated in the scientific literature (Roberts *et al.*, 2016), including variation at individual tree level (e.g. Berni *et al.*, 2009; Santini *et al.*, 2019b). By contrast, thermal data have been used at the tree level as proxies of tree transpiration and water status (e.g. González-Dugo *et al.*, 2013; Ludovisi *et al.*, 2017). Hence, the approach developed in this study could allow for easy routine analysis of barely investigated phenotypic traits.

In a previous study performed in the same common garden, vegetation indices and canopy temperature derived at the plot level revealed interpopulation differentiation in vegetation characteristics (Santini *et al.*, 2019a). The present study provided tree-level information which is suited for genotype–phenotype association studies and the assessment of relationships among phenotypic traits at the individual level. In this regard, the PCA indicated strong relationships between indices related to leaf area, canopy temperature and above-ground growth traits. Trees having high leaf area and reduced canopy temperature (indicative of high transpiration; González-Dugo *et al.*, 2013) showed high growth, indicated by large height and crown area. These results confirm the strong dependence of growth to variation in photosynthetic surface and total transpiration in *P. halepensis* under drought conditions (Voltas *et al.*, 2008; Santini *et al.*, 2019a). By contrast, variation in leaf biochemistry seemed not to affect tree growth to the same degree (Santini *et al.*, 2019a).

Table 3 Annotation of the single nucleotide polymorphisms (SNPs) detected in genome-wide association study (GWAS).

SNP ID	Accession	E-values BLASTX	Annotation	Species	Molecular function	Biological function
SNP9	XP_006826240.1	2E-10	Heat shock factor protein HSF24	<i>Amborella trichopoda</i>	DNA-binding transcription factor activity; RNA polymerase II <i>cis</i> -regulatory region; sequence-specific DNA	Cellular response to heat; regulation of transcription from RNA polymerase II promoter in response to stress
SNP18	ACJ09662.1	9E-45	Putative calcium-dependent protein kinase	<i>Cupressus sempervirens</i>	Kinase; transferase	–
SNP67	XP_031397970.1	1E-14	Leaf rust 10 disease-resistance locus receptor	<i>Punica granatum</i>	ATP binding; polysaccharide binding; protein serine/threonine kinase activity	–
SNP91	ABF73316.1	4E-14	Clavata-like receptor	<i>Picea glauca</i>	ATP binding; protein kinase activity	–
SNP108	TKS08810.1	2E-41	DNAJ heat shock N-terminal domain-containing family protein	<i>Populus alba</i>	–	–
SNP133	ABG34278.1	6E-58	Polygalacturonase	<i>Eucalyptus globulus</i>	Polygalacturonase activity	Carbohydrate metabolic process; cell wall organization
SNP151	XP_002320762.1	3E-13	Peroxisomal membrane protein 11D	<i>Populus trichocarpa</i>	Identical protein binding	Peroxisome fission; peroxisome organization; regulation of eproxisome size
SNP159	ACJ09662.1	9E-45	Putative calcium-dependent protein kinase, partial	<i>Cupressus sempervirens</i>	Kinase; transferase	–
SNP206	ABR15469.1	0.00	GDP-mannose pyrophosphorylase	<i>Pinus taeda</i>	Nucleotidyltransferase activity	Biosynthetic processes
SNP241	ATP71577.1	0.004	Hypothetical protein	<i>Pinus pinaster</i>	–	–
SNP258	AJP06341.1	0.00	PIN2	<i>Pinus tabuliformis</i>	–	Auxin-activated signalling pathway; transmembrane transport
SNP273	AAQ63936.1	8E-15	Cellulose synthase, partial	<i>Pinus radiata</i>	Cellulose synthase (UDP-forming); metal ion binding	–
SNP340	ABR15469.1	0.00	GDP-mannose pyrophosphorylase	<i>Pinus taeda</i>	Nucleotidyltransferase activity	Biosynthetic process
SNP350	AEX11975.1	2E-78	Hypothetical protein	<i>Pinus taeda</i>	–	–

Association genetics of growth traits

Despite the low number of markers considered, the results of this study showed the effectiveness of combining genotypic information with UAV imagery to characterize the genomic basis of phenotypic differentiation in forest species. Indeed, we identified relevant genotypic–phenotypic associations for several traits related to growth and vegetation indices. Such associations provide first insight into adaptive genomic variation in *P. halepensis*, although an intrinsic limitation of our approach is the lack of strong evidence on how variation at the cellular and tissue levels is coordinated with changes at the whole-organism level. Regardless, the detected loci deserve further attention in relation to their potential adaptive role for this and other closely related pines.

GWAS identified 12 SNPs significantly associated with UAV-derived crown area and tree height, explaining a proportion of phenotypic variance of 5–10%. Because linkage disequilibrium decays rapidly in conifers (De La Torre *et al.*, 2014; Plomion *et al.*, 2016), these SNPs are probably quantitative trait nucleotides (i.e. single polymorphisms influencing the phenotype) or are located in close proximity to the causative polymorphisms. Our results indicate that these SNPs do not produce protein change or alteration, being probably involved in changes of gene expression. Most of the relevant SNPs were annotated with known proteins, even though a correct annotation was not feasible in some cases. This may be partially due to the relatively scarce information on conifer genomes that is available to date, which emphasizes the need of further studies to characterize tree genomes (De La Torre *et al.*, 2014).

Table 4 Results of the multilocus mixed model (MLMM) analysis.

Trait	Indicator	SNPs as cofactors in the best fitting model	PVE (%)
Leaf area	NDVI	183	2.13
	OSAVI	183	3.05
	RDVI	108, 183	3.48
	EVI	16, 38, 106, 108, 222, 256, 264, 269, 326, 350	18.34
	MCARI	133, 222	5.17
	TCARI	38, 140, 222, 312, 350, 368	10.04
Photosynthetic pigments	TCARI/OSAVI	315	2.24
	ARI2	261	2.83
	CRI2	8	4.09
Water content	WBI	140, 156, 205, 216, 304, 312, 319, 343, 350	19.83
Canopy temperature	T	128, 140, 180, 215, 248, 258, 275, 319, 375	16.51
Growth traits	Crown area	21, 49	8.48
	H_{UAV}	18, 350	15.74

For each trait, the single nucleotide polymorphisms (SNPs) included as cofactor in the best fitting model are reported, as well as the percentage of variance explained (PVE) explained by the combination of SNPs.

NDVI, normalized difference vegetation index; OSAVI, optimized soil adjusted vegetation index; RDVI, renormalized difference vegetation index; EVI, enhanced vegetation index; MCARI, modified chlorophyll absorption reflectance index; TCARI, transformed chlorophyll absorption reflectance index; ARI2, anthocyanin reflectance index 2; CRI2, carotenoid reflectance index 2; WBI, water band index; T, canopy temperature; H_{UAV} , tree height derived from drone imagery.

Growth-associated SNPs that could be annotated included genomic regions encoding for a disparate number of proteins. In particular, differences in crown area were associated with a gene marker (SNP 9) coding for a heat stress transcription factor, which is part of a family of proteins involved in response to heat stress (Scharf *et al.*, 2012). In the case of tree height, significant associations were found with genes encoding for: calcium-dependent kinase (SNP 18 and SNP 159), which influences plant responses to endogenous and environmental cues (Romeis, 2001); GDP-mannose pyrophosphorylase (SNP 340) involved in ascorbic acid metabolism (Keller *et al.*, 1999); cellulose synthase (SNP 273), which influences cell wall synthesis and organization (Richmond & Somerville, 2000); and polygalacturonase (SNP 133 and SNP 206) involved in tissue development and stomatal dynamics (Rui *et al.*, 2017).

These results suggest that the genetic control of complex traits such as growth is associated with a large number of interacting genes with very diverse functions. However, the multilocus association analysis revealed that a combination of only two SNPs explained most phenotypic variation (16%) in tree height. This outcome could be partially due to the inability of the MLMM to identify loci with small effects in the presence of loci with large effects on the phenotype. By contrast, the result of the MLMM may also indicate that the effects of a single locus of tree height identified by GWAS are largely overlapping. Finally, it must be noted that differences in the number of SNPs detected through each approach could originate, at least in part, from different statistical treatments of missing data.

Association genetics of vegetation properties

GWAS also revealed several SNPs significantly associated with phenotypic variation in vegetation indices. These associations were inconsistent across indices related to leaf area, however.

Some indices (i.e. NDVI, RDVI and OSAVI) are known to saturate at high leaf areas, being potentially unable to intercept fine interindividual differences in this trait (Roberts *et al.*, 2016). By contrast, several marker-trait associations emerged with EVI, which is less sensitive to saturation. Two of the identified SNPs related to EVI were annotated to known proteins. SNP 91 codes for a clavata-like receptor kinase, which is involved in tissue development and meristem differentiation (Clark, 2001). SNP 273 is found in a gene coding for a cellulose synthase, which is part of a family of genes with a crucial role in cell wall synthesis and organization, in turn influencing tissue growth and development (Richmond & Somerville, 2000). Interestingly, proteins of the cellulose synthase family are associated with the control of leaf size in *Arabidopsis* (Horiguchi *et al.*, 2006) and maize (Li *et al.*, 2018). Another gene marker (SNP 151) was associated with TCARI, an index partly related to leaf area. SNP 151 is found in a gene coding for PEX11C protein involved in peroxisome metabolism. Among other functions, proteins of the PEX family have been shown to regulate photomorphogenesis through a light-mediated pattern of induction of peroxisome proliferation, with potential implications on leaf development (Hu *et al.*, 2002; Kaur *et al.*, 2013). Interestingly, this SNP was reported to be under selection in *P. halepensis* and associated with variation in summer precipitation at the origin of natural populations (Ruiz Daniels *et al.*, 2018). In other Mediterranean pines (*P. pinaster* and *P. pinea*), the expression of a homologous gene was induced by drought stress (Perdiguero *et al.*, 2013).

Although TCARI is also sensitive to changes in leaf Chl content, no significant marker-trait associations emerged with the index TCARI/OSAVI, which is specifically related to Chl content (Zarco-Tejada *et al.*, 2004). This may be due to weak genetic differentiation in photosynthetic capacity of *P. halepensis* (Santini *et al.*, 2019a). By contrast, four SNPs were associated with carotenoid or anthocyanin content, and two of them were

annotated with homologous proteins described in other species. In particular, SNP 258 is located in a gene coding for a PIN2 protein involved in auxin transport and auxin-mediated signalling, while the protein associated with SNP 67 is involved in pathogen resistance. Anthocyanins and carotenoids are involved in a myriad of plant processes related to stress response (Havaux, 2014; Van der Ende & El-Esawe, 2014). Interestingly, some studies have reported that carotenoid accumulation is, in some cases, mediated by auxin balance (Du *et al.*, 2012; Su *et al.*, 2015). In this regard, SNP 258 was also identified in a recent study as a candidate gene involved in adaptation to local climate (Ruiz Daniels *et al.*, 2019).

The WBI index, indicative of leaf water content, was associated with five SNPs. Two of them are located in genes of known functions. SNP 133 is found in a gene encoding the polygalacturonase isoform X3 protein, which has been widely studied in the model plant *Arabidopsis*. This protein influences tissue development and is involved in stomatal dynamics of mature leaves, which might explain the association found with leaf water content (Rui *et al.*, 2017). SNP 91 codes for a clavata-like receptor kinase and it was associated also with leaf area indices (Clark, 2001).

However, the identified SNPs had small effects on phenotypes, explaining individually less than 6% of the variation in vegetation indices or canopy temperature. Genetic variation in single markers usually accounts for a small proportion of phenotypic variance in forest species (Eckert *et al.*, 2009; Prunier *et al.*, 2013; Baison *et al.*, 2019). Indeed, many adaptive traits are probably under polygenic control, suggesting that many loci with minor effect might modify their expression (Savolainen *et al.*, 2007; White *et al.*, 2007). Moreover, the single contribution of many other genes to complex traits such as leaf architecture or water use may be too small to be detected by GWAS (Resende *et al.*, 2017). In this regard, the multilocus association test showed that combinations of several markers are able to explain a relatively large proportion of the phenotypic variation in leaf area and transpiration. This result suggests that association analyses carried out considering a larger number of markers simultaneously can capture a sizeable part of the variation of complex traits in forest species (Savolainen *et al.*, 2007; Grattapaglia *et al.*, 2009).

Even considering a polygenic model, however, a large part of the phenotypic variation both in vegetation indices and in growth traits remained unexplained. We speculate that this may be related to inaccuracies in UAV-derived records and, also, to low marker coverage, which stresses the importance of allowing for the largest possible number of markers across the genome (Resende *et al.*, 2017). Nevertheless, it is worth noting that a high phenotypic variance is seldom explained by genetic variation even in organisms for which dense, genome-wide sets of markers are available (Manolio *et al.*, 2009; Bourrat *et al.*, 2017). This issue, known as 'missing heritability', suggests that phenotypic variation may be broadly influenced by largely unknown underlying factors such as epistatic interactions or epigenetic modifications of gene expression (Manolio *et al.*, 2009; Trerotola *et al.*, 2015; Bourrat *et al.*, 2017).

Pleiotropy

Some SNPs showed significant associations with multiple correlated traits, a possible indication of pleiotropic effects with genes influencing different traits simultaneously (Wu *et al.*, 2000). However, these SNPs probably concur to determine only one trait, showing multiple associations because of correlated traits. For example, three SNPs (108, 273 and 350) were associated with both tree height and some indices related to leaf area, traits that were associated among individuals. By contrast, SNPs 133 and 350 were associated with both tree height and WBI, traits that were weakly correlated among individuals. Such associations involving more than one trait could be indicative of true pleiotropy, with gene markers influencing both leaf water content and tree growth concurrently.

Conclusions

In this study, we developed a workflow for (semi-)automatic phenotyping of trees growing in a common garden. In the era of genomics, retrieving meaningful phenotypic information with time- and cost-effective tools has the potential to boost the characterization of the genetic basis of fundamental evolutionary processes involved in phenotypic differentiation of nonmodel organisms (Großkinsky *et al.*, 2015). Although the scope of our inferences could be undoubtedly broadened by using a significantly larger number of markers and individuals, this study highlights the type of information that can be obtained through high-throughput phenotyping approaches. Indeed, our results provide insight into the molecular processes controlling phenotypic differentiation in a widespread conifer, underlining the potential of widely available new technologies to fill the gap between genetic variability and individual phenotypes (Houle *et al.*, 2010).

Acknowledgements


This work was supported by the Spanish Government, grant number RTI2018-094691-B-C31 (MCIU/AEI/FEDER, EU). We thank Ricardo Alía (CIFOR-INIA) for providing the genotypic data. We thank three anonymous reviewers whose insightful comments and suggestions helped to improve the manuscript. The authors declare no conflicts of interest.


Author contributions


JV and FS conceived and designed the research. FS, SCK, VRdD, DG and JV collected the data; FS and SMG analysed the data; FS and JV wrote the manuscript, with contributions from SCK, JLA, VRdD, SMG and DG.

ORCID

José Luis Araus  <https://orcid.org/0000-0002-8866-2388>
Delphine Grivet  <https://orcid.org/0000-0001-8168-4456>
Shawn Carlisle Kefauver  <https://orcid.org/0000-0002-1687-1965>

Víctor Resco de Dios  <https://orcid.org/0000-0002-5721-1656>

Filippo Santini  <https://orcid.org/0000-0002-5681-6039>

Jordi Voltas  <https://orcid.org/0000-0003-4051-1158>

References

- Baison J, Vidalis A, Zhou L, Chen Z, Li Z, Sillanpää MJ, Bernhardsson C, Scofield D, Forsberg N, Grahn T *et al.* 2019. Genome-wide association study identified novel candidate loci affecting wood formation in Norway spruce. *The Plant Journal* **100**: 83–100.
- Berni J, Zarco-Tejada PJ, Suarez L, Fereres E. 2009. Thermal and narrowband multispectral remote sensing for vegetation monitoring from an unmanned aerial vehicle. *IEEE Transactions on Geoscience and Remote Sensing* **47**: 722–738.
- Bourrat P, Lu Q, Jablonka E. 2017. Why the missing heritability might not be in the DNA. *BioEssays* **39**: 1700067.
- Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics* **23**: 2633–2635.
- Chen J, Chen Z. 2008. Extended Bayesian information criteria for model selection with large model spaces. *Biometrika* **95**: 759–771.
- Clark SE. 2001. Cell signalling at the shoot meristem. *Nature Reviews Molecular Cell Biology* **2**: 276–284.
- Climent J, Prada MA, Calama R, Chambel MR, de Ron DS, Alía R. 2008. To grow or to seed: ecotypic variation in reproductive allocation and cone production by young female Aleppo pine (*Pinus halepensis*, Pinaceae). *American Journal of Botany* **95**: 833–842.
- Daughtry C. 2000. Estimating corn leaf chlorophyll concentration from leaf and canopy reflectance. *Remote Sensing of Environment* **74**: 229–239.
- De La Torre AR, Birol I, Bousquet J, Ingvarsson PK, Jansson S, Jones SJM, Keeling CI, MacKay J, Nilsson O, Ritland K *et al.* 2014. Insights into conifer giga-genomes. *Plant Physiology* **166**: 1724–1732.
- Du H, Wu N, Fu J, Wang S, Li X, Xiao J, Xiong L. 2012. A GH3 family member, OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and cold tolerance in rice. *Journal of Experimental Botany* **63**: 6467–6480.
- Eckert AJ, Bower AD, Wegrzyn JL, Pande B, Jermstad KD, Krutovsky KV, St. Clair JB, Neale DB. 2009. Association genetics of coastal Douglas fir (*Pseudotsuga menziesii* var. *menziesii*, Pinaceae). I. Cold-hardiness related traits. *Genetics* **182**: 1289–1302.
- Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using the software structure: a simulation study. *Molecular Ecology* **14**: 2611–2620.
- Filipescu CN, Groot A, MacIsaac DA, Cruickshank MG, Stewart JD. 2012. Prediction of diameter using height and crown attributes: a case study. *Western Journal of Applied Forestry* **27**: 30–35.
- Gitelson AA, Merzlyak MN, Chivkunova OB. 2001. Optical properties and nondestructive estimation of anthocyanin content in plant leaves. *Photochemistry and Photobiology* **74**: 38–45.
- Gitelson AA, Zur Y, Chivkunova OB, Merzlyak MN. 2002. Assessing carotenoid content in plant leaves with reflectance spectroscopy. *Photochemistry and Photobiology* **75**: 272–281.
- González-Dugo V, Zarco-Tejada P, Nicolás E, Nortés PA, Alarcón JJ, Intrigliolo DS, Fereres E. 2013. Using high resolution UAV thermal imagery to assess the variability in the water status of five fruit tree species within a commercial orchard. *Precision Agriculture* **14**: 660–678.
- Grattapaglia D, Plomion C, Kirst M, Sederoff RR. 2009. Genomics of growth traits in forest trees. *Current Opinion in Plant Biology* **12**: 148–156.
- Grattapaglia D, Resende MDV. 2011. Genomic selection in forest tree breeding. *Tree Genetics & Genomes* **7**: 241–255.
- Großkinsky DK, Svendsgaard J, Christensen S, Roitsch T. 2015. Plant phenomics and the need for physiological phenotyping across scales to narrow the genotype-to-phenotype knowledge gap. *Journal of Experimental Botany* **66**: 5429–5440.
- Haboudane D, Miller JR, Tremblay N, Zarco-Tejada PJ, Dextraze L. 2002. Integrated narrow-band vegetation indices for prediction of crop chlorophyll content for application to precision agriculture. *Remote Sensing of Environment* **81**: 416–426.
- Hardy O, Vekemans X. 2015. *SPAGeDi 1.5. a program for spatial pattern analysis of genetic diversity*. Brussels, Belgium: Université Libre de Bruxelles.
- Havaux M. 2014. Carotenoid oxidation products as stress signals in plants. *The Plant Journal* **79**: 597–606.
- Horiguchi G, Ferjani A, Fujikura U, Tsukaya H. 2006. Coordination of cell proliferation and cell expansion in the control of leaf size in *Arabidopsis thaliana*. *Journal of Plant Research* **119**: 37–42.
- Houle D, Govindaraju DR, Omholt S. 2010. Phenomics: the next challenge. *Nature Reviews Genetics* **11**: 855–866.
- Hu J, Aguirre M, Peto C, Alonso J, Ecker J, Chory J. 2002. A role for peroxisomes in photomorphogenesis and development of *Arabidopsis*. *Science* **297**: 405–409.
- Huete A, Didan K, Miura T, Rodriguez EP, Gao X, Ferreira LG. 2002. Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment* **83**: 195–213.
- Jakobsson M, Rosenberg NA. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**: 1801–1806.
- Jaramillo-Correa JP, Prunier J, Vázquez-Lobo A, Keller SR, Moreno-Letelier A. 2015. Molecular signatures of adaptation and selection in forest trees. *Advances in Botanical Research* **74**: 265–306.
- Kaur N, Li J, Hu J. 2013. Peroxisomes and photomorphogenesis. In: Luis A, Rio D, eds. *Peroxisomes and their key role in cellular signaling and metabolism*. Dordrecht, the Netherlands: Springer, 195–211.
- Keller R, Renz FSA, Kossmann J. 1999. Antisense inhibition of the GDP-mannose pyrophosphorylase reduces the ascorbate content in transgenic plants leading to developmental changes during senescence. *The Plant Journal* **19**: 131–141.
- Khan MA, Korban SS. 2012. Association mapping in forest trees and fruit crops. *Journal of Experimental Botany* **63**: 4045–4060.
- Klein T, Di Matteo G, Rotenberg E, Cohen S, Yakir D. 2013. Differential ecophysiological response of a major Mediterranean pine species across a climatic gradient. *Tree Physiology* **33**: 26–36.
- Li W, Yang Z, Yao J, Li J, Song W, Yang X. 2018. Cellulose synthase-like D1 controls organ size in maize. *BMC Plant Biology* **18**: 239.
- Lobos GA, Camargo AV, del Pozo A, Araus JL, Ortiz R, Doonan JH. 2017. Editorial: plant phenotyping and phenomics for plant breeding. *Frontiers in Plant Science* **8**: 2181.
- Lockhart BR, Weih RC, Smith KM. 2005. Crown radius and diameter at breast height relationships for six bottomland hardwood species. *Journal of the Arkansas Academy of Science* **59**: 110–115.
- Loiselle BA, Sork VL, Nason J, Graham C. 1995. Spatial genetic structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of Botany* **82**: 1420–1425.
- Ludovisi R, Tauro F, Salvati R, Khoury S, Mugnozza Scarascia G, Harfouche A. 2017. UAV-based thermal imaging for high-throughput field phenotyping of black poplar response to drought. *Frontiers in Plant Science* **8**: 1681.
- Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorf LA, Hunter DJ *et al.* 2009. Finding the missing heritability of complex diseases. *Nature* **461**: 747–753.
- McGaughey RJ. 2012. *FUSION/LDV: Software for LiDAR data analysis and visualization—V3. 10*. Portland, OR, USA: USDA Forest Service, Pacific Northwest Research Station.
- McKown AD, Klápště J, Guy RD, Gerales A, Porth I, Hannemann J, Friedmann M, Muchero W, Tuskan GA, Ehlting J *et al.* 2014. Genome-wide association implicates numerous genes underlying ecological trait variation in natural populations of *Populus trichocarpa*. *New Phytologist* **203**: 535–553.
- Meyer F, Beucher S. 1990. Morphological segmentation. *Journal of Visual Communication and Image Representation* **1**: 21–46.
- Nevalainen O, Honkavaara E, Tuominen S, Viljanen N, Hakala T, Yu X, Hyyppä J, Saari H, Polonen I, Imai NN *et al.* 2017. Individual tree detection and classification with UAV-based photogrammetric point clouds and hyperspectral imaging. *Remote Sensing* **9**: 185.

- Peakall R, Smouse PE. 2006. Genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
- Peñuelas J, Filella I, Biel C, Serrano L, Save R. 1993. The reflectance at the 950–970 nm region as an indicator of plant water status. *International Journal of Remote Sensing* 14: 1887–1905.
- Perdiguerio P, del Carmen Barbero M, Cervera MT, Collada C. 2013. Molecular response to water stress in two contrasting Mediterranean pines (*Pinus pinaster* and *Pinus pinea*). *Plant Physiology and Biochemistry* 67: 199–208.
- Pinosio S, González-Martínez SC, Bagnoli F, Cattonaro F, Grivet D, Marroni F, Lorenzo Z, Pausas JG, Verdú M, Vendramin GG. 2014. First insights into the transcriptome and development of new genomic tools of a widespread circum-Mediterranean tree species, *Pinus halepensis* Mill. *Molecular Ecology Resources* 14: 846–856.
- Plomion C, Bartholomé J, Lesur I, Boury C, Rodríguez-Quilón I, Lagraret H, Ehrenmann F, Bouffier L, Gion JM, Grivet D *et al.* 2016. High-density SNP assay development for genetic analysis in maritime pine (*Pinus pinaster*). *Molecular Ecology Resources* 16: 574–587.
- Plowright A. 2018. *Forestools: analyzing remotely sensed forest data*. R package v.0.2.0 [WWW document] URL <https://cran.r-project.org/web/packages/ForestTools/index.html>.
- Popescu SC, Wynne RH. 2004. Seeing the trees in the forest. *Photogrammetric Engineering & Remote Sensing* 70: 589–604.
- Pretzsch H, Biber P, Uhl E, Dahlhausen J, Rötzer T, Caldentey J *et al.* 2015. Crown size and growing space requirement of common tree species in urban centres, parks, and forests. *Urban Forestry & Urban Greening* 14: 466–479.
- Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155: 945–959.
- Prunier J, Pelgas B, Gagnon F, Despots M, Isabel N, Beaulieu J, Bousquet J. 2013. The genomic architecture and association genetics of adaptive characters using a candidate SNP approach in boreal black spruce. *BMC Genomics* 14: 368.
- QGIS Development Team. 2019. *QGIS geographic information system*. Open Source Geospatial Foundation Project. [WWW document] URL <http://qgis.osgeo.org/> [accessed 20 January 2020].
- R Core Team. 2019. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing [WWW document] URL <https://www.R-project.org/> [accessed 20 January 2020].
- Resende RT, Resende MDV, Silva FF, Azevedo CF, Takahashi EK, Silva-Junior OB, Grattapaglia D. 2017. Regional heritability mapping and genome-wide association identify loci for complex growth, wood and disease resistance traits in *Eucalyptus*. *New Phytologist* 213: 1287–1300.
- Richmond TA, Somerville CR. 2000. The cellulose synthase superfamily. *Plant Physiology* 124: 495–498.
- Roberts DA, Roth KL, Perroy RL. 2016. Hyperspectral vegetation indices. In: Huete A, Lyon JG, Thenkabail PS, eds. *Hyperspectral remote sensing of vegetation*. Boca Raton, FL, USA: CRC Press, 309–328.
- Rodríguez Quilón I. 2017. *Ecological and association genetics in two Mediterranean pine species*. PhD thesis, Universidad Politécnica de Madrid, Madrid, Spain.
- Romeis T. 2001. Calcium-dependent protein kinases play an essential role in a plant defence response. *EMBO Journal* 20: 5556–5567.
- Rondeaux G, Steven M, Baret F. 1996. Optimization of soil-adjusted vegetation indices. *Remote Sensing of Environment* 55: 95–107.
- Roujean J-L, Breon F-M. 1995. Estimating PAR absorbed by vegetation from bidirectional reflectance measurements. *Remote Sensing of Environment* 51: 375–384.
- Rouse J, Haas RH, Schell JA, Deering DW. 1974. Monitoring vegetation systems in the Great Plains with ERTS. *NASA Special Publication* 351: 309.
- Roussel JR, Auty D. 2018. *lidR: airborne LiDAR data manipulation and visualization for forestry applications*. R package v.1.0.0 [WWW document] URL <https://rdr.io/cran/lidR/>.
- Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large datasets. *Molecular Biology and Evolution* 34: 3299–3302.
- Rui Y, Xiao C, Yi H, Kandemir B, Wang JZ, Puri VM, Anderson CT. 2017. Polygalacturonase involved in expansion3 functions in seedling development, rosette growth, and stomatal dynamics in *Arabidopsis thaliana*. *The Plant Cell* 29: 2413–2432.
- Ruiz Daniels R, Taylor RS, González-Martínez SC, Vendramin GG, Fady B, Oddou-Muratorio S, Piotti A, Simioni G, Grivet D, Beaumont MA. 2019. Looking for local adaptation: convergent microevolution in Aleppo Pine (*Pinus halepensis*). *Genes* 10: 673.
- Ruiz Daniels R, Taylor RS, Serra-Varela MJ, Vendramin GG, González-Martínez SC, Grivet D. 2018. Inferring selection in instances of long-range colonization: the Aleppo pine (*Pinus halepensis*) in the Mediterranean Basin. *Molecular Ecology* 27: 3331–3345.
- Santini F, Kefauver SC, Resco de Dios V, Araus JL, Voltas J. 2019a. Using unmanned aerial vehicle-based multispectral, RGB and thermal imagery for phenotyping of forest genetic trials: a case study in *Pinus halepensis*. *Annals of Applied Biology* 174: 262–276.
- Santini F, Serrano L, Kefauver SC, Abdullah-Al M, Aguilera M, Sin E, Voltas J. 2019b. Morpho-physiological variability of *Pinus nigra* populations reveals climate-driven local adaptation but weak water use differentiation. *Environmental and Experimental Botany* 166: 103828.
- Savolainen O, Pyhäjärvi T, Knürr T. 2007. Gene flow and local adaptation in trees. *Annual Review of Ecology, Evolution and Systematics* 38: 595–619.
- Scharf KD, Berberich T, Ebersberger I, Nover L. 2012. The plant heat stress transcription factor (Hsf) family: structure, function and evolution. *Biochimica et Biophysica Acta (BBA) – Gene Regulatory Mechanisms* 1819: 104–119.
- Schiller G, Atzmon N. 2009. Performance of Aleppo pine (*Pinus halepensis*) provenances grown at the edge of the Negev desert: a review. *Journal of Arid Environments* 73: 1051–1057.
- Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, Nordborg M. 2012. An efficient multi-locus mixed-model approach for genome-wide association studies in structured populations. *Nature Genetics* 44: 825.
- Serra-Varela MJ, Aliá R, Daniels RR, Zimmermann NE, Gonzalo-Jiménez J, Grivet D. 2017. Assessing vulnerability of two Mediterranean conifers to support genetic conservation management in the face of climate change. *Diversity and Distributions* 23: 507–516.
- Storey JD, Bass AJ, Dabney A, Robinson D. 2019. *qvalue: Q-value estimation for false discovery rate control*. R package v.2.16.0 [WWW document] URL <https://rpubs.com/jmr1990/eiccqvalue>.
- Storey JD, Tibshirani R. 2003. Statistical significance for genomewide studies. *Proceedings of the National Academy of Sciences, USA* 100: 9440–9445.
- Su L, Diletto G, Purgatto E, Danoun S, Zouine M, Li Z, Roustan JP, Bouzayen M, Giuliano G, Chervin C. 2015. Carotenoid accumulation during tomato fruit ripening is modulated by the auxin–ethylene balance. *BMC Plant Biology* 15: 114.
- Tognetti R, Michelozzi M, Giovannelli A. 1997. Geographical variation in water relations, hydraulic architecture and terpene composition of Aleppo pine seedlings from Italian provinces. *Tree Physiology* 17: 241–250.
- Trerotola M, Relli V, Simeone P, Alberti S. 2015. Epigenetic inheritance and the missing heritability. *Human Genetics* 9: 17.
- Van den Ende W, El-ESawe SK. 2014. Sucrose signaling pathways leading to fructan and anthocyanin accumulation: a dual function in abiotic and biotic stress responses? *Environmental and Experimental Botany* 108: 4–13.
- de Villemereuil P, Gaggiotti OE, Mouterde M, Till-Bottraud I. 2016. Common garden experiments in the genomic era: new perspectives and opportunities. *Heredity* 116: 249–254.
- Voltas J, Chambel MR, Prada MA, Ferrio JP. 2008. Climate-related variability in carbon and oxygen stable isotopes among populations of Aleppo pine grown in common-garden tests. *Trees* 22: 759–769.
- Voltas J, Lucabaugh D, Chambel MR, Ferrio JP. 2015. Intraspecific variation in the use of water sources by the circum-Mediterranean conifer *Pinus halepensis*. *New Phytologist* 208: 1031–1041.
- Voltas J, Shestakova TA, Patsiou T, di Matteo G, Klein T. 2018. Ecotypic variation and stability in growth performance of the thermophilic conifer *Pinus halepensis* across the Mediterranean basin. *Forest Ecology and Management* 424: 205–215.
- Wallace L, Lucieer A, Malenovsky Z, Turner D, Vopěnka P. 2016. Assessment of forest structure using two uav techniques: a comparison of airborne laser scanning and structure from motion (SfM) point clouds. *Forests* 7: 62.

- Weiss M, Baret F. 2017. Using 3D point clouds derived from UAV RGB imagery to describe vineyard 3D macro-structure. *Remote Sensing* **9**: 111.
- White TL, Adams WT, Neale DB. 2007. *Forest genetics*. Cambridge, MA, USA: CABI.
- Wu R, Zeng ZB, McKeand SE, OMalley DM. 2000. The case for molecular mapping in forest tree breeding. *Plant Breeding Reviews* **19**: 41–68.
- Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB *et al.* 2006. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nature Genetics* **38**: 203–208.
- Zarco-Tejada PJ, Diaz-Varela R, Angileri V, Loudjani P. 2014. Tree height quantification using very high resolution imagery acquired from an unmanned aerial vehicle (UAV) and automatic 3D photo-reconstruction methods. *European Journal of Agronomy* **55**: 89–99.
- Zarco-Tejada PJ, Miller JR, Morales A, Berjón A, Agüera J. 2004. Hyperspectral indices and model simulation for chlorophyll estimation in open-canopy tree crops. *Remote Sensing of Environment* **90**: 463–476.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 Aerial view of the common garden in July 2016.

Fig. S2 Geographical locations of the populations tested in the common garden.

Fig. S3 Schematized process for estimation of tree height and crown area, and for obtaining crown multispectral and thermal images.

Fig. S4 Boxplots of intertree variation in growth data, canopy temperature and vegetation indices.

Fig. S5 Correlations between ground measured and UAV-derived growth data.

Fig. S6 ΔK results from STRUCTURE analysis.

Fig. S7 Plot of the first and second coordinates of PCoA performed on a matrix of genetic distances between populations.

Methods S1 Multispectral camera calibration and validation of reflectance data.

Methods S2 Retrieval and validation of data retrieval.

Methods S3 Description of vegetation indices considered in the study.

Methods S4 Description of the SNPs markers considered in the study.

Table S1 Geographical origin of the 56 *Pinus halepensis* populations tested in this study.

Table S2 Correlation matrix between phenotypic traits.

Table S3 First step of the SNP annotation.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.