

Geomorphological and sedimentary processes of the glacially influenced northwestern Iberian continental margin and abyssal plains

Estefanía Llave^{a,*}, Gloria Jané^a, Adolfo Maestro^{a,b}, Jerónimo López-Martínez^b, F. Javier Hernández-Molina^c, Sandra Mink^a

^a Dept. Investigación y Prospectiva Geocientífica, Instituto Geológico y Minero de España, IGME, Calera 1, 28760 Tres Cantos, Madrid, Spain

^b Dept. Geología y Geoquímica, Facultad de Ciencias, Universidad Autónoma de Madrid, Carretera de Colmenar, Viejo km 15, 28049, Madrid, Spain

^c Dept. Earth Sciences, Royal Holloway University London, Egham TW20 0EX, UK



ARTICLE INFO

Article history:

Received 12 April 2017

Received in revised form 13 March 2018

Accepted 23 March 2018

Available online 5 April 2018

Keywords:

Morphological features

Along-slope and down-slope processes

Neotectonic

Northwestern Iberia

ABSTRACT

The offshore region of northwestern Iberia offers an opportunity to study the impacts of along-slope processes on the morphology of a glacially influenced continental margin, which has traditionally been conceptually characterised by predominant down-slope sedimentary processes. High-resolution multibeam bathymetry, acoustic backscatter and ultrahigh-resolution seismic reflection profile data are integrated and analysed to describe the present-day and recent geomorphological features and to interpret their associated sedimentary processes. Seventeen large-scale seafloor morphologies and sixteen individual echo types, interpreted as structural features (escarpments, marginal platforms and related fluid escape structures) and depositional and erosional bedforms developed either by the influence of bottom currents (moats, abraded surfaces, sediment waves, contourite drifts and ridges) or by gravitational features (gullies, canyons, slides, channel-levee complexes and submarine fans), are identified for the first time in the study area (spanning ~90,000 km² and water depths of 300 m to 5 km). Different types of slope failures and turbidity currents are mainly observed on the upper and lower slopes and along submarine canyons and deep-sea channels. The middle slope morphologies are mostly determined by the actions of bottom currents (North Atlantic Central Water, Mediterranean Outflow Water, Labrador Sea Water and North Atlantic Deep Water), which thereby define the margin morphologies and favour the reworking and deposition of sediments. The abyssal plains (Biscay and Iberian) are characterised by pelagic deposits and channel-lobe systems (the Cantabrian and Charcot), although several contourite features are also observed at the foot of the slope due to the influence of the deepest water masses (i.e., the North Atlantic Deep Water and Lower Deep Water). This work shows that the study area is the result of Mesozoic to present-day tectonics (e.g. the marginal platforms and structural highs). Therefore, tectonism constitutes a long-term controlling factor, whereas the climate, sediment supply and bottom currents play key roles in the recent short-term architecture and dynamics. Moreover, the recent predominant along-slope sedimentary processes observed in the studied northwestern Iberian Margin represent snapshots of the progressive stages and mixed deep-water system developments of the marginal platforms on passive margins and may provide information for a predictive model of the evolution of other similar margins.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

Continental margins are built up by several sedimentary processes driven by plate tectonic evolution and environmental changes (Einsele, 2000; Pickering and Hiscott, 2016). The dominant sedimentary processes mainly originate from the combined action of down-slope movements that are mainly driven by gravity, and along-slope sedimentary processes that are forced by bottom currents (e.g., Faugères et al., 1999; Weaver et al., 2000; Stow et al., 2002; Rebesco and Camerlenghi, 2008). Understanding how these processes build continental margins

provides new insights into the origins of different morphologies and their relative importance in the evolution of passive margins.

In locations where down-slope processes are dominant, continental slope successions comprise turbidites, debrites, and submarine landslides (Weaver et al., 2000; Pickering and Hiscott, 2016). Where along-slope processes are dominant, a contourite depositional system (CDS) or contourite sedimentary system, comprising both depositional (mainly “drifts”) and erosional features (e.g., moats, channels, and furrows), is generated (Stow et al., 2002; Hernández-Molina et al., 2003, 2011; Rebesco, 2005). Although the dominant process regime on the continental slope can have temporal and spatial variations, there are many examples of continental margins, especially those described between 26°N and 56°N in the northeastern Atlantic, that have

* Corresponding author.

E-mail address: e.llave@igme.es (E. Llave).

been considered to be predominantly built up by down-slope sedimentary processes and were then defined as “glacially influenced margins” (e.g., Weaver et al., 2000; Wynn et al., 2000; Benetti, 2006, among many others). This type of margin comprises mainly linear pathways of canyons and channels (e.g., Kenyon et al., 1978; Crémer, 1983; Faugères et al., 1998; Zaragosi et al., 2000; Mulder et al., 2001; Bourillet et al., 2006; Gonthier et al., 2006; Gaudin et al., 2007). On these margins, along-slope processes have been considered to play a minor role, except in the Gulf of Cádiz and at the southern Portuguese margins (Stow et al., 2002; Alves et al., 2003; Hernández-Molina et al., 2006a, 2016a, 2016b; García et al., 2009; Roque et al., 2012, among others).

The northwestern Iberian continental margin is located within the eastern sector of the central North Atlantic (Fig. 1A), an area included within the classification of a “glacially influenced margin” (Weaver et al., 2000; Zaragosi et al., 2001a; Mojtabid et al., 2005). Geomorphological studies have characterised the area by both their down-slope and morphostructural features, such as the numerous submarine canyons, large marginal platforms (Ortegal, Pardo Bazán and Castro) and several structural highs (Maestro et al., 2015). In proximal areas, such as the Galicia Bank to the south (Ercilla et al., 2008a; Llave et al., 2008) and the Cantabrian Margin to the east (Ercilla et al., 2008b; Iglesias, 2009), previous geomorphological studies have also considered the predominance of those morphologies related to tectonics and down-slope sedimentary processes. The major goal of this study is to reveal the impacts of both along-slope and down-slope oriented sediment transport on the northwestern Iberian Margin, establishing the importance of the dynamic oceanographic settings and assessing the impacts of their interactions on reshaping the inherited structures such as marginal platforms and structural highs. A novel and detailed description of the submarine morphological features

is presented for a better understanding of the associated sedimentary processes along the continental margin and the abyssal plains off northwestern Iberia.

2. Geological and oceanographic settings

2.1. Geological setting

The northwestern Iberian continental margin comprises the Galicia continental margin and the westernmost sector of the Cantabrian Margin (Fig. 1A). The rift structures, characterised by horsts and grabens bounded by N-S, NE-SW, E-W and NW-SE faults, control the present-day morphology (Montadert et al., 1979; Groupe Galice, 1979; Boillot et al., 1979; Maestro et al., 2017) (Fig. 2). These faults have developed during the geodynamic evolution of the margin from the late-Variscan period until the present day. From the Upper Miocene until the present day, those faults with NE-SW direction have been reverse faults, the NW-SE have been normal faults and the N-S and E-W have been left-lateral and right-lateral strike-slip faults, respectively (González-Casado and Giner, 2000; De Vicente et al., 2004; Martín-González et al., 2010). The Galicia Bank is one of the most important structural highs in the Galicia margin and creates the intraslope basin called the Galicia inner basin (Boillot et al., 1975; Vanney et al., 1979). Other distinct morphostructural features in the northwestern Iberia margin are the Ortegal, Pardo Bazán and Castro Marginal Platforms and the Charcot, Coruña and Finisterre High (Maestro et al., 2015) (Fig. 2).

This area is considered to be a starved non-volcanic passive margin with only a thin sedimentary cover. The Mesozoic to Neogene sedimentary cover of the northwestern Iberian Margin is approximately 1300 m thick (Groupe Cybere, 1984; Vigneaux, 1974; Derégnaucourt




Fig. 1. A) Digital bathymetric model of the study area. B) Regional circulation of the main water masses (updated and modified from Hernández-Molina et al., 2011).




Fig. 2. Geological and tectonic setting of the northwestern Iberian continental margin and the adjacent abyssal domains, modified from Rodríguez-Fernández et al. (2014). The bathymetric map was derived from satellite and ship track data (Smith and Sandwell, 1997).

and Boillot, 1982; Thion et al., 2001; Gallastegui et al., 2002). The recent sedimentary units (Plio-Quaternary) show a very irregular distribution overlying the Mesozoic-Neogene sedimentary succession. This unit consists of alternating deposits of silt and clay laminas with interbedded coarse sediments. These sediments mainly include turbidites, debrites, contourites, and pelagites/hemipelagites (Vanney et al., 1979; Boillot et al., 1987; Comas and Maldonado, 1988; Alonso et al., 2008; Ercilla et al., 2006, 2008a). The low sediment thickness is mainly due to two reasons: a) the redistribution of sediments by longshore bottom currents and b) the shortness of the rivers of the study area as well as their steep water courses, reflecting their low sediment supply (Jané et al., 2010). Only 4.2% of the overall catchment area of the Iberian Peninsula drains to the north coast, between the Galicia-Asturias border and the French border (Uriarte et al., 2004).

2.2. Oceanographic setting

In the study area, most of the water masses are of North Atlantic origin (Pollard et al., 1996; González-Pola, 2006) and are described below from shallowest to deepest (Fig. 1B).

- a) The Eastern North Atlantic Central Water (ENACW) extends down to depths of 400–600 m and flows towards the southwest (Ambar and Fiuza, 1994; Fiúza et al., 1998; Pérez et al., 2001; González-Pola, 2006). This water mass generally flows at a velocity of 1 cm/s, although it can occasionally reach velocities of 10 cm/s (Pingree and Le Cann, 1990).
- b) The Mediterranean Outflow Water (MOW) flows along the middle slope of the Portuguese margin towards the Galicia Margin and the Bay of Biscay and extends to depths of approximately 1500 m. This water mass has two distinct cores, centred at depths of 800 and nearly 1200 m (Ambar and Howe, 1979; Mazé et al., 1997; Iorga and Lozier, 1999; Ambar et al., 2002; Slater, 2003; González-Pola, 2006), one of which flows to the west of the Galicia Bank plateau and then continues northward, while the other flows eastwards along the Cantabrian Margin slope. On its southwestern slope, the MOW exhibits high speeds and variabilities at 1100 m depth (with a mean of 18 cm/s and peaks of >40 cm/s, Ruiz-Villarreal et al., 2006). Moreover, several observations of the meddies (eddies from MOW) along the Galician margin have been published (Fiúza et al., 1998; Paillet et al., 1999, 2002; Zhang et al., 2016).
- c) The deep-water masses, i.e., those below a 1500 m water depth (wd), consist of the southward-flowing North Atlantic Deep Water (NADW), flowing between 1500 and 3000 m wd, below the northward-flowing Lower Deep Water (LDW) (Van Aken, 2000). The NADW flows below the LSW from the Labrador Sea eastwards towards the Bay of Biscay (Pingree and Le Cann, 1990) and reaches the Galicia margin. This water mass may exhibit a north-westward return flow over the Celtic continental slope (Paillet et al., 1998), and its core, initially defined by Saunders (1986), is located in this margin, between approximately 2500–3000 m wd. On the Galicia margin, the LDW is characterised by a near-bottom northward-directed flow at approximately 4000 m wd (Paillet and Mercier, 1997; Van Aken, 2000). A cyclonic recirculation cell over the Biscay Abyssal Plain has been identified as having a characteristic poleward velocity near the continental margin of 1.2 (± 1.0) cm/s (Dickson et al., 1985; Paillet and Mercier, 1997).

Internal tides and waves have been locally reported along the slopes of the Bay of Biscay as being due to a combination of favourable water mass stratification, steep topography and strong barotropic tidal currents (Pingree and Le Cann, 1989, 1990; Fiúza et al., 1998; Apel, 2002; Jackson, 2004; Varela et al., 2005; García-Lafuente et al., 2006; Krahnmann et al., 2008; Prieto et al., 2013; Zhang et al., 2016).

3. Data and methodology

The study area is located between the latitudes of 44°57' N–43°10' N and the longitudes of 13°50' W–05°29' W, extending across an area of approx. 90,000 km² at water depths of 300–5000 m (Fig. 1A). This work has been conducted in the framework of the 'Scientific Research Programme of the Economic Exclusive Zone of Spain', which is coordinated and directed by the Defence Ministry of Spain. During the oceanographic cruises conducted aboard the R/V Hespérides in 2003 and from 2006 to 2009, bathymetry and reflectivity data as well as ultrahigh-resolution seismic profiles were acquired.

A high-resolution bathymetric map was obtained using the Simrad EM-12 S120 multibeam echosounder, which allows for the simultaneous collection of high-resolution seafloor bathymetry and backscatter strength measurements. This system covers a sector of the seafloor that is approximately three times the water depth in which it operates. The seismic records comprise ultrahigh-resolution single-channel profiles that were obtained with a TOPAS (TOPographic PARametric Sonar) PS 018 system. This system is a hull-mounted seabed and sub-bottom echosounder that is based on a parametric acoustic array and that operates using the nonlinear acoustic properties of the water (Dybedal and Boe, 1994). The penetration of the acoustic signal achieved with the TOPAS system varies from between 0 and 200 ms (twtt) at full oceanic depths.

The data analysis includes: (1) the classification and cartography of the main morphologies, (2) echo-character/backscatter analyses and mapping, and (3) a correlation between the bathymetry, superficial acoustic facies and acoustic backscatter to interpret and discuss the possible sedimentary processes and their shaping of the seafloor. The nomenclature for the echo-character analysis was based on the classic guidelines established by Damuth (1980) and Pratson and Laine (1989) but was adapted for the acoustic response of the study area. High-resolution bathymetry and backscatter data analyses have been very useful for the marine research community as they reflect the many studies undertaken for the characterisation of surficial morphologies, sedimentary composition and regional deep-sea environments (Damuth, 1978; Damuth et al., 1988; Pratson and Laine, 1989; Jackson and Briggs, 1992; Hernández-Molina, 1993; Hughes Clarke et al., 1996; Pudsey and Howe, 1998; Wynn et al., 2000; Lee et al., 2002, among others). However, to avoid interpretation errors based on these indirect methods, the information should be complemented with sedimentological data from surface samples (ground truths), near-surface stratigraphy and, if possible, information of near-bottom currents, benthic organism reworking, temperature and salinity (Damuth, 1980; Mc Clennen, 1989; Pratson and Laine, 1989). The scarcity of these types of data in our study area has been compensated for via the correlation of our results with the published results from other works on the sedimentological and sedimentary processes in other nearby areas (e.g., the Armorican system, Zaragosi et al., 2000, 2001a, 2001b, 2006; the southwestern Galicia Bank, Ercilla et al., 2008a, 2011; Hernández-Molina et al., 2008; the Cantabrian continental margin, Crémer, 1983; Ercilla et al., 2008b; Iglesias, 2009; along the southern Iberian Peninsula margin, Gràcia et al., 2003; Terrinha et al., 2003) and in similar systems, such as those located in northwestern Africa (Wynn et al., 2002; Talling et al., 2007).

The nomenclature for the general morphological features and classification criteria is based on Emery (1980), Weaver et al. (2000) and Posamentier et al. (2000). For more specific morphologies, such as those related to the along-slope sedimentary processes, the classifications for the contourite depositional features (*drifts*) proposed by Faugères et al. (1999), Rebesco and Camerlenghi (2008) and García et al. (2009) for contourite erosive features have been used. The mounded, elongated and separated drifts have been referred as *separated drifts*. Those deposits that show the intercalations of along- and down-slope and/or hemipelagic/pelagic sedimentary processes have been called *mixed systems*. In addition, those bathymetric




Fig. 3. A) Digital bathymetric model of the study area and the locations of some main physiographic and morphosedimentary elements: (1) Ortega Marginal Platform, (2) Pardo Bazán Marginal Platform, (3) Castro Marginal Platform, (4) Avilés Canyon, (5) La Frouseira Canyon, (6) San Jorge Canyon, (7) El Ferrol Canyon, (8) A Coruña Canyon, (9) Lage Canyon, (10) Inner Basin, (11) Fernando High, (12) Ordoño High, (13) Finisterre High, (14) A Coruña High, (15) Charcot High, (16) Theta Gap, (17) Biscay Abyssal Plain, (18) Iberian Abyssal Plain. B) Gradient map. C) Morphological map.




Fig. 4. A) Echo-character map and locations of some main physiographic elements (see the legend of the main physiographic and morphosedimentary elements in Fig. 3A). Legend of the different echo-type descriptions in Table 2. B) Backscatter map.

elevations atop the marginal platforms between submarine canyons, within which sedimentary waves do not show the same wavy expressions as on the seafloor, have been called *relict sediment waves*. The morphological nomenclature related to the down-slope sedimentary processes is based on the works of Mutti and Normark (1987), Carter (1988), Alonso and Ercilla (2000), Einsele (2000), Mayall and Stewart (2000), and Posamentier and Kolla (2003), among others. The nomenclature for the echo-character analysis was based on the classic guidelines established by Damuth (1975, 1978, 1980), Damuth and Hayes (1977), Mullins et al. (1979), Mc Clennen (1989), and Pratson and Laine (1989).

4. Main morphosedimentary features

On the continental slopes and abyssal plains of the study area, numerous morphological features have been identified based on their seafloor expressions, echo and backscatter characteristics (Figs. 3, 4), as summarised in Tables 1 and 2.

4.1. Physiographic provinces

The northwestern Iberian margins are generally characterised by narrow shelves (approx. 30 km wide) with slopes that extend down to 5000 m wd and widths that range from between 22 and 45 km and increase eastwards. Their gradients are highly variable and range from 1° to 12° (Figs. 3A, B).

Westwards of 9°W, the average width of the continental slope is 35 km and extends down to 5000 m wd. This zone shows an average

gradient of 6° and is crossed by three main submarine canyons that are, from west to east, the Lage, A Coruña and El Ferrol canyons. Additionally, three marginal platforms, namely, the Ortegá, Pardo Bazán and Castro platforms, are located between 200 and 2400 m wd (Fig. 3A, B).

Eastwards of 9°W, the continental slope extends down to 4800 m wd and is narrower and steeper (being 22 km wide with a gradient of 12°). This zone is characterised by numerous submarine canyons, the most significant of which are the San Jorge, La Frouseira and Avilés canyons (Figs. 3A, B).

The Galicia and Cantabrian continental slopes abruptly pass into the Biscay and Iberian abyssal plains (Figs. 3A, B), which extend down to 5000 and 5300 m wd, respectively, and are characterised by the E-W Charcot High to the north and the A Coruña and Finisterre highs to the west, which show a NE-SW trend. A deep passage, the Theta Gap, connects these two abyssal plains (Figs. 3A, B). The Biscay Abyssal Plain, which is characterised by a gentle relief with a 0.03° gradient towards the SW, extends down to 5100 m wd near the Theta Gap but is found to reach approximately 4900 m wd near the Cantabrian Margin. The Iberian Abyssal Plain shows a gradient of 0.05° towards the southwest (Figs. 3A, B).

































4.2. Morphologies related to tectonics

Four main morphologies related to tectonic have been described: (1) *marginal platforms*, (2) *structural highs*, (3) *scarps*, and (4) *fluid escape-related features* (Fig. 3C). The main morphological features, such as the lengths, widths, directions, gradients and depths are

Table 1
Main morphological features identified in the study area based on their geneses. * Length and width measured only within the study area. ** Gradient of the surface where a feature is developed. *** Gradient of the steepest flank.

Morphology	Name	Length (km)	Width (km)	Direction	Gradient (°)	Depth (m)		
Structural features	Structural highs	Ordoño	11,6*	18,6*	NW-SE	16–30	2000	
		Fernando	17,8	4,7–0,7	NW-SE	23–30	2200	
		Charcot	324*	44*	E-W	8–17	3200	
		A Coruña	87*	96*	NNE-SSW	8–18	4200	
		Finisterre	75*	20–30*	NNE-SSW	3–17	4300	
	Scarps		5–20	0,5–2	NW-SE, NE-SW, NNW-SSE	14–27	600–2700	
	Marginal platforms	Ortegá	160	22	NE-SW	0,4	200–800	
		Pardo Bazán	80	25	NE-SW	2,5	900–2000	
		Castro	32	17	NE-SW	1,5	2200–2700	
	Fluid escapes-related features		0,03–0,45	0,03–0,45	NNW-SSE, NW-SE, NE-SW		200–800	
	Depositional features	Plastered drifts	7–55	10–20	NE-SW, NW-SE, E-W	0,5–0,8**	1000–2600	
		Mounded elongated and separated drifts	10–15	2–5	NE-SW, NW-SE	2,2–4,6**	500–4500	
		Mixed systems	32	17	NE-SW	0,8–1,1**	2200–2700	
Relict sediment waves		20–30	7–8	NE-SW	2–4	2400–2700		
Sediment waves		0,2	2	E-W	3***	600–650		
Mass transport deposits		5–16	3–12	NE-SW, E-W	1,7**	4000–4200		
Submarine fans		30–300	5–70	NE-SW, NW-SE, N-S	0,6–2,2	4600–4900		
Levees		35–150	4–13	NE-SW, E-W	0,03** 0,2–0,8*** 0,04**	4800		
Erosive features		Moats	5–15	0,5–2	NE-SW, NW-SE	1–5	500–4500	
		Abraded surface	90	15	NE-SW	1–2	300–650	
	Canyons	Lage	80	5–20	NE-SW, E-W	11–17	2200–4800	
		A Coruña	72	4–6	NE-SW, NW-SE	4–9	500–4800	
		El Ferrol	64	2–3	E-W, NW-SE	4–12	500–4800	
		San Jorge	35	2–3	N-S, NE-SW	11	500–4500	
		La Frouseira	26	1,5–4	NE-SW, NNE-SSW	10	900–4700	
		Gullies	5–12	1,5	NW-SE	2,8	2200–3500	
	Submarine channels	Cantabrian System	Southern	90	2–8,5	E-W	0,1	4700–4800
			Ortegá	306	1–7	NE-SW, E-W	0,03	4900–5100
			Vizcaya	242	2–12,5	NE-SW, E-W	0,02	4900–5000
		Charcot System	Charcot	170	3–17,5	NE-SW, E-W	0,02	4900
			Hespérides	358	3–12	NE-SW, E-W	0,04	4900–5100
Interplains channel			97	2,2–5	ENE-WSW, NW-SE, NE-SW	0,10	5000–5300	

Table 2
Characteristics of the main differentiated echo-character types and subtypes.

Echo type	Characteristics	Distribution	Backscatter	Interpretation	Sedimentary processes
<i>Distinct echo</i>					
	Continuous echo with absence of sub-bottom reflectors	Continental slope		Abraded surface	Erosive contourite process
	Continuous echo with stratified sub-bottom reflectors	Continental slope and abyssal plain		Mixed system	Depositional turbiditic or pelagic/hemipelagic process
	Continuous echo with prograding stratified sub-bottom reflectors	Continental slope		Mounded drift	Depositional contourite process
	Continuous echo with transparent fill	Abyssal plain		Debrite	Depositional mass flow process
	Continuous echo and no sub-bottom reflectors in the first few meters followed by zones of parallel sub-bottom reflectors and intermittent transparent layers	Abyssal plain		Channel infill	Depositional pelagic/hemipelagic process
	Continuous echo with oblique sub-bottom reflectors	Continental slope		Abraded surface	Erosive contourite process
	Continuous echo with wavy and outcropping sub-bottom reflectors with columnar disturbances	Continental slope		Pockmarks	Tectonic process related to fluids escape
	Continuous echo with wavy, outcropping and parallel sub-bottom reflectors	Continental slope		Relict sediment waves	Erosive contourite process
	Erosive bottom surface with parallel and truncated sub-bottom reflectors	Abyssal plain		Channel	Erosive turbiditic process
<i>Irregular echo</i>					
	Irregular bottom, transparent and semi-prolonged layer followed by parallel and continuous reflectors	Continental slope		Mass-transport deposits	Depositional gravitational process
<i>Hyperbolic echo</i>					
	Irregular hyperbolae overlapping with varying vertex elevations	Continental slope and abyssal plain		Erosion or outcrop	Erosive gravitational process or basement outcrop
	Regular hyperbolae overlapping with varying vertex elevations and with few or no sub-bottom reflectors	Continental slope		Slides and/or turbidites	Depositional gravitational process
	Small regular hyperbolae overlapping tangent to the seafloor without sub-bottom reflectors	Continental slope		Sediment waves	Depositional turbiditic and/or contourite processes
<i>Wavy echo</i>					
	Wavy echo with parallel sub-bottom reflectors	Continental slope and abyssal plain		Mixed system	Depositional turbiditic and/or contourite processes
	Wavy echo with no parallel sub-bottom reflectors	Abyssal plain		Sediment waves	Depositional turbiditic process
	Wavy echo with absence of sub-bottom reflectors	Continental slope		Sediment waves	Depositional contourite process




Fig. 5. A) Digital bathymetric model of the study area and locations of some main physiographic and morphosedimentary elements (see the legend of the main physiographic elements in Fig. 3A). B) and C) Bathymetric profiles. TOPAS (D–I) and digital bathymetric examples (D'–I') of fluid escape-related features. See text for detailed descriptions.

summarised in Table 1. The echo types are shown in Fig. 4, and the acoustic and reflectivity characteristics are summarised in Table 2.

- *Marginal platforms* have considerable spatial extents (approximately 1600 km²), relatively gentle slopes (0.2° to 2.5°) and are characterised as having Echo-Type 3A in its distal parts. These include the Ortegale, Pardo Bazán and Castro platforms (Fig. 5A).
- *Structural highs*. The main structural highs are a) the Ordoño and Fernando Highs of the northern Galicia Bank zone, with up to 650 m of vertical relief, which are delimited by several NNW-SSE and NE-SW scarps (Figs. 5A–C); and b) those located on the abyssal plain comprising the E-W trending Charcot High, which is located at 3200 m wd, and the predominantly NNW-SSE trending A Coruña and Finisterre highs, which are found at approximately 4200–4300 m wd. These three highs are characterised by reliefs varying from 650 to 1800 m and by Echo-Type 3A.
- *Scarps*. On the Ortegale and Pardo Bazán Marginal platforms, there are several scarps with slopes of 14–27° and 150–700 m reliefs (Fig. 5C). These scarps are acoustically characterised by Echo-Type 3A.
- *Fluid escape-related features* are located mainly in the Ortegale Marginal Platform and are characterised by Echo-Type 1G. These features have

been observed as having dome-shaped cross-sections with heights of 1 m and diameters of 150 m (Fig. 5D). Other positive relief features with circular and/or elliptical shapes that correspond to coralline mounds with heights of 3 m and diameters of 150 m are shown (Fig. 5E). In addition, almost 340 pockmarks (collapse craters) up to 16 m deep with 30–450 m diameters have been identified (Figs. 5D, E).

4.3. Morphologies related to bottom currents

Five main depositional contourite features have been observed in the study area (Figs. 3, 6A): (a) *plastered drifts*, (b) *mounded elongated and separated drifts*, (c) *mixed systems*, (d) *relict sediment waves* and (e) *sediment waves*. Two erosive contourite features comprising (f) *moats* and (g) *abraded surfaces* have also been identified. The main morphological features, such as length, width, direction, gradient and depth are summarised in Table 1. The echo types are shown in Fig. 4 and the acoustic and reflectivity characteristics are summarised in Table 2.

- *Plastered drifts*. From south to north, several upslope-prograding plastered drifts (Echo-Type 1C), with thicknesses ranging from




Fig. 6. A) Digital bathymetric model of the study area and the locations of some main physiographic and morphosedimentary elements (see the legend of the main physiographic elements in Fig. 3A). B–I) TOPAS examples of the main contourite features. Depositional contourite features comprising plastered drifts (B), separated drifts (C and D) and sediment waves (G and H). Erosive contourite features are represented by moats (C and D), abraded surfaces (I) and the erosion of relict sediment waves (F). Mixed drifts occur where alternations of contourite and turbidite features are shown (E).

between 20 and 80 ms (twtt), have been mapped as follows, (a) along the flanks of structural highs (northern zone of the Galicia Bank) at 2600 m wd (Fig. 6B), (b) on the Pardo Bazán Marginal Platform at 1200–1700 m wd, (c) at the head of the A Coruña Canyon at 400–500 m wd, and (d) in nearby canyons, such as to the north of the El Ferrol Canyon (1000–1200 m wd), south of the A Coruña Canyon (1200 m wd), and around the Lage Canyon (down to 2500 m wd).

- *Mounded, elongated and separated drifts.* From south to north, several separated drifts have been described as being characterised by Echo-Type 1C and showing average thicknesses of 40 ms (twtt). They have been identified at the following locations: (a) to the southwest of the Fernando High at 2700 m wd, (b) to the west of the Pardo Bazán Marginal Platform at 1600 m wd, (c) to the northeast of the Pardo Bazán Marginal Platform at 1200–1400 m wd, (d) to the northwest of the Ortegal Marginal Platform at 600–800 m wd, (Fig. 6C) and (e) to the north of the Galicia Bank and Castro Marginal Platform at 5000 m wd in the Biscay Abyssal Plain (Fig. 6D).
- *Mixed systems.* The whole Castro Marginal Platform is a zone comprising mixed systems that are approximately 50 ms (twtt) thick and are found over several parts of the structural highs. Acoustically, these deposits are characterised by well-stratified facies interbedded with transparent and chaotic layers (Fig. 6E), corresponding to Echo-Type 1B.
- *Relict sediment waves.* The western distal parts of the Pardo Bazán and Castro marginal platforms are characterised by sediment waves 2 km long and 15–25 m high that do not show the same morphological expression. These deposits have been classified as Echo-Type 1H (Fig. 6F).
- *Sediment waves.* These are observed only at the head of the El Ferrol Canyon and are characterised by N-S alignments, heights of 5 m, lengths of 200 m, wavelengths of 400–600 m and Echo-Type 4C. These waves are asymmetric with a steeper (3°) eastern side (Figs. 6G, H).
- *Moats.* Several moats have been described as associated with the previously reported separated drifts. They are characterised by seismic reflections truncated by the depression walls with generally symmetric V-shaped cross-sections (Fig. 6C). They are identified as follows: (a) south of the Fernando High, at 2700 m wd with a thickness of 50 m; (b) in the southwestern and northeastern zones of the Pardo Bazán Marginal Platform, at 1800 and 1500 m wd, respectively, with incisions that range from between 4 and 15 m deep; (c) in the Ortegal Marginal Platform, at the heads of the A Coruña and El Ferrol canyons, located at 450–800 m and showing incisions that are 5–20 m deep (Fig. 6C); and (d) in the Biscay Abyssal Plain, where they are characterised by 10 m deep incisions (Fig. 6D).
- *Abraded surfaces.* An extensive area characterised by an erosional surface is located to the southwest of the Ortegal Marginal Platform with a NE-SW trend. The surface defines a very high reflectivity subhorizontal plain, described as characterised by Echo-Type 1A or Echo-Type 1F (Figs. 6C, I). Also, small sectors in the shallowest part of the Ortegal Marginal Platform are described by Echo-Type 1F as abraded surfaces.

4.4. Morphologies related to gravitational processes

Four main depositional gravitational features have been observed in the study area, including (a) *submarine slides* and (b) *debris-flow deposits*, both of which are included as mass-transport deposits in Table 1, as well as (c) *submarine fans* and (d) *levees*. In addition, three erosive gravitational features have been identified, including (e) *canyons*, (f) *submarine channels* and (g) *gullies* (Fig. 7A). The main morphological features are summarised in Table 1. The echo types are

shown in Fig. 4, and the acoustic and reflectivity characteristics are summarised in Table 2.

- *Slides* are common at the borders of the canyons walls, and several isolated slides have been identified along the steepest zones of the continental slope, such as the northern zone of the Galicia Bank (17°) and within the transition of the Ortegal to Pardo Bazán marginal platforms (5°). These slides are characterised by slightly to highly deformed, back-rotated, stratified and chaotic masses resting at the base of a steep scarp (Fig. 7B) as well as by Echo-Type 3B. The heads of the larger slides display scars on the seafloor or show a rugged seafloor when many slides coalesce.
- *Debris-flow deposits.* Extensions of debris-flow deposits of up to 200 km² have developed along the lower continental slopes in the distal parts of the El Ferrol and A Coruña canyons at 4800–4900 m wd (Fig. 7C). Moreover, on the flanks of the Charcot High, these morphologies are generally 5–16 km long, 3–12 km wide and associated with steep slopes (Fig. 7D). These deposits are characterised by Echo-Type 1D. Also, another mass-transport deposit has been identified in the shallowest sector of the Ortegal Marginal Platform and is characterised by Echo-Type 2A.
- *Submarine fans.* Several submarine fans have been described at the termination of the main submarine canyons located on the Cantabrian Margin and to the north of the Inner Basin (Fig. 7A). These fans are described from east to west as follows: a) the first group comprises the three fans (i, ii and iii in Fig. 7A) that cover the most extensive span of the study area (approx. 300 km long and 70 km wide) and show a dominantly E-W orientation as well as a SE-NW trend with fingers-like shapes in their distal regions; b) the second group comprises three fans located perpendicular to the Cantabrian and Galicia margins that are approximately 5–45 km wide and 29 to 80 km long (iv to vi in Fig. 7A). Acoustically, these fans are mainly characterised by Echo-Type 1B, but fan iii in Fig. 7A is also characterised by Echo-Type 3B.
- *Channel-levees.* A group of channel-levees is described in the Biscay Abyssal Plain between the Ortegal Spur and the Charcot High and is called the Charcot Mid-Oceanic Channel System (Fig. 7E). Two more levees are observed along the right margin of the Ortegal Channel. These levees form part of a second mid-oceanic channel system called the Cantabrian Mid-Oceanic Channel System (Fig. 7F). These levees are approximately 5–18 m thick and are characterised by Echo-Type 1B.
- *Submarine canyons.* Numerous submarine canyons cross the continental slope of the study area, displaying very steep walls and U- or V-shaped cross-sections with high degrees of incision. The main canyons, described from west to east, are as follows (Fig. 8A): (a) Lage Canyon, which begins in the Castro Marginal Platform (2200 m wd) (Fig. 8B); (b) A Coruña Canyon, which begins in the Ortegal Marginal Platform (500 m wd) (Fig. 8B); (c) El Ferrol Canyon, which begins in the Ortegal Marginal Platform (500 m wd) (Fig. 8B); (d) San Jorge Canyon, which is located to the west of the Ortegal Spur (500 m wd) (Fig. 8C); and (e) La Frouseira Canyon, which begins at a 900 m wd (Fig. 8C).

Numerous minor canyons to the east at 1200–1500 m wd have been described as being primarily characterised by perpendicular and rectilinear trajectories and are U- and V-shaped with 200 m deep incisions. Note that there are several of these minor canyons located in the northern sector of the Ortegal Marginal Platform and to the east of the La Frouseira Canyon that change their orientations from S-N to SW-NE in the distal areas (Fig. 8A). These erosive features are characterised by Echo-Type 3A.

- *Submarine channels.* The channels comprising the Charcot Mid-Oceanic Channel System are characterised by three main channels that are approximately 2–10 m deep: the Biscay, Hespérides and Charcot channels, with several tributaries that are approximately




Fig. 7. A) Digital bathymetric model of the study area and the locations of some main physiographic and morphosedimentary elements (see the legend of the main physiographic elements in Fig. 3A). B–F) TOPAS examples of the main depositional gravitational features. Depositional gravitational features comprise slides (B), debris (C and D) and levees (E and F). Also, note that several erosive features are represented by the turbidite channels (E and F).

3–14 km wide, 100 km long, 3–10 m deep and have a NE-SW trend (Figs. 8A, F, E, G). However, the mid-oceanic channels of the Cantabrian System comprise two main channels (the Southern and Ortegual channels) that are 20 to 15 m deep (Fig. 8A). These channels are mainly characterised by Echo-Type 1 L but are also characterised by Echo-Type 4B in the proximal zones of the Biscay Channel and by Echo-Type 1E in some parts of the Charcot System and extensive zones of the Ortegual and Interplains Channels. All of these submarine channels converge and are constrained through the Theta Gap, where the interplain channel is formed (Figs. 8A, H, I). This channel adapts its flow to the A Coruña and Finisterre highs and is characterised by a V-shaped cross-sectional profile that is 200 m deep.

- **Gullies.** Several V-shaped gullies that are approximately 30–70 m deep have been observed in the distal parts of the Pardo Bazán and Castro marginal platforms at 2200–2700 m wd and in the northern part of the Galicia Bank at 3500 m wd, although these are 25 m deep (Figs. 8A, D).

5. Interpretation of the results and insights into continental margin sedimentary processes

5.1. Geomorphology and links with morphostructures and sedimentary processes

The integration and comparison of the results provides some practical applications for indirect near-surface and seafloor interpretations. Even though occasionally smooth or subdued bathymetric reliefs characterise the study area, especially in the abyssal plains (Figs. 3A, B), backscatter maps help to better differentiate the heterogeneities and diversities of the seafloor deposits (Fig. 4B). The characteristics and

regional distributions of the morphological features (Fig. 3C and Table 1) and echo types (Figs. 4A, 9 and Table 2) show that the seafloor in the study area is highly variable and complex.

The continental slope is predominantly characterised by highly reflective erosional features that result in *Echo-Types 1A, 1F, 1H and 3A*. The areas characterised by *Echo-Types 1A and 1F* are located along an extended SW-NE surface of the Ortegual Marginal Platform and of the Pardo Bazán Marginal Platform. The near-surface sediments obtained in this area have been studied and found to consist of clean, coarse sands and silty sands comprising surrounded and well-sorted quartz and glauconite grains as well as abundant bioclastic fragments (Alejo et al., 2012). This is evidence of high-energy environments and evolving erosion. The area characterised by *Echo-Type 3A* is located along submarine canyons, on the steep marginal platform edges and on structural highs and could be associated with the outcropping of sedimentary rocks or consolidated sediments derived from both erosion and/or tectonics. The area characterised by *Echo-Type 1H* is observed on the lower continental slope and shows similar characteristics to that of *Echo-Type 1C* but is superficially eroded. This pattern can be interpreted as contourite drifts comprising sediment waves that have recently been partially reshaped or eroded. On the other hand, the depositional features mainly located at the heads of A Coruña and El Ferrol canyons (*Echo-Type 4C*), between the Ortegual and Pardo Bazán marginal platforms and on the flanks of the structural highs (*Echo-Type 1C*), represent sediment waves and widespread contourite drifts.

Likewise, the area characterised by *Echo-Type 3B* in submarine canyons could be associated with slides and those characterised by *Echo-Type 2A* in the shallowest zones of the Ortegual Platform are possibly associated with mass-transport deposits from depositional gravitational processes. The areas characterised by *Echo-Type 3C*, recorded on the




Fig. 8. A) Digital bathymetric model of the study area and the locations of some main physiographic and morphosedimentary elements (see the legend of the main physiographic elements in Fig. 3A). B) and C) Digital bathymetric model of the main studied submarine canyons. D–I) TOPAS examples of the main erosional gravitational features, such as submarine canyons (B and C), gullies (D) and mid-oceanic channels (E to I).

floors of several channels of *Echo-Type 1B*, are observed on the Castro Marginal Platform. *Echo-Type 4A* is observed in the proximal zones of structural highs and could be associated with the interplay of turbidites or contourite sedimentary processes. Finally, a sector characterised by

Echo-Type 1G is locally observed in a distal sector of the Ortegal Marginal Platform and is related to fluid escape processes, making this process very characteristic of zones where a continuous supply of fluid flows towards the surface from below (Hovland and Judd, 1988).




Fig. 9. Main morphologies interpreted via backscatter analyses.

These results indicate that gravitational processes are predominant on the abyssal plains. The depositional gravitational features are mainly described as the levees (*Echo-Type 1B*) along several deep-sea channels and as the debrites at the base of several structural highs (*Echo-Type 1D*). Deposits from recent hemipelagic/pelagic sedimentation (*Echo-Type 1E*) infilling several areas of the deep-sea channels suggest the inactivity of those channels. In addition, despite sediment waves being a common feature of many turbidite systems found in areas crossed by unconfined turbidity currents, the observed sediment waves on several levees and submarine lobes (*Echo-Type 4B*) as well as the undulating deposits on proximal zones of the structural highs (*Echo-Type 4A*) cannot be definitively attributed to either turbidity or contour current influence. There is no clear evidence to identify a specific sedimentary process, so we considered that the origins of these deposits and sediment waves can be influenced by both down- and along-slope.

The erosive gravitational processes are represented by several deep-sea channels (*Echo-Type 1I*), especially those located along the westernmost part of the study area (i.e., Hespérides, Charcot and Biscay channels). Another erosive feature of the gravitational processes or the tectonic-related basement outcrops is described the steep structural highs (*Echo-Type 3A*).

5.2. Consideration of tectonic processes

The main morphostructural features of the northwestern Iberian Margin (marginal platforms, scarps and structural highs, Fig. 5 and Table 2) are controlled by several basement structural highs and depressions as well as asymmetric ridges produced by extensional rotated blocks, which are the result of the complex geodynamic evolution of this area that spans the Mesozoic to the present. Currently, it is possible to observe the recent rejuvenation of the escarpments via the reactivation of the NW-SE to N-S and NW-SE normal faults, the tilting of the recent sedimentary sequences above the structural highs, the development of gentle folds related to basement horst and grabens, the sudden changes in the lateral thicknesses of the sedimentary units, the development of growth faults affecting the sedimentary deposits, and the development of sedimentary wedges at the feet of escarpments related to the erosional processes of the fault scarps or sedimentary instabilities (Vázquez et al., 2008). The fault reactivation is evidenced by low to moderate seismic activity that is mainly distributed along the NE-SW and NW-SE fractures (Giner et al., 1999; López-Fernández et al., 2002).

Morphological features generated by fluid escape processes in the Ortegal Marginal are observed to be aligned in two directions, NE-SW and NW-SE (Jané et al., 2010). These directions are parallel to the orientations of the local fractures and are coincident with the locations of the master basement faults. In the same way, the strikes of the maximum axes of the elongated pockmarks are NW-SE to N-S, NE-SW and ESE-WNW, which are also parallel to the main tectonic structures. Moreover, the presence of mound-type deposits with conical or elongated morphologies on the surrounding seabed close to the pockmarks suggest violent eruptions that were sufficiently intense to instantaneously lift huge sedimentary ejecta volumes and were probably related to earthquakes (Clifton et al., 1971; Hasiotis et al., 1996; Tsunogai et al., 1998; Nikonov, 2002; Hieke, 2004; Kusçu et al., 2008; Lomtev and Gurinov, 2009; Vologina et al., 2012; Fischer et al., 2013). It has been demonstrated that earthquakes may suddenly increase fluid emission rates when the magnitude of the seismic event is greater than or equal to a 5 on the Richter scale. This is consistent with the calculations of the maximum magnitudes carried out by Martín-González et al. (2010) from a fault analysis of the northwestern sector of the Iberian Peninsula, which reported earthquake magnitudes between 6.1 and 6.8 Mw.

On the Biscay Abyssal Plain, the two large morphostructural features described, namely, the Charcot and A Coruña Highs, are related to the uplift of the oceanic crust spanning the Alpine Orogeny to the present

day (Medialdea et al., 2009) and follow an orientation established by a cortical anisotropy. The oceanic crust typically displays a tectonic spreading fabric parallel to the spreading ridge at which the crust was generated (Lonsdale, 1977; Laughton and Searle, 1979). The extension in this sector led to the formation of the Bay of Biscay with an E-W direction and the North Atlantic Ocean with a NE-SW direction. Thus, the Charcot High, which exhibits an E-W orientation, and the A Coruña High, which shows a NE-SW trend, are the result of the reactivation and uplift of the oceanic crust under the Cenozoic stress field favouring tectonic spreading. The deformation and progressive uplift of the Galicia continental margin produces the tilting of the Biscay Abyssal Plain, increasing the seafloor slope towards the north and northwest by approximately 2.5° on average.

Moreover, the orientations of the submarine canyons coincide with the main structural trends defined in the northwestern sector of the Iberian Peninsula (Arthaud and Matte, 1975; De Vicente and Vegas, 2009; Maestro et al., 2017). The Lage, A Coruña and El Ferrol canyons in the Galician margin have NE-SW, NW-SE and EW orientations (Fig. 8B), and the San Jorge and Frouseria canyons in the Cantabrian Margin have predominantly N-S to NNE-SSW and NE-SW orientations (Fig. 8C). The easternmost part of the study area includes the Avilés Canyon, which is related to the offshore continuation of the Ventaniella Fault, has a N137°E orientation and was identified from early bathymetric studies of the Biscay Margin (Boillot et al., 1979; Derégnaucourt and Boillot, 1982).

The presence of steep slopes that commonly exceed 30° in the study of the continental slope of the northern part of the Galicia Margin is caused by tectonic processes that occurred during the Alpine Orogeny, leading to the uplift of the Galicia and Cantabria margins due to the subduction of the Biscay oceanic crust under the Iberian continental crust (Gallastegui, 2000; Gallastegui et al., 2002). Some studies have shown that these uplift processes are still active, as seen via those emerged areas with episodic elevations of coastal terraces (Flor, 1983; Mary, 1983; Álvarez-Marrón et al., 1997, 2008) and by the presence of reverse faults in the Quaternary materials located on the coastal terraces (Gutiérrez-Claverol et al., 2006).

5.3. Down-slope sedimentary processes

Several submarine canyons, turbidites, slides and debrites are locally distributed on the upper and lower slopes (Fig. 7A and Table 1). They represent the products of both erosional and depositional gravitational sedimentary processes, but most of these features are also caused by tectonics (see Section 5.2). This is the case for the main submarine canyons in this study, which are often deeply incised. Many of these canyons are obliquely aligned with the slope and are isolated from areas with fluvial input. Thus, their morphologies are related to their underlying basement topography and faulting. Likewise, the occurrence of slope failures seems to be largely associated with the steep slope gradients of the canyon walls and the tectonic activity that can trigger seismic events. There are numerous studies that relate the mass-transport processes with well-known earthquakes (Sultan et al., 2004; Biscontin and Pestana, 2006; Shanmugam, 2006; Mulder, 2011; Argnani et al., 2012; Dugan, 2012; Lindhorst et al., 2012; Vargas et al., 2012).

The abyssal plains are characterised by the presence of two deep-sea channels systems that are the products of gravitational sedimentary processes. For the first time, this study reveals the courses and morphologies of two independent mid-oceanic channel systems in the Biscay Abyssal Plain: the *Cantabrian and Charcot turbidite systems* (Fig. 10A). The origins of these systems seem to be associated with the distal parts of the Cap Ferret (Figs. 10B, C) and the Armorican deep-sea fans (Fig. 11A), respectively. These systems can extend over long distances, which may be due to an increase in the slope gradient along the abyssal plain that sustains the energies of turbidity currents, as was observed by other authors in different regions (Laughton, 1960; Carter, 1988). Moreover, another factor in their development could be the same as that





Fig. 10. A) Digital bathymetric model of the study area and the locations of some of the main physiographic and morphosedimentary elements (see the legend of the main physiographic elements in Fig. 3A). B) Down-slope sedimentary processes described in this study. C) Correlation of this study within the Cantabrian Margin and Abyssal Plain interpretation by Iglesias (2009).

observed in other similar systems, which is associated with landslide-generated sediment gravity flows that are capable of spreading over extensive abyssal plains and submarine fans (Piper and Normark, 1982; Weaver et al., 1992; Wynn et al., 2002, 2010; Skene and Piper, 2006, 2003; Talling et al., 2007; Hunt et al., 2011), which are often low-energy and non-erosive environments (Weaver and Thomson, 1993; Weaver, 1994).

- The *Cantabrian Turbidite System*, comprising several channels and lobate bodies beyond the ends of these channels (*Lobes-i to iii* in Fig. 10B), shows similar planform sizes and features to those described by Prélat et al. (2010). The similarities of the study Cantabrian System to other systems described in the literature (Deptuck et al., 2008; Prélat et al., 2009) suggest that this system constitutes a nice example of compensational stacking. The depositional architecture of a basin floor fan is determined in large part by flow discharge, sand-to-mud ratio, slope length, slope gradient, and seafloor rugosity (Posamentier and Kolla, 2003; Deptuck et al., 2008; Prélat et al., 2009, 2010). Thus, depositional elements can change based on location and through time as environmental parameters change (Prélat et al., 2010). For lobes where backscatter data have been collected along with cores, a relationship was proposed between the low (light) backscatter intensity and the presence of coarse-grained deposits (Unterseh et al., 1998; Zaragosi et al., 2000; Kenyon et al., 2002). Therefore, taking this premise into consideration and given the observed backscatter and geometry in the studied lobes (Figs. 3, 4, 9), we suggest that Lobe-i is the most reflective and homogeneous and may have originated in a channel

that propagates into the basin from the complex system of the Avilés Canyon (Fig. 10B). As this lobe is very similar to frontal lobes described in other areas (e.g., Posamentier and Kolla, 2003; Morris et al., 2014; Ortiz-Karpf et al., 2015) and seems to be dominated by deposits from high density turbidity currents and other high concentration flows. Its connection with the channel is covered by the development of two more lobes: Lobe-ii, the southernmost lobe in the system, is superimposed on some parts of the higher-backscatter Lobe-i and exhibits adaptations of its morphology to the previous Lobe-i (Fig. 10B); and Lobe-iii, which developed in the northern sector of Lobe-i, also adjusted its distribution depending on the previous lobes. These two lobes comprise many elongated finger-like or finger bodies spanning across the lobes (Fig. 10B). These features are commonly attributed to flow transformations from turbidity currents to strongly density stratified flows (e.g., Klauke et al., 2004; Talling et al., 2010; Groenenberg et al., 2010; Kane et al., 2017; Spychala et al., 2017). Lobe-iii also has sediment waves on its surface that can be migrated upslope and are roughly parallel to the regional slope (Fig. 10B). These waves seem to represent primary depositional features rather than being the product of slope failures or other types of post-sedimentary deformation. Similar bedforms located in the proximal areas of the submarine fan systems and channel-lobe transitions have been described and interpreted as sediment wave fields formed by unconfined turbidity currents (Wynn et al., 2002). These features have been observed in the overall Cantabrian drainage system, comprising lobes and channels corresponding to a channel-lobe transition zone from the Cap Ferret system, which could be considered as a bypass-dominated

Fig. 11. A) Locations of the Celtic and Armorican turbidity systems related to the development of the Charcot Mid-Oceanic Channel. B–D) Evolution of the Charcot Mid-Oceanic Channel in the western part of the Biscay Abyssal Plain.



area following the classification by [Stevenson et al. \(2015\)](#). The study system could be mainly connected to the Cap Ferret system ([Crémer, 1983](#); [Ercilla et al., 2008b](#); [Iglesias, 2009](#)) and the Avilés Canyon area ([Gómez-Ballesteros et al., 2013](#)).

- In the *Charcot Turbidite System*, prominent deep-sea channels parallel to the bathymetry that comprises numerous distributaries and several levees have been described ([Figs. 7A, 8A and Table 1](#)). Primarily based on backscatter and seismic profile data, this system is characterised by a low sinuosity ([Fig. 9](#)) and seems to be the oceanic continuation of the large Armorican Fan system ([Fig. 11A](#)) that is the result of sediment-laden turbidity currents that have eroded and deposited sediment several hundreds of kilometres basinward. The large extension of the Charcot System can be explained following the results of studies of other high-latitude regions where the Coriolis forces has been observed to influence the gravity currents within these submarine channels due to the very large channel-length scales ([Wells and Cossu, 2013](#)). There are numerous factors that can control the developments and sinuosities of submarine channels, such as the slope gradients, flow interactions with the seafloor topography, the calibre of sediment the channel is transporting and eroding into, and the channel maturity (e.g., [Clark and Pickering, 1996](#); [Kane et al., 2008](#); [Peakall et al., 2012](#), among others). However, channel sinuosity is also affected by other factors, such as tectonic controls and sediment supply (e.g., [Cronin, 1995](#); [Clark and Cartwright, 2011](#); [Babonneau et al., 2002](#)), and several authors have also suggested that channel sinuosity increases with time, with meander cut-offs forming during this process (e.g., [Deptuck et al., 2003](#); [Gee et al., 2007](#); [Wynn et al., 2007](#); [Babonneau et al., 2010](#); [Maier et al., 2013](#)) or being linked to the Coriolis Force (e.g., [Peakall et al., 2012](#); [Wells and Cossu, 2013](#)). The Charcot System shares several affinities with the Equatorial Atlantic Mid-marine channel ([Baraza et al., 1997](#)), and the Northwest Atlantic Mid-Ocean Channel ([Klaucke et al., 1997](#)) as well as with the more recently identified and cored Agadir system ([Wynn et al., 2002](#); [Talling et al., 2007](#); [Masson et al., 2011](#); [Stevenson et al., 2014](#)) and Tanzania channel system ([Bourget et al., 2008](#)). The examples recently described in the literature have had origins of disintegrative submarine landslides, which can generate long run-out sediment gravity flows, or have been linked to the strong structural control of a sediment pathway associated with a massive sediment transfer towards the ocean related to tectonic activity. The presence or absence of levees in the Charcot System could be related to changes in the density stratification of its associated flows, as is discussed in other places, such that if there is a very narrow grain-size range, the construction of levees is unlikely ([Kane and Hodgson, 2011](#)). The observed levees can be classified as external levees (outer levees *sensu* [Deptuck et al., 2003, 2007](#); [Kane and Hodgson, 2011](#)). Moreover, several levees show sediment waves that are commonly identified in other regions of the modern seabed (e.g., [Wynn et al., 2002, 2007](#); [Posamentier and Kolla, 2003](#)), marking constructional crests and/or silt-prone deposits on the outside outer bends of the external levees (e.g., [Normark et al., 1993](#); [Migeon et al., 2010](#); [Wynn and Stow, 2002](#)). Pelagic/hemipelagic sedimentation could produce these semi-transparent facie units with uniform thicknesses that mimic the previous morphologies (including the valley axes), indicating that the channels were inactive (e.g., [Damuth et al., 1988](#); [Flood et al., 1995](#); [Baraza and Ercilla, 1997](#); [Wynn et al., 2007](#)).

5.4. Along-slope sedimentary processes

Several erosional (abraded surfaces and moats) and depositional (mounded and plastered drifts and sediment waves) features have been described as occurring mainly along the continental margin, e.g., the Ortegá, Pardo Bazán and Castro marginal platforms (at 0.4–0.8, 0.9–2.0 and 2.2–2.7 km wd, respectively) ([Fig. 12](#)) and to a lesser extent along the Biscay Abyssal Plain at a 5 km wd at the base of the

slope ([Fig. 6 and Table 1](#)). Their distribution demonstrates the dominant control of the bottom currents on the sedimentary processes since the distribution coincides with the depths presently under the influence of several bottom currents (ENACW-MOW-NADW along the continental slope and LDW along the abyssal plain) and bounded by density contrasts (pycnoclines) ([Figs. 1, 12](#)).

The preliminary results comprising numerous examples of contourite features confirm the interaction of the impinging MOW from west with the Ortegá Marginal Platform and the generation of a CDS. The Ortegá Spur can be considered an obstacle to the MOW into the Bay of Biscay ([Fig. 1](#)), which locally enhances its flow velocity ([Hernández-Molina et al., 2011](#)) and generates contourite drifts containing fine sands to muddy sandy on their surfaces ([Hernández-Molina et al., 2009a](#)). The depths of influence of the MOW ranges between 600 m to 1500 m wd in the study area ([Fig. 12](#)), so the interaction of the lower part of the MOW can explain many of the identified erosional and depositional features on the Pardo Bazán Marginal Platform and the upper MOW on the distal parts of the Ortegá Marginal Platform ([Fig. 12](#)). Moreover, the irregularities of the seafloor can locally force the MOW to circulate at shallower depths of the slope. In general, the interaction of the contour currents with the steep slopes causes enhanced and focused bottom-current flows evolving erosion along the moat and on the upslope flank, and deposition of separated mounded drifts on the other flank ([McCave, 1982](#)). However, the plastered drifts located on the Pardo Bazán Marginal Platform ([Fig. 12](#)) follow the typical contourite accumulation scenario proposed by [Faugères et al. \(1999\)](#) and [Rebesco \(2005\)](#) for their development, consisting of a broad, non-focused current on a gentle slope at low current velocities. The drifts located at the base of the Fernando and Ordoño highs, which are influenced by the NADW, are evidence of the effects of obstacles on an impinging flow ([Fig. 13](#)). As has been described by other authors for different water masses, the water flow in the northern hemisphere seems to accelerate to the left of a seamount (looking downstream) and decelerate to the right ([Roberts et al., 1974](#); [Gould et al., 1981](#); [Hernández-Molina et al., 2006b](#)). This standard behaviour in bottom currents supports the development of moats/mounded drifts on the left flanks of the Fernando and Ordoño highs and plastered drifts located on the right flank and/or around structural highs ([Fig. 13](#)). Moreover, the broad described erosional and abraded surfaces showing low slope angles in the study area could correlate to the influence of less confined tabular bottom currents as [Rebesco et al. \(2014\)](#) define in similar areas.

This conceptual sedimentary model describe the generation of most of the major studied contourite features but do not answer the development of minor features such as the deposition of sediment waves in Ferrol and A Coruña submarine canyons. In this case their construction can be associated to the origination of meddies, internal waves and internal baroclinic tides at the ENACW-MOW interface ([Fig. 12](#)). These oceanographic processes are commonly recognised to be originated at the pycnoclines ([Cacchione et al., 2002](#); [Lamb, 2014](#)) and there are several studies about the generation of sediment waves as products derived from their interactions with steep continental margins (e.g., [Hernández-Molina et al., 2011](#); [Pomar et al., 2012](#); [Preu et al., 2013](#); [Shanmugam, 2013](#); [Hanebuth et al., 2015](#); [Collart et al., 2018](#)). On the Galician upper slope there are oceanographic measurements obtained by [García-Lafuente et al. \(2006\)](#) that show baroclinic bottom currents of up to 19.4 cm/s. Moreover, [Balmforth and Young \(2001\)](#), [Zhang et al. \(2008\)](#), [Quaresma and Pichon \(2013\)](#), [Lamb \(2014\)](#) and [Collart et al. \(2018\)](#) consider the formation of consistent internal tide on the Galician middle slope. This type of interaction of the seafloor with the filaments of water masses forced to flow upslope could develop these kinds of sedimentary waves, as has been reported for other slopes (e.g., [Van Rooij et al., 2003, 2010](#); [Iglesias, 2009](#)).

On the other hand, those relict drifts and sediment waves described at 2400–2700 m wd ([Table 1](#)) are formed by water masses strong enough to develop contourites and sediment waves. Although the present depth of the AABW flow is below 3500 m ([Fig. 1](#)), it is believed




Fig. 12. Vertical distribution of the water masses flowing in the eastern part of the study area in the digital terrain model (DTM). B) Seismic profile crossing the terraces (line position in panel A), where the different water masses and their interfaces are indicated for the present day, and the theoretical position of the MOW is indicated during glacial stages (in black) (Sánchez-González, 2013). C) Water mass percentages in the area adjacent to the contourite terraces (from Sánchez-González, 2013).

that this current was enhanced (Sarnthein et al., 1982; Duplessy et al., 1988; Maslin et al., 1995; Zahn et al., 1997; Elliot et al., 2002) and shoaled at ~2500 m (Keigwin, 2004) during the past, especially during the glacial periods. Several studies carried out along the Atlantic margin have discussed the variations in depth of the AABW in the deep Atlantic basin at subtropical Iberian latitudes during the glacial/stadial periods (Sarnthein et al., 2001; Curry and Oppo, 2005; Voelker et al., 2010;

Friedrich et al., 2014; Rodríguez-Tovar et al., 2015; Howe et al., 2016) as well as the importance of its influence on developing contourite features (Roque et al., 2015).

Finally, the distal parts of several submarine canyons, e.g., the very noteworthy features to the east of La Frouseira Canyon (Fig. 4A) and the turbidite fans situated at the base of the slope (i.e., turbidite fan iv in Fig. 7A), show a change in their direction from S-N to SW-NE,




Fig. 13. A) Digital terrain model (DTM) of the northern part of the Galicia Bank showing the NADW bottom interaction that triggers the development of contourite deposits related to structural highs. Examples of high-resolution seismic profiles showing mounded, elongated and separated drifts in the western flank of the Ordoño (B) and Fernando (C) highs and D) a plastered drift on the western flank of the Ordoño High.

which is interpreted as a consequence of the interplay of the S-N trending gravitational influence through the canyon with the W-S LDW circulation (Fig. 1).

6. Discussion: architectural sedimentary changes and evolution

6.1. Significance of the along-slope sedimentary processes in changing the conceptual glacially influenced margin model and its comparison with other glacially influenced margins

When considering all described bottom-current products, especially those described for the Ortegá, Pardo Bazán, and Castro marginal platforms (Fig. 5A, B and Table 1), the first noticeable aspect is the relationship between the positions of the water masses (Fig. 1B) and the developments of both depositional and erosive contourite features (Fig. 3C). These features are characteristics of *contourite terraces*, which are formed by the interplay between margin physiography and local current regimes. Marginal platforms or *marginal plateaus* are defined in the submarine morphology as “flat (subhorizontal) but deep (deeper than the shelf break) domains within the continental slope” (Mercier de Lépinay et al., 2016). These features occur on all passive margins, occupying areas of >18,000,000 km² and comprising 5.11% of the oceans (Harris et al., 2014). The polyphase tectonic evolution, subsidence evolution and the sedimentary settings controlled by deep currents make the marginal plateaus a target for furthering our understanding of the

continental margin and hydrocarbon exploration. Our study reflects that the bottom currents have overemphasized the tectonic processes responsible for the initial development of the marginal platforms and structural highs (Figs. 5, 12). In this new scenario, the Ortegá, Pardo Bazán and Castro marginal platforms, where both erosional and depositional contourite features are described (Fig. 12B) and where the water masses show remarkable density contrasts (Fig. 12C), can be considered to constitute large *contourite terraces* (hereafter named the *Ortegá, Pardo Bazán and Castro terraces*) (Fig. 14). Therefore, the study area constitutes an excellent example for the study of the control of inherited structures and the influence of the topography of marginal plateaus on the local and global circulations of intermediate and deep oceanic waters. Based on the correlations of these morphologies and the ocean regimes, we propose that terrace genesis is strongly connected to the turbulent current patterns typical of water mass interfaces. Furthermore, this energetic pattern might be enforced by internal tides and internal waves, which are known to be generated along strong density gradients typical of water mass interfaces and which have tremendous effects on sediment dynamics, resulting in erosion and resuspension (Dickson and McCave, 1986; van Raaphorst et al., 2001; McCave, 2001; Cacchione et al., 2002; Hosegood and van Haren, 2004). Several contourite terraces have been defined at other continental margins at different latitudes, with the difference with respect to our study area being related to the absence of a structural control on the terrace formation (e.g., Gulf of Cádiz, Hernández-Molina et al., 2016a;




Fig. 14. Sketch of the conceptual sedimentary model proposed in this study. The main tectonic, along- and down-slope sedimentary processes and the principal depositional/erosional products generated by these processes are shown.

Argentine margin, Hernández-Molina et al., 2009b; Gruetzner et al., 2012; Preu et al., 2013; the Uruguayan margin, Hernández-Molina et al., 2016c; Alboran Sea, Ercilla et al., 2016).

This contribution proposes that although there are similarities with respect to the previous conceptual “glacially influenced margin” related to the development of numerous submarine canyons and turbidity currents that deliver sediment to abyssal plains (e.g., Kenyon et al., 1978; Lebreiro et al., 1997; Wynn et al., 2000), there is a remarkable difference in the interplay of the down- and along-slope processes, including the presence of 1) three marginal platforms on the continental slope dominated by along-slope processes evolving into contourite terraces and 2) down-slope processes that are only minimally active on the upper slope but are dominant in the lower slope and abyssal plain regions (Fig. 14). These are two major and peculiar characteristics that suggest another perspective of the study area wherein the contour currents rework and/or redistribute the gravity deposits.

6.2. Recent changes in the architectural sedimentary evolution

Down-slope and along-slope sediment transports are the dominant processes during the evolution of continental margins and have major impacts on the shapes of these margins (Stow and Faugères, 1998; Faugères et al., 1999; Stow et al., 2002). Previous studies of glacially influenced margins have suggested that during full-glacial periods, sediment delivery onto the slope was mainly achieved by down-slope mass-wasting processes, while along-slope processes that deposited contourites dominated during interglacial times when the bottom currents were generally stronger and on the abyssal plains (Weaver et al.,

2000). In the study area, we propose that the contourite processes were dominant over the middle slope, mainly caused by the MOW circulation and associated with the enhancement of this current during the glacial cycles (Cacho et al., 2000; Schönfeld and Zahn, 2000; Llave et al., 2006, 2010, 2011; Voelker et al., 2006; Hernández-Molina et al., 2011, 2016a; Toucanne et al., 2007, 2012; Rogerson et al., 2012). However, the down-slope processes, which are mainly focused on the upper and lower slopes, are also considered to be more active in the study area during regressive and lowstand stages (Zaragosi et al., 2001a, 2001b; Toucanne et al., 2012) but are dissipated by along-slope processes on the middle slope where the along-slope processes are dominant. This scenario could explain the absence of extensive turbidity systems off the contourite terraces (Fig. 14), and nevertheless they only develop to the east and west of the terraces and associated to the Cap Ferret and Armorican turbidity systems, respectively. Therefore, water masses circulating in the study area show another example within modern oceans of their importance in re-suspending and advecting sea-floor sediments or pirate sediments from other sedimentary processes (e.g., gravity flows).

The contourite terraces may have been controlled by the following key factors, starting with a) the local morphology of the margin: inherited geological features conditioned the study area as having pelagic platforms located on its middle/upper slopes. These structures produced changes in the regional slope trends, providing the best areas for local water mass interactions with the seabed. The next factor is b) sediment supply: superfacial samples of the drifts are mainly characterised as biogenic sandy contourites (Hernández-Molina et al., 2009a; Alejo et al., 2012). However, a wide variety of sediment inputs

in CDS generation could also be considered. Contour currents can redistribute gravity supplied sediments or simply rework gravity deposits and transport any sedimentary drift bodies downward. Additionally, a greater direct sediment supply from the shelf and hinterland sources should be expected during the regressive and lowstand periods. The third factor is c) oceanographic changes: a vertical variability of the contourite features is observed, including erosive features close to the interface depths, plastered drifts between the interface depths and the upslope progradation of the mounded drifts and sediment waves, suggesting a vertical upward/downward migration or expansion, which observed most clearly at the depths of the MOW. Glacioeustatic changes could have facilitated the absence of the sedimentary imprints of the studied contourite terraces. Palaeoceanographic models of the MOW and its interfaces in the Western Mediterranean Sea and the Gulf of Cádiz assume that intermediate and deep Mediterranean water masses strengthened and sank to approximately 600–700 m deeper than its modern counterpart during the glacial periods of the Quaternary (Cacho et al., 2000; Schönfeld and Zahn, 2000; Llave et al., 2006, 2010, 2011; Voelker et al., 2006; Hernández-Molina et al., 2011, 2016a; Toucanne et al., 2007, 2012; Rogerson et al., 2012). In the study area, this deepening reflects a shift of the upper interface to below 1100 m wd and of the lower interface down to 2200 m wd (Sánchez-González, 2013). Our sedimentary model suggests that during colder periods, the deeper ENACW/MOW and MOW/LSW pycnoclines influenced the distal sector of the Ortegale terrace, the upper part and distal sector of the Pardo Bazán terrace and the upper part of the Castro terrace, mainly by developing erosive features (Fig. 12B) that could have masked the development of depositional features, including those that originated from gravity sedimentary processes. Those sectors located between the pycnoclines were more favourable for depositional features, which could be relict or even have been eroded by successive variations of the interface depths. It is possible that the interplay between the along- and down-slope processes is less evident along the Castro Marginal Platform as a result of the smaller sensitivities to local changes in the LSW compared to those in the MOW, which would lead to alternating periods of dominant along- and down-slope sedimentary processes as well as the generation of the described mixed systems.

On the other hand, the Iberian Abyssal Plain is mainly characterised by two turbidite systems shaping the seafloor, which are the main contributors to both the outbuilding of the distal continental margin and basin infilling (Fig. 10). The Charcot System records multiple phases of incision and several pelagic/hemipelagic sediment infills that are evidence of intervals of active and inactive channel-levee systems (Figs. 7E, F, 8E–I). This fact has sometimes been related to local or regional changes in sediment supply rates or has been considered as being relative to variations of the sea-level. The systems that are connected to the Armorican and Celtic systems in the north (Fig. 11A) are assumed to have increased activity when increased sediment delivery is produced during lowstand sea levels (Zaragosi et al., 2001a, 2001b; Toucanne et al., 2012). In contrast, one can expect that the complete burial and abandonment of a large and long-persistent deep-sea channel could be a consequence of first-order tectonic processes. Our study proposed that there is an evolution in this development/abandonment of several channels and distributaries that is mainly associated with the continental margin deformation and consequent tilting with a progressively westward increased gradient of the abyssal plain during the Pleistocene. The confinement by the structural highs of the flow along with the glacioeustatic changes and influences on the sediment supply are especially important in the submarine landside and the turbidity current activity during the Quaternary (Auffret et al., 2002). The progressive changes in the recent evolution of the system can be synthesized as follows: (a) the first stage consists of an erosive phase when the Charcot, Biscay and Hespérides channels were incised into the seafloor following the topography of the Charcot High and were intensified by the constricted conditions between the Charcot High and the Ortegale Spur (Fig. 11B); (b) the second stage, marking an

intensification of the turbidity currents, is observed along the Hespérides channel, which progrades towards the Charcot High, and the development of additional tributaries and levees is observed (Fig. 11C); (c) the third and final stage, consisting of the inactivity of the tributaries and the predominant intensification of the Hespérides channel, is then observed. In this final stage, an extensive levee also develops along the Hespérides channel to the south of the Charcot High (Fig. 11D).

The Biscay Abyssal Plain regionally slopes in a southward direction and (in the area close to the A Coruña High) its deepest section favours the convection of all of these channel/levee systems through the Theta Gap (Fig. 8A). This connection narrows to approximately 5 km wide between the Biscay Abyssal Plain (average water depth of 5100 m) and the Iberian Abyssal Plain (approximately 5300 m deep) and could have acted as a deep-ocean overflow between oceanic basins of different depths (Legg et al., 2009). Active erosion occurs approximately 15 km upslope of the narrowing between the structural highs and creates initial incisions of up to 200 m and later incisions of approximately 30 m at points further along the current's course (Figs. 8H, I and Table 1). The straightening of the LDW water mass (Fig. 1) due to this constriction and a topographic barrier could be responsible for an overflow of the water mass generation that favours the excavation of the interplain channel. A change in sedimentation in the more distal part of the interplain channel system would then occur, and turbidites would progressively be replaced by the pelagic/hemipelagic deposits caused by the switch-off of turbidity flow activity (Fig. 8I).

These sedimentary modes and their recent evolution comprising the interplays of the structural and along- and down-slope sedimentary processes require further investigation (i.e., regarding their stratigraphy and sedimentary characterisation) since they have important implications for regional and global climate dynamics, morphostructural evolution and hydrocarbon reservoirs. Thus, the interplay of down-slope and along-slope processes represents a critical and little known topic in the study of deep-marine sedimentation (e.g., Hernández-Molina et al., 2009b, 2016a, 2016b; Rebesco et al., 2008; Mulder et al., 2008; Mutti et al., 2009; Marches et al., 2010; Talling et al., 2012; Brackenkridge et al., 2013; Stow et al., 2013).

7. Conclusions

A comprehensive sedimentary evolution of the recent depositional processes operating along the northwestern Iberian Margin and its adjacent abyssal plains is proposed herein. Our evolutionary and conceptual model provides new perspectives of the interplays between down- and along-slope sedimentary processes in shaping this glacially influenced margin and the important role of bottom currents in obscuring the tectonic process signatures in this area.

Down-slope processes are dominant on the upper and lower slope and within the abyssal plain regions but have minimal impact on the middle slope, which is strongly influenced by along-slope processes. These processes show minimal concurrent interactions.

Therefore, this study provides a new perspective of the early continental margin model based on two main and peculiar characteristics. A) The northwestern Iberian continental slope is affected by down-slope gravitational processes (mass-transport and turbidity) that mainly occur within canyons. However, the dominance of contourite features (contourite drifts, sediment waves, moats and abraded surfaces), described mainly along the inherited marginal platforms, is evidence of the influence of the intermediate bottom currents and their importance in shaping the seafloor and the evolution of contourite terraces. B) The abyssal plains are dominated by pelagic/hemipelagic deposits as well as by debris flows and turbidity currents where two complex channel-lobe systems develop (i.e., the Cap Ferret and Armorican turbidite systems). Nevertheless, in this abyssal plain down-slope scenario, several contourite features are also observed at the foot of the continental slope as a result of the influence of the deepest water masses.

This recent and/or present-day sedimentary stacking pattern and the sediment dynamics in the study area are the result of both the structural framework and the variability of sedimentary systems and processes influenced by glacioeustatic changes. Tectonic processes are considered to be the controlling factors over the long-term, while sediment supply and climatic conditions have critical influences on the present-day configuration of the area at the basin scale, having played major roles over the short-term by influencing the variabilities of different sedimentary processes. Note that the tectonic processes have mainly conditioned the inherited marginal platforms as well as the structural highs, which have then influenced the interactions with the bottom currents, even modifying the signatures of down-slope activity. In the abyssal plains, tectonic processes have also controlled the westward migration of the channels/levee systems by tilting the continental margin. In addition, the lateral and vertical shifts of the water masses influence the depth, and their interfaces have left imprints on the reshaping and evolution of the recent sedimentary architecture of the continental slope, which was conditioned by the climatic variations during the Pleistocene.

Future work in this area should aim to provide a better understanding of the formation, timing and evolution of the morphostructural and morphosedimentary features of this region and to better constrain the interactions between tectonic, gravitational and bottom-current processes as well as sediment supply and climate variations.

Acknowledgements

We are especially grateful to those who participated in the research cruises of the ZEEE (Spanish Economic Exclusive Zone) project, which was headed by the Instituto Hidrográfico de la Marina (Hydrographic Institute of the Spanish Navy) and the Real Instituto y Observatorio de la Armada (Spanish Navy Royal Observatory), as well as to the captains and crews of the R/V Hespérides. This research was funded by the Instituto Geológico y Minero de España (Geological Survey of Spain) Research scholarship (2007–2011). This research is supported through the PROGEZEE (CTM2011-13367-E) and CGL2016-80445-R (AEI/FEDER, UE) SCORE projects. This research was partially conducted within the framework of “The Drifters Research Group” of the Royal Holloway University of London (UK). The authors also wish to express gratitude to the Editor, Dr. Plater and Reviewers, Dr. Hodgson and Anonymous, for their constructive feedback.

References

- Alejo, I., Nombela, M.A., Ercilla, G., Brackenridge, R.E., Francés, G., Pérez-Arlucea, M., Mena, A., Hernández-Molina, F.J., Stow, D.A.V., Medialdea, T., 2012. Caracterización de los sedimentos superficiales en tres sistemas deposicionales contorníticos (Golfo de Cádiz, Cabo Ortegal y El Cachucho): implicaciones conceptuales. *Resúmenes extendidos del VIII Congreso Geológico de España. Geo-Temas* 13, 1781–1784.
- Alonso, B., Ercilla, G., 2000. Valles Submarinos y Sistemas Turbidíticos Modernos. In: Alonso, B., Ercilla, G. (Eds.), *Consejo Superior de Investigaciones Científicas Barcelona* (292 pp.).
- Alonso, B., Ercilla, G., Casas, D., Estrada, F., Farrán, M., García, M., Rey, D., Rubio, B., 2008. Late Pleistocene and Holocene sedimentary facies on the SW Galicia Bank (Atlantic NW Iberian Peninsula). *Mar. Geol.* 249, 46–63.
- Álvarez-Marrón, J., Rubio, E., Torné, M., 1997. Subduction related structures of North Iberian Margin. *J. Geophys. Res.* 102 (10), 22245–22511.
- Álvarez-Marrón, J., Niedermann, S., Hetzel, R., Menéndez, R., Marquinez, J., 2008. Origin, structure and exposure history of a wave-cut platform more than 1 Ma in age at the coast of northern Spain: a multiple cosmogenic nuclide approach. *Geomorphology* 93, 316–334.
- Alves, T.M., Gawthorpe, R.L., Hunt, D.W., Monteiro, J.H., 2003. Cenozoic tectono-sedimentary evolution of the western Iberian margin. *Mar. Geol.* 195, 75–108.
- Ambar, I., Fiúza, A., 1994. Some features of the Portuguese current system: a poleward slope undercurrent, an upwelling related southward flow and an autumn-winter poleward coastal surface current, 2nd International Conference on Air-Sea Interaction and on Meteorology and Oceanography of the coastal zone, preprints, September 22–27. *Am. Meteorol. Soc.* 286–287.
- Ambar, I., Howe, M.R., 1979. Observations of the Mediterranean outflow. Deep circulation in the vicinity of the Gulf of Cadiz. *Deep-Sea Res.* 26 (5), 535–554.
- Ambar, I., Serra, N., Brogueira, M.J., Cabeçadas, G., Abrantes, F., Freitas, P., Gonçalves, C., Gonzalez, N., 2002. Physical, chemical and sedimentological aspects of the Mediterranean outflow off Iberia. *Deep-Sea Res. II Top. Stud. Oceanogr.* 49 (19), 4163–4177.
- Apel, J.R., 2002. *Oceanic Internal Waves and Solitons. An Atlas of Oceanic Internal Solitary Waves*. pp. 1–40.
- Argnani, a., Armigliato, a., Pagnoni, G., Zaniboni, F., Tinti, S., Bonazzi, C., 2012. Active tectonics along the submarine slope of South-Eastern Sicily and the source of the 11 January 1693 earthquake and tsunamis. *Nat. Hazards Earth Syst. Sci.* 12, 1311–1319.
- Arthaud, F., Matte, P., 1975. Les décrochements tardihercyniens du sudouest de l'Europe. *Géometrie et essai de reconstitution des conditions de la déformation. Tectonophysics* 25, 139–171.
- Auffret, G., Zaragoza, S., Dennielou, B., Cortijo, E., Van Rooij, D., Grousset, F., Pujol, C., Eynaud, F., Siebert, M., 2002. Terrigenous fluxes at the Celtic margin during the last glacial cycle. *Mar. Geol.* 188 (1–2), 79–108.
- Babonneau, N., Savoye, B., Cremer, M., Klein, B., 2002. Morphology and architecture of the present canyon and channel system of the Zaire deep-sea fan. *Mar. Pet. Geol.* 19, 445–467.
- Babonneau, N., Savoye, B., Cremer, M., Bez, M., 2010. Sedimentary architecture in meanders of a submarine channel: detailed study of the present Congo turbidite channel (Zaiango project). *J. Sediment. Res.* 80, 852–866.
- Balmforth, N., Ierley, G., Young, W., 2002. Tidal conversion by subcritical topography. *J. Phys. Oceanogr.* 32 (10), 2900–2914.
- Baraza, J., Ercilla, G., Farrán, M., Casamor, J.L., Sorribas, J., Flores, J.A., Wersteeg, W., 1997. The Equatorial Atlantic Mid-Ocean Channel: An Ultra High-Resolution Image of Its Burial History Based on TOPAS Profiles. *Mar. Geophys. Res.* 19 (2), 15–135.
- Benetti, S., 2006. Late Quaternary Sedimentary Processes along the Western North Atlantic Margin. (Unpublished Ph.D. Thesis). University of Southampton, UK (188 pp.).
- Biscontin, G., Pestana, J.M., 2006. Factors affecting seismic response of submarine slopes. *Nat. Hazards Earth Syst. Sci.* 6 (1), 97–107.
- Boillot, G., Dupeuble, P.A., Musellec, P., 1975. Carte géologique du plateau continental nord-portugais au nord du cap Carvoeiro, Bulletin de la Société Géologique de France. vol. 4 pp. 462–480.
- Boillot, G., Dupeuble, P.A., Malod, J., 1979. Subduction and tectonics on the continental margin off northern Spain. *Mar. Geol.* 32, 53–70.
- Boillot, G., Malod, J.A., Dupeuble, P.A., Cybere Group, 1987. Mesozoic evolution of Ortegal Spur, North Galicia margin: comparison with adjacent margins. In: Boillot, G., Winterer, E.L., Meyer, A.W. (Eds.), *Proceedings of the ODP, Initial Reports 103, A*, pp. 107–119.
- Bourget, J., Zaragoza, S., Garlan, T., Gabelotaud, I., Guyomard, P., Dennielou, B., Ellouzi-Zimmermann, N., Schenider, J.L., 2008. *Mar. Geol.* 255, 179–185.
- Bourillet, J.F., Zaragoza, S., Mulder, T., 2006. The French Atlantic margin and deep-sea submarine systems. *Geo-Mar. Lett.* 26, 311–315.
- Brackenridge, R.E., Hernández-Molina, F.J., Stow, D.A.V., Llave, E., 2013. A Pliocene mixed contourite-turbidite system offshore the Algarve Margin, Gulf of Cadiz: seismic response, margin evolution and reservoir implications. *Mar. Pet. Geol.* 46, 36–50.
- Cacchione, D.A., Pratson, L.F., Ogston, A.S., 2002. The shaping of continental slopes by internal tides. *Science* 296 (5568), 724–727.
- Cacho, I., Grimalt, J.O., Sierro, F.J., Shackleton, N., Canals, M., 2000. Evidence for enhanced Mediterranean thermohaline circulation during rapid climatic coolings. *Earth Planet. Sci. Lett.* 183, 417–429.
- Carter, R.M., 1988. The nature and evolution of deep-sea channel systems. *Basin Res.* 1, 41–54.
- Clark, R.I., Cartwright, A.J., 2011. Key controls on submarine channel development in structurally active settings. *Mar. Pet. Geol.* 28, 1333–1349.
- Clark, J.D., Pickering, K.T., 1996. *Submarine Channels: Process and Architecture*. Vallis Press, London (231 pp.).
- Clifton, H., Greene, H., Moore, G., Phillips, R., 1971. Methane seep off Malibu Point following San Fernando earthquake. *Geol. Surv. Prof. Pap. (United States)* 733, 112–116.
- Collart, T., Verreydt, W., Hernández-Molina, F.J., Llave, E., León, R., Gómez-Ballesteros, M., Pons-Branchu, E., Stewart, H., Van Rooij, D., 2018. Sedimentary processes and cold-water coral mini-mounds at the Ferrol canyon head, NW Iberian margin. *Prog. Oceanogr.* <https://doi.org/10.1016/j.pcean.2018.02.027> In Press.
- Comas, M.C., Maldonado, A., 1988. Late Cenozoic sedimentary facies and processes in the Iberian Abyssal Plain, Site 637, ODP Leg 103. In: Boillot, G., Winterer, E.L., et al. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results, College Station, TX (Ocean Drilling Program)*. 103, pp. 635–655.
- Crémer, M., 1983. Approches sédimentologique et géophysique des accumulations turbiditiques. L'éventail profond du Cap Ferret (Golfe de Gascogne), la Série des Gres d'Annot (Alpes de Haute Provence). (Ph.D. Thesis). Univ Bordeaux, France, p. 344.
- Cronin, B.T., 1995. Structurally-controlled deep sea channel courses: examples from the Miocene of southeast Spain and the Alboran Sea, southwest Mediterranean. *Geol. Soc. Lond. Spec. Publ.* 94, 115–135.
- Curry, W.B., Oppo, D.W., 2005. Glacial water mass geometry and the distribution of d13C of SumCO2 in the western Atlantic Ocean. *Paleoceanography* 20 (1) (PA1017).
- Damuth, J.E., 1975. Echo character of the western equatorial Atlantic floor and its relationship to the dispersal and distribution of terrigenous sediments. *Mar. Geol.* 18, 17–45.
- Damuth, J.E., 1978. Echo character of the Norwegian–Greenland Sea: relationship to Quaternary sedimentation. *Mar. Geol.* 28, 1–36.
- Damuth, J.E., 1980. Use of high-frequency (3.5–12 kHz) echo-grams in the study of near-bottom sedimentation processes in the deep-sea: a review. *Mar. Geol.* 38, 51–75.
- Damuth, J.E., Hayes, D.E., 1977. Echo character of the East Brazilian continental margin and its relationship to sedimentary processes. *Mar. Geol.* 24, 73–95.
- Damuth, J.E., Flood, R.D., Kowsmann, R.O., Belderson, R.H., Gorini, M.A., 1988. Anatomy and growth pattern of Amazon deep-sea fan as revealed by long-range side-scan

- sonar (GLORIA) and high resolution seismic studies. *Bull. Am. Assoc. Pet. Geol.* 72, 885–911.
- De Vicente, G., Vegas, R., 2009. Large-scale distributed deformation controlled topography along the western Africa–Eurasia limit: tectonic constrains. *Tectonophysics* 474, 124–143.
- De Vicente, G., Vegas, R., Muñoz Martín, A., González-Casado, J.M., Carbó, A., Álvarez, J., Cloetingh, S., Andriessen, P., Elorza, F.J., Olaiz, A., 2004. El Sistema Central. In: Vera, J.A. (Ed.), *Geología de España*, Instituto Geológico y Minero de España, Madrid, pp. 621–626.
- Deptuck, M.E., Steffens, G.S., Barton, M., Pirmez, C., 2003. Architecture and evolution of upper fan channel belts on the Niger Delta slope and in the Arabian Sea. *Mar. Pet. Geol.* 20, 649–676.
- Deptuck, M.E., Sylvestre, Z., Pirmez, C., O'Byrne, C., 2007. Migration/aggradation history and 3-D seismic geomorphology of submarine channels in the Pleistocene Benin-major canyon, western Niger Delta slope. *Mar. Pet. Geol.* 24, 406–433.
- Deptuck, M.E., Piper, D.J.W., Savoye, B., Gervais, A., 2008. Dimensions and architecture of late Pleistocene submarine lobes off the northern margin of East Corsica. *Sedimentology* 55 (4), 869–898.
- Derégnaucourt, D., Boillot, G., 1982. Nouvelle carte structurale du golfe de Gascogne. *Comptes Rendus de l'Académie des Sciences de Paris II* 294, 219–222.
- Dickson, R.R., McCave, I.N., 1986. Nepheloid layers on the continental slope west of Porcupine Bank. *Deep-Sea Res.* 33 (6), 791–818.
- Dickson, R.R., Gould, W.J., Muller, T.J., Maillard, C., 1985. Estimates of the mean circulation in the deep (>2000 m) layer of the eastern North Atlantic. *Prog. Oceanogr.* 14, 103–127.
- Dugan, B., 2012. A review of overpressure, flow focusing, and slope failure. In: Yamada, Y., Kawamura, K., Ikehara, K., Ogawa, Y., Urgeles, R., Mosher, D., Chaytor, J., Strasser, M. (Eds.), *Submarine Mass Movements and their Consequence*, Springer, Netherlands, pp. 267–276.
- Duplessy, J.-C., Shackleton, N.J., Fairbanks, R.G., Labeyrie, L., Oppo, D., Kallel, N., 1988. Deepwater source variations during the last climatic cycle and their impact on the global deepwater circulation. *Paleoceanography* 3, 343–360.
- Dybedal, J., Boe, R., 1994. Ultra high resolution sub-bottom profiling for detection of thin layers and objects. *Oceans'94/Osates*, Brest, France.
- Einsele, G., 2000. *Sedimentary Basins. Evolution, Facies, and Sediment Budget*. second ed. Springer, Berlin (792 pp.).
- Elliot, M.L., Labeyrie, L., Duplessy, J.C., 2002. Changes in North Atlantic deep-water formation associated with the Dansgaard-Oeschger temperature oscillations (60–10 ka). *Quat. Sci. Rev.* 21 (10), 1153–1165.
- Emery, K.O., 1980. Continental margins - classification and petroleum prospects. *Am. Assoc. Pet. Geol. Bull.* 64, 297–315.
- Ercilla, G., Córdoba, D., Gallart, J., Gracia, E., Muñoz, J.A., Somoza, L., Vázquez, J.T., Vilas, F., 2006. Geological characterization of the prestige sinking area. *Mar. Pollut. Bull.* 53, 208–219.
- Ercilla, G., García-Gil, S., Estrada, F., Gràcia, E., Vizcaíno, A., Vázquez, T., Díaz, S., Vilas, F., Casas, D., Alonso, B., Dañobeitia, J., Farrán, M., 2008a. High resolution seismic stratigraphy of the Galicia Bank region and neighbouring abyssal plains (NW Iberian continental margin). *Mar. Geol.* 249, 108–127.
- Ercilla, G., Casas, D., Estrada, F., Vázquez, J.T., Iglesias, J., García, M., Gómez, M., Acosta, J., Gallart, J., Maestro-González, A., Team, Marconi, 2008b. Morpho-sedimentary features and recent depositional architectural model of the Cantabrian continental margin. *Mar. Geol.* 247, 61–83.
- Ercilla, G., Casas, D., Vázquez, J.T., Iglesias, J., Somoza, L., Juan, C., Medialdea, T., León, R., Estrada, F., García-Gil, S., Farrán, M., Bohoyo, F., García, M., Maestro, A., ERGAP Project and Cruise Teams, 2011. Imaging the recent sediment dynamics of the Galicia Bank region (Atlantic, NW Iberian Peninsula). *Mar. Geophys. Res.* 32, 99–126.
- Ercilla, G., Juan, C., Hernández-Molina, F.J., Bruno, M., Estrada, F., Alonso, B., Casas, D., Farran, M., Llave, E., García, M., Vázquez, J.T., D'Acromont, E., Gorini, C., Palomino, D., Valencia, J., El Moumni, B., Ammar, A., 2016. Significance of bottom currents in deep-sea morphodynamics: an example from the Alboran Sea. *Mar. Geol.* 378, 157–170.
- Faugères, J.C., Imbert, P., Mézerais, M.L., Crémer, M., 1998. Seismic patterns of a muddy contourite fan (Vema Channel, south Brazilian Basin) and a sandy distal turbidite deep-sea fan (cap ferret system, Bay of Biscay): a comparison. *Mar. Geol.* 115, 81–110.
- Faugères, J.C., Stow, D.A.V., Imbert, P., Viana, A.R., 1999. Seismic features diagnostic of contourite drifts. *Mar. Geol.* 162, 1–38.
- Fischer, D., Mogollón, J.M., Strasser, M., Pape, T., Bohrmann, G., Fekete, N., Spiess, V., Kasten, S., 2013. Subduction zone earthquake as potential trigger of submarine hydrocarbon seepage. *Nat. Geosci.* 6, 647–651.
- Fiúza, A.F.G., Hamann, M., Ambar, I., Díaz del Río, G., González, N., Cabanas, J.M., 1998. Water masses and their circulation off western Iberia during May 1993. *Deep-Sea Res.* 1 Oceanogr. Res. Pap. 45, 1127–1160.
- Flood, R.D., Piper, D.J.W., Klaus, A., et al., 1995. Proceedings of the ocean drilling program. *Initial Report Leg. Vol. 155*. ODP, College Station, Texas (1233 pp.).
- Flor, G., 1983. Las rasas asturianas: ensayos de correlación y emplazamiento. *Trab. Geol.* 13, 65–82.
- Friedrich, T., Timmermann, A., Stichel, T., Pahnke, K., 2014. Ocean circulation reconstructions from eNd: a model-based feasibility study. *Paleoceanography* 29. <https://doi.org/10.1002/2014PA002658>.
- Gallastegui, J., 2000. Estructura cortical de la cordillera y margen continental cantábricos: perfiles ESCI-N. *Trabajos de Geología de la Universidad de Oviedo* 22, 9–231.
- Gallastegui, J., Pulgar, J.A., Gallart, J., 2002. Initiation of an active margin at the north Iberian continent-ocean transition. *Tectonics* 21, 15.1–15.13.
- García, M., Hernández-Molina, F.J., Llave, E., Stow, D.A.V., León, R., Fernández-Puga, M.C., Díaz del Río, V., Somoza, L., 2009. Contourite erosive features caused by the Mediterranean outflow water in the Gulf of Cadiz: quaternary tectonic and oceanographic implications. *Mar. Geol.* 257 (24–10).
- García-Lafuente, J., Díaz-del-Río, G., Sánchez-Berrolcal, C., 2006. Vertical structure and bottom-intensification of tidal currents off northwestern Spain. *J. Mar. Syst.* 62, 55–70.
- Gaudin, M., Mulder, T., Cirac, P., Berné, S., Imbert, P., 2007. Past and present sedimentary activity in the Capbreton canyon, southern Bay of Biscay. *Geo-Mar. Lett.* 26, 331–345.
- Gee, M.J.R., Gawthorpe, R.L., Bakke, K., Friedmann, S.J., 2007. Seismic geomorphology and evolution of submarine channels from the Angolan Continental Margin. *J. Sediment. Res.* 77, 433–446.
- Giner, J., González-Casado, J.M., Muñoz-Martín, A., De Vicente, G., 1999. Recent Tectonic in the Northwest Región of the Iberian Peninsula, an Intraplate Deformation. 11th Meeting of the 'Association of European Geological Societies' MAEGS-11. *European Palaeogeography and Geodynamics, A multidisciplinary approach. Abstracts*, p. 17.
- Gómez-Ballesteros, M., Druet, M., Muñoz, A., Arrese, B., Rivera, J., Sánchez, F., Cristobo, J., Parra, S., García-Alegre, A., González-Pola, C., Gallastegui, J., Acosta, J., 2013. Geomorphology of the Avilés canyon system, Cantabrian Sea (Bay of Biscay). *Deep-Sea Res. II Top. Stud. Oceanogr.* 106, 99–117.
- Gonthier, E., Cirac, P., Faugères, J.C., Gaudin, M., Crémer, M., Bourillet, F.F., 2006. Instabilities and deformation in the sedimentary cover on the upper slope of the southern Aquitaine continental margin, north of the Capbreton canyon (Bay of Biscay). *Sci. Mar.* 70 (S1), 89–100.
- González-Casado, J.M., Giner, J., 2000. Relaciones entre fallas y sismicidad en el noreste peninsular. *Geogaceta* 28, 71–74.
- González-Pola, C., 2006. Variabilidad Climática en la Región Sureste del Golfo de Vizcaya. Ph.D. Thesis, Univ. Oviedo, Spain, p. 192.
- Gould, W.J., Hendry, R., Huppert, H.E., 1981. An abyssal topography experiment. *Deep Sea Res. Part A Oceanographic Research Papers* 28, 409–440.
- Gràcia, E., Dañobeitia, J., Vergés, J., Team, P.A.R.S.I.F.A.L., 2003. Mapping active faults offshore Portugal (36°N–38°N): implications for seismic hazard assessment along the southwest Iberian margin. *Geology* 31 (1), 83–86.
- Groeneweg, R.M., Hodgson, D.M., Prelat, A., Luthi, S.M., Flint, S.S., 2010. Flow-deposit interaction in submarine lobes: insights from outcrop observations and realizations of a process-based numerical model. *J. Sediment. Res.* 80, 252–267.
- Groupe Cybere, 1984. La marge déformée du Nord Ouest de l'Espagne. Campagne Cybere du submersible Cyana (Août 1982). Résultats des Campagnes à la Mer 29. CNEXO, Paris, p. 135.
- Groupe Galice, 1979. The continental margin off Galicia and Portugal: acoustical stratigraphy, dredge stratigraphy and structural evolution. In: Sibuet, J.C., Ryan, W.B.F., et al. (Eds.), *Initial Reports Deep Sea Drilling Project*, Washington D.C., 47B, 2, pp. 633–662.
- Gruetzner, J., Uenzelmann-Neben, G., Franke, D., 2012. Variations in sediment transport at the central Argentine continental margin during the Cenozoic. *Geochim. Geophys. Geosyst.* 13 (10) (Q10003).
- Gutiérrez-Claverol, M., López-Fernández, C., Alonso, J.L., 2006. Procesos neotectónicos en los depósitos de rasa en la zona de Cane-ro (Occidente de Asturias). *Geogaceta* 40, 75–78.
- Hanebuth, T.J.J., Zhang, W., Hofmann, A.L., Löwemarker, L.A., Schwenk, T., 2015. Oceanic density fronts steering bottom-current induced sedimentation deduced from a 50 ka contourite-drift record and numerical modeling (off NW Spain). *Quat. Sci. Rev.* 112, 207–225.
- Harris, P.T., McMillan-Lawler, M., Rupp, J., Baker, E.K., 2014. Geomorphology of the oceans. *Mar. Geol.* 352, 4–24.
- Hasiotis, T., Papatheodorou, G., Kastanos, N., Ferentinos, G., 1996. A pockmark bed in the Patras Gulf (Greece) and its activation during the 4/7/93 seismic event. *Mar. Geol.* 130, 333–344.
- Hernández-Molina, F.J., 1993. Dinámica sedimentaria y evolución durante el Pleistoceno terminal-Holoceno del margen noroccidental del Mar de Alborán. Modelo de estratigrafía secuencial de muy alta resolución en plataformas continentales. (Ph.D. Thesis). Univ. Granada, Spain (617 pp.).
- Hernández-Molina, F.J., Llave, E., Somoza, L., Fernández-Puga, M.C., Maestro, A., León, R., Barnolas, A., Medialdea, T., García, M., Vázquez, J.T., Díaz del Río, V., Fernández-Salas, L.M., Lobo, F., Alveirinho Dias, J.M., Roderio, J., Gardner, J., 2003. Looking for clues to paleoceanographic imprints: a diagnosis of the gulf of Cádiz contourite depositional systems. *Geology* 31 (1), 19–22.
- Hernández-Molina, F.J., Llave, E., Stow, D.A.V., García, M., Somoza, L., Vázquez, J.T., Lobo, F.J., Maestro, A., Díaz del Río, V., León, R., Medialdea, T., Gardner, J., 2006a. The contourite depositional system of the Gulf of Cadiz: a sedimentary model related to the bottom current activity of the Mediterranean outflow water and its interaction with the continental margin. *Deep Sea Res., Part II* 53, 1420–1463.
- Hernández-Molina, F.J., Larer, R.D., Rebeco, M., Maldonado, A., 2006b. Miocene reversal of bottom water flow along the Pacific Margin of the Antarctic Peninsula: stratigraphic evidence from a contourite sedimentary tail. *Mar. Geol.* 228, 93–116.
- Hernández-Molina, F.J., Llave, E., Ercilla, G., Maestro, A., Medialdea, T., Ferrin, A., Somoza, L., Gràcia, E., Masson, D.G., García, M., Vizcaíno, A., León, R., 2008. Recent sedimentary processes in the Prestige site area (Galicia Bank, NW Iberian Margin) evidenced by high-resolution marine geophysical methods. *Mar. Geol.* 249 (1–2), 21–45.
- Hernández-Molina, F.J., Matias, H., Llave, E., Stow, D.A.V., 2009a. Onset of contourite deposition in the Gulf of Cádiz: preliminary results. 6th Symposium on the Atlantic Iberian Margin (MIA 09), Oviedo, Abstracts Volume, pp. 277–280.
- Hernández-Molina, F.J., Paterlini, M., Violante, R., Marshall, P., de Isasi, M., Somoza, L., Rebeco, M., 2009b. Contourite depositional system on the argentine slope: an exceptional record of the influence of Antarctic water masses. *Geology* 37, 507–510.
- Hernández-Molina, F.J., Serra, N., Stow, D.A.V., Ercilla, G., Llave, E., Van Rooij, D., 2011. Along-slope oceanographic processes and sedimentary products around Iberia. In: Hernández-Molina, F.J., Stow, D.A.V., Llave, E., et al. (Eds.), *Deep Water Circulation: Processes and Products*. *Geo-Marine Letters*, 31, pp. 315–341.

- Hernández-Molina, F.J., Sierro, F.J., Llave, E., Roque, C., Stow, D.A.V., Williams, T., Lofi, J., Van der Schee, M., Arnáiz, A., Ledesma, S., Rosales, C., Rodríguez-Tovar, F.J., Pardo-Igúzquiza, E., Brackenkridge, R.E., 2016a. Evolution of the Gulf of Cadiz margin and Southwest Portugal contourite depositional system: tectonic, sedimentary and paleoceanographic implications from IODP expedition 339. *Mar. Geol.* 377, 7–39.
- Hernández-Molina, F.J., Wählin, A., Bruno, M., Ercilla, G., Llave, E., Serra, N., Roson, G., Puig, P., Rebesco, M., Van Rooij, D., Roque, C., González-Pola, C., Sánchez, F., Gómez, M., Preu, B., Schwenk, T., Hanebuth, T.J.J., Sánchez-Leal, R.F., García-Lafuente, J., Brackenkridge, R.E., Juan, C., Stow, D.A.V., Sánchez-González, J.M., 2016b. Oceanographic processes and products around the Iberian margin: a new multidisciplinary approach. *Mar. Geol., Special Issue: The Contourite Log-Book 378*, 127–156.
- Hernández-Molina, F.J., Soto, M., Piola, A.R., Tomasini, J., Preu, B., Thompson, P., Badalini, G., Creaser, A., Violante, R.A., Morales, E., Paterlini, M., De Santa Ana, H., 2016c. A contourite depositional system along the Uruguayan continental margin: sedimentary, oceanographic and paleoceanographic implications. *Mar. Geol., Special Issue: The Contourite Log-Book 378*, 333–349.
- Hieke, W., 2004. The August 27, 1886 earthquake in Messenia (Peloponnesus) and reported flames over the Ionian Sea—a Mediterranean ridge gas escape event? *Mar. Geol.* 207, 259–265.
- Hosegood, P., van Haren, H., 2004. Near-bed solibores over the continental shelf in the Faeroe–Shetland channel. *Deep-Sea Res.* 51 (25), 2943–2971.
- Hovland, M., Judd, A., 1988. Seabed Pockmarks and Seepages: Impact on Geology, Biology, and the Marine Environment. Vol. 293. Graham and Trotman, London (565 pp.).
- Howe, J.N.W., Piotrowski, A.M., Noble, T.L., Mulitza, S., Chiessi, C.M., Bayon, G., 2016. North Atlantic deep water production during the last glacial maximum. *Nat. Commun.* 7, 11765.
- Hughes Clarke, J.E., Mayer, L.A., Wells, D.E., 1996. Shallow-water imaging multibeam sonars: a new tool for investigating seafloor processes in the coastal zone and on the continental shelf. *Mar. Geophys. Res.* 18 (6), 607–629.
- Hunt, J.E., Wynn, R.B., Masson, D.G., Talling, P.J., Teagle, D.A.H., 2011. Sedimentological and geochemical evidence for multistage failure of volcanic island landslides: a case study from Icod landslide on North Tenerife. *Geochem. Geophys. Geosyst.* 12 (Q12007).
- Iglesias, J., 2009. Sedimentation on the Cantabrian Continental Margin From Late Oligocene to Quaternary. Univ. Vigo, Spain (Ph.D. Thesis, 185 pp.).
- Iorga, M., Lozier, M.S., 1999. Signatures of the Mediterranean outflow from a North Atlantic climatology: salinity and density fields. *J. Geophys. Res.* 194, 25985–26029.
- Jackson, C.R. (Ed.), 2004. An Atlas of Internal Solitary-like Waves and their Properties (Global Ocean Associates).
- Jackson, D.R., Briggs, K.B., 1992. High-frequency bottom backscattering: roughness versus sediment volume scattering. *J. Acoust. Soc. Am.* 92 (2), 962–977.
- Jané, G., Maestro, A., Ercilla, G., López-Martínez, J., De Andrés, J.R., Casas, D., González-Aller, D., Catalán-Morollón, M., 2010. Occurrence of pockmarks on the Ortelgal Spur continental margin, Northwestern Iberian Peninsula. *Mar. Pet. Geol.* 27, 1551–1564.
- Kane, I., Hodgson, D., 2011. Sedimentological criteria to differentiate submarine channel levee subenvironments: exhumed examples from the Rosario Fm. (upper cretaceous) of Baja California, Mexico, and the Fort Brown Fm. (Permian), Karoo Basin, S. Africa. *Mar. Pet. Geol.* 28, 807–823.
- Kane, I.A., McCaffrey, W.D., Peakall, J., 2008. Controls on sinuosity evolution within submarine channels. *Geology* 36, 287–290.
- Kane, I.A., Pontén, A.S.M., Vangdal, B., Eggenhuisen, J.T., Hodgson, D.M., Sychala, Y.T., 2017. The stratigraphic record and processes of turbidity current transformation across deep-marine lobes. *Sedimentology* 64, 1236–1273.
- Keigwin, L.D., 2004. Radiocarbon and stable isotope constraints on Last Glacial Maximum and Younger Dryas ventilation in the western North Atlantic. *Paleoceanography* 19 (PA4012).
- Kenyon, N.H., Belderson, R.H., Stride, A.H., 1978. Channels, canyons and slump folds on the continental slope between south-West Ireland and Spain. *Oceanol. Acta* 1 (3), 369–380.
- Kenyon, N.H., Klauke, I., Millington, J., Ivanov, M.K., 2002. Sandy submarine canyon-mouth lobes on the western margin of Corsica and Sardinia, Mediterranean Sea. *Mar. Geol.* 184, 69–84.
- Klauke, I., Hesse, R., Ryan, W.B.F., 1997. Flow parameters of turbidity currents in a low-sinuosity, giant deep-sea channel. *Sedimentology* 44, 1093–1102.
- Klauke, I., Masson, D.G., Kenyon, N.H., Gardner, J.V., 2004. Sedimentary processes of the lower Monterey Fan channel and channel-mouth lobe. *Mar. Geol.* 206, 181–198.
- Krahmann, G., Brandt, P., Klaeschen, D., Reston, T., 2008. Mid-depth internal wave energy off the Iberian Peninsula estimated from seismic reflection data. *J. Geophys. Res.* Oceans 113 (C12).
- Kuscu, I., Halbach, P., Inthorn, M., Kuhn, T., Seifert, R., 2008. The R/V Meteor Cruise Leg M44/1 in February 1999 in the sea of Marmara: the first multibeam bathymetric study and analysis of methane in sediment and water columns. *Turk. J. Earth Sci.* 17, 461–480.
- Lamb, K.G., 2014. Internal wave breaking and dissipation mechanisms on the continental slope/shelf. In: Davis, S.H., Moin, P. (Eds.), *Annual Review of Fluid Mechanics*, Vol. 46. Annual Review of Fluid Mechanics. Annual Reviews, Palo Alto, pp. 231–254.
- Loughton, A.S., 1960. An interplain deep-sea channel system. *Deep-Sea Res.* 7, 75–88.
- Loughton, A.S., Searle, R.C., 1979. Tectonic processes on slow spreading ridges. In: Talwani, M., Harrison, C.G., Hayes, D.E. (Eds.), *Deep Drilling Results in the Atlantic Ocean: Ocean Crust: Washington (Am. Geophys. Union)*, Maurice Ewing Series 2, pp. 15–32.
- Lebreiro, S.M., McCave, I.N., Weaver, P.P.E., 1997. Late quaternary emplacement of turbidites on the horseshoe abyssal plain. *J. Sediment. Res.* 67, 856–870.
- Lee, S.H., Chough, S.K., Back, G.G., Kim, Y.B., 2002. Chirp (2–7-kHz) echo characters of the South Korea plateau, East Sea: Syles of mass movement and sediment gravity flow. *Mar. Geol.* 184, 227–247.
- Legg, S., Briegleb, B., Chang, Y., Chassignet, E.P., Danabasoglu, G., Ezer, T., Gordon, A.L., Griffies, S., Hallberg, R., Jackson, L., Large, W., Özgökmen, T.M., Peters, H., Price, J., Riemenschneider, U., Wu, W., Xu, X., Yang, J., 2009. Improving oceanic overflow representation in climate models. The gravity current entrainment climate process team. *Bull. Am. Meteorol. Soc.* 657–670.
- Lindhorst, K., Gruen, M., Krastel, S., Schwenk, T., 2012. Hydroacoustic analysis of mass wasting deposits in Lake Ohrid (FYR Macedonia/Albania). In: Yamada, Y., Kawamura, K., Ikehara, K., Ogawa, Y., Urgeles, R., Mosher, D., Chaytor, J., Strasser, M. (Eds.), *Submarine Mass Movements and Their Consequences*, Springer, Netherlands, pp. 245–253.
- Llave, E., Schönfeld, J., Hernández-Molina, F.J., Mulder, T., Somoza, L., Díaz del Río, V., Sánchez-Almazo, I., 2006. High-resolution stratigraphy of the Mediterranean outflow contourite system in the Gulf of Cadiz during the late Pleistocene: the impact of Heinrich events. *Mar. Geol.* 227, 241–262.
- Llave, E., García, M., Pérez, C., Sayago, M., Farrán, M., Ercilla, G., Somoza, L., León, R., Maestro, A., Medialdea, T., Hernández-Molina, F.J., Álvarez, R., Durán, R., Mohamed, K., 2008. Morphological feature analyses of the prestige half-graben on the SW Galicia Bank. *Mar. Geol.* 249, 7–20.
- Llave, E., Matias, H., Hernández-Molina, F.J., Ercilla, G., Stow, D.A.V., Medialdea, T., 2010. Pliocene and quaternary seismic stacking pattern and distribution of contourites in the Algarve margin (Northern Gulf of Cadiz, Spain). *Geo-Temas* 11, 103–104.
- Llave, E., Matias, H., Hernández-Molina, F.J., Ercilla, G., Stow, D.A.V., Medialdea, T., 2011. Pliocene and quaternary seismic stacking pattern and distribution of contourites in the Algarve margin (Northern Gulf of Cadiz, Spain). *Geo-Mar Letters* 31 (5–6), 377–390.
- Lomtev, V.L., Gurinov, M.G., 2009. The tectonic conditions of the august 2, 2007, M ~ 6.1 Nevsk earthquake. *Russian J. Pac. Geol.* 3 (5), 450–459.
- Lonsdale, P., 1977. Structural geomorphology of a fast-spreading rise crest: the East Pacific rise near 3°25'S. *Mar. Geophys. Res.* 3, 251–293.
- López-Fernández, C., Pulgar, J.A., Gallart, J., González-Cortina, J.M., Díaz, J., Ruíz, M., 2002. Actividad sísmica reciente en el noroeste de la Península Ibérica. 3ª Asamblea Hispano Portuguesa de Geodesia y Geofísica, Valencia.
- Maestro, A., Bohoyo, F., López-Martínez, J., Acosta, J., Gómez-Ballesteros, M., Llave, E., Muñoz, A., Terrinha, P.G., Dominguez, M., Fernández, F., 2015. Influencia de los procesos tectónicos y volcánicos en la morfología de los márgenes continentales ibéricos. *Bol. Geol. Min.* 126 (2–3), 427–482.
- Maestro, A., Jané, G., Llave, E., López-Martínez, J., Bohoyo, F., Druet, M., 2017. The role of tectonic inheritance in the morphostructural evolution of the Galicia continental margin and adjacent abyssal plains from digital bathymetric model (DBM) analysis (NW Spain). *Int. J. Earth Sci.* <https://doi.org/10.1007/s00531-017-1531-4>.
- Maier, K., Fildani, A., Paull, C., McHargue, T., Graham, S., Caress, D., 2013. Deep-sea channel evolution and stratigraphic architecture from inception to abandonment from high-resolution Autonomous Underwater Vehicle surveys offshore central California. *Sedimentology* 60, 935–960.
- Marches, E., Mulder, T., Gonthier, E., Cremer, M., Hanquiez, V., Garlan, T., Lecroart, R., 2010. Perched lobe formation in the Gulf of Cadiz: interactions between gravity processes and contour currents (Algarve Margin, Southern Portugal). *Sediment. Geol.* 229, 81–94.
- Martín-González, F., Insua, J.M., Martínez-Díaz, J., Giner, J., 2010. Estudio preliminar de las fallas sismogénicas en el NO de la Península Ibérica. Resúmenes de la 1ª Reunión Ibérica sobre Fallas Activas y Paleosismología. Sigüenza, Spain, pp. 57–60.
- Mary, G., 1983. Evolución del margen costero de la cordillera Cantábrica en Asturias desde el Mioceno. *Trabajos de Geología, Universidad de Oviedo* 13, 3–35.
- Maslin, M.A., Shackleton, N.J., Pflaumann, U., 1995. Surface water temperature, salinity, and density changes in the northeast Atlantic during the last 45,000 years: Heinrich events, deep water formation, and climatic rebounds. *Paleoceanography* 10 (3), 527–544.
- Masson, D.G., Huvenne, V.A.I., Stigter, H.C., Arzola, R.G., LeBas, T.P., 2011. Sedimentary processes in the middle Nazare canyon. *Deep-Sea Res.* 58, 2369–2387.
- Mayall, M., Stewart, I., 2000. The architecture of turbidite slope channels. In: Weimer, P., Slatt, R.M., Coleman, J., Rosen, N.C., Nelson, H., Bouma, A.H., Styzen, M.J., Lawrence, D.T. (Eds.), *Deep-Water Reservoirs of the World*. Gulf Coast Section SEPM 20th Bob F. Perkins Research Conference, pp. 578–586.
- Mazé, J.P., Arhan, M., Mercier, H., 1997. Volume budget of the eastern boundary layer off the Iberian Peninsula. *Deep-Sea Res.* 44, 1543–1574.
- Mc Clennen, C.E., 1989. Microtopography a surficial sediment patterns in the Central Gulf of Maine: a 3.5 kHz survey and interpretation. *Mar. Geol.* 89, 69–85.
- McCave, I.N., 1982. Erosion and deposition by currents on submarine slopes. *Bull. Inst. Géol. Bassin d'Aquitaine* 31, 47–55.
- McCave, I.N., 2001. Nepheloid layers. In: Steele, J.H., Thorpe, S.A., Turekian, K.K. (Eds.), *Encyclopedia of Ocean Sciences* 4. Academic Press, London, pp. 1861–1870.
- Medialdea, T., Somoza, L., Bohoyo, F., Vázquez, J.T., Vegas, R., Patriat, M., Thinon, I., Ercilla, G., León, R., 2009. Compresión cenozoica en la Cuenca de la Unión y Monte Submarino Armoricano (Llanura Abisal de Vizcaya). 6º Simposio sobre el Margen Ibérico Atlántico, MIA09, 1–5 de Diciembre de 2009, Oviedo.
- Mercier de Lépinay, M., Loncke, L., Basile, C., Roest, W.R., Patriat, M., Maillard, A., De Clarens, P., 2016. Transform continental margins—part 2: a worldwide review. *Tectonophysics* 693, 96–115.
- Migeon, S., Ducassou, E., Le Gonidec, Y., Rouillard, P., Mascle, J., Revel-Rolland, M., 2010. Lobe construction and sand/mud segregation by turbidity currents and debris flows on the western Nile deep-sea fan (eastern Mediterranean). *Sediment. Geol.* 229, 124–143.
- Mojtahid, M., Eynaud, F., Zaragosi, S., Scourse, J., Bourillet, J.F., Garlan, T., 2005. Palaeoclimatology and palaeohydrography of the glacial stages on Celtic and Armorican margins over the last 360,000 yrs. *Mar. Geol.* 224, 57–82.
- Montadert, L., Roberts, D.G., Auffret, G.A., Bock, W.D., Dupeuble, P.A., Hailwood, E.A., Harrison, W.E., Kagami, H., Lumsden, D.N., Müller, C.M., Schnitker, D., Thompson, T.L., Timofeev, P.P., 1979. Annotated Record of the Detailed Examination of Mn Deposits from DSDP Leg 48 (Holes 400A, 401, 402). Initial Reports of the Deep Sea Drilling Project. U. S. Government Printing Office, XLVIII (1183 pp.).

- Morris, E.A., Hodgson, D.M., Flint, S.S., Brunt, R.L., Butterworth, P.J., Verhaeghe, J., 2014. Sedimentology, stratigraphic architecture and depositional context of submarine frontal lobe complexes. *J. Sediment. Res.* 84, 763–780.
- Mulder, T., 2011. Gravity processes on continental slope, rise and abyssal plains. In: Huneke, H., Mulder, T. (Eds.), *Deep-sea Sediments. Developments in Sedimentology*, 63. Elsevier, Amsterdam, pp. 25–148.
- Mulder, T., Weber, O., Abschutz, P., Jorissen, J., Jouanneau, M., 2001. A few months old storm generated turbidite deposited in the Capbreton canyons (Bay of Biscay, SW France). *Geo-Mar. Lett.* 21, 149–156.
- Mulder, T., Faugères, J.-C., Gonthier, E., 2008. Chapter 21 mixed turbidite–contourite systems. *Dev. Sedimentol.* 60. [https://doi.org/10.1016/S0070-4571\(08\)10021-8](https://doi.org/10.1016/S0070-4571(08)10021-8).
- Mullins, H.T., Bordman, M.R., Neumann, A.C., 1979. Echo character of off-platform carbonates. *Mar. Geol.* 32, 251–268.
- Mutti, E., Normark, W.R., 1987. Comparing examples of modern and ancient turbidite systems: problems and concepts. In: Leggett, J.R., Zuffa, G.G. (Eds.), *Marine Clastic Sedimentology: Concepts and Case Studies*. G. Graham and Trotman, London, pp. 1–37.
- Mutti, E., Bernoulli, D., Lucchi, F.R., Tinetti, R., 2009. Turbidites and turbidity currents from Alpine ‘flysch’ to the exploration of continental margins. *Sedimentology* 56, 267–318.
- Nikonov, A.A., 2002. The Crimean earthquakes of 1927: unknown phenomena at sea. *Priroda* 9, 13–20.
- Normark, W.R., Posamentier, H., Mutti, E., 1993. Turbidite systems: state of the art and future directions. *Rev. Geophys.* 31, 91–116.
- Ortiz-Karpp, A., Hodgson, D.M., McCaffrey, W.D., 2015. The role of mass-transport complexes in controlling channel avulsion and the subsequent sediment dispersal patterns on an active margin: the Magdalena fan, offshore Colombia. *Mar. Pet. Geol.* 64, 58–75.
- Paillet, J., Mercier, H., 1997. An inverse model of the eastern North Atlantic general circulation and thermocline ventilation. *Deep Sea Res., Part I* 44, 1293–1328.
- Paillet, J., Arhan, M., MacCartney, M.S., 1998. The spreading of Labrador Sea water in the eastern North Atlantic. *J. Geophys. Res.* 103, 10223–10239.
- Paillet, J., Le Cann, B., Serpette, A., Morel, Y., Carton, X., 1999. Real-time tracking of a Galician Meddy. *Geophys. Res. Lett.* 26 (13), 1877–1880.
- Paillet, J., Le Cann, B., Carton, X., Morel, Y., Serpette, A., 2002. Dynamics and evolution of a northern meddy. *J. Phys. Oceanogr.* 32 (1), 55–79.
- Peakall, J., Kane, I.A., Masson, D.G., Keevil, G., McCaffrey, W.D., Corney, R., 2012. Global (latitudinal) variation in submarine channel sinuosity. *Geology* 40, 11–14.
- Pérez, F.F., Castro, C.G., Álvarez-Salgado, X.A., Ríos, A.F., 2001. Coupling between the Iberian basin-scale circulation and the Portugal boundary current system. A chemical study. *Deep Sea Research* 48, 1519–1533.
- Pickering, K.T., Hiscott, R.N., 2016. *Deep Marine Systems: Processes, Deposits, Environments, Tectonics and Sedimentation*. American Geophysical Union and Wiley (672 pp.).
- Pingree, R.D., Le Cann, B., 1989. Celtic and Armerican slope and shelf residual currents. *Progress in Oceanography* 23, 303–338.
- Pingree, R.D., Le Cann, B., 1990. Structure, strength and seasonality of the slope currents in the Bay of Biscay region. *J. Mar. Biol. Assoc. UK* 70, 857–885.
- Piper, D.J.W., Normark, W.R., 1982. Acoustic interpretation of Quaternary sedimentation and erosion on the channelled upper Laurentian Fan, Atlantic margin of Canada. *Can. J. Earth Sci.* 19, 1974–1984.
- Pollard, R.T., Griffiths, M.J., Cunningham, S.A., Read, J.F., Pérez, F.F., Ríos, A.F., 1996. Vivaldi 1991 – a study of the formation, circulation and ventilation of eastern North Atlantic Central Water. *Prog. Oceanogr.* 37, 167–192.
- Pomar, L., Morsilli, M., Hallock, P., Bádenas, B., 2012. Internal waves, an under-explored source of turbulence events in the sedimentary record. *Earth Sci. Rev.* 111 (1–2), 56–81.
- Posamentier, H.W., Kolla, V., 2003. Seismic geomorphology and stratigraphy of depositional elements in deep-water settings. *J. Sediment. Res.* 73 (3), 367–388.
- Posamentier, H.W., Mezarwin, Wisman, P.S., Plawman, T., 2000. Deep water depositional systems–Ultra-deep Makassar Strait, Indonesia. In: Weimer, P., Slatt, R.M., Coleman, J., Rosen, N.C., Nelson, H., Bouma, A.H., Styzen, M.J., Lawrence, D.T. (Eds.), *Deep-Water Reservoirs of the World: Gulf Coast Society of the Society of Economic Paleontologists and Mineralogists Foundation, 20th Annual Research Conference*, pp. 806–816.
- Pratson, L.F., Laine, E.P., 1989. The relative importance of gravity-induced versus current-controlled sedimentation during the Quaternary along the Mideast U.S. outer continental margin revealed by 3.5 kHz echo character. *Mar. Geol.* 89, 87–126.
- Prélat, A., Hodgson, D.M., Flint, S.S., 2009. Evolution, architecture and hierarchy of distributary deep-water deposits: a high-resolution outcrop investigation from the Permian Karoo Basin, South Africa. *Sedimentology* 56 (2132–U25).
- Prélat, A., Covault, J.A., Hodgson, D.M., Fildani, A., Flint, S.S., 2010. Intrinsic controls on the range of volumes, morphologies, and dimensions of submarine lobes. *Sediment. Geol.* 232 (1), 66–76.
- Preu, B., Hernández Molina, F.J., Vilonate, R., Piola, A.R., Paterlini, C.M., Schwenk, T., Voigt, I., Krastel, S., Spiess, V., 2013. Morphosedimentary and hydrographic features of the northern Argentine margin: the interplay between erosive, depositional and gravitational processes and its conceptual implications. *Deep Sea Res., Part I* 75, 157–174.
- Prieto, E., González-Pola, C., Lavín, A., Sánchez, R.F., Ruiz-Villarreal, M., 2013. Seasonality of intermediate waters hydrography west of the Iberian Peninsula from an 8 yr semi-annual time series of an oceanographic section. *Ocean Sci.* 9 (2), 411–429.
- Pudsey, C.J., Howe, J.A., 1998. Quaternary history of the Antarctic Circumpolar Current: evidence from the Scotia Sea. *Mar. Geol.* 148, 83–112.
- Quaresma, L.S., Pichon, A., 2013. Modelling the barotropic tide along the west-Iberian margin. *J. Mar. Syst.* 109–110, S3–S25.
- Rebesco, M., 2005. Contourites. In: Richard, C., Selley, R.C., Cocks, L.R.M., Plimer, I.R. (Eds.), *Encyclopedia of Geology* 4. Elsevier, London, pp. 513–527.
- Rebesco, M., Camerlenghi, A., 2008. *Contourites. Developments in Sedimentology*. vol. 60. Elsevier (688 pp.).
- Rebesco, M., Camerlenghi, A., VanLoon, A.J., 2008. Contourite research: a field in full development. In: Rebesco, M., Camerlenghi, A. (Eds.), *Contourites*. Elsevier, Amsterdam, pp. 1–10.
- Rebesco, M., Hernández-Molina, F.J., Van Rooij, D., Wählin, A., 2014. Contourites and associated sediments controlled by deep-water circulation processes: state of the art and future considerations. *Mar. Geol.* 352, 111–154.
- Roberts, D.G., Hogg, N.G., Derek, G., Bishop, G., Flewelling, C.G., 1974. Sediment distribution around moated seamounts in the Rockall trough. *Deep Sea Res. Part A Oceanographic Research Papers* 21, 175–184.
- Rodríguez-Fernández, L.R., López-Olmedo, F., Oliveira, J.T., Medialdea, T., Terrinha, P., Matas, J., Martín-Serrano, A., Martín-Parra, L.M., Rubio, F., Montes, M., Nozal, F., 2014. Mapa Geológico de España y Portugal a escala 1:1.000.000. Instituto Geológico y Minero de España 978-84-7840-949-5.
- Rodríguez-Tovar, F.J., Dorador, J., Grunert, P., Hodell, D., 2015. Deep-sea trace fossil and benthic foraminiferal assemblages across glacial terminations 1, 2 and 4 at the “Shackleton Site” (IODP Expedition 339, site U1385). *Glob. Planet. Chang.* 133, 359–370.
- Rogerson, M., Rohling, E.J., Bigg, G.R., Ramirez, J., 2012. Paleocirculation of the Atlantic-Mediterranean exchange: overview and first quantitative assessment of climatic forcing. *Rev. Geophys.* 50 RG2003.
- Roque, C., Duarte, H., Terrinha, P., Valadares, V., Noiva, J., Cahão, M., Ferreira, J., Legoinha, P., Zitellini, N., 2012. Pliocene and Quaternary depositional model of the Algarve margin contourite drifts (Gulf of Cadiz, SW Iberia): seismic architecture, tectonic control and paleoceanographic insights. *Mar. Geol.* 303–306, 42–62.
- Roque, C., Hernández-Molina, F.J., Ercilla, G., Casas, D., Quartau, R., Llave, E., Alonso, B., Ferran, M., Mena, A., Francés, G., Cruise Party, M.O.W.E.R., 2015. Slope failure and mass movements in the Sines Contourite Drift (West Portuguese Margin): preliminary results. In: Díaz del Río, V., Bárcenas, P., Fernández-Salas, L.M., López-González, N., Palomino, D., Rueda, J.L., Sánchez-Guillamón, O., Vázquez, J.T. (Eds.), *Volumen de Comunicaciones presentadas en el VIII Simposio sobre el Margen Ibérico Atlántico, MIA15*. Ediciones Sia Graf, Málaga, Spain, pp. 579–582.
- Ruiz-Villarreal, M., González-Pola, C., Otero, P., Díaz del Río, G., Lavín, A., Cabanas, J.M., 2006. Circulation in Galicia–Southern Bay of Biscay: reanalysis of the circulation influencing the prestige oil spill. *European Operational Oceanography: present and future* 342–348.
- Sánchez-González, J.M., 2013. *Oceanographic and Morphosedimentary Features in the North-western Sector of the Galicia Margin: Implications*. MSc Work Univ. Vigo, Spain.
- Sarnthein, M., Thiede, J., Pflaumann, U., 1982. Atmospheric and ocean circulation patterns off Northwest Africa during the past 25 million years. In: von Rad, U., Hinz, K., Sarnthein, M., Seibold, E. (Eds.), *Geology of the Northwest African Continental Margin*. Springer, Berlin, pp. 545–604.
- Sarnthein, M., Karl Statterger, D.D., Erlenkeuser, H., Schulz, M., Seidov, D., Simstich, J., Van Kreveld, S., 2001. Fundamental Modes and Abrupt Changes in North Atlantic Circulation and Climate over the last 60 ky. In: Schafer, P., Ritzrau, W., Scghluter, M., Thiede, J. (Eds.), *The Northern North Atlantic: A Changing Environment*. Springer, Berlin, pp. 365–410.
- Saunders, P.M., 1986. The accuracy of measurement of salinity, oxygen and temperature in the deep ocean. *J. Phys. Oceanogr.* 16, 189–195.
- Schönfeld, J., Zahn, R., 2000. Late glacial to Holocene history of the Mediterranean outflow. Evidence from benthic foraminiferal assemblages and stable isotopes at the Portuguese margin. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 159, 85–111.
- Shanmugam, G., 2006. Deep-water processes and facies models. Implications for Sandstone Petroleum Reservoirs, 5. Elsevier, Amsterdam.
- Shanmugam, G., 2013. Modern internal waves and internal tides along oceanic pycnoclines, challenges and implications for ancient deep-marine baroclinic sands. *AAPG Bull.* 97 (5), 767811.
- Skene, K.I., Piper, D.J.W., 2003. Late Quaternary stratigraphy of Laurentian Fan: a record of events off the eastern Canadian continental margin during the last deglacial period. *Quat. Int.* 99–100, 135–152.
- Skene, K.I., Piper, D.J.W., 2006. Late Cenozoic evolution of Laurentian Fan: development of a glacially-fed submarine fan. *Mar. Geol.* 227 (1–2), 67–92.
- Slater, D.R., 2003. *The Transport of Mediterranean Water in the North Atlantic Ocean*. Univ. Southampton, PhD Thesis.
- Smith, W.H.F., Sandwell, D.T., 1997. Global seafloor topography from satellite altimetry and ship depth soundings. *Science* 277, 1957–1962.
- Spychala, Y.T., Hodgson, D.M., Prélat, A., Kane, I.A., Flint, S.S., Mountney, N.P., 2017. Frontal and lateral submarine lobe fringes: comparing sedimentary facies, architecture and flow processes. *J. Sediment. Res.* 87, 75–96.
- Stevenson, C.J., Talling, P.J., Masson, D.G., Sumner, E.J., Frenz, M., Wynn, R.B., 2014. The spatial and temporal distribution of grain-size breaks in turbidites. *Sedimentology* 61, 1120–1156.
- Stevenson, C.J., Jackson, C.A.L., Hodgson, D.M., Hubbard, S.M., Eggenhuisen, J.T., 2015. Deep-water sediment bypass. *J. Sediment. Res.* 85, 1058–1081.
- Stow, D.A.V., Faugères, J.C., 1998. Contourites, turbidites and processes interaction. *Sediment. Geol., Special Issue* 82, 1–310.
- Stow, D.A.V., Pudsey, C.J., Howe, J.A., Faugères, J.C., Viana, A.R., 2002. Deep-water Contourite Systems: Modern Drifts and Ancient Series, Seismic and Sedimentary Characteristics, Memoirs 22. Geological Society of London, London (464 pp.).
- Stow, D.A.V., Hernández-Molina, F.J., Alvarez Zarikian, C.A., the Expedition 339 Scientists, 2013. *Proceedings IODP Exp. 339*. Tokyo (Integrated Ocean Drilling Program Management International, Inc.). <https://doi.org/10.2204/iodp.proc.339.2013>.
- Sultan, N., Cochonat, P., Canals, M., Cattaneo, A., Dennielou, B., Hafliadason, H., Laberg, J.S., Long, D., Mienert, J., Urgeles, R., Vorren, T., Wilson, C., 2004. Triggering mechanisms of slope instability processes and sediment failures on continental margins: a geotechnical approach. *Mar. Geol.* 213, 291–321.

- Talling, P.J., Wynn, R.B., Masson, D.G., Frenz, M., Cronin, B.T., Schiebel, R., Akhmetzhanov, A.M., Dallmeier-Tiessen, S., Benetti, S., Weaver, P.P.E., Georgiopolou, A., Zuhlsdorff, C., Amy, L.A., 2007. Onset of submarine debris flow deposition far from original giant landslide. *Nature* 450, 541–544.
- Talling, P.J., Wynn, R.B., Rixon, R., Schmidt, D., Sumner, E., Amy, L.A., 2010. How did submarine flows transport boulder sized mud clasts to the fringes of the Mississippi Fan? *J. Sed. Res.* 80, 829–851.
- Talling, P.J., Masson, D.G., Sumner, E.J., Malgesini, G., 2012. Subaqueous sediment density flows: depositional processes and deposit types. *Sedimentology* 59, 1937–2003.
- Terrinha, P., Pinheiro, L.M., Henriët, J.-P., Matias, L., Ivanov, M.K., Monteiro, J.H., Akhmetzhanov, A., Volkonskaya, A., Cunha, T., Shaskin, P., Rovere, M., 2003. Tsunamigenic–seismogenic structures, neotectonics, sedimentary processes and slope instability on the southwest Portuguese margin. *Mar. Geol.* 195 (1–4), 55–73.
- Thinon, I., Fidalgo-González, L., Réhault, J.P., Olivet, J.L., 2001. Pyrenean deformations in the Bay of Biscay. *Comptes Rendus de l'Académie des Sciences, Series IIA: Science Terre Planets* 332, 561–568.
- Toucanne, S., Mulder, T., Schönfeld, J., Hanquiez, V., Gonther, E., Duprat, J., Cremer, M., Zaragosi, S., 2007. Contourites of the Gulf of Cadiz: a high-resolution record of the paleocirculation of the Mediterranean outflow water during the last 50,000 years. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 246, 354–366.
- Toucanne, S., Jouet, G., Ducassou, E., Bassetti, M.A., Dennielou, B., Minto'o, C.M.A., Lahmi, M., Touyet, N., Charlier, K., Lericolais, G., Mulder, T., 2012. A 130,000-year record of Levantine intermediate water flow variability in the Corsica trough, western Mediterranean Sea. *Quat. Sci. Rev.* 33, 55–73.
- Tsunogai, U., Ishibashi, J., Wakita, H., Gamoto, T., 1998. Methane-rich plumes in the Suruga trough (Japan) and their carbon isotopic characterization. *Earth Planet. Sci. Lett.* 160, 97105.
- Unterseh, S., Cochonot, P., Voisset, M., Ollier, G., Harmegnies, F., 1998. Acoustic Data Ground Truthing Through In-situ Measurements of Physical Properties. International Offshore and Polar Engineering Conference (ISOPE). ISOPE, Montreal, Canada. pp. 689–692.
- Uriarte, A., Collins, M., Cearreta, A., Bald, J., Evans, G., 2004. Chapter 5 Sediment supply, transport and deposition: contemporary and Late Quaternary evolution. In: Collins, B.A.M. (Ed.), *Oceanography Series*. Elsevier, pp. 97–131.
- Van Aken, H.M., 2000. The hydrography of the mid-latitude Northeast Atlantic Ocean I: The deep water masses. *Deep Sea Res., Part I* 47, 757–788.
- van Raaphorst, W., Malschaert, H., van Haren, H., Boer, W., Brummer, G.-J., 2001. Cross-slope zonation of erosion and deposition in the Faeroe–Shetland Channel, North Atlantic Ocean. *Deep Sea Res. Part 1 Oceanogr. Res. Pap.* 48(2), 567–591.
- Van Rooij, D., De Mol, B., Huvenne, V., Ivanov, M., Henriët, J.P., 2003. Seismic evidence of current-controlled sedimentation in the Belgica mound province, upper porcupine slope, southwest of Ireland. *Mar. Geol.* 195, 31–53.
- Van Rooij, D., Iglesias, J., Hernández-Molina, F.J., Ercilla, G., Gómez-Ballesteros, M., De Hauwere, A., Casas, D., Llave, E., García-Gil, S., Henriët, J.P., Acosta, J., 2010. The Le Danois contourite depositional system off the north Iberian margin, Spain. *Mar. Geol.* 274, 1–20.
- Vanney, J.R., Auxietre, J.L., Dunand, J.P., 1979. Geomorphic provinces and the evolution of the northwestern Iberian continental margin. *Annales de l'Institut Océanographique Paris* 55, 5–20.
- Varela, R.A., Roson, G., Herrera, J.L., Torres-Lopez, S., Fernandez-Romero, A., 2005. A general view of the hydrographic and dynamical patterns of the Rias Baixas adjacent sea area. *J. Mar. Syst.* 54, 97–113.
- Vargas, C.A., Mann, P., Gómez, C., 2012. Morphologic expression of accretionary processes and recent submarine landslides along the southwestern Pacific margin of Colombia. In: Yamada, Y., Kawamura, K., Ikehara, K., Ogawa, Y., Urgeles, R., Mosher, D., Chaytor, J., Strasser, M. (Eds.), *Submarine Mass Movements and Their Consequences*. Springer, Netherlands, pp. 365–377.
- Vázquez, J.T., Medialdea, T., Ercilla, G., Somoza, L., Estrada, F., Fernández Puga, M.C., Gallart, J., Gràcia, E., Maestro, A., Sayago, M., 2008. Cenozoic deformational structures on the Galicia Bank region (NW Iberian continental margin). *Mar. Geol.* 249, 128–149.
- Vigneaux, M., 1974. The geology and sedimentation history of the Bay of Biscay. In: Nairn, A.E.M., Francis, G., Stehli, F.G. (Eds.), *Ocean Basins and Margins*. Vol. 2. Plenum Press, New York-London, pp. 273–374.
- Voelker, A., Lebreiro, S., Schönfeld, J., Cacho, I., Exlenkenser, H., Abrantes, F., 2006. Mediterranean outflow strengthenings during Northern Hemisphere Coolings: a salt sources for the glacial Atlantic? *Earth Planet. Sci. Lett.* 245, 39–55.
- Voelker, A.H.L., Rodrigues, T., Billups, K., Oppo, D.W., McManus, J.F., Stein, R., Hefter, J., Grimalt, J.O., 2010. Variations in mid-latitude North Atlantic surface water properties during the mid-Brunhes (MIS 9–14) and their implications for the thermohaline circulation. *Climate of the Past* 6 (4), 531–552.
- Vologina, E.G., Sturm, M., Radziminovich, Y.B., Vorob'eva, S.S., Schetnikov, A.A., 2012. The 1912 earthquake in South Baikal: traces in bottom sediments and gas release into the water column. *Russ. Geol. Geophys.* 53, 1342–1350.
- Weaver, P.P.E., 1994. Determination of turbidity current erosional characteristics from reworked coccolith assemblages, Canary Basin, North-east Atlantic. *Sedimentology* 41, 1025–1038.
- Weaver, P.P.E., Thomson, J., 1993. Calculation erosion by deep-sea turbidity currents during initiation and flow. *Nature* 364, 136–138.
- Weaver, P.P.E., Rothwell, R.G., Ebbing, J., Gunn, D., Hunter, P.M., 1992. Correlation, frequency of emplacement and source directions of megaturbidites on the Madeira Abyssal Plain. *Mar. Geol.* 109, 1–20.
- Weaver, P.P.E., Wynn, R.B., Kenyon, N.H., Evans, J., 2000. Continental margin sedimentation, with special reference to the north-East Atlantic margin. *Sedimentology* 47, 239–256.
- Wells, M., Cossu, R., 2013. The possible role of Coriolis forces in structuring large-scale sinuous patterns in submarine channel-levee systems. *Phil. Trans. R. Soc. A* 371, 20120366.
- Wynn, R.B., Stow, D.A.V., 2002. Classification and characterisation of deep-water sediment waves. *Mar. Geol.* 192, 7–22.
- Wynn, R.B., Masson, D.G., Stow, D.A.V., Weaver, P.P., 2000. Turbidity current sediment waves on the submarine slopes of the western Canary Islands. *Mar. Geol.* 163, 185–198.
- Wynn, R.B., Kenyon, N.H., Masson, D.G., Stow, D.A.V., Weaver, P.P.E., 2002. Characterization and recognition of deep-water channel-lobe transition zones: American Association of Petroleum Geologists. *Bulletin* 86, 1441–1462.
- Wynn, R.B., Cronin, B.T., Peakall, J., 2007. Sinuous deep-water channels: genesis, geometry and architecture. *Mar. Pet. Geol.* 24, 341–387.
- Wynn, R.B., Talling, P.J., Masson, D.G., Stevenson, C.J., Cronin, B.T., Le Bas, T.P., 2010. Investigating the timing, processes and deposits of one of the world's largest submarine gravity flows: The “Bed 5 Event” off Northwest Africa. In: Mosher, D.C., Shipp, R.C., Moscardelli, L., Chaytor, J.D., Baxter, C.D.P., Lee, H.J., Urgeles, R. (Eds.), *Submarine Mass Movements and their Consequences: Advances in Natural and Technological Hazards Research*. Springer, Dordrecht, pp. 463–474.
- Zahn, R., Schönfeld, J., Kudrass, H.-R., Park, M.-H., Erlenkeuser, H., Grootes, P., 1997. Thermohaline instability in the North Atlantic during meltwater events: stable isotope and ice-rafted detritus records from core S075-26KL, Portuguese margin. *Paleoceanography* 12, 696–710.
- Zaragosi, S., Auffret, G.A., Faugères, J.C., Garlan, T., Pujol, C., Cortijo, E., 2000. Physiography and recent sediment distribution of the Celtic deep-sea fan, Bay of Biscay. *Mar. Geol.* 169, 207–237.
- Zaragosi, S., Eynaud, F., Pujol, C., Auffret, G.A., Turon, J.L., Garlan, T., 2001a. Initiation of the European deglaciation as recorded in the northwestern Bay of Biscay slope environments (Meriadzek terrace and Trevelyan escarpment): a multi-proxy approach. *Earth Planet. Sci. Lett.* 188, 493–507.
- Zaragosi, S., Le Suave, R., Bourillet, J.F., Auffret, G., Faugères, J.C., Pujol, C., Garlan, T., 2001b. The deep-sea Armorican depositional system (Bay of Biscay), a multiple source, ramp model. *Geo-Mar. Lett.* 20, 219–232.
- Zaragosi, S., Bourillet, J.F., Eynaud, F., Toucanne, S., Denhard, B., Van Toer, A., Lanfume, V., 2006. The impact of the last European deglaciation on the deep-sea turbidite systems of the Celtic-Armorican margin (Bay of Biscay). *Geo-Mar. Lett.* 26, 317–329.
- Zhang, H., King, B., Swinney, H.L., 2008. Resonant generation of internal waves on a model continental slope. *Phys. Rev. Lett.* 100 (24), 244504.
- Zhang, W., Hanebuth, T.J., Stöber, U., 2016. Short-term sediment dynamics on a meso-scale contourite drift (off NW Iberia): impacts of multi-scale oceanographic processes deduced from the analysis of mooring data and numerical modelling. *Mar. Geol.* 378, 81–100.