

# Update and latest advances in antiretroviral therapy

Luis Menéndez-Arias<sup>1</sup> and Rafael Delgado<sup>2</sup>

*1 Centro de Biología Molecular “Severo Ochoa”, Consejo Superior de Investigaciones Científicas & Universidad Autónoma de Madrid, 28049 Madrid, Spain.*

*2 Laboratory of Molecular Microbiology. Instituto de Investigación Hospital 12 de Octubre (Imas12) and The University Complutense School of Medicine, Madrid, Spain.*

*Corresponding authors: E-mail: [lmenendez@cbm.csic.es](mailto:lmenendez@cbm.csic.es), [rafael.delgado@salud.madrid.org](mailto:rafael.delgado@salud.madrid.org)*

## **Abstract**

Since the first cases of AIDS appeared in 1981, human immunodeficiency virus type 1 (HIV-1) infection has reached pandemic proportions. Forty years later, research has led to the approval of more than thirty antiretroviral drugs, while combination therapies have turned HIV-1 infection into a chronic, but manageable disease. Still, drug toxicity and acquired and transmitted drug resistance remain as major threats to therapy success. In this review, we provide an overview on currently available anti-HIV drugs and the latest developments in antiretroviral therapy focused on new antiretroviral agents acting on known and unexploited antiviral targets, prevention therapies aimed to improve available drug combinations, and research on new long-acting therapies, particularly those involving novel drug candidates such as lenacapavir or islatravir.

**Keywords:** HIV, antiretroviral drug, drug resistance, toxicity, combination therapy, long-acting agents

## **Historical perspective and current HIV/AIDS situation**

Forty years ago, in June and July 1981, the Centers for Disease Control and Prevention in the U.S. reported clusters of Kaposi's sarcoma and *Pneumocystis* pneumonia among gay men in California and New York City. Patients were immunosuppressed and affected by a deadly disease whose etiological agent was a retrovirus identified two years later and designated as human immunodeficiency virus (HIV) [1].

Although azidothymidine (3'-azido-3'-deoxythymidine, zidovudine, or AZT) was approved in 1987 as the first anti-HIV drug, the lethality of the disease remained high until therapies based on drug combinations emerged in the mid-1990s [1]. These combination therapies included three or more drugs directed against at least two different targets. With the introduction of highly active antiretroviral therapy (HAART), the number of AIDS-related deaths in the U.S. and Europe decreased by half in the span of three years. Although HAART was lifesaving, the early regimens were far from perfect. The side effects were burdensome, and their daily dosing was complex. Drug toxicities, the difficulties of maintaining long-term adherence and the development of drug resistance frequently led to virological failure and clinical progression of the disease. In 2006, the approval of a combination of tenofovir

disoproxil fumarate (TDF), emtricitabine and efavirenz (Atripla) as the first once-daily single tablet regimen was a major therapeutic advance (reviewed in [2]). The efficacy of combination therapies (currently referred to as cART instead of HAART) increased from 43% in the mid-90s to 78% in 2010, measured as their ability to maintain undetectable viral loads for a minimum of 48 weeks, when administered to patients with less than 100,000 copies/mL at the initiation of treatment [3]. These data, obtained from a large meta-analysis, reflect the impact of Atripla and other novel therapies developed after the introduction of HAART.

Today, HIV infection has become a manageable chronic health condition, enabling people living with HIV, and under virological suppression, a good quality of life that is in many aspects comparable to the general population. However, HIV/AIDS remains as a major global health threat. Worldwide, an estimated 27.2-47.8 million people have already died of AIDS [<https://www.unaids.org/en/resources/fact-sheet>]. About 38 million are living with HIV/AIDS, but half of them do not know their HIV status. The World Health Organization estimated that in 2020 there were 680,000 AIDS-related deaths and 1.5 million new infections worldwide [<https://www.unaids.org/en/resources/fact-sheet>]. At present, HIV/AIDS is the leading cause of death among young people (aged 10-24) in Africa, and after road injuries, remains as the second largest cause of death in the world for the 25-49 year age group [4].

A cure for HIV infection, meaning that the virus is completely eradicated in the patient is still an elusive goal. However, sustained antiretroviral therapy-free remission (also known as ‘functional cure’) could be a feasible accomplishment in coming years, although HIV persistence due to the proliferation of infected cells constitutes a major challenge for a successful intervention [5,6].

Combination therapies with at least three different antiretroviral medicines is nowadays standard treatment for all people newly diagnosed with HIV. In addition, potent dual therapies are also gaining support, most notably as a choice for maintenance treatment. Currently, there are 27.5 million people globally on HIV therapy [<https://www.unaids.org/en/resources/fact-sheet>]. Although promising, these numbers are still far from the goals needed to achieve HIV eradication across the world. In addition, adverse side effects, long-term drug toxicity and the potential appearance of resistance remain as major threats to current therapies.

In this review, we provide an update on available antiretroviral therapies, and recent efforts aimed at the development of long-acting drugs, treatment simplification and HIV prevention often associated to novel antiretroviral drugs with different mechanisms of action.

### **Approved antiretroviral drugs**

The aim of HIV treatments is to reduce viral loads to undetectable levels. Antiretroviral drugs act by different mechanisms of action, while targeting various steps in the HIV replication process [Figure 1]. Thus, fostemsavir is a precursor of temsavir, a drug that binds the HIV envelope glycoprotein gp120 and blocks virus attachment to CD4+ T cells [8]. Fostemsavir was found to be effective in patients with multidrug-resistant HIV-1 infection and limited therapy options [9]. In addition, it shows no in vitro cross-resistance with currently available antiretroviral drugs. Binding to the cellular receptor CD4 can be also inhibited by ibalizumab. This is a non-immunosuppressive humanised monoclonal antibody with broad specificity that neutralizes many HIV-1 strains [10]. Chemokine receptors CXCR4 and CCR5 function along with CD4 to facilitate viral entry into target cells. Maraviroc is a negative allosteric modulator of the CCR5 receptor, and unique among antiretroviral drugs as a molecule targeting a host protein. A fourth mechanism of action among drugs blocking HIV entry is represented by enfuvirtide. This is a synthetic peptide (of 36 amino acids) with an acetylated N-terminus and a carboxamide at the C-terminus. Enfuvirtide blocks the conformational changes affecting the HIV complex gp120/gp41 that facilitate fusion of the viral envelope and the host cell membrane.

More than half of the approved drugs are reverse transcriptase (RT) inhibitors. These molecules block the conversion of the viral single-stranded genomic RNA into double-stranded DNA (dsDNA), necessary for integration into host genomic DNA. HIV-1 RT is a heterodimeric enzyme composed of subunits of 66 and 51 kDa. The large subunit contains catalytic sites of DNA polymerase and RNase H activities [11,12]. RTs can use RNA or DNA as templates during DNA polymerization.

Approved drugs targeting the RT's DNA polymerase activity include nucleoside analogues (NRTIs) and non-nucleoside RT inhibitors (NNRTIs). NRTIs are prodrugs that have to be converted to triphosphorylated derivatives before being incorporated into proviral DNA. However, approved NRTIs lack a 3'OH in the ribose ring and act as chain terminators,

blocking DNA synthesis. Within this class, tenofovir, an acyclic analogue of deoxyadenosine 5'-monophosphate, is usually included in clinical guidelines as part of recommended first-line therapies [13]. Currently, the prodrug tenofovir disoproxil fumarate (TDF) is being substituted by tenofovir alafenamide (TAF), which ensures higher active metabolite concentrations in peripheral blood mononuclear cells (PBMCs) and lower plasma tenofovir exposure [14]. NNRTIs are small hydrophobic compounds that inhibit DNA polymerization through binding to an allosteric site located a few angstroms away from the polymerase catalytic site (reviewed in [15]). Nevirapine was the first drug of this class approved for clinical use, and efavirenz has been widely used for many years. First-generation NNRTIs have been replaced in recent formulations by second-generation inhibitors such as etravirine, rilpivirine and doravirine.

Retroviral integrases catalyze the insertion of viral dsDNA into the host cell's DNA. Four identical copies of the integrase grab the two ends of the viral DNA, creating a stable complex called an intasome. The intasome binds to the cellular DNA and performs an integration reaction. First, two or three nucleotides are removed from one or both 3' ends of the proviral DNA to expose a conserved CA dinucleotide at both 3' ends (3'-processing), and then a strand transfer reaction ligates the processed 3' ends of the viral DNA to the host chromosomal DNA [16]. The HIV-1 integrase (32 kDa) has three domains and the central core contains the active site. The N- and C-terminal domains of the enzyme contain an HH-CC zinc-binding domain and a DNA-binding domain with an SH3 fold, respectively. Approved HIV-1 integrase inhibitors (raltegravir, elvitegravir, dolutegravir, cabotegravir and bictegravir) bind to the catalytic domain and block the strand transfer reaction [17,18].

Antiretroviral drugs acting in the late phase of the replicative cycle of HIV are represented by HIV protease inhibitors (saquinavir, lopinavir, darunavir and others, see [Figure 1]). Although ritonavir, indinavir and nelfinavir have been replaced by less toxic and more convenient to take protease inhibitors, ritonavir is widely used at low dose as a pharmacokinetic enhancer (booster) of protease inhibitor-based combination therapies. HIV protease cleaves viral precursor polyproteins (i.e. Gag and Gag-Pol) to generate the mature protein components of an HIV virion. The HIV-1 protease is a homodimer containing subunits of 99 amino acids each. All approved protease inhibitors except tipranavir are peptidomimetics that contain a nonhydrolyzable hydroxyethylene core instead of a peptide linkage which would otherwise be cleaved by hydrolysis [19].

## **Combination antiretroviral therapies**

After the approval of saquinavir in 1996 and the introduction of HAART regimens, combination therapies including three or more drugs directed against at least two different targets, were widely adopted for treating HIV infection. Ten years later pharmacological advances led to the approval of the first one-pill combination therapy, that contained TDF, emtricitabine and efavirenz (i.e., two NRTIs and one NNRTI). During the last 15 years, new one-pill combinations have been approved (Table 1). In general, new formulations show an improved safety profile (less adverse effects), more potent antiviral effects, and more resilience to the acquisition of drug resistance. Examples illustrating these advances include the replacement of TDF by TAF that reduces renal and bone toxicity [14], the incorporation of second-generation NNRTIs (e.g. rilpivirine or doravirine) effective against efavirenz-resistant viruses [15], or the introduction of integrase inhibitors such as raltegravir, dolutegravir or bictegravir that do not require pharmacological boosting. In addition, dolutegravir and bictegravir show a very high genetic barrier to resistance development [15]. Clinical trials demonstrated that recently approved first-line regimens achieve viral suppression rates close to 90% after 48 weeks of treatment [20,21]. However, major threats to the efficacy of antiretroviral therapies are toxicity and drug resistance, and major efforts in drug design and discovery are focused in these areas of research.

## **Drug resistance and antiretroviral drug toxicity as threats to therapy success**

Despite the rapid turnover and high mutation rate of HIV (around  $10^{-4}$  to  $10^{-5}$  mutations per nucleotide and cycle of replication), emergence of resistance has been largely contained in the clinical setting due to the increased potency of current combination regimens. However, high levels of transmitted drug resistance have been reported for antiretroviral drugs with low genetic barriers of resistance, but prescribed for many years [22,23]. Examples are M184I/V, associated with NRTI resistance, and K103N/S, Y181C/I and G190A/S, associated with resistance to first-generation NNRTIs [15] [Figure 2]. The prevalence of these mutations seems to be decreasing as dolutegravir and new drugs with high genetic barriers to resistance development are included in the prescribed regimens. Medicinal chemistry efforts are currently focused on the discovery of new drugs targeting unexploited antiviral mechanisms of action and/or effective against drug-resistant HIV strains [7].

Drug toxicity is also very important considering the life-long duration of antiretroviral therapies. Despite being safe, current treatments are not benign. We mentioned above the effects of TDF on bone and renal toxicity, and how TAF ensures lower plasma tenofovir exposure [14]. Novel prodrugs (e.g. octadecyloxyethyl benzyl tenofovir) facilitating a slow release of the active metabolite could improve safety in tenofovir-containing regimens. On the other hand, recently developed integrase inhibitors induce adipogenesis, lipogenesis, oxidative stress, fibrosis and insulin resistance, resulting in weight gain and obesity [26,27]. This has been shown in clinical trials measuring the efficacy of dolutegravir-containing regimens, particularly in patients receiving TAF as part of the antiretroviral treatment [28]. For dolutegravir, obesity and metabolic disturbances are a matter of concern for patients from African descent which would be at a higher risk of cardiovascular disease [29]. Among integrase inhibitors, dolutegravir also bears a burden of neuropsychiatric side effects [30]. Preliminary data from clinical trials and non-trial settings show that bictegravir has lower rates of adverse events in comparison with dolutegravir [29,31,32].

### **Treatment simplification and long-acting antiretroviral drugs**

Apart from being highly effective and strong suppressors of HIV replication, novel antiretroviral drugs should be safe, tolerable and facilitate medication adherence through convenient dosing and administration. Simplifying an HIV regimen can include reducing the number of antiretroviral drugs. This is acceptable as a maintenance therapy for virologically suppressed people, i.e. those patients with viral load in blood below 50 copies/mL for at least six months. Suitable two-drug combinations for maintenance therapy include dolutegravir and rilpivirine (Juluca) [33], dolutegravir and lamivudine (Dovato) [34], boosted darunavir and lamivudine and boosted atazanavir and lamivudine [35]. Combinations containing lamivudine should be avoided if there is evidence of resistance (M184I/V detection or virologic failure while on lamivudine), as well as in patients suffering chronic hepatitis B.

Interestingly, one of those therapies (dolutegravir/lamivudine) has been approved by regulatory agencies in Europe and the U.S. for the initial treatment of HIV infection, based on the results of the GEMINI 1 and 2 clinical studies that enrolled more than 1,400 people around the world [36]. However, lamivudine resistance can be relatively frequent and if the

M184I/V mutation is present, development of dolutegravir resistance could be facilitated, ruling out future treatments with integrase inhibitors.

Long-acting or extended release nanoproducts of antiretroviral therapies can be administered monthly or less frequently. These formulations can be oral, parenteral (intramuscular or subcutaneous), transdermal or implantable technologies [37]. In January 2021, Cabenuva™ became the first FDA-approved injectable intended as a complete antiretroviral regimen for adults [38]. It contains cabotegravir and rilpivirine both as extended-release injectable suspensions, that should be given intramuscularly every month. Cabotegravir is an integrase inhibitor similar in structure to dolutegravir, while rilpivirine is a second-generation NNRTI [Figure 3]. At the same time, the regulatory agency also approved a single-pill cabotegravir/rilpivirine combination (named Vocabria™). Vocabria™ should be administered for one month prior to starting treatment with Cabenuva™, in order to ensure that medications are well-tolerated before switching to the extended-release injectable formulation. In addition, ATLAS and FLAIR clinical trials showed that once-monthly intramuscular injections of long-acting cabotegravir/rilpivirine was non-inferior to conventional oral antiretroviral therapy for maintenance of HIV-RNA suppression [39,40].

### **Pre-exposure prophylaxis (PrEP) and HIV prevention**

PrEP has been shown to be highly effective in preventing HIV infections. Currently recommended formulations to be used as PrEP for HIV/AIDS include combinations of NRTIs such as TDF/emtricitabine, TDF/lamivudine and TAF/emtricitabine. Unfortunately, use of pre-exposure prophylaxis is still low among women, particularly in vulnerable populations, and recent efforts have been concentrated on the development of strategies to improve adherence in both men and women.

In this scenario, HIV topical microbicides applied to the vagina or anus and rectum have been developed to prevent HIV infection through sexual intercourse. A flexible silicone matrix vaginal ring containing dapivirine (an NNRTI), worn for a month at a time, was found to reduce HIV infection in women by  $\square$ 30% compared to placebo, in two phase 3 placebo-controlled trials [41,42], and last year the World Health Organization and the European Medicines Agency supported its use in HIV prevention for women in high HIV burden settings [43]. Besides, novel long-acting and extended-release formulations of antiretroviral



drugs are expected to facilitate a widespread use of PrEP, particularly in the most vulnerable populations.

The safety and efficacy of a cabotegravir injection every 8 weeks for HIV prevention among cisgender women has been compared to daily oral TDF/emtricitabine in a large clinical study carried out in several African countries. Although both drugs were highly effective at preventing HIV acquisition, long-acting cabotegravir injection showed statistical significant superiority in the study population [44]. Long-acting cabotegravir has low aqueous solubility and a high melting point, but requires a 2-ml dosing volume to elicit injection site reactions. Its approval by the U.S. FDA is anticipated in the end of 2021. Animal studies showed that nanocrystals of lipophilic fatty acid ester derivatives of cabotegravir improve the pharmacokinetic properties of the drug, thereby extending its apparent half-life and making yearly-dosing feasible [45].

Other antiretroviral therapies considered in PrEP studies and HIV prevention include the combination of dapivirine and maraviroc [46] and neutralizing antibodies such as VRC01. Although the results of a recent clinical trial showed that this antibody alone did not prevent overall HIV-1 acquisition more effectively than placebo, interestingly it showed 75% protection against VRC01-sensitive HIV-1 isolates, providing proof-of-concept that neutralizing antibodies prophylaxis can be effective [47].

### **Novel antiretroviral drugs in advanced clinical trials**

High potency, low toxicity and effectiveness against drug-resistant variants are desirable characteristics for newly developed antiretroviral drugs, that should be also amenable to long-acting therapy and increased dosing intervals. Islatravir (a novel NRTI) and lenacapavir (an HIV capsid assembly inhibitor) are the most advanced candidates, and their antiretroviral potential is being evaluated in phase 3 clinical trials, both in classical and long-acting formulations [48,49].

#### **1. Islatravir**

Islatravir (4'-ethynyl-2-fluoro-2'-deoxyadenosine, EFdA, or MK-8591) is an investigational NRTI [Figure 3], in development for extended administration in a subdermal drug-eluting implant [50]. After conversion to islatravir-triphosphate, the nucleotide analogue is

incorporated into the DNA chain. Unlike currently approved NRTIs, islatravir-triphosphate acts as a translocation inhibitor due to its 4'-ethynyl group, that in combination with the 3'-hydroxyl group, results in chain termination [51,52]. Islatravir inhibited HIV-1 replication in activated PBMCs with an EC<sub>50</sub> of 0.05-0.3 nM depending on the viral strain, a potency several orders of magnitude better than any of the currently prescribed NRTIs. It also shows a strong effect on multidrug-resistant strains [51,53] [Box 1]. The active triphosphate of islatravir has an intracellular half-life of up to 128 hours [54] that allows for flexible-dose treatment including once per week dosing or more spaced administration intervals. In clinical trials, single doses of 0.5 mg of islatravir suppressed HIV-1 RNA by more than one log at day 7 in treatment-naive adults with HIV-1 infection, and were generally well tolerated with no evidence of emergence of drug-resistance [54].

In phase 2b clinical trials, daily administration of islatravir in combination with doravirine/lamivudine or doravirine/TDF showed high levels of virological suppression in HIV-infected individuals [55]. In the context of this trial, researchers are evaluating the possibility of using islatravir and doravirine as a maintenance two-drug regimen, for those individuals achieving an HIV-1 RNA concentration lower than 50 copies/mL after 24 weeks of treatment. Due to its long half-life, islatravir is also an excellent candidate for long-acting formulations. The efficacy of islatravir in preexposure prophylaxis has been shown in the rhesus macaque simian/human immunodeficiency virus (SHIV) rectal challenge model [56]. Moreover, islatravir-eluting implants for yearly HIV PrEP have been evaluated for safety, tolerability and pharmacokinetics in healthy HIV-negative men and women. The implants inserted with a Nexplanon applicator were well tolerated, and specifically the 56-mg implant was found to be effective in maintaining islatravir-triphosphate concentrations above the pharmacokinetics threshold [57].

## **2. Lenacapavir**

The HIV capsid protein (CA) forms a conical shell that protects the viral genomic RNA [Box 1]. Lenacapavir [Figure 3], formerly known as GS-6207 or GS-CA2, inhibits HIV-1 replication mainly by stabilizing and thereby preventing capsid disassembly in infected cells [58]. Currently in phase 3 clinical trials, lenacapavir is the most advanced HIV capsid binding inhibitor. Its development resulted from the optimization of PF-3450074 (PF74), a peptidomimetic compound built around a phenylalanine core and capped with indole-3-

acetic acid and aniline moieties at the amino and carboxylate ends, respectively [Figure 4]. PF74 interferes with capsid assembly and disassembly as well as binding to the cellular HIV-1 cofactors Nup153 and CPSF6 that mediate nuclear import of preintegration complexes [59]. Lenacapavir is a tight inhibitor that binds two contiguous capsid subunits while promoting distal intra- and inter-hexamer interactions that stabilize the curved capsid lattice [58]. The compound inhibits HIV-1 replication in T cells and PBMCs with picomolar activity. Resistance to lenacapavir is associated with HIV-1 CA amino acid substitutions L56I, M66I, Q67H, K70N, N74D, N74S and T107N (alone and in different combinations). These substitutions confer reduced susceptibility to the inhibitor, as well as reduced viral fitness [60,61]. Q67H and N74D are the most frequently selected mutations in HIV-1 grown in cell culture, while in some patients Q67H was identified after one-week exposure to low doses of lenacapavir [62]. These experiments showed no evidence of cross-resistance with protease, RT or integrase inhibitors. Furthermore, mutations at the Gag polyprotein cleavage sites (as found in response to protease inhibitor therapies or bevirimat and other maturation inhibitors) had no effect on lenacapavir susceptibility [63].

Lenacapavir has a half-life of 7 to 11 weeks when administered subcutaneously [64], and evidence collected from experiments with non-human primates support its further development as a long-acting agent for PrEP [65]. Lenacapavir can be administered orally [55] and its highly favorable antiviral properties have stimulated efforts towards the development of second-generation inhibitors, including compounds with different binding sites in HIV-1 CA [Figure 4].

### **3. Other targets and drug candidates in development**

The most advanced antiviral agents in clinical trials are maturation inhibitors and neutralizing antibodies. Maturation inhibitors block a late protease cleavage event occurring between CA and the spacer polypeptide SP1. When CA-SP1 is not properly processed, an abnormal eccentric capsid is formed and the virus is not infectious [66]. The first representative of this class of agents was bevirimat, a betulinic acid-like compound, whose development was interrupted due to numerous resistance-related problems. Common Gag polymorphisms such as V362I and variability at positions 369-370 of the polyprotein precursor limit the efficacy of bevirimat and second-generation maturation inhibitors (e.g.

GSK3532795). GSK3640254 is active against viruses containing these polymorphisms. Currently in clinical trials, this compound showed favorable pharmacokinetics when combined with TAF/emtricitabine [67]. Still, emergence of resistance remains as a major concern in the development of these drugs.

Broadly neutralizing antibodies (bNAbs) provide a new approach to HIV-1 prevention and treatment, as inhibitors of viral entry. Clinical trials have shown the efficacy of some of those bNAbs in reducing viremia and maintaining viral suppression. Combination of bNAbs demonstrated antiviral activity in small clinical trials, but their dose, route of administration, and bioavailability are important concerns hampering their further development [68]. Besides, bNAbs could be used alone or in combination as long-acting antiretroviral drugs, since their administration every 3 to 6 months could be feasible. Unfortunately, the evaluation of VRC01 as a PrEP agent in a clinical trial has shown that the bNAb did not prevent overall HIV-1 acquisition more effectively than placebo [47]. Nevertheless, new clinical trials with antibodies showing increased potency and breadth, also in combination with other antiretroviral drugs such as long-acting cabotegravir or lenacapavir, are currently in progress (e.g., clinical trials NCT03739996 and NCT04811040, respectively).

Other antiviral agents in development (mostly at preclinical stage) include molecules expanding the currently available antiretroviral drugs, as well as compounds directed against unexploited viral targets. In the first group, there are representatives of different classes of antiretroviral agents: (i) NRTIs, such as rosofosvir etalafenamide (GS-9131), a prodrug of GS-9148 [(5-(6-amino-purin-9-yl)-4-fluoro-2,5-dihydro-furan-2-yl)oxymethyl]phosphonic acid] that after phosphorylation acts as a chain terminator of DNA polymerization [69]; (ii) NNRTIs (e.g. elvitegravir, IQP-0528, MIV-150 and MK-8507 among others [70]); (iii) protease inhibitors (e.g. GRL-001, GRL-003, GRL-121 and GRL-142, some of them effective against highly resistant HIV mutants [19,71]); (iv) integrase inhibitors (targeting the catalytic site such as MK-2048 [72,73], or allosteric inhibitors that affect integrase multimerization and HIV morphogenesis) [16,74,75]; and (v) entry inhibitors (including novel bNAbs, the humanized IgG1 monoclonal antibody UB-421 and other agents interfering with CD4 binding, CXCR4 antagonists, and fusion inhibitors) [76].

On the other hand, other druggable targets include the ribonuclease activity of HIV-1 RT [77], the viral nucleocapsid protein [78], HIV-1 Nef [79], and relevant interactions between host factors and viral proteins (e.g. HIV-1 Vpu and host BST-2, or HIV-1 Vif-APOBEC3

complexes) [80]. Despite efforts of the scientific community, suitable drug candidates targeting those biological processes are still unavailable.

### **Conclusions and perspectives**

More than 30 antiretroviral drugs have been licensed and drug combinations demonstrated high potency and minimal toxicity. However, long-term toxicity and acquired and transmitted drug resistance remain as major threats to therapy success. Continued efforts are needed to improve current formulations and facilitate adherence. Newly developed antiviral agents have interesting advantages towards this goal: low toxicity, high potency (often very efficient against resistant HIV strains frequently found in the infected population), and convenient dosing. There are multiple options for administration while very long-acting therapies facilitate their dosing every 3, 6 or even 12 months. Although drug resistance might still be problematic for islatravir, its combination with lenacapavir as a dual long-acting combination treatment holds promise for future convenient treatment of HIV infection. Besides, although not covered in this review, a number of efforts towards eradication of HIV [81], including the use of latency reversal agents [81,82], are being addressed and probably we will witness significant advances in the coming years.

### **Acknowledgements**

This work was supported in part by the Spanish Ministry of Science and Innovation (grant PID2019-104176RB-I00/AEI/10.13039/501100011033), and an institutional grant of Fundación Ramón Areces (Madrid, Spain).

### **Disclaimer Statement**

L.M.-A. declares that he has no conflicts of interest with the contents of this article.

R.D. has received conference fees from GSK, Gilead and MSD

### **References**

- [1] Fauci, A.S. (2003) HIV and AIDS: 20 years of science. *Nat. Med.* 9, 839-843
- [2] De Clercq, E. (2009) Anti-HIV drugs: 25 compounds approved within 25 years after the discovery of HIV. *Int. J. Antimicrob. Agents* 33, 307-320

- [3] Lee, F.J. *et al.* (2014) Efficacy of initial antiretroviral therapy for HIV-1 infection in adults: a systematic review and meta-analysis of 114 studies with up to 144 weeks' follow-up. *PLoS One* 9, e97482
- [4] GBD 2019 Diseases and Injuries Collaborators (2019) Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 396, 1204-1222. Erratum in: *Lancet* 2020, 396, 1562
- [5] Bailon, L. *et al.* (2020) Novel approaches towards a functional cure of HIV/AIDS. *Drugs* 80, 859-868
- [6] Bale, M.J. and Kearney, M.F. (2019) Review: HIV-1 phylogeny during suppressive antiretroviral therapy. *Curr. Opin. HIV AIDS* 14, 188-193
- [7] Ma, Y. *et al.* (2021) Medicinal chemistry strategies for discovering antivirals effective against drug-resistant viruses. *Chem. Soc. Rev.* 50, 4514-4540
- [8] Meanwell, N. A. *et al.* (2018) Inhibitors of HIV-1 attachment: the discovery and development of temsavir and its prodrug fostemsavir. *J. Med. Chem.* 61, 62-80
- [9] Kozal, M. *et al.* (2020) Fostemsavir in adults with multidrug-resistant HIV-1 infection. *N. Engl. J. Med.* 382, 1232-1243
- [10] Pace, C.S. *et al.* (2013) Anti-CD4 monoclonal antibody ibalizumab exhibits breadth and potency against HIV-1, with natural resistance mediated by the loss of a V5 glycan in envelope. *J. Acquir. Immune Defic. Syndr.* 62, 1-9
- [11] Menéndez-Arias, L. *et al.* (2017) Viral reverse transcriptases. *Virus Res.* 234, 153-176
- [12] Xavier Ruiz, F. and Arnold, E. (2020) Evolving understanding of HIV-1 reverse transcriptase structure, function, inhibition, and resistance. *Curr. Opin. Struct. Biol.* 61, 113-123
- [13] Saag, M.S. *et al.* (2020) Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2020 Recommendations of the International Antiviral Society-USA Panel. *JAMA* 324, 1651-1669
- [14] Tao, X. *et al.* (2020) Efficacy and safety of the regimens containing tenofovir alafenamide versus tenofovir disoproxil fumarate in fixed-dose single-tablet regimens for initial treatment of HIV-1 infection: A meta-analysis of randomized controlled trials. *Int. J. Infect. Dis.* 93, 108-117
- [15] Menéndez-Arias, L. (2013) Molecular basis of human immunodeficiency virus type 1 drug resistance: overview and recent developments. *Antiviral Res.* 98, 93-120

- [16] Maertens, G.N. *et al.* (2021) Structure and function of retroviral integrase. *Nat. Rev. Microbiol.*, doi: 10.1038/s41579-021-00586-9
- [17] Smith, S.J. *et al.* (2021) Integrase strand transfer inhibitors are effective anti-HIV drugs. *Viruses* 13, 205
- [18] Józwiak, I.K. *et al.* (2020) Structural biology of HIV integrase strand transfer inhibitors. *Trends Pharmacol. Sci.* 41, 611-626
- [19] Weber, I.T. *et al.* (2021) HIV protease: historical perspective and current research. *Viruses* 13, 839
- [20] Gallant, J. *et al.* (2017) Bictegravir, emtricitabine, and tenofovir alafenamide versus dolutegravir, abacavir, and lamivudine for initial treatment of HIV-1 infection (GS-US-380-1489): a double-blind, multicentre, phase 3, randomised controlled non-inferiority trial. *Lancet* 390, 2063-2072
- [21] Sax, P.E. *et al.* (2017) Coformulated bictegravir, emtricitabine, and tenofovir alafenamide versus dolutegravir with emtricitabine and tenofovir alafenamide, for initial treatment of HIV-1 infection (GS-US-380-1490): a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet* 390, 2073-2082
- [22] Baxter, J.D. *et al.* (2015) Global HIV-1 transmitted drug resistance in the INSIGHT Strategic Timing of AntiRetroviral Treatment (START) trial. *HIV Med.* 16 (Suppl 1), 77-87
- [23] Inzaule, S.C. *et al.* (2020) Tenofovir resistance in early and long-term treated patients on first-line antiretroviral therapy in eight low-income and middle-income countries. *AIDS* 34, 1965-1969
- [24] Wensing, A.M. *et al.* (2019) 2019 update of the drug resistance mutations in HIV-1. *Top. Antivir. Med.* 27, 111-121
- [25] Asante-Appiah, E. *et al.* (2021) Impact of HIV-1 resistance-associated mutations on susceptibility to doravirine: analysis of real-world clinical isolates. *Antimicrob. Agents Chemother.*, doi:10.1128/AAC.01216-21
- [26] Gorwood, J. *et al.* (2020) The integrase inhibitors dolutegravir and raltegravir exert proadipogenic and profibrotic effects and induce insulin resistance in human/simian adipose tissue and human adipocytes. *Clin. Infect. Dis.* 71, e549-e560
- [27] Scarsi, K.K. *et al.* (2020) HIV-1 integrase inhibitors: a comparative review of efficacy and safety. *Drugs* 80, 1649-1676
- [28] Venter, W.D.F. *et al.* (2019) Dolutegravir plus two different prodrugs of tenofovir to treat HIV. *N. Engl. J. Med.* 381, 803-815

- [29] Kolakowska, A. *et al.* (2019) Update on adverse effects of HIV integrase inhibitors. *Curr. Treat. Options Infect. Dis.* 11, 372-387
- [30] Apostolova, N. *et al.* (2015) Efavirenz and the CNS: what we already know and questions that need to be answered. *J. Antimicrob. Chemother.* 70, 2693-2708
- [31] Stellbrink, H.J. *et al.* (2019) Co-formulated bictegravir, emtricitabine, and tenofovir alafenamide versus dolutegravir with emtricitabine and tenofovir alafenamide for initial treatment of HIV-1 infection: week 96 results from a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet HIV* 6, e364-e372
- [32] Wohl, D.A. *et al.* (2019) Bictegravir combined with emtricitabine and tenofovir alafenamide versus dolutegravir, abacavir, and lamivudine for initial treatment of HIV-1 infection: week 96 results from a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet HIV* 6, e355-e363
- [33] Aboud, M. *et al.* (2019) Efficacy and safety of dolutegravir-rilpivirine for maintenance of virological suppression in adults with HIV-1: 100-week data from the randomised, open-label, phase 3 SWORD-1 and SWORD-2 studies. *Lancet HIV* 6, e576-e587
- [34] van Wyk, J. *et al.* (2020) Efficacy and safety of switching to dolutegravir/lamivudine fixed-dose 2-drug regimen vs continuing a tenofovir alafenamide-based 3- or 4-drug regimen for maintenance of virologic suppression in adults living with human immunodeficiency virus type 1: phase 3, randomized, noninferiority TANGO study. *Clin. Infect. Dis.* 71, 1920-1929
- [35] Perez-Molina, J.A. *et al.* (2018) Individual patient data meta-analysis of randomized controlled trials of dual therapy with a boosted PI plus lamivudine for maintenance of virological suppression: GeSIDA study 9717. *J. Antimicrob. Chemother.* 73, 2927-2935
- [36] Cahn, P. *et al.* (2019) Dolutegravir plus lamivudine versus dolutegravir plus tenofovir disoproxil fumarate and emtricitabine in antiretroviral-naive adults with HIV-1 infection (GEMINI-1 and GEMINI-2): week 48 results from two multicentre, double-blind, randomised, non-inferiority, phase 3 trials. *Lancet* 393, 143-155
- [37] Flexner, C. *et al.* (2021) Long-acting drugs and formulations for the treatment and prevention of HIV infection. *Int. J. Antimicrob. Agents* 57, 106220
- [38] Orkin, C. *et al.* (2021) Long-acting cabotegravir plus rilpivirine for treatment in adults with HIV-1 infection: 96-week results of the randomised, open-label, phase 3 FLAIR study. *Lancet HIV* 8, e185-e196
- [39] Swindells, S. *et al.* (2020) Long-acting cabotegravir and rilpivirine for maintenance of HIV-1 suppression. *N. Engl. J. Med.* 382, 1112-1123



- [40] Rizzardini, G. *et al.* (2020) Long-acting injectable cabotegravir + rilpivirine for HIV maintenance therapy: week 48 pooled analysis of phase 3 ATLAS and FLAIR trials. *J. Acquir. Immune Defic. Syndr.* 85, 498-506
- [41] Baeten, J.M. *et al.* (2016) Use of a vaginal ring containing dapivirine for HIV-1 prevention in women. *N. Engl. J. Med.* 375, 2121-2132
- [42] Nel, A. *et al.* (2016) Safety and efficacy of a dapivirine vaginal ring for HIV prevention in women. *N. Engl. J. Med.* 375, 2133-2143
- [43] Baeten, J.M. *et al.* (2021) Safety, uptake, and use of a dapivirine vaginal ring for HIV-1 prevention in African women (HOPE): an open-label, extension study. *Lancet HIV* 8, e87-e95
- [44] Landovitz, R.J. *et al.* (2021) Cabotegravir for HIV prevention in cisgender men and transgender women. *N. Engl. J. Med.* 385, 595-608
- [45] Gautam, N. *et al.* (2021) Lipophilic nanocrystal prodrug-release defines the extended pharmacokinetic profiles of a year-long cabotegravir. *Nat. Commun.* 12, 3453
- [46] Srinivas, N. *et al.* (2020) Translational approach to predicting the efficacy of maraviroc-based regimens as HIV preexposure prophylaxis. *Antimicrob. Agents Chemother.* 64, e01729-19
- [47] Corey, L. *et al.* (2021) Two randomized trials of neutralizing antibodies to prevent HIV-1 acquisition. *N. Engl. J. Med.* 384, 1003-1014
- [48] Gu, S.X. *et al.* (2020) Recent discoveries in HIV-1 reverse transcriptase inhibitors. *Curr. Opin. Pharmacol.* 54, 166-172
- [49] Zhang, D.W. *et al.* (2019) A HTRF based competitive binding assay for screening specific inhibitors of HIV-1 capsid assembly targeting the C-terminal domain of capsid. *Antiviral Res.* 169, 104544
- [50] Barrett, S.E. *et al.* (2018) Extended-duration MK-8591-eluting implant as a candidate for HIV treatment and prevention. *Antimicrob. Agents Chemother.* 62, e01058-18
- [51] Michailidis, E. *et al.* (2014) 4'-Ethyne-2'-fluoro-2'-deoxyadenosine (EFdA) inhibits HIV-1 reverse transcriptase with multiple mechanisms. *J. Biol. Chem.* 289, 24533-24548
- [52] Salie, Z.L. *et al.* (2016) Structural basis of HIV inhibition by translocation-defective RT inhibitor 4'-ethyne-2'-fluoro-2'-deoxyadenosine (EFdA). *Proc. Natl. Acad. Sci. U. S. A.* 113, 9274-9279
- [53] Takamatsu, Y. *et al.* (2018) The high genetic barrier of EFdA/MK-8591 stems from strong interactions with the active site of drug-resistant HIV-1 reverse transcriptase. *Cell Chem. Biol.* 25, 1268-1278.e3

- [54] Schürmann, D. *et al.* (2020) Safety, pharmacokinetics, and antiretroviral activity of islatravir (ISL, MK-8591), a novel nucleoside reverse transcriptase translocation inhibitor, following single-dose administration to treatment-naive adults infected with HIV-1: an open-label, phase 1b, consecutive-panel trial. *Lancet HIV* 7, e164-e172
- [55] Molina, J.M. *et al.* (2021) Islatravir in combination with doravirine for treatment-naive adults with HIV-1 infection receiving initial treatment with islatravir, doravirine, and lamivudine: a phase 2b, randomised, double-blind, dose-ranging trial. *Lancet HIV* 8, e324-e333
- [56] Markowitz, M. *et al.* (2020) Once-weekly oral dosing of MK-8591 protects male rhesus macaques from intrarectal challenge with SHIV109CP3. *J. Infect. Dis.* 221, 1398-1406
- [57] Matthews, R.P. *et al.* (2021) Next-generation islatravir implants projected to provide yearly HIV prophylaxis. 2021 CROI Conference, Abstract 88. Available at: [https://www.natap.org/2021/CROI/croi\\_36.htm](https://www.natap.org/2021/CROI/croi_36.htm) (accessed September 29, 2021)
- [58] Bester, S.M. *et al.* (2020) Structural and mechanistic bases for a potent HIV-1 capsid inhibitor. *Science* 370, 360-364
- [59] Price, A.J. *et al.* (2014) Host cofactors and pharmacologic ligands share an essential interface in HIV-1 capsid that is lost upon disassembly. *PLoS Pathog.* 10, e1004459
- [60] Link, J.O. *et al.* (2020) Clinical targeting of HIV capsid protein with a long-acting small molecule. *Nature* 584, 614-618
- [61] Marcelin, A.G. *et al.* (2020) Frequency of capsid substitutions associated with GS-6207 in vitro resistance in HIV-1 from antiretroviral-naive and -experienced patients. *J. Antimicrob. Chemother.* 75, 1588-1590
- [62] Margot, N. *et al.* (2020) Lenacapavir resistance analysis in a phase 1b clinical proof-of-concept study. HIV Glasgow 2020 Virtual Congress, Poster O-324. Available at: <http://www.hivglasgow.org/2020-posters> (accessed June 24, 2021)
- [63] Margot, N. *et al.* (2021) Absence of lenacapavir (GS-6207) phenotypic resistance in HIV Gag cleavage site mutants and in isolates with resistance to existing drug classes. *Antimicrob. Agents Chemother.* 65, e02057-20
- [64] Begley, R. *et al.* (2020) GS-6207 sustained delivery formulation supports 6-month dosing interval. 23rd International AIDS Conference virtual, Abstract PEB0265
- [65] Bekerman, E. *et al.* (2021) Long-acting HIV capsid inhibitor effective as prep in a SHIV rhesus macaque model. 2021 CROI Conference. Abstract 717. Available at: [https://www.natap.org/2021/CROI/croi\\_06.htm](https://www.natap.org/2021/CROI/croi_06.htm) (accessed September 29, 2021)

- [66] Kleinpeter, A.B. and Freed, E.O. (2020) HIV-1 maturation: lessons learned from inhibitors. *Viruses* 12, 940
- [67] Pene Dumitrescu, T. *et al.* (2021) A phase I evaluation of the pharmacokinetics and tolerability of the HIV-1 maturation inhibitor GSK3640254 and tenofovir alafenamide/emtricitabine in healthy participants. *Antimicrob. Agents Chemother.* 65, e02173-20
- [68] Mahomed, S. *et al.* (2021) Clinical trials of broadly neutralizing monoclonal antibodies for human immunodeficiency virus prevention: a review. *J. Infect. Dis.* 223, 370-380
- [69] Cihlar, T. *et al.* (2008) Design and profiling of GS-9148, a novel nucleotide analog active against nucleoside-resistant variants of human immunodeficiency virus type 1, and its orally bioavailable phosphonoamidate prodrug, GS-9131. *Antimicrob. Agents Chemother.* 52, 655-665
- [70] Wang, Y. *et al.* (2019) Current and emerging non-nucleoside reverse transcriptase inhibitors (NNRTIs) for HIV-1 treatment. *Expert Opin. Drug Metab. Toxicol.* 15, 813-829
- [71] Hattori, S.I. *et al.* (2019) Halogen bond interactions of novel HIV-1 protease inhibitors (PI) (GRL-001-15 and GRL-003-15) with the flap of protease are critical for their potent activity against wild-type HIV-1 and multi-PI-resistant variants. *Antimicrob. Agents Chemother.* 63, e02635-18
- [72] Liu, A.Y. *et al.* (2019) Phase 1 pharmacokinetic trial of 2 intravaginal rings containing different dose strengths of vicriviroc (MK-4176) and MK-2048. *Clin. Infect. Dis.* 68, 1129-1135
- [73] Hoesley, C.J. *et al.* (2019) Phase 1 safety and pharmacokinetics study of MK-2048/vicriviroc (MK-4176)/MK-2048A intravaginal rings. *Clin. Infect. Dis.* 68, 1136-1143
- [74] Sun, J. *et al.* (2021) Optimized binding of substituted quinoline ALLINIs within the HIV-1 integrase oligomer. *J. Biol. Chem.* 296, 100363
- [75] Gupta, K. *et al.* (2021) Allosteric HIV integrase inhibitors promote formation of inactive branched polymers via homomeric carboxy-terminal domain interactions. *Structure* 29, 213-225.e5
- [76] Xiao, T. *et al.* (2021) HIV-1 entry and membrane fusion inhibitors. *Viruses* 13, 735
- [77] Tramontano, E. *et al.* (2019) Ribonuclease H, an unexploited target for antiviral intervention against HIV and hepatitis B virus. *Antiviral Res.* 171, 104613
- [78] Iraci, N. *et al.* (2018) NCp7: targeting a multitask protein for next-generation anti-HIV drug development part 2. Noncovalent inhibitors and nucleic acid binders. *Drug Discov. Today* 23, 687-695

- [79] Lv, Y. *et al.* (2020) Research progress of HIV-1 Nef inhibitors. *AIDS Rev.* 22, 221-226
- [80] Duan, S. *et al.* (2020) A novel HIV-1 inhibitor that blocks viral replication and rescues APOBEC3s by interrupting vif/CBF $\beta$  interaction. *J. Biol. Chem.* 295, 14592-14605
- [81] Margolis, D.M. *et al.* (2020) Curing HIV: seeking to target and clear persistent infection. *Cell* 181, 189-206
- [82] Spivak, A.M. and Planelles, V. (2018) Novel latency reversal agents for HIV-1 cure. *Annu. Rev. Med.* 69, 421-436
- [83] Markowitz, M. and Sarafianos, S.G. (2018) 4'-Ethynyl-2-fluoro-2'-deoxyadenosine, MK-8591: a novel HIV-1 reverse transcriptase translocation inhibitor. *Curr. Opin. HIV AIDS* 13, 294-299
- [84] Schürmann, D. *et al.* (2020) Safety, pharmacokinetics, and antiretroviral activity of islatravir (ISL, MK-8591), a novel nucleoside reverse transcriptase translocation inhibitor, following single-dose administration to treatment-naive adults infected with HIV-1: an open-label, phase 1b, consecutive-panel trial. *Lancet HIV* 7, e164-e172
- [85] Sun, L. *et al.* (2021) An insight on medicinal aspects of novel HIV-1 capsid protein inhibitors. *Eur. J. Med. Chem.* 217, 113380
- [86] Chia, T. *et al.* (2021) A small molecule, ACAi-028, with anti-HIV-1 activity targets a novel hydrophobic pocket on HIV-1 capsid. *Antimicrob. Agents Chemother.* 65, e0103921

## Highlights

Current antiretroviral therapies are highly efficacious in maintaining undetectable viral loads for a long period of time.

Efforts to improve current therapies focus on treatment simplification (i.e. reducing the number of antiretroviral drugs and their doses) and the development of long-acting drugs for treatment and pre-exposure prophylaxis.

Lenacapavir (a capsid-binding agent) and islatravir (a nucleoside reverse transcriptase inhibitors) are currently in advanced clinical trials and are suitable for extended-release formulations.

Low toxicity, high potency and efficiency against wild-type and drug-resistant strains of HIV, and convenient dosing are desirable properties to be met by newly developed antiretroviral drugs.

## Glossary

**Allosteric inhibitors:** molecules that bind to an enzyme changing its shape and activity, by interacting in location different from that of the catalytic site. In antiretroviral therapy, NNRTIs are allosteric inhibitors of HIV-1 RT.

**Boosting:** in combination antiretroviral therapies, this term describes the use of a small amount of a drug (typically ritonavir or cobicistat) that competes in the liver for the degradation of protease inhibitors, thereby increasing the concentration of antiretroviral drugs in the bloodstream, by reducing the speed at which the drugs are metabolized and cleared.

**cART or combination antiretroviral therapy:** a synonym of HAART, implying the use of at least three antiretroviral drugs directed against two or more molecular targets.

**gp120/gp41:** envelope glycoprotein of HIV-1. It is a homotrimer composed of subunits gp120 (surface glycoprotein) and gp41 (transmembrane glycoprotein).

**HAART or highly active antiretroviral therapy:** Term used since the mid 90' to describe the combination of antiretroviral drugs to achieve high antiviral effect.

**Latency reversal agents:** agents that activate HIV latently infecting resting CD4 lymphocytes, contributing to their destruction and subsequent reduction of the latent HIV reservoir after administration of conventional antiretroviral therapies.

**Neutralizing antibodies:** immunoglobulins that bind to the viral particle, typically to the envelope glycoprotein, in such a way that cell infection is blocked.

**Nexplanon:** single, radiopaque, rod-shaped implant, traditionally used in birth control for women when containing contraceptives. The implant can be loaded in the needle of a disposable applicator.

**Preintegration complex:** ribonucleoprotein form after viral entry and reverse transcription, that is transported into the nucleus for integration. Although its specific composition and structure is still unknown, it contains double-stranded DNA, integrase, and viral proteins, most notably, MA (matrix protein), Vpr, and probably CA and NC proteins.

**Proviral DNA:** stands for the genetic information of retroviruses, integrated as DNA in the chromosome of infected cells with the capability to be activated and produce new infective viral particles.

**Reverse transcriptase (RT):** retrovirus-encoded enzyme responsible for the production of double-stranded proviral DNA from single-stranded genomic RNA.

**Viral load:** term used to indicate the quantity of genetic material, commonly RNA, of a virus present in the blood. Usually, in HIV-infected patients, the measurement is given as HIV-1 RNA copies per mL of plasma.

**Virological suppression:** in treated HIV-infected individuals, it describes the achievement of complete inhibition of viral replication, measured by viral load in plasma as less than 50 HIV-1 RNA copies/mL.

### **Outstanding Questions**

Could long-acting strategies for pre-exposure prophylaxis (PrEP) have a significant global impact on the AIDS pandemic?

Could the new highly active antiretroviral drugs circumvent the inexorable development of resistance mutations?

Are there any new antiviral targets still worth exploring for the treatment of HIV infection?

Could monoclonal neutralizing antibodies be part of standard prophylaxis or therapies against HIV-1 infection?

### **Box 1**

**Islatravir resistance.** The combination of amino acid substitutions I142V/T165R/M184V was selected *in vitro* after successive passage of HIV-1 in the presence of 2'-deoxy-4'-ethynyl-adenosine, a nonfluorinated analogue of islatravir. M184V, the first mutation that appeared in those experiments conferred a 7.5-fold reduction in islatravir susceptibility. M184V contributes to islatravir resistance particularly in combination with P119S and T165A (reviewed in [83]). The relatively high prevalence of M184I/V in heavily-treated populations due to the extensive use of lamivudine and emtricitabine in combination therapies is a matter of concern for the use of islatravir. However, the drug candidate retains activity against a wide range of NRTI-resistant mutants, including M41L/T69-insertion/T215Y mutants characterized by their very high-level resistance to AZT. Furthermore, common drug resistance mutations found in the HIV-1 RT-coding region (e.g. K65R, L74V and Q151M) confer hypersusceptibility to the drug, and many studies have confirmed that M184I/V have a negative impact on the HIV-1 replication capacity [15]. So far, emergence of resistance in patients has not been reported in ongoing clinical trials for the treatment of HIV-1 infection with islatravir in combination with other antiretroviral drugs, as well as for pre-exposure prophylaxis (PrEP) of HIV-1 infection as a single agent [84].

**HIV-1 CA structure and CA-binding inhibitors.** HIV-1 CA originates from cleavage of the 55-kDa Gag polyprotein by the viral protease. It has two independently folded domains connected by a 5-residue flexible linker [Figure 4]. The N-terminal domain includes residues 1-145, while the C-terminal domain contains amino acids 151 to 231. Less than two dozen CA-targeting compounds have been described, including small molecules, peptides and a specific antibody [49]. PF74 was extensively studied and a lead compound of lenacapavir. It inhibits both early and late steps of viral replication. According to their mechanism of action and target CA-targeting compounds can be classified in different groups: (i) molecules promoting CA multimerization and targeting the N-terminal domain, around positions 67, 70 and 107 (e.g. lenacapavir, PF74 and GS-CA1); (ii) covalent agents targeting the C-terminal

region, at Cys198 and Cys218, and promoting multimerization, such as ebselen; (iii) compounds decreasing multimerization and acting in the late phase of the viral replication cycle (e.g. CAP-1, BD-1 and BM-1, binding around residues 28-32); and (iv) agents that decrease CA multimerization through binding a pocket defined by residues Glu98, His120 and Ile124 (e.g. C-1 and I-XW-053) [85]. ACAi-028 represents a fifth class of compounds binding to the N-terminal domain of CA. This compound, identified after virtual screening using *in silico* docking simulations, interacts with amino acid residues Gln13, Ser16, and Thr19 through hydrogen bonds. ACAi-028 acts in the early phase of HIV-1 replication and reduces CA multimerization [86].



## LEGENDS TO FIGURES

**Figure 1.** Approved antiretroviral drugs and HIV replication cycle. Less-frequently used or discontinued drugs are shown in purple and smaller font size. Abbreviations: /c, boosted with cobicistat; /r, boosted with ritonavir; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate. Cobicistat and ritonavir are pharmacokinetic enhancers administered with specific protease inhibitors or the integrase inhibitor, elvitegravir, to optimize their therapeutic concentrations. Their mechanism of action involves the inhibition of cytochrome P450 3A (CYP3A4) enzymes. Adapted from ref. [7] with permission from the Royal Society of Chemistry.

**Figure 2.** Amino acid substitutions associated with resistance to approved antiretroviral drugs. Major resistance mutations for each drug are indicated by red boxes, while accessory mutations are highlighted with yellow boxes. Data were taken from ref. [24], and the Stanford University HIV Drug Resistance Database (<http://hivdb.stanford.edu>), with additional data on doravirine susceptibility from ref. [25].

**Figure 3.** Chemical structures of antiretroviral drugs approved as long-acting formulations or in advanced clinical trials.


**Figure 4.** HIV-1 CA monomer showing the location of binding sites of representative assembly and disassembly inhibitors in pre-clinical development. The upper right panel shows a cartoon representation of the structure of a hexameric complex. One of the subunits is represented in cyan. Images were obtained with PyMol using PDB coordinates 6VKV and 6OBH.

**Table 1.** FDA-approved single-tablet drug combinations of two or more antiretroviral drugs for antiretroviral therapy.<sup>1</sup>


	Commercial name	Combination	FDA approval date
Three-drug combinations for initial therapy			
	Atripla	TDF/emtricitabine/efavirenz	July 12, 2006
	Complera	TDF/emtricitabine/rilpivirine	August 10, 2011
	Stribild	TDF/emtricitabine/elvitegravir (cobicistat)	August 27, 2012
	Triumeq	Abacavir/lamivudine/dolutegravir	August 22, 2014
	Genvoya	TAF/emtricitabine/elvitegravir (cobicistat)	November 5, 2015
	Odefsey	TAF/emtricitabine/rilpivirine	March 1, 2016
	Biktarvy	TAF/emtricitabine/bictegravir	February 7, 2018
	Symfi Lo & Symfi	TDF/lamivudine/efavirenz	February 5/March 22, 2018
	Symtuza	TAF/emtricitabine/darunavir (cobicistat)	July 17, 2018
	Delstrigo	TDF/lamivudine/doravirine	August 30, 2018
Two-drug combinations for maintenance therapy <sup>2</sup>			
	Juluca	Dolutegravir/rilpivirine	November 21, 2017
	Dovato	Dolutegravir/lamivudine	April 8, 2019
Two-drug combinations for initial therapy			
	Cabenuva	Cabotegravir/rilpivirine (both supplied as extended-release injectable suspensions)	January 22, 2021

<sup>1</sup> Information taken from: <https://aidsinfo.nih.gov/understanding-hiv-aids/fact-sheets/21/58/fda-approved-hiv-medicines> (accessed September 30, 2021).

<sup>2</sup> To be used only in patients with undetectable viral load in blood (less than 50 copies/mL) for at least 6 months.




## RT inhibitors (NRTIs and NNRTIs)



### Mutation complexes conferring high-level resistance:

- All approved NRTIs:  
M41L/A62V/T69(insert)/K70R/L210W/T215(F/Y)
- All except tenofovir:  
A62V/V75I/F77L/F116Y/Q151M
- All except lamivudine and emtricitabine:  
M41L/D67N/K70R/L210W/T215(F/Y)/K219(E/Q)

## Integrase inhibitors



## Entry inhibitors

### Fostemsavir

Resistance to temsavir associated with amino acid substitutions in gp120:  
S375H/I/N/M/T, M426L/P, M434I/K and M475I.

### Ibalizumab-uyyk

Resistance acquired through the loss of glycosylation sites in the V5 region of gp120 (i.e. N460Q and N464Q alone or in combination).


### Maraviroc


Resistance through selection of viruses using the CXCR4 coreceptor, or mutations in V2, V3 and V4 loops of gp120.

### Enfuvirtide


Resistance conferred by several amino acid substitutions in the Env transmembrane subunit gp41:G36D/E/S, I37T/N/V, V38A/E/M, Q39R, Q40H, N42T and N43D/K/S.

## Protease inhibitors







**Rilpivirine**





**Islatravir**



**Cabotegravir**



**Lenacapavir**



N-terminal  
domain

C-terminal  
domain

