

Nomenclatural changes in a taxonomically complex group (Amphipoda: Gammaridae)

MARIO GARCÍA-PARÍS^{1,4}, PILAR JURADO-ANGULO^{1,5}, PAULA C. RODRÍGUEZ-FLORES² &
NATALIA ROSAS-RAMOS^{1,3}

¹Departamento de Biodiversidad y Biología Evolutiva. Museo Nacional de Ciencias Naturales, MNCN-CSIC. c/ José Gutiérrez Abascal, 2. 28006, Madrid, Spain

²Department of Organismic and Evolutionary Biology, Museum of Comparative Zoology, Harvard University, 26 Oxford St., Cambridge MA 02138, USA.  paularodriguezflores@g.harvard.edu;  <https://orcid.org/0000-0003-1555-9598>

³Departamento de Biología Animal (Área de Zoología), Facultad de Biología (Edificio de Farmacia, planta 5), Universidad de Salamanca, Campus Miguel de Unamuno s/n, 37007 Salamanca, Spain.

 nataliarosasr@usal.es;  <https://orcid.org/0000-0002-8653-0306>

⁴ mparis@mncn.csic.es;  <https://orcid.org/0000-0002-9361-9405>

⁵ si.pilarjurado@gmail.com;  <https://orcid.org/0000-0003-1144-9076>

Abstract

The family Gammaridae is a large monophyletic group with a complicated taxonomy. A nomenclatural review of the taxonomic status of *Pectenogammarus* Reid, 1940 was conducted based on Hou & Sket (2016) and Sket & Hou (2018) phylogenetic proposals. These authors used the name *Homoeogammarus* Schellenberg, 1937 for a set of morphologically diverse taxa previously included mainly within *Echinogammarus* Stebbing, 1899. Unfortunately, the use of the name *Homoeogammarus* Schellenberg, 1937 is problematic. As a result of a detailed bibliographic review, we propose the name *Pectenogammarus* Reid, 1940 (**stat. nov.**) for the “*Homoeogammarus*” clade (Sket & Hou 2018), which results in 20 new combinations. In addition, two genera are here formally synonymised as follows: *Pectenogammarus* Reid, 1940 = *Homoeogammarus* Barnard et Barnard, 1983, **syn. nov.** = *Laurogammarus* G.S. Karaman, 1984, **syn. nov.** A synonymic catalogue is provided.

Key words: Amphipoda, *Homoeogammarus*, *Laurogammarus*, new synonymy, nomenclature, *Pectenogammarus*, taxonomy

Introduction

In a recent series of papers, Hou & Sket (2016) and Sket & Hou (2018) proposed a phylogenetic framework that solved some of the long-standing taxonomic problems within Gammaridae. Some of the taxonomic changes derived from Hou & Sket (2016) and Sket & Hou (2018) phylogenetic proposals, were the restriction of the genus *Echinogammarus* Stebbing, 1899 to the *E. berilloni* (Catta, 1878a) species-group (*sensu* Pinkster 1969, 1973) and the transferring of the other species formerly included in *Echinogammarus* to the genera *Chaetogammarus* Martynov, 1925, *Marinogammarus* Schellenberg, 1937, *Parhomoeogammarus* Schellenberg, 1943, and most of them to *Homoeogammarus* Schellenberg, 1937. By doing so, Hou & Sket (2016) and Sket & Hou (2018) rendered well supported monophyletic genera, showing that many of the morphological characters used for the diagnosis of the former genera are largely homoplastic. In this paper, a nomenclatural review of Hou & Sket (2016) and Sket & Hou (2018) “*Homoeogammarus*” is conducted.

Results

Hou & Sket (2016) and Sket & Hou (2018) used the name *Homoeogammarus* Schellenberg, 1937 for a large clade

that included sets of morphologically diverse taxa, most of them previously included within *Echinogammarus*, as *E. simoni* (Chevreux, 1894), and *E. pungens* (Milne Edwards, 1840) species-groups (*sensu* Stock 1968; Pinkster & Stock 1972). Unfortunately, the use of the name *Homoeogammarus* Schellenberg, 1937, is problematic. Schellenberg (1937) proposed *Homoeogammarus* as a subgenus of *Gammarus* Fabricius, 1775, without designating a type species among the two species he included within it: *Gammarus simoni* Chevreux, 1894 and *G. tacapensis* Chevreux & Gauthier, 1924. Therefore, as previously pointed out by Stock (1968), because of the articles 13.3 and 67.4.1 of the International Code of Zoological Nomenclature (International Commission on Zoological Nomenclature—ICZN, 1999), *Homoeogammarus* Schellenberg, 1937 is not an available name. The name *Homoeogammarus* was later made available by Barnard & Barnard (1983: p. 487) who designated *Gammarus simoni* as its type species.

In addition to *Homoeogammarus* Barnard & Barnard, 1983, four other genera names were applied to species-groups included in Sket and Hou's (2018) "*Homoeogammarus*" clade: *Ostiogammarus* S. Karaman, 1931, *Neogammarus* Ruffo, 1937, *Pectenogammarus* Reid, 1940, and *Laurogammarus* G.S. Karaman, 1984.

Ostiogammarus was described including a large number of species, but none of them was designated as type species; therefore, according to articles 13.3 and 67.4.1 of the International Code of Zoological Nomenclature (International Commission on Zoological Nomenclature—ICZN, 1999) *Ostiogammarus* is also an unavailable name. Stock (1968), followed by Barnard & Barnard (1983) and Pinkster (1993), indicated erroneously that Karaman (1931) selected *G. berilloni* as type species for *Ostiogammarus*; but Karaman (1931) rather said that *G. berilloni* could be the type species of *Echinogammarus*: "Und wenn das ganze Genus *Echinogammarus* auf diesem Merkmal beruht und *E. berilloni* als Typus der Gruppe betrachtet werden sollte (wie dies Chevreux-Fage anführen), müsste die ganze Gruppe *Echinogammarus* als solche gestrichen werden" (Karaman 1931: p. 62). Some of the species included by Karaman (1931) in the original description of *Ostiogammarus* (i.e., *G. olivii* Milne Edwards, 1830, *G. pungens* Milne Edwards, 1840, *G. scutarensis* Schäferna, 1923 and *G. thoni* Schäferna, 1923) are part of Sket & Hou's (2018) "*Homoeogammarus*" clade. Nevertheless, the name *Ostiogammarus* cannot be used for any of these taxa.

Neogammarus Ruffo, 1937, described as a subgenus of *Gammarus* for the species *G. festai* Ruffo, 1937 and *G. rhipidiophorus* Catta, 1878b, is unavailable because it was described without fixation of type species. Dahl (1958) reserved *Neogammarus* for *G. festai*, but as indicated in article 67.5 of the International Code of Zoological Nomenclature (ICZN 1999) this action does not constitute a designation of type species (contrary to the indication of Stock (1971)), and consequently *Neogammarus* Dahl, 1958 is also an unavailable name. *Neogammarus* was made subsequently available by Karaman (1969: p. 57; 1971: p. 206) who designated *G. rhipidiophorus* Catta, 1878b as its type species. *Gammarus rhipidiophorus* (type species of *Neogammarus* G.S. Karaman, 1969) is not included in Sket & Hou's (2018) "*Homoeogammarus*" clade, and therefore *Neogammarus* cannot be used to name this clade.

Pectenogammarus was described as a subgenus of *Gammarus* for *G. (Pectenogammarus) planicrurus* Reid, 1940 (type species by monotypy). It is an available name and has nomenclatural precedence over *Homoeogammarus* Barnard & Barnard, 1983 and also over *Neogammarus* G.S. Karaman, 1969 according to the principle of priority (Art. 23.1 ICZN, 1999).

Laurogammarus was described for *Carinogammarus scutarensis* Schäferna, 1923 (type species by original designation). It is an available name, and its type species is included in Sket and Hou's (2018) "*Homoeogammarus*" clade, but it does not have priority over *Pectenogammarus* nor *Homoeogammarus* Barnard & Barnard, 1983.

Among the available names, *Pectenogammarus* is the oldest, and thus has nomenclatural priority for Sket & Hou's (2018) "*Homoeogammarus*" clade. The type species of *Pectenogammarus*, *G. (P.) planicrurus*, was included by Sket & Hou's (2018) within their "*Homoeogammarus*" clade. *Pectenogammarus planicrurus* is deeply nested within the "*Homoeogammarus*" clade (as shown in the supplementary material of Sket and Hou (2018)), confirming the applicability of the name *Pectenogammarus* for the entire clade.

Based on this information, the synonymous list of *Pectenogammarus* (including unavailable names) remains as:

***Pectenogammarus* Reid 1940**

Ostiogammarus S. Karaman, 1931: p. 61 (**unavailable name**)

Type species: none designated originally; some of the species included originally by Karaman (1931) are included in Sket & Hou's (2018) "*Homoeogammarus*" clade.

Homoeogammarus Schellenberg, 1937: p. 272 (described as subgenus of *Gammarus*) (**unavailable name**)

Type species: none designated originally; *Gammarus simoni* and *Gammarus tacapensis* were included originally in *Homoeogammarus*, but none of them was designated as type species (Schellenberg 1937). Synonymized with *Echinogammarus* Stebbing, 1899 by Stock (1968 as an invalid name).

Neogammarus Ruffo, 1937: p. 442 (described as a subgenus of *Gammarus*) (**unavailable name**)

Type species: none designated originally. *Gammarus festai* Ruffo, 1937 and *G. rhipidiophorus* Catta, 1878b were included originally in *Neogammarus*, but none of them was designated as type species (Ruffo 1937).

Pectenogammarus Reid, 1940: p. 288 (described as a subgenus of *Gammarus*) (**stat. nov.**)

Type species: *Gammarus (Pectenogammarus) planicrurus* Reid, 1940, by monotypy. Raised to genus by Stock (1982) and retained as such by Duhig & Humphries (1955), Kant *et al.* (1968), Morgan (1970), and Van Maren (1975a, b), among others. Included in the synonymy of *Echinogammarus* Stebbing, 1899 by Karaman (1976) and of *Homoeogammarus* Schellenberg, 1937 by Sket & Hou (2018).

Homoeogammarus Barnard & Barnard, 1983: p. 487 (as a synonym of *Echinogammarus*) (**new synonymy**)

Type species: *Gammarus simoni* Chevreux, 1894, by original designation. Treated as a synonym of *Echinogammarus* Stebbing, 1899 by Barnard & Barnard (1983). Restored as valid by Hou & Sket (2016) and Sket & Hou (2018), but erroneously attributing authorship to Schellenberg, 1937.

Laurogammarus G.S. Karaman, 1984: p. 29 (**new synonymy**)

Type species: *Carinogammarus scutarensis* Schäferna, 1923, by original designation. Treated as an independent genus by Pinkster (1993) and Grabowski *et al.* (2014) among others. Synonymized with *Homoeogammarus* Schellenberg, 1937 by Sket & Hou (2018), because its type species *C. scutarensis* is nested within the clade.

Included species in *Pectenogammarus* based on Hou & Sket (2016) and Sket & Hou (2018) are:

Pectenogammarus antalyae (G.S. Karaman, 1971) **new combination**

Pectenogammarus atlanticus (Dahl, 1958) **new combination**

Pectenogammarus cyrtus (Pinkster & Platvoet, 1986) **new combination**

Pectenogammarus festai (Ruffo, 1937) **new combination**

Pectenogammarus foxi (Schellenberg, 1928) **new combination**

Pectenogammarus monomerus (Stock, 1977) **new combination**

Pectenogammarus nudus (Stock, 1971) **new combination**

Pectenogammarus olivii (Milne Edwards, 1830) **new combination**

Pectenogammarus planicrurus (Reid, 1940) (see Stock (1968); Pinkster (1993)).

Pectenogammarus platvoeti (Pinkster, 1993) **new combination**

Pectenogammarus pungens (Milne Edwards, 1840) **new combination**

Pectenogammarus ruffoi (Pinkster & Stock, 1970) **new combination**

Pectenogammarus simoni (Chevreux, 1894) **new combination**

Pectenogammarus scutarensis (Schäferna, 1923) **new combination**

Pectenogammarus spinulicornis (Pinkster & Stock, 1971) **new combination**

Pectenogammarus stammeri (S. Karaman, 1931) **new combination**

Pectenogammarus stocki (G.S. Karaman, 1970) **new combination**

Pectenogammarus tabu (G.S. Karaman, 1971) **new combination**

Pectenogammarus thoni (Schäferna, 1923) **new combination**

Pectenogammarus tibaldii (Pinkster & Stock, 1970) **new combination**

Pectenogammarus veneris (Heller, 1865) **new combination**

Discussion

According to Hou & Sket (2016) and Sket & Hou (2018) the genus *Echinogammarus* should be restricted to the species formerly included in the *E. berilloni* species-group (*sensu* Pinkster 1973). Since *Echinogammarus* in Sket & Hou (2018) is monophyletic and well characterized morphologically (Pinkster 1973, 1979; Platvoet & Pinkster 1980; Fernández Lop 1987; Hou & Sket 2016; Sket & Hou 2018), all remaining species previously assigned to *Echinogammarus* but not included in the *E. berilloni* species group (see Pinkster 1993) should be transferred to other genera. Hou & Sket (2016) and Sket & Hou (2018) already transferred most of the species previously included in *Echinogammarus* (but not in the *E. berilloni* species group) to *Chaetogammarus*, *Marinogammarus*, *Parhomoeogammarus*, and specially to *Pectenogammarus* ("*Homoeogammarus* Schellenberg, 1937"); nevertheless,

to avoid breaking the well-funded monophyly of *Echinogammarus*, those species of *Echinogammarus* not included in the *E. berilloni* species group, but not yet assigned with molecular information to any particular genus, should be included temporarily in *Pectenogammarus*.

Pectenogammarus as discussed here, remains in need of a deep taxonomic revision based on molecular and morphological data combined. We hope that widely used data bases (e.g., Horton *et al.* 2022) will keep-up with the most recent phylogenetic information and its taxonomic consequences (e.g., Hou & Sket 2016; Sket & Hou 2018), rather than retaining species in non-monophyletic assemblages, most of them based on homoplastic character states. By using the clades resulting from the mentioned phylogenies, the internal taxonomy of proposed monophyletic groups (i.e., *Pectenogammarus sensu nov.*, *Dinarogammarus*, *Echinogammarus*, etc.) can be worked-out at a much closer scale (taxonomic and geographic), facilitating identification of useful morphological taxonomic characters, although homoplastic at other scales, and species recognition.

Acknowledgements

We thank Miguel A. Alonso Zarazaga for a critical review of the nomenclatural acts included in this work. We also thank the librarians of the Museo Nacional de Ciencias Naturales (especially Ignacio Pino) for their constant help.

This work has been possible thanks to the project grant PID2019-110243GB-100 /AEI/10.13039/50110001103 3 (Ministerio de Ciencia, Innovación y Universidades, Spain) (PI: M.G.-P.). PCR-F is supported by a Biodiversity postdoctoral fellowship from the MCZ (Museum of Comparative Zoology) at Harvard University.

References

- Barnard, J.L. & Barnard, C.M. (1983) *Freshwater Amphipoda of the world. Vol. II*. Hayfield Associates, Mt. Vernon, Virginia, 830 pp.
- Catta, J.D. (1878a) Note sur le *Gammarus berilloni* (n. sp.). *Bulletin de la Société de Borda, Dax*, 1878 (1er trimestre), 69–73.
- Catta, J.D. (1878b) Sur un Amphipode nouveau, le *Gammarus rhipidiophorus*. *Actes de la Société Helvétique des Sciences Naturelles Reunie a Bex les 20, 21 et 22 Aout 1877*, 60e Session, Compte-Rendu, 1876/1877, 256–263.
- Chevreux, E. (1894) *Gammarus simoni*, nov. sp., amphipode des eaux douces d'Algérie et de Tunisie. *Bulletin de la Société Zoologique de France*, 19, 171–176.
<https://doi.org/10.5962/bhl.title.15972>
- Chevreux, E. & Gauthier, H. (1924) Descriptions d'un nouveau Gammare de Tunisie. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 15, 53–55.
- Dahl, E. (1958) Fresh and brackish water amphipods from the Azores and Madeira. *Boletim do Museu Municipal do Funchal*, 11 (27), 5–25.
- Duhig, M. & Humphries, C. (1955) *Amphilochus brunneus* Della Valle, an amphipod new to Britain and Ireland and notes on other amphipods not previously recorded in Irish coastal waters. *Proceedings of the Royal Irish Academy, Section B*, 57 (8), 123–129.
- Fabricius, J.C. (1775) *Systema Entomologiae, sistens Insectorum Classes, Ordines, Genera, Species, adjectis Sysnonymis, Locis, Descriptionibus, Observationibus*. Kortii, Flensburgi et Lipsiae, 832 pp.
<https://doi.org/10.5962/bhl.title.36510>
- Fernández Lop, A. (1987) *Echinogammarus navaensis* (Crustacea, Amphipoda) a new species of the *berilloni*-group from Asturias (northern Spain). *Bulletin Zoologisch Museum*, 11 (1), 1–7.
- Grabowski, M., Bacela-Spychalska, K. & Pešić, V. (2014) Reproductive traits and conservation needs of the endemic gammarid *Laurogammarus scutarensis* (Schäferna, 1922) from the Skadar Lake system, Balkan Peninsula. *Limnologica*, 47, 44–51.
<https://doi.org/10.1016/j.limno.2014.04.001>
- Horton, T., Lowry, J., De Broyer, C., Bellan-Santini, D., Coleman, C.O., Corbari, L., Costello, M.J., Daneliya, M., Dauvin, J.-C., Fišer, C., Gasca, R., Grabowski, M., Guerra-García, J.M., Hendrycks, E., Hughes, L., Jaume, D., Jazdzewski, K., Kim, Y.-H., King, R., Krapp-Schickel, T., LeCroy, S., Lörz, A.-N., Mamos, T., Senna, A.R., Serejo, C., Sket, B., Souza-Filho, J.F., Tandberg, A.H., Thomas, J.D., Thurston, M., Vader, W., Väinölä, R., Vonk, R., White, K. & Zeidler, W. (2022) World Amphipoda Database. *Echinogammarus* Stebbing, 1899. Available from: <https://www.marinespecies.org/amphipoda/aphia.php?p=taxdetails&id=101535> (accessed 2 January 2022)
- Hou, Z. & Sket, B. (2016) A review of Gammaridae (Crustacea: Amphipoda): the family extent, its evolutionary history, and taxonomic redefinition of genera. *Zoological Journal of the Linnean Society*, 176 (2), 323–348.
<https://doi.org/10.1111/zoj.12318>

- ICZN (1999) *International Code of Zoological Nomenclature (ICZN)*. International Commission on Zoological Nomenclature, London, 306 pp.
- Kant, P., Pinkster, S. & Stock, J. (1968) Sur la présence en la Méditerranée de *Pectenogammarus planicrurus* Reid, 1940. *Vie et Milieu, Série A, Biologie Marine*, 19 (1-A), 207–209.
- Karaman, G.S. (1969) XXVI. Beitrag zur Kenntnis der Amphipoden. Taxonomie und Verbreitung der Art *Neogammarus rhipidiophorus* (Catta) in Mittelmeerbassin. *Glasnik Republickog Zavoda Za Zastitu Prirode I Prirodnjake Zbirke u Titograd*, 2, 47–58.
- Karaman, G.S. (1970) XXV. Prilog poznavanju Amphipoda Kriticka zapazanja o vrstama *Echinogammarus acarinatus* (S. Kar. 1931) i *Echinogammarus stocki* n. sp. *Poljoprivreda I Sumarstvo, Titograd*, 16 (1–2), 193–214.
- Karaman, G.S. (1971) XIX. Beitrag zur Kenntnis der Amphipoden. Eine neue Art der Gattung *Sarothrogammarus* (Gammaridae) aus Afghanistan, *S. ruffoi* n. sp. *Crustaceana*, 20 (2), 199–207.
<https://doi.org/10.1163/156854069X00222>
- Karaman, G.S. (1984) 135. Contribution to the knowledge of the Amphipoda. The Genus *Laurogammarus*, n. gen. (Fam. Gammaridae) in Yugoslavia. *Bilten Društva ekologa Bosne i Hercegovine*, Serie B, Naučni skupovi i savjetovanja, No. 3: III kongres ekologa Jugoslavije, Sarajevo, Knj. II, 29–35. Not available
- Karaman, S. (1931) III. Beitrag zur Kenntnis der Amphipoden Jugoslaviens, sowie einiger Arten aus Griechenland. *Prirodoslovne Rasprave*, 1, 31–66.
- Martynov, A.V. (1925) Études sur les Crustacees de mer du bassin du bas Don et leur distribution ethologique. *Akademii Nauk SSSR, Leningrad, Ezhegodnik Zoologicheskogo Muzeja*, 25 (1924), 1–115, pl. 1.
- Milne Edwards, H. (1830) Extrait de recherches pour servir à l'histoire naturelle des crustacés amphipodes. *Annales des Sciences Naturelles*, 20, 353–399, pls. 10–11.
<https://doi.org/10.5962/bhl.part.12300>
- Milne Edwards, H. (1840) *Histoire naturelle des crustacés: comprenant l'anatomie, la physiologie et la classification de ces animaux. Vol. 3*. Collection Suites a Buffon. Roret, Paris, 638 pp.
- Morgan, E. (1970) The effect of environmental factors on the distribution of the amphipod *Pectenogammarus planicrurus*, with particular reference to grain size. *Journal of the Marine Biological Association of the United Kingdom*, 50 (3), 769–785.
<https://doi.org/10.1017/S0025315400005038>
- Pinkster, S. (1969) A new *Echinogammarus* of the berilloni-group, *E. aquilifer* nov.sp., from the Pyrenees (Crustacea, Amphipoda). *Bulletin Zoologisch Museum Universiteit van Amsterdam*, 1 (11), 137–150.
- Pinkster, S. (1973) The *Echinogammarus berilloni*-group, a number of predominantly Iberian amphipod species (Crustacea). *Bijdragen tot de Dierkunde*, 43 (1), 1–38.
<https://doi.org/10.1163/26660644-04301001>
- Pinkster, S. (1993) A revision of the genus *Echinogammarus* Stebbing, 1899 with some notes on related genera (Crustacea, Amphipoda). *Memorie del Museo Civico di Storia Naturale*, Serie 2^a, Sezione Scienze della Vita, Verona, 10, 1–185.
- Pinkster, S. & Platvoet, D. (1986) Remarks on the identity of *Echinogammarus thoni* (Schäferna, 1922) with description of a new species, *Echinogammarus cyrtus*, from southern France (Crustacea, Amphipoda). *Bulletin Zoologisch Museum Universiteit van Amsterdam*, 10 (20), 173–181.
- Pinkster, S. & Stock, J.H. (1970) On three new species of *Echinogammarus*, related to *E. veneris* (Heller, 1865), from Italy and Switzerland (Crustacea, Amphipoda). *Beaufortia*, 17 (228), 85–104.
- Pinkster, S. & Stock, J.H. (1971) Sur deux *Echinogammarus limniques* nouveaux du Bassin d'Aquitaine. *Bijdragen tot de Dierkunde*, 41, 37–51.
<https://doi.org/10.1163/26660644-04101006>
- Platvoet, D. & Pinkster, S. (1980) A new species of *Echinogammarus* from Northern Spain, *Echinogammarus pseudoaquilifer* nov. spec. (Crustacea, Amphipoda). *Bulletin Zoologisch Museum Universiteit van Amsterdam*, 7 (2), 13–16.
- Pinkster, S. & Stock, J.H. (1972) Members of the *Echinogammarus simoni*-group and the genus *Eulimnogammarus* (Crustacea - Amphipoda) from Iberian Peninsula and North Africa, with descriptions of a new species. *Bulletin Zoologisch Museum Universiteit van Amsterdam*, 2 (10), 85–115.
- Reid, D. (1940) On *Gammarus (Pectenogammarus) planicrurus* subg. et sp. n. (Crust. Amph.). *Annals and Magazine of Natural History*, Series 11, 6, 287–292.
<https://doi.org/10.1080/03745481.1940.9723680>
- Ruffo, S. (1937) Una nuova specie di *Gammarus* del Mar Ligure. *Annali del Museo Civico di Storia Naturale Giacomo Doria*, 59, 438–446.
- Schäferna, K. (1923) Amphipoda balcanica, spolu s poznámkami o jiných sladkovodních Amphipodech. *Mémoires de la Société Royale des Sciences de Bohême [Věstník Královské České Společnosti Nauk], Classe des Sciences, Année 1921–1922*, 12, 1–111, pls. 1–11. [<https://www.biodiversitylibrary.org/page/42558753#page/1/mode/1up>]
- Schellenberg, A. (1928) Report on Amphipoda. Zoological results of the Cambridge Expedition to the Suez Canal. *Transactions of the Zoological Society of London*, 22, 633–692.
<https://doi.org/10.1111/j.1096-3642.1928.tb00209.x>
- Schellenberg, A. (1937) Schlüssel und Diagnosen der dem Süsswasser-*Gammarus* nahestehenden Einheiten ausschließlich der Arten des Baikalsees und Australiens. *Zoologischer Anzeiger*, 117, 267–280.

- Schellenberg, A. (1943) Portugiesische Süßwasser-Amphipoden. *Memorias e estudos do Museu Zoológico da Universidade de Coimbra*, 139, 1–6.
- Sket, B. & Hou, Z. (2018) Family Gammaridae (Crustacea: Amphipoda), mainly its *Echinogammarus* clade in SW Europe. Further elucidation of its phylogeny and taxonomy. *Acta Biologica Slovenica*, 61, 93–102.
- Stebbing, T.R.R. (1899) Amphipoda from the Copenhagen Museum and other sources II. *Transactions of the Linnean Society of London, Zoology*, 7 (8), 395–432, pls. 30–35.
<https://doi.org/10.1111/j.1096-3642.1899.tb00202.x>
- Stock, J.H. (1968) A revision of the European species of the *Echinogammarus pungens* - group (Crustacea, Amphipoda). *Beaufortia*, 16, 13–78.
- Stock, J.H. (1971) A revision of the *Sarathrogammarus*—Group (Crustacea, Amphipoda). *Bijdragen tot de Dierkunde*, 41 (2), 94–129.
<https://doi.org/10.1163/26660644-04102002>
- Stock, J.H. (1977) The non-marine Gammarids of the Balearic Islands. *Boletín de la Sociedad de Historia Natural de Baleares*, 22, 17–47.
- Stock, J.H. (1982) Validité du genre *Pectenogammarus* Reid, 1940 et distribution de son espèce-type, *P. planicrurus* (Reid, 1940) (Crustacea, Amphipoda, Gammaridae). *Cahiers de Biologie Marine*, 23, 325–329.
- van Maren, M.J. (1975a) Some notes on the intertidal gammarids (Crustacea, Amphipoda) from the Atlantic coast of the Iberian peninsula. *Beaufortia*, 23 (305), 153–168.
- van Maren, M.J. (1975b) The biology of *Chaetogammarus marinus* (Leach) and *Eulimnogammarus obtusatus* (Dahl), with some notes on other intertidal gammarid species (Crustacea, Amphipoda). *Bijdragen tot de Dierkunde*, 45 (2), 205–224.
<https://doi.org/10.1163/26660644-04502005>