USE OF THERMAL UNITS TO ESTIMATE CORN CROP COEFFICIENTS UNDER SEMIARID CLIMATIC CONDITIONS

A. Martínez-Cob

Estación Experimental Aula Dei (CSIC), Apdo. 202, 50080 Zaragoza (Spain). Phone: +34 976716086. Fax: +34 976716145. E-mail: macoan@eead.csic.es

Short running title

Estimation of corn crop coefficient using thermal units

ABSTRACT

Two crop coefficient equations were derived as a function of fraction of thermal units from lysimeter measured corn evapotranspiration (ET_{c-lys}) during 1997 and 1998, and reference evapotranspiration obtained from: a) lysimeter measurements ($K_{c_{mes}}$) or FAO Penman-Monteith (ET_{o-PM}) estimates ($K_{c_{est-PM}}$). For validation, corn evapotranspiration (ET_{c-est}) was estimated in 2005 and 2006 from ET_{o-PM} and: a) the equation for $K_{c_{mes}}$ with ($ET_{c-est-lyslc}$) or without ($ET_{c-est-lys}$) locally calibrated ET_{o-PM} ; b) the equation for $K_{c_{est-PM}}$; and c) the FAO approach ($ET_{c-est-FAO}$). The $ET_{c-est-lys}$ estimates showed the lowest bias (0.09 mm day^{-1}); the $ET_{c-est-PM}$ and $ET_{c-est-FAO}$, the highest ($0.50\text{-}0.51 \text{ mm day}^{-1}$). However, the root mean square error (RMSE, $1.23\text{-}1.27 \text{ mm day}^{-1}$) and the index of agreement (IA, around 0.94) of the $ET_{c-est-lys}$, $ET_{c-est-lyslc}$ and $ET_{c-est-PM}$ were similar. Therefore, $ET_{c-est-lys}$ is recommended although the $ET_{c-est-lyslc}$ was almost as accurate. The $ET_{c-est-PM}$ is less recommended due to poorer bias and systematic mean square error, and a general underestimation except for low corn ET values. For real time irrigation scheduling, the $ET_{c-est-FAO}$

should be avoided as RMSE (1.35 mm day^{-1}), IA (0.93) and bias were slightly worse, corn ET was overestimated but for high values, and the length of the four phenological stages must be known in advance.

1. INTRODUCTION

When estimating crop water requirements, it is common to compute crop coefficient (K_c) values as a function of the length of the four phenological stages in which crop development is divided following guidelines from the Food and Agriculture Organization of the United Nations (FAO) (Allen et al., 1998). Once these phenological stages are established, three K_c values are estimated to define the K_c curve. This estimation requires that these phenological stages, and the average soil wetting conditions (initial stage) and minimum relative humidity and wind speed (mid-season and final stages), be defined early in the season, which is an important limitation to the use of the FAO approach for estimation of crop water requirements on a real-time basis. A common practice is to use the same K_c curve for different years without taking into consideration the influence of environment and other factors over crop development for a particular year. Subsequently, crop evapotranspiration estimates becomes a function only of estimated reference evapotranspiration (ET_o).

Other alternative approaches have been proposed over the last years to estimate K_c curves for annual crops as a function of time in terms of days after sowing (DAS) or month of the year (Wright, 1982; Wright, 1991; Devitt et al., 1992; Steele et al., 1996; Nielsen and Hinkle, 1996; Tyagi et al., 2000a, b, 2003; Sepaskhah and Andam, 2001). This approach is easy to implement but, as with the FAO methodology, it does not take into account the influence of environmental and cultural factors on the rate of canopy development. For this reason, several attempts

1 have been made to present Kc curves as a function of other variables more related to
2 crop development: leaf area index (Kang et al., 2003), percent canopy that shades
3 the ground (Grattan et al., 1998) or thermal-based index expressed as cumulative
4 growing degree days or thermal units (TU) (Amos et al., 1989; Nielsen and Hinkle,
5 1996; Steele et al., 1996; Hunsaker, 1999; Sepaskhah and Andam, 2001).

6 The use of TU to estimate Kc curves has the advantage that air temperature
7 data is readily available and there is enough evidence of the influence of such
8 variable on crop development (Ritchie and NeSmith, 1991). Equations to estimate Kc
9 curves for corn as a function of TU have been developed in previous studies where
10 the required experimental Kc values were obtained from measured corn
11 evapotranspiration (ET) by using the neutron probe, and ET_o estimates from
12 meteorological data using several methods: Jensen-Haise (Amos et al., 1989; Steele
13 et al., 1996), Penman-Monteith (Nielsen and Hinkle, 1996; Kang et al., 2003),
14 Cuenca-Nicholson Penman (Sammis et al., 1985), and Allen Penman (Steele et al.,
15 1996). A constraint of these studies is the uncertainty linked to each ET_o estimation
16 method which may require previous local calibration. Subsequently, different Kc
17 curves have been obtained for the same corn ET datasets depending on the method
18 used to estimate ET_o (Steele et al., 1996). The maximum value that a given Kc curve
19 can attain is affected by the accuracy of the method used for estimation of ET_o . For a
20 given corn ET dataset, those ET_o methods that have the largest overestimates of the
21 true ET_o will lead to lower maximum Kc values when compared to those ET_o methods
22 in closer agreement with true ET_o . The opposite would occur for ET_o estimation
23 methods underestimating true ET_o . In addition, the use of neutron probe readings did
24 not provide measured corn ET values at time scales shorter than weekly or 10-day
25 periods (Amos et al., 1989; Nielsen and Hinkle, 1996; Steele et al., 1996). For real-

time irrigation scheduling of corn in semiarid climates, it would be more appropriate to develop Kc curves from daily experimental values.

Sammis et al. (1985), Steele et al. (1996) and Kang et al. (2003) used the absolute values of cumulative TU to develop their corn Kc estimation equations. However, Amos et al. (1989) and Nielsen and Hinkle (1996) used the fraction of TU (FTU), i.e. the ratio of cumulative TU for day *i* to total TU from emergence to physiological maturity. The use of FTU allows a general application of the Kc curve across cultivars requiring different TU totals from emergence to physiological maturity (Amos et al., 1989). In any case, the equations from Amos et al. (1989) and Nielsen and Hinkle (1996) may not be adequate for some climatic conditions due to the low peak estimated Kc values (about 1.0 or less); these equations were developed for sub-humid and water-limitation conditions and that peak value of 1.0 is well below the expected value under semi-arid and well irrigated conditions (Allen et al., 1998).

The goals of this paper were: 1) to develop two equations to estimate daily corn Kc values as a function of FTU from either lysimeter measured or FAO Penman-Monteith estimated reference evapotranspiration; 2) to validate these equations using additional lysimeter measured corn evapotranspiration values; and 3) to evaluate whether the use of Kc values obtained as a function of FTU instead of using the FAO methodology improved the accuracy of the corn ET estimations.

2. MATERIAL AND METHODS

2.1. Development of crop coefficient curves

The research was conducted on an experimental farm located in the middle Ebro River Valley (NE Spain), along the terraces of Gállego River (41°43'09" N latitude, 0°49'11" W longitude, altitude 225 m), about 8 km from where both rivers

meet. This is an irrigated area of about 5-7 km width stretching along the Gállego River for about 20-25 km. Soils in the experimental site are Typic Xerofluvent. Most crops in the area are corn, alfalfa, other pastures, fruit tree orchards, vegetables, and natural riparian vegetation. The climate is semiarid Mediterranean continental with an average annual precipitation of about 330 mm (Faci et al., 1994). Air temperature averages 5.5 °C in January and 24.4 °C in July, with 14.6 °C as the annual average.

Measurements were taken over two adjacent plots, A (1.3 ha, 130 m x 100 m) and B (1.0 ha, 100 m x 100 m). During 1997 and 1998, plot A was uniformly covered with grass (*Festuca arundinacea* Moench. cv. Demeter), while plot B was cultivated with corn (*Zea mays* cv. Juanita) sown on May 8 (1997), and May 14 (1998). Distance within adjacent rows was 0.75 m and average plant density was about 80.000 plants ha⁻¹. Corn was harvested on October 24 (1997) and October 22 (1998). A weighing lysimeter, 1.7 m depth and 6.3 m² effective surface area, was located in the center of each plot (Martínez-Cob, 2001). Identical management practices (sprinkler irrigation, fertilization, grass clippings) were regularly performed in both the lysimeter and surrounding plot, as required to keep both crops as close as possible to the optimum growing conditions (Allen et al., 1998). Martínez-Cob (2001) provides more detailed description of the lysimeters, including how 30-min ET rates were obtained from lysimeter mass losses. Half-hour ET rates were summed up to obtain daily lysimeter measured reference (ET_{o-lys}) and corn (ET_{c-lys}) ET rates.

Days during which the above mentioned management practices were performed were discarded. Thus, only those days (62 for 1997 and 63 for 1998), with both ET_{c-lys} and ET_{o-lys} available (both expressed in mm day⁻¹), were retained to obtain measured crop coefficient (K_{c_{mes}}) values (Allen et al., 1998):

$$Kc_{mes} = \frac{ET_{c-lys}}{ET_{o-lys}} \quad (1)$$

An automatic weather station (CR10 Campbell Scientific) was located next to the grass lysimeter. Daily averages of air temperature and relative humidity (at 1.5 m above ground), solar radiation and wind speed (at 2.0 m above ground) were recorded during the 1997 and 1998 growing periods, using a Vaisala probe (model HMP35C), a Skye pyranometer (model SP1110), and a Vector switching anemometer (model A100R), respectively. Daily ET_o values were estimated in both years using the FAO Penman-Monteith method (ET_{o-PM}) from those daily meteorological averages. The FAO Penman-Monteith method is completely described in Allen et al. (1998). Thus, a second set of experimental (estimated) crop coefficient (Kc_{est-PM}) values were derived (both ET_{c-lys} and ET_{o-PM} expressed in mm day⁻¹) as:

$$Kc_{est-PM} = \frac{ET_{c-lys}}{ET_{o-PM}} \quad (2)$$

For this set of experimental Kc values, 91 and 98 observations of daily ET_{c-lys} and ET_{o-PM} were retained in 1997 and 1998, respectively. The FAO Penman-Monteith method has become the reference crop definition (Allen et al., 1998) and it is being implemented in most of the automatic weather station networks used to provide real-time crop water requirement recommendations to farmers throughout the world. For this reason, it was considered relevant in this work to study the differences between the Kc curves obtained using this method and those obtained using measured ET_o .

The daily averages of air temperature were used to compute cumulative thermal units as follows (Ritchie and NeSmith, 1991):

$$\begin{aligned} \text{1} \quad \text{TU}_i &= \text{TU}_{i-1} + (T_a - T_b)_i && \text{if } T_a > T_b \\ \text{2} \quad \text{TU}_i &= \text{TU}_{i-1} && \text{if } T_a \leq T_b \end{aligned} \quad (3)$$

2 where: TU_i , cumulative thermal units for day i ($^{\circ}\text{C}$); TU_{i-1} , cumulative thermal units for
 3 day $i-1$ ($^{\circ}\text{C}$); $(T_a - T_b)_i$, difference between air and basal temperatures for day i ($^{\circ}\text{C}$).
 4 Basal temperature was assumed to be 8°C (Kiniry, 1991). The total cumulative
 5 thermal units for the 1997 and 1998 corn seasons (sowing to harvest) were 2108 and
 6 2028 $^{\circ}\text{C}$, respectively. Cavero et al. (1999) reported cumulative TU of 2250, 2116,
 7 and 1938 $^{\circ}\text{C}$ during the 1994, 1995 and 1996 corn (cv. Juanita) growing periods,
 8 respectively, at the same study area. The average of these five values 2088 $^{\circ}\text{C}$
 9 (coefficient of variation, 5.5 %), a value about 10 % higher of the cumulative TU
 10 reported as typical for corn in Spain by Boons-Prins et al. (1993). So that it was
 11 assumed in this experiment that total TU of 2090 $^{\circ}\text{C}$ is a good estimation for corn
 12 growing period. Subsequently, values of TU_i (Eq. 3) were expressed as the fraction of
 13 total cumulative thermal units for that day i (FTU_i): $\text{FTU}_i = \text{TU}_i / 2090$.

14 Next, experimental crop coefficients, Kc_{mes} and $\text{Kc}_{\text{est-PM}}$, were plotted against
 15 FTU_i values for both years for those days with available data. The plots suggested
 16 polynomial regressions would be adequate:

$$\text{17} \quad \text{Kc} = C'_0 + C'_1 \text{FTU}_{(\text{tr})i} + C'_2 \text{FTU}_{(\text{tr})i}^2 + \dots + C'_m \text{FTU}_{(\text{tr})i}^m \quad (4)$$

18 where: C'_0 to C'_m are the regression parameters; $\text{FTU}_{(\text{tr})i} = \text{FTU}_i - \overline{\text{FTU}}$. Note that the
 19 independent variable was expressed as a deviation around its mean $\overline{\text{FTU}}$. The
 20 reason for doing so was to substantially reduce the multicollinearity problem, i.e.
 21 correlation between independent variables (Neter et al., 1983). Average ($\overline{\text{FTU}}$) of the
 22 125 experimental FTU values available for Kc_{mes} was 0.518, while the corresponding
 23 $\overline{\text{FTU}}$ of the 189 experimental FTU values available for $\text{Kc}_{\text{est-PM}}$ was 0.548. The

criterion of the change in the adjusted coefficient of determination (R_{aj}^2) as the polynomial order increased was used to select the appropriate m^{th} order of the polynomial fit for each set of experimental Kc values (Neter et al., 1983). Values of R_{aj}^2 were used instead of coefficient of determination (R^2) to take into account the sample size (Neter et al., 1983). Thus, two Kc equations were developed, the first one from the $K_{c_{mes}}$ values and the second one from the $K_{c_{est-PM}}$ values. Once a polynomial regression equation was fit to the experimental values, the parameters C'_0 to C'_m were back-transformed in terms of the original units as described by Neter et al. (1983). The back-transformation equations varied depending upon the m^{th} order polynomial selected. Thus, those back-transformation equations are described in section 3.1, once an appropriate m^{th} order was selected in each case.

2.2. Validation of crop coefficient curves

The validation of the two developed Kc equations was performed with corn evapotranspiration data gathered during 2005 and 2006 in the experimental plot A used for development of the equation, and sown with corn (cultivar Pioneer PR34N43) on April 27, for both years. Distance within adjacent rows also was 0.75 m and average plant density was about 80.000 plants ha^{-1} . Corn was harvested on October 5 (2005), and October 2 (2006). Identical management practices (sprinkler irrigation, fertilization, etc.) were regularly performed in both the lysimeter and surrounding plot as required to keeping the crop as close as possible to the optimum growing conditions (Allen et al., 1998). The weighing lysimeter of plot A was used to record half-hour corn ET values that were summed up to get daily values (ET_{c-mes}). Only days without incidences (irrigations, fertilization, etc.) were retained for further analyses (98 days in each year).

For the validation, grass lysimeter values were not available. Therefore, ET_o estimates were obtained using the FAO Penman-Monteith method (ET_{o-PM}) (Allen et al., 1998) from daily averages of meteorological data (air temperature and relative humidity, wind speed and global solar radiation) recorded at an automatic weather station located over a grass plot next to the experimental plot. This station was equipped with a CR10X Campbell Scientific datalogger, a Vaisala probe (HMP45AC) for air temperature and relative humidity, a Skye SP1110 pyranometer, and a wind monitor Young 05103, all instruments located at 2.0 m height. These ET_{o-PM} estimates were applied in both Kc_{mes} and Kc_{est-PM} equations to get estimates of corn ET. The use of these ET_{o-PM} estimates could introduce some additional uncertainty during the validation of the Kc_{mes} equation if the FAO Penman-Monteith method would not exactly reproduce the measured ET_{o-lys} values. Although this method has been reported as highly accurate (Allen et al., 1998, 2005; Lecina et al., 2003), in this work, a previous comparison between ET_{o-lys} and ET_{o-PM} was performed using simple linear regression analysis ($y = b_0 + b_1 x$) and data gathered in 1997 and 1998 in order to calibrate the FAO Penman-Monteith equation for the study area if required during the validation of the Kc_{mes} equation. For this regression analysis, the measured ET_{o-lys} values were taken as the independent variable x , and the estimated ET_{o-PM} values were taken as the dependent variable y .

Likewise, the FAO methodology (Allen et al., 1998) also was applied to compute Kc estimates, $Kc_{est-FAO}$. To apply this methodology, it was necessary to divide the corn season in four phenological stages: 1) initial, from sowing to 10 % ground cover; 2) development, from 10 % to effective full (about 80 %) ground cover; 3) mid season, from full ground cover to senescence; and 4) final, from senescence to harvest. Then, ground cover data was required to define the end of the stages 1

1 and 2. Ground cover was taken as the average ratio of absorbed (at ground level) to
2 incident (above canopy) photosynthetically active radiation (PAR), expressed as
3 percentage. PAR values at each spot were measured weekly at 10 spots of 4
4 sampling areas within the experimental plot A using a SunScan Canopy Analysis
5 System (Delta-T Devices, Cambridge, UK) (Potter et al., 1996). These readings were
6 taken up to 90-95 % ground cover was attained. Senescence was detected by
7 weekly visual inspection of the four areas where ground cover was measured.

8 The procedures described by Allen et al. (1998) to compute single K_c values
9 were used to derive the three values defining a FAO $K_{c\text{est-FAO}}$ curve: 1) initial crop
10 coefficient, $K_{c\text{FAO-ini}}$; 2) mid-season crop coefficient, $K_{c\text{FAO-mid}}$; and 3) final crop
11 coefficient, $K_{c\text{FAO-end}}$. Number of irrigations and precipitation events and their
12 corresponding intensities were used to get a $K_{c\text{FAO-ini}}$ value for both 2005 and 2006.
13 Average minimum relative humidity and wind speed during the mid-season stage
14 were used to adjust tabulated $K_{c\text{FAO-mid}}$ value for corn to take into account local
15 climatic conditions. A value of $K_{c\text{FAO-end}} = 0.35$ as $K_{c\text{FAO-end}}$ should not be adjusted
16 when tabulated value is below 0.45 (Allen et al., 1998). Values of $K_{c\text{FAO-ini}}$, $K_{c\text{FAO-mid}}$
17 and $K_{c\text{FAO-end}}$ were used to derive daily $K_{c\text{est-FAO}}$ values for the whole season as
18 described by Allen et al. (1998).

19 Therefore, four sets of estimates of daily corn evapotranspiration ($ET_{c\text{-est}}$) were
20 obtained for 2005 and 2006 from the computed K_c values and the $ET_{o\text{-PM}}$ estimates:
21 1) $ET_{c\text{-est-lys}}$, using the $K_{c\text{mes}}$ equation (Figure 2); 2) $ET_{c\text{-est-lyslc}}$, using the $K_{c\text{mes}}$
22 equation (Figure 2) and the local calibration of the $ET_{o\text{-PM}}$ estimates (Figure 3); 3)
23 $ET_{c\text{-est-PM}}$, using the $K_{c\text{est-PM}}$ equation (Figure 2); and 4) $ET_{c\text{-est-FAO}}$, using a K_c curve
24 obtained applying the FAO methodology ($K_{c\text{est-FAO}}$). The corresponding daily
25 estimates of corn ET were compared against the measured corn ET values to

1 validate the Kc equations. Simple linear regression analyses ($y = b_0 + b_1 x$) were
 2 performed where the ET_{c-mes} and the ET_{c-est} values were the independent (x) and the
 3 dependent variables (y), respectively.

4 Regression is useful but not ideal when the goal is to quantify the agreement
 5 between estimates and measurements rather than fitting of a model to the
 6 measurements (Willmott, 1982; Kobayashi and Us Salam, 2000). Subsequently,
 7 additional statistics were computed to measure the difference between estimated and
 8 measured values (Willmott, 1982): mean estimation error (MEE), root mean square
 9 error (RMSE), systematic mean square error (MSE_s) and index of agreement (IA):

$$10 \quad MEE = \frac{\sum_{i=1}^N y_i - x_i}{N} \quad (5)$$

$$11 \quad RMSE = \left[\frac{\sum_{i=1}^N (y_i - x_i)^2}{N} \right]^{0.5} \quad (6)$$

12 where N is the number of days with available data.

$$13 \quad MSE_s = \frac{\sum_{i=1}^N (\hat{y}_i - x_i)^2}{N} \quad (7)$$

14 where $\hat{y}_i$ is the predicted value of corn ET using the corresponding simple linear
 15 regression between measured and estimated corn ET values.

$$16 \quad IA = 1 - \frac{\sum_{i=1}^N (y_i - x_i)^2}{\sum_{i=1}^N (|y_i - \bar{x}| + |x_i - \bar{x}|)^2} \quad (8)$$

17 where $\bar{x}$ is the average of the estimated corn ET values.

3. RESULTS AND DISCUSSION

3.1. Development of crop coefficient curves

Meteorological conditions in 1997 and 1998 were different (Figure 1). Averages of air temperature for the whole corn-season (9 May to 23 October in 1997, and 15 May to 21 October in 1998) were quite similar in both years, 20.9 °C in 1997, and 21.0 °C in 1998. Nevertheless, air temperature was somewhat higher in 1998 for late June to middle July, and somewhat lower late in the season. Minimum relative humidity was lower in 1998 for most of the season, and so whole corn-season averages were 46 % in 1997 and 41 % in 1998. Wind speed and estimated ET_{o-PM} were higher in 1998 for most of the period. Thus, whole corn-season averages of wind speed were 2.2 m s⁻¹ in 1997 and 2.8 m s⁻¹ in 1998, while whole corn-season averages of estimated ET_{o-PM} were 4.4 mm day⁻¹ in 1997 and 5.3 mm day⁻¹ in 1998.

In accordance with these differences in meteorological conditions, there also were differences between measured lysimeter ET values. Thus, average measured ET_{o-lys} was 4.3 mm day⁻¹ in 1997 and 5.0 mm day⁻¹ in 1998, while average measured ET_{c-lys} was 4.0 mm day⁻¹ in 1997 and 4.7 mm day⁻¹ in 1998. However, in terms of crop coefficient, differences between both years were smaller: average measured Kc_{mes} was 0.88 in 1997, and 0.94 in 1998, i.e. only 6 % higher in 1998, while ET rates were about 17 % higher. Minimum measured Kc_{mes} values were quite similar, 0.18 in 1997 and 0.17 in 1998, while maximum measured Kc_{mes} values only were about 5 % higher in 1998 (1.54 against 1.47 in 1997).

The values of R_{aj}^2 for the 2nd, 3rd, and 4th order polynomials fit to the Kc_{mes} values as a function of fraction of total cumulative thermal units (FTU) were 0.741, 0.771 and 0.778, respectively; correspondingly, the values of R_{aj}^2 for the 2nd, 3rd, and

4th order polynomials fit to the $Kc_{\text{est-PM}}$ values as a function of FTU were 0.647, 0.664 and 0.662, respectively. These R_{aj}^2 values were moderately high and represent that there still is an important fraction of Kc variability (about 25-35 %) that is not explained by FTU variability. This result should be expected as several factors affecting Kc variability have not been taken into account; for instance, the intensity and frequency of soil moistening during the initial stages of the crop when soil evaporation is more important than transpiration. In this work, attempts have been made to include other meteorological factors as relative humidity and wind speed but there were no improvements of the estimation equation of Kc in terms of R_{aj}^2 . It is likely that there was not enough variability in those meteorological variables as to have a significant effect in the Kc variability. Further research should consider the development of Kc curves including experimental data from several locations showing higher variability in those meteorological conditions.

The change of R_{aj}^2 as the polynomial order increased was used to select the appropriate m^{th} order of the polynomial fit for each set of experimental Kc values (Neter et al., 1983): a 3rd order polynomial in the case of Kc_{mes} , and a 2nd order polynomial, in the case of $Kc_{\text{est-PM}}$. In a later step, the regression parameters were back-transformed in terms of the original units according to the back-transformation equations described by Neter et al. (1983):

a) In the case of a 3rd polynomial regression fit:

$$\begin{aligned}
 C_0 &= C'_0 - C'_1 \overline{FTU} + C'_2 \overline{FTU}^2 - C'_3 \overline{FTU}^3 \\
 C_1 &= C'_1 - 2C'_2 \overline{FTU} + 3C'_3 \overline{FTU}^2 \\
 C_2 &= C'_2 - 3C'_3 \overline{FTU} \\
 C_3 &= C'_3
 \end{aligned} \tag{9}$$

1 b) In the case of a 2nd polynomial regression fit:

$$\begin{aligned} C_0 &= C'_0 - C'_1 \overline{FTU} + C'_2 \overline{FTU}^2 \\ C_1 &= C'_1 - 2C'_2 \overline{FTU} \\ C_2 &= C'_2 \end{aligned} \quad (10)$$

3 Figure 2 shows the experimental values of Kc_{mes} and Kc_{est-PM} plotted against
4 FTU as well as the polynomials fit in each case after back-transforming the
5 regression parameters to the original units using Eqs. (9) and (10). Scattering of
6 experimental values was relatively similar to that reported in previous works (Sammis
7 et al., 1985; Amos et al., 1989; Steele et al., 1996), the coefficients of determination
8 were slightly worse although it should be considered that this work used daily values
9 while previous works used 10-days or monthly averages so the time variability of
10 experimental Kc values was smoothed. The shape of both Kc curves (Figure 2) was
11 different as the fit polynomial order was different. In the case of Kc_{mes} , the period
12 before maximum Kc was reached was longer than the period after that maximum.
13 Maximum Kc value was 1.37, attained for FTU ranging from 0.57 to 0.65. In the case
14 of Kc_{est-PM} , the length of periods before and after maximum Kc value was more even.
15 That maximum was 1.17, attained for FTU ranging from 0.49 to 0.59. The duration of
16 maximum Kc value was slightly longer in the case of Kc_{est-PM} . Then, different Kc
17 curves and different fit Kc values were obtained depending upon the ET_o approach
18 used to derive experimental Kc values as pointed by Steele et al. (1996).

19 The FAO Penman-Monteith method has been shown to be very accurate for
20 estimation of daily ET_o in different climatic conditions (Allen et al., 1998; Allen et al.,
21 2005). This method has also been accurate for estimation of ET_o in semi-arid and
22 advective conditions as those occurring in the study area (Lecina et al., 2003;
23 Berengena and Gavilán, 2005). However, in this work, some overestimation has

1 been observed for the study period (May to October 1997 and 1998) as shown in
2 Figure 3. Averages of ET_{o-lys} and ET_{o-PM} values shown in Figure 3 were 4.7 and 5.1
3 $mm\ day^{-1}$, respectively. Therefore, the FAO Penman-Monteith method overestimated
4 lysimeter ET_o by about 9 % on average. MEE and RMSE values of this comparison
5 were 0.41 and 0.84 $mm\ day^{-1}$, while index of agreement (IA) was quite high, 0.936.
6 Likewise, standard deviations of measured and estimated ET_o were also quite
7 similar, 1.69 and 1.67 $mm\ day^{-1}$. These statistics suggested that the agreement
8 between ET_{o-lys} and ET_{o-PM} was quite high (Willmott, 1982) although some
9 overestimation was still observed. The regression coefficient shown in Figure 3 was
10 used for local calibration of the FAO Penman-Monteith by solving ET_{o-PM} as function
11 of ET_{o-lys} .

12 This overestimation of the FAO Penman-Monteith method partially explains
13 the difference between maximum Kc value obtained with the two fit equations (Figure
14 2). This difference also was due to the fact that the number of available days used to
15 fit the equations was different in both cases, Kc_{mes} and Kc_{est-PM} . If the same sample
16 size would have been used, maximum Kc value obtained with the fit equation in the
17 case of Kc_{est-PM} would have been 1.20. The different responses of Kc and ET_o to
18 other climatological conditions (relative humidity, wind speed, etc.) were also likely to
19 be responsible for the difference between maximum Kc values obtained with the fit
20 equations shown in Figure 2. If the FAO methodology would have been followed to
21 estimate Kc values for corn in the study area, a maximum value of 1.23 would have
22 been obtained as the average wind speed and minimum relative humidity were 2.3 m
23 s^{-1} and 40 %, respectively, for the mid-season stage, approximately from the third
24 decade of July to the first decade of September.

Maximum values of K_c obtained with the $K_{c_{mes}}$ equation were similar to those reported by Kang et al. (2003) for corn in a semiarid area of China. However, these authors obtained a K_c equation as a function of days after sowing (DAS) and the equation was developed for corn cultivars requiring about 30-40 days less to complete their cycle than the cultivar used in this work. As Kang et al. (2003) did not use fraction of DAS, the K_c equation developed by these authors is only limited to corn cultivars of similar cycle lengths. Tyagi et al. (2003) reported maximum K_c values of 1.23 for corn in semiarid areas of India when the FAO Penman-Monteith method was used to estimate the ET_o values required for computing K_c . These authors also reported other maximum values of K_c obtained with other ET_o methods and the same corn ET dataset, values that were higher or lower than 1.23, depending upon the method underestimated or overestimated the FAO Penman-Monteith method. Sammis et al. (1985) reported maximum K_c values of 1.11 for corn in semiarid areas of New Mexico (USA) when the Penman method was used to estimate ET_o . This method has been reported as highly overestimating ET_o (Jensen et al., 1990; Smith et al., 1991). Other studies on the development of K_c curves for corn as a function of thermal units have reported maximum values of 1.0, when local climatic conditions were sub-humid or humid (Nielsen and Hinkle, 1996; Steele et al., 1996) or corn was grown under water limiting conditions (Amos et al., 1989).

3.2. Validation of crop coefficient curves

Figure 4 shows the average meteorological conditions for 2005 and 2006. Average air temperature was warmer for these two years than 1997 and 1998. Thus, the corn whole-season averages of air temperature were 21.5 °C for 2005 (28 April to 3 October) and 21.7 °C for 2006 (28 April to 1 October). Likewise, whole-season averages of minimum relative humidity were lower (28 % for 2005, and 30 % for

2006) than those for 1997 and 1998. Whole-season averages of global solar radiation for 2005 and 2006 were intermediate to those recorded for 1997 and 1998. Whole-season average of wind speed for 2005 (2.3 m s^{-1}) was slightly higher to that of 1997 and 2006 (2.2 m s^{-1}), and noticeably lower than that of 1998 (2.8 m s^{-1}). Finally, whole-season averages of estimated $\text{ET}_{\text{o-PM}}$ for 2005 and 2006 (5.5 and 5.4 mm day^{-1} , respectively) were quite similar to that of 1998 (5.6 mm day^{-1}) but higher than that of 1997 (4.6 mm day^{-1}). Summarizing, 2005 and 2006 were quite similar to 1998 in terms of estimated $\text{ET}_{\text{o-PM}}$ but were warmer, drier and less windy. Compared to 1997, 2005 and 2006 were warmer and drier, with higher estimated $\text{ET}_{\text{o-PM}}$ and similar windy conditions.

Percentage ground cover during the validation seasons (2005 and 2006), as determined from PAR measurements, and visual inspection for senescence were used to divide the corn cycle in the four phenological stages defined by the FAO methodology (Allen et al., 1998): 1) initial stage up to 10 % ground cover, attained at 28 DAS in 2005, and 31 DAS in 2006; 2) development stage up to 80 % ground cover, attained at 58 DAS in both years; 3) mid-season stage up to senescence, attained at 128 DAS in both years; and 4) final stage up to harvest, 161 DAS in 2005, and 158 DAS in 2006. Initial crop coefficient ($\text{Kc}_{\text{FAO-ini}}$) was computed taken into account the following factors (Allen et al., 1998): 1) average water depth of wetting events (irrigation and precipitation) during the initial stage, about 16 mm in 2005, and 7 mm in 2006; 2) frequency of wetting events, about 6 days in 2005, and 3 days in 2006; and 3) average estimated $\text{ET}_{\text{o-PM}}$ during the initial stage, about 5.0 mm day^{-1} in both years. Thus, values of $\text{Kc}_{\text{FAO-ini}}$ of 0.48 and 0.59 were computed for 2005 and 2006, respectively. For the mid-season Kc ($\text{Kc}_{\text{FAO-mid}}$), the corresponding averages of wind speed (2.4 m s^{-1} in 2005 and 2.3 m s^{-1} in 2006) and minimum relative humidity

(27 % in both years) were used to adjust the tabulated value of 1.20 (Allen et al., 1998). Thus, $K_{cFAO-mid}$ was 1.28 in 2005 and 1.27 in 2006. Finally, the value of final stage crop coefficient ($K_{cFAO-end}$) was 0.35 in both years (section 2.2).

Figure 5 shows the graphs of estimated (ET_{c-est}) versus measured (ET_{c-lys}) corn ET for the four cases considered, $ET_{c-est-lys}$, $ET_{c-est-lyslc}$, $ET_{c-est-PM}$, and $ET_{c-est-FAO}$. Coefficients of determination were relatively high, about 80 %. These values suggest that there still was an important fraction of variability, about 20 %, not explained by the estimation methods. This result should be expected as crop development is highly but not completely affected by thermal units; other climatic, plant, soil and management factors should be considered to estimate Kc curves. On average, estimates of corn ET obtained using the K_{cmes} equation showed a high agreement (in the case of $ET_{c-est-lys}$ estimates) or a slight underestimation of measured ET_{c-lys} (in the case of $ET_{c-est-lyslc}$ estimates) (Figure 5). Corn ET was underestimated when using the $K_{c-est-PM}$ equation except for low ET values, and it was overestimated when using the $K_{c-est-FAO}$ approach except for high ET values (Figure 5).

Table 1 lists the error analysis statistics computed for the comparison of the four sets of estimated against the measured corn ET. The ratio of means, $\bar{y} / \bar{x}$, and the MEE values indicated that use of the K_{cmes} equation without local calibration for the ET_{o-PM} estimates provided the lowest bias (case $ET_{c-est-lys}$). The MEE was not significantly different than 0 ($\alpha = 0.95$). The close agreement between estimated and measured corn ET was also indicated by the intercept and the slope of the corresponding simple linear regression, which were not significantly different than 0 and 1 ($\alpha = 0.95$), respectively (Figure 5). Using these statistics as criteria, the $ET_{c-est-lys}$ could be considered as the best method for estimation of ET_{c-lys} of the four approaches being evaluated.

The use of the $K_{c_{mes}}$ equation with local calibration for the ET_{o-PM} estimates (case $ET_{c-est-lyslc}$) was also in close agreement with measured corn ET as indicated by an intercept and slope of the corresponding simple linear regression that were not different than 0 and 1 ($\alpha = 0.95$), respectively. However, the MEE value, although lower than 0.2 mm day^{-1} , was significantly different than 0 ($\alpha = 0.95$) and the ratio of means, $\bar{y} / \bar{x}$, suggested an average underestimation of almost 4 % (Table 1). In general terms, it could be concluded that both approaches $ET_{c-est-lys}$ and $ET_{c-est-lyslc}$ provided similar estimates of corn ET, $ET_{c-est-lys}$ being slightly better. At first glance, it would have been expected that the accuracy of the $ET_{c-est-lyslc}$ estimates would have been higher than that of the $ET_{c-est-lys}$ estimates. However, this did not occur probably due to the different local climatic conditions of the validation years (2005 and 2006) and those of 1997 and 1998 (Figures 1 and 4). Lecina et al. (2003) reported a slightly lower ET_o overestimation (about 5 %) by the FAO Penman-Monteith method for the years 1999 and 2000 in the same study area. Therefore, ET_o overestimation by this method observed for 1997 and 1998 (Figure 3) likely was somewhat higher than the 'true' overestimation of this method for the local climatic conditions of the study area.

Applying the $K_{c_{est-PM}}$ equation provided slightly worse corn ET estimates according to $\bar{y} / \bar{x}$ and MEE, which also was significantly different than 0 (Table 1). At first glance, it was expected that these two statistics would have been similar for the two approaches, $ET_{c-est-lyslc}$ and $ET_{c-est-PM}$. However, the different response of K_c and ET_o to climatic, crop and management conditions has likely been responsible for this higher bias of the $ET_{c-est-PM}$ estimates. Applying the $K_{c_{est-FAO}}$ estimates led to significant ($\alpha = 0.95$) overestimation as indicated by $\bar{y} / \bar{x}$ and MEE values, similar in magnitude to the underestimation observed for the $ET_{c-est-PM}$ approach. In terms of the systematic MSE, the two approaches using the $K_{c_{mes}}$ equation also seemed to be

the best methods (Table 1) although MSE_s values were also relatively small for the other two evaluated approaches, $ET_{c-est-PM}$ and $ET_{c-est-FAO}$. The quite low values of the MSE_s for the $ET_{c-est-lys}$ and $ET_{c-est-lyslc}$ approaches indicate that very little improvement of their accuracy can be attained without significant changes in the structure of the methods (Willmott, 1982).

When the other two error analysis statistics, RMSE and IA were considered, the performance of the four corn ET estimation approaches was more similar than suggested by $\bar{y} / \bar{x}$, MEE and MSE_s values. Thus, RMSE and IA values were almost the same for the $ET_{c-est-lys}$, $ET_{c-est-lyslc}$ and $ET_{c-est-PM}$ estimates (Table 1), while the $ET_{c-est-FAO}$ estimates were slightly worse according to these two statistics. When the performance of different methods is evaluated, the $\bar{y} / \bar{x}$ and MEE statistics have the problem that they may be small because of cancelling errors of similar magnitude but opposite sign. RMSE is more suited to overall measurement of model performance as it summarizes the mean difference in the units of the evaluated variable (Willmott, 1982; Kobayashi and Us Salam, 2000). However, RMSE do not provide information about the relative size of the average difference between estimated and measured values and about the nature of those differences. Relative difference measures, such as $RMSE/\bar{x}$ sometimes have been reported but this kind of indices are questionable because they are unbounded and unstable for $\bar{x}$ and/or N close to 0 (Willmott, 1982). This author has suggested the use of the index of agreement, which is both a relative and bounded measure. Likewise, Willmott (1982) has suggested computing the systematic and unsystematic portions of the mean square error (the root square of RMSE) as the systematic difference should approach 0 for a “good” method. Therefore, the interpretation of the differences between measured and estimated

values, and the performance of different methods should rely on the use of several statistics as they may offer complementary information each other.

Thus, taking into account the different statistics listed in Table 1 and Figure 5, it could be assumed that the use of the Kc_{mes} equation without local calibration for the FAO Penman-Monteith method has been the best method to estimate corn ET for the climatic conditions of the study area. The $ET_{c-est-lyslc}$ approach provided estimates having slightly higher bias but showing a quite similar overall difference to measured values, so the use of this approaches seems at least to be almost as reasonable as the use of the $ET_{c-est-lys}$ approach. The $ET_{c-est-PM}$ approach provided estimates having similar overall difference to measured values than the two above mentioned approaches, but the results of Figure 5 and the MEE and MSE_s statistics suggest that the $ET_{c-est-PM}$ was less accurate.

Regarding the $ET_{c-est-FAO}$ estimates, the RMSE and the IA values were slightly worse and the bias was relatively high (Table 1). An overall overestimation of the measured values was observed except for high corn ET values (Figure 5) which can be used as a criterion, together with the MEE and MSE_s statistics, to consider this method as the worst of the four evaluated. In any case, it should be realized that the $Kc_{est-FAO}$ curve in this study was computed after the experimental campaign was finished, once measured values of ground cover and senescence were available to divide corn development in the four phenological stages established by the FAO methodology (Allen et al., 1998). However, in general, for real time irrigation scheduling, this division in four phenological stages must be established before the crop attains them so it must be based in long-term average values that may not well represent the specific conditions for a given year. For this reason, it should be expected that the performance of the $ET_{c-est-FAO}$ estimates would have been worse in

this study if that 'long-term average phenological stage lengths' would have been taken. Thus, it could be concluded that the use of the $K_{c_{est-FAO}}$ approach for real time irrigation scheduling should be avoided for climatic conditions similar to those seen in this study when there is the possibility of using the other approaches evaluated.

4. CONCLUSIONS

A 3rd degree polynomial fit has been obtained to estimate crop coefficient (K_c) for corn as a function of fraction thermal units when experimental K_c values ($K_{c_{mes}}$) were obtained from measured corn and reference evapotranspiration using weighing lysimeters in 1997 and 1998. When experimental K_c values ($K_{c_{est-PM}}$) were derived using estimated reference evapotranspiration (FAO Penman-Monteith method), a 2nd degree polynomial fit was obtained. Adjusted coefficients of determination were relatively high, 0.77 in the case of $K_{c_{mes}}$ and 0.66 in the case of $K_{c_{est-PM}}$.

The two developed equations for K_c estimation were validated using measured corn evapotranspiration and estimated reference evapotranspiration values gathered in 2005 and 2006 in the same experimental plot. The error analysis statistics RMSE and IA indicate that the use of the equation developed for $K_{c_{mes}}$ provided similar estimates than the use of the equation developed for $K_{c_{est-PM}}$; the use of a local calibration factor to take into account the observed overestimation of the FAO Penman-Monteith equation did not improve the estimates. Based on the bias MEE, the MSE_s and the simple linear regression results, it would be suggested that the use of the $K_{c_{mes}}$ equation would provide slightly better corn evapotranspiration estimates particularly when using it without local calibration of the FAO Penman-Monteith. But the use of the $K_{c_{est-PM}}$ equation should also be assumed

as reasonable based on RMSE and IA, although likely less accurate based on MEE, MSE_s and the coefficients of the simple linear regression against measured corn ET.

The use of the FAO methodology to estimate corn K_c showed slightly worse results in terms of the various statistics computed. For real time irrigation scheduling, the use of this methodology should be avoided if it is possible to use FTU to estimate K_c as the FAO methodology requires dividing the corn development season in four phenological stages in advance and so the possible variations of corn development due to different climatic conditions for a particular year can not be taken into account. Therefore, the use of FTU to estimate corn K_c would slightly improve the uncertainty of the FAO methodology.

Acknowledgments

This research has been funded through projects HID96-1295-C04-04 and AGL-2004-06675-CO3-02 (Spanish Ministry of Education), and PIP090/2005 (Regional Government of Aragón). Special thanks are due to: Enrique Playán and José M. Faci for the design of the weighing lysimeters; José Caverro, Nery Zapata, Miguel Izquierdo, Jesús Gaudó, Daniel Mayoral and Juan Manuel Acín for their assistance in the maintenance of the experimental plot; M^a Pilar Paniagua and Eva Medina for their assistance to collect ground cover data.

REFERENCES

Allen R.G., Pereira L.S., Raes D., Smith M. 1998. Crop evapotranspiration: guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56. FAO, Rome, Italy.

1 Allen R.G., Walter I.A., Elliott R.L., Howell T.A., Itenfisu D., Jensen M.E., Snyder R.L.
2 (eds.). 2005. The ASCE Standardized Reference Evapotranspiration Equation.
3 Am. Soc. Civil Engrs., Reston, Virginia, USA.

4 Amos B., Stone L.R., Bark L.D. 1989. Fraction of thermal units as the base for
5 evapotranspiration crop coefficient curve for corn. Agron. J. 81: 713-717.

6 Berengena J., Gavilán P. 2005. Reference ET estimation in a highly advective semi-
7 arid environment. J. Irrig. Drain. Engr. ASCE. 131 (2): 147-163.

8 Boons-Prins E.R., De Koning G.H.J., Van Diepen C.A., Penning de Vries F.W.T.
9 1993. Crop specific simulation parameters for yield forecasting across the
10 European Community. Simulation Reports CABO-TT, N° 32. Centre for
11 Agrobiological Research (CABO-DLO) and Winand Staring Centre (SC-DLO),
12 Wageningen Agricultural University, Wageningen, Netherlands.

13 Cavero J., Zaragoza C., Suso M.L., Pardo A. 1999. Competition between maize and
14 *Datura stramonium* in an irrigated field under semi-arid conditions. Weed
15 Research. 39: 225-240.

16 Devitt D.A., Morris R.L., Bowman D.C. 1992. Evapotranspiration, crop coefficients,
17 and leaching fractions of irrigated desert turfgrass systems. Agron. J. 84: 717-723.

18 Faci J.M., Martínez-Cob A., Cabezas-Andrade A. 1994. Agroclimatología de los
19 regadíos del bajo Gállego: doce años de observaciones diarias en Montañana
20 (Zaragoza). Departamento de Agrícola, Ganadería y Montes, Diputación
21 General de Aragón, Zaragoza, Spain.

22 Grattan S.R., Bowers W., Dong A., Snyder R.L., Carroll J.J., George W. 1998. New
23 crop coefficients estimate water use of vegetables, row crops. California
24 Agriculture. 52 (1): 16-21.

- 1 Hunsaker D.J. 1999. Basal crop coefficients and water use for early maturity cotton.
2 Trans. ASAE. 42 (4): 927-936.
- 3 Jensen M.E., Burman R.D., Allen R.G. 1990. Evapotranspiration and Irrigation Water
4 Requirements. ASCE Manuals and Reports on Engineering Practice, No. 70. Am.
5 Soc. Civil Engrs., New York, USA.
- 6 Kang S.Z., Gu B.J., Du T.S., Zhang J.H. 2003. Crop coefficient and ratio of
7 transpiration to evapotranspiration of winter wheat and maize in a semi-humid
8 region. *Agricul. Water Manage.* 59 (3): 239-254.
- 9 Kiniry J.R. 1991. Maize Phasic Development. In: *Modeling Plant and Soil Systems*.
10 Hanks J., Ritchie J.T. (eds.). Series Agronomy N° 31. 55-70. American Society of
11 Agronomy, Crop Science Society of America, Soil Science Society of America,
12 Madison, WI, USA.
- 13 Kobayashi K., Us Salam M. 2000. Comparing simulated and measured values using
14 mean squared deviation and its components. *Agron. J.* 92: 345-352.
- 15 Lecina S., Martínez-Cob A., Pérez P.J., Villalobos F.J., Baselga J.J. 2003. Fixed
16 versus variable bulk canopy resistance for reference evapotranspiration estimation
17 using the Penman-Monteith equation under semiarid conditions. *Agric. Water*
18 *Manage.* 60 (3): 181-198.
- 19 Martínez-Cob A. 2001. Adequacy of Villalobos method to adjust eddy covariance
20 latent heat flux. *Irrig. Sci.* 20: 175-188.
- 21 Neter J., Wasserman W., Kutner M.H. 1983. *Applied linear regression models*. Irwin,
22 Homewood, Illinois, USA.

- 1 Nielsen D.C., Hinkle S.E. 1996. Field evaluation of basal crop coefficients for corn
2 based on growing degree days, growth stage, or time. Trans. ASAE. 39 (1): 97-
3 103.
- 4 Potter E., Wood J., Nicholl C. 1996. SunScan Canopy Analysis System: User
5 Manual. Document SS1-UM-1.05. Delta-T Devices Ltd, Cambridge, UK.
- 6 Ritchie J.T., NeSmith D.S. 1991. Temperature and crop development. In: Modeling
7 Plant and Soil Systems. Hanks J., Ritchie J.T. (eds.). Series Agronomy N° 31. 5-
8 29. American Society of Agronomy, Crop Science Society of America, Soil
9 Science Society of America, Madison, WI, USA.
- 10 Sammis T.W., Mapel C.L., Lugg D.G., Lansford R.R., McGuckin J.T. 1985.
11 Evapotranspiration crop coefficients predicted using growing-degree-days. Trans.
12 ASAE. 28 (3): 773-780.
- 13 Sepaskhah A.R., Andam M. 2001. Crop coefficient of sesame in a semiarid region of
14 I.R. Iran. Agricul. Water Manage. 49: 51-63.
- 15 Smith M., Allen R.G., Monteith J.L., Perrier A., Pereira L.S., Segeren A. 1991. Report
16 on the expert consultation on revision of FAO methodologies for crop water
17 requirements. FAO Land and Water Development Division, FAO, Rome, Italy.
- 18 Steele D.D., Sajid A.H., Prunty L.D. 1996. New corn evapotranspiration crop curves
19 for southeastern North Dakota. Trans. ASAE. 39 (3): 931-936.
- 20 Tyagi N.K., Sharma D.K., Luthra S.K. 2000a. Determination of evapotranspiration
21 and crop coefficients of rice and sunflower with lysimeter. Agricul. Water Manage.
22 45: 41-54.
- 23 Tyagi N.K., Sharma D.K., Luthra S.K. 2000b. Evapotranspiration and crop
24 coefficients of wheat and sorghum. J. Irrig. Drain. Engr. 126 (4): 215-222.

- 1 Tyagi N.K., Sharma D.K., Luthra S.K. 2003. Determination of evapotranspiration for
2 maize and berseem clover. *Irrig. Sci.* 21: 173-181.
- 3 Willmott C.J. 1982. Some comments on the evaluation of model performance. *Bull.*
4 *Am. Meteorol. Soc.* 63 (11): 1309-1313.
- 5 Wright J.L. 1982. New evapotranspiration crop coefficients. *J. Irrig. Drain. Div., Proc.*
6 *ASCE.* 108 (IR2): 57-74.
- 7 Wright J.L. 1991. Using weighing lysimeters to develop evapotranspiration crop
8 coefficients. In: *Lysimeters for evapotranspiration and environmental*
9 *measurements.* Allen R.G., Howell T.A., Pruitt W.O., Walter I.A., Jensen M.E.
10 (eds.). 191-199. *Proc. International Symposium on Lysimetry.* 23-25 July 1991.
11 Honolulu, Hawaii. Am. Soc. Civil Engrs., New York, USA.

Table 1. Error analysis statistics computed for the comparison between measured, (ET_{c-lys} , variable x), and estimated (ET_{c-est} , variable y) corn evapotranspiration for the validation years, 2005 and 2006. $\bar{x}$, mean of variable x; $\bar{y}$, mean of variable y; MEE, mean estimation error; RMSE, root mean square error; MSE_s , systematic mean square error; IA, index of agreement. Sample size N = 196.

ET_{c-est}	$\bar{y}$ (mm day ⁻¹)	$\bar{y} / \bar{x}^{(a)}$	MEE (mm day ⁻¹)	RMSE (mm day ⁻¹)	MSE_s (%)	IA
$ET_{c-est-lys}$	5.44	1.017	0.09 ^{ns0}	1.235	0.6	0.944
$ET_{c-est-lyslc}$	5.16	0.964	-0.19 ^{s0}	1.269	2.4	0.942
$ET_{c-est-PM}$	4.84	0.904	-0.51 ^{s0}	1.228	29.5	0.936
$ET_{c-est-FAO}$	5.86	1.094	0.50 ^{s0}	1.348	21.2	0.925

^(a) $\bar{x} = 5.35$ mm day⁻¹.

^{ns0} no significantly different than 0; ^{s0} significantly different than 0 ($\alpha = 0.95$).

Figure 1. Average meteorological conditions during 10-day periods between May to October 1997 and 1998. a) mean air temperature; b) minimum relative humidity; c) wind speed at 2.0 m height; and d) estimated reference evapotranspiration.

1

2 Figure 2. Polynomial equations fitted to experimental values of Kc_{mes} and Kc_{est-PM} ,
 3 obtained for 1997 and 1998, as a function of fraction of thermal units (FTU).

4

1

2 Figure 3. Regression of estimated against measured lysimeter (ET_{o-lys}) reference
 3 evapotranspiration using the FAO Penman-Monteith method (ET_{o-PM}) for 1997 and
 4 1998.

5

Figure 4. Average meteorological conditions during 10-day periods between May to September 2005 and 2006. a) mean air temperature; b) minimum relative humidity; c) wind speed at 2.0 m height; and d) estimated reference evapotranspiration.

Figure 5. Daily measured (ET_{c-lys}) versus estimated (ET_{c-est}) corn evapotranspiration using four approaches: 1) $ET_{c-est-lys}$, using equation for $K_{c_{mes}}$ (Figure 2); 2) $ET_{c-est-lyslc}$, using equation for $K_{c_{mes}}$ (Figure 2) and a calibration factor for the FAO Penman-Monteith method (Figure 3); 3) $ET_{c-est-PM}$, using equation for $K_{c_{est-PM}}$ (Figure 4); and 4) $ET_{c-est-FAO}$, using the FAO K_c curve ($K_{c_{est-FAO}}$).