

# A surface ocean trajectories visualization tool and its initial application to the Galician coast

Otero<sup>1\*</sup>, P., N. S. Banas<sup>2</sup> and M. Ruiz-Villarreal<sup>1</sup>.

<sup>1</sup>Instituto Español de Oceanografía, C. O. A Coruña. Paseo Marítimo Alcalde Francisco Vázquez, 10, 15001, A Coruña, Spain, Phone Number: 0034981205362, Fax Number: 0034981229077

<sup>2</sup>Joint Institute for the Study of the Atmosphere and Ocean, Box 355672, University of Washington, Seattle, WA 98195, USA

\*Corresponding author: otero\_pablo@hotmail.com

**Submitted to *Environmental Modelling & Software***

**July 2013**

**Revised**

**December 2013**

**Published**


**January 2015**

Please cite:

**Otero et al., 2015. A surface ocean trajectories visualization tool and its initial application to the Galician coast. *Environmental, Modelling and Software*, 66, pp. 12-16. doi:10.1016/j.envsoft.2014.12.006**

## Abstract

Alavai is a free tool built with Processing to visualize forecasts of drift trajectories in the sea in a very fast and user-friendly graphic environment. It plots trajectories of both passive particles trapped near the ocean surface and oil slicks. Input data come from an operational configuration of the ROMS ocean model focused on the Euroregion Galicia (Spain) – North Portugal, however, this tool is easily adaptable to run with outputs coming from any operational high resolution ocean model configuration. The Prestige accident is used here in hindcast mode to demonstrate the main characteristics of Alavai.



## Highlights

- A free, fast and friendly tool to visualize ocean surface drift is disseminated.
- It is easily adaptable to any operational high resolution ocean model configuration.
- 3-days forecast oil spill and passive surface particles trajectories are plotted.
- The usefulness is demonstrated in hindcast mode during the Prestige accident.

**Keywords:** ROMS, Alavai, ocean model, drift, oil spill, Iberia, Prestige

## **Software and/or data availability**

Name: Alavai

Code developers: Pablo Otero and Neil Banas.

Contact: Instituto Español de Oceanografía, C. O. A Coruña. Paseo Marítimo Alcalde Francisco Vázquez, 10, 15001, A Coruña, Spain, Phone Number: 0034981205362, Fax Number: 0034981229077, e-mail: otero\_pablo@hotmail.com

Year: 2013.

Hardware requirements: 512 Mb of RAM memory. Internet connection.

Software requirements: Java Runtime Environment (JRE) 1.7 or later.

Code availability: <https://github.com/PabloOtero/Alavai>

Web: (Spanish version) <http://centolo.co.ieo.es:8080/alavai/>

(English version) [http://centolo.co.ieo.es:8080/alavai\\_en/](http://centolo.co.ieo.es:8080/alavai_en/)

Costs: Free. Under Creative Commons Licence Attribution-NonCommercial-ShareAlike 3.0

Unported.

Program language: Built with Processing (<http://www.processing.org/>)

Program size: ~20Mb.

Data repository: <http://centolo.co.ieo.es:8080/thredds/catalog.html>

Size of archives: 4Mb

Access form: Distributed Oceanographic Data System (DODS) protocols for client/server data access.

## **1. Introduction**

The computation of drift trajectories in the sea is a hard although necessary task, useful to predict the evolution of oil spills, harmful algae blooms, or to help in fish egg and larvae dispersion studies, among others.

In particular with pollution contingencies, these computations must be done in predictive mode, taking advantage of the present state-of-the-art in operational oceanographic models, in order to prevent impacts on coastal areas or environmental sensitive areas (Hackett et al., 2006).

Similarly, forecasting drift trajectories helps Search and Rescue teams in case of maritime accidents. For all these cases, the base information comes from oceanographic and meteorological numerical predictions. Many operational services disseminate currents and wind speed predictions as vector maps through their websites. But it is of more interest to the public

and decision-makers to know how these physical variables affect the drifting of objects in the sea. In spite of that, trajectories summarizing the drift history are rarely shown in operational services. Most of the tools available through the web that do allow visualization of drift trajectories are Lagrangian models that need to be run offline by a technician (e.g., the General NOAA Operational Modeling Environment, GNOME; MEDSLINK-II, De Dominicis et al., 2013; ROFF, Carr et al., 2008), have a restricted list of users (e.g., POSEIDON-OSM System, implemented at the Hellenic Centre for Marine Research) or are run on demand (e.g., NCSU forecast system at <http://omgsrv1.meas.ncsu.edu:8080/ocean-circulation/trajectory.jsp>; MOTHY implemented by Météo-France; Daniel et al., 2005, 2004). In addition, some of these tools require very complex input information, or their outputs are simply hard to understand by a non-expert.

Previous difficulties are part of a wider problem, which is illustrated by the growing need within the management community and the general public for easy access to the wealth of environmental data coming from monitoring systems and models, and also for effectively communicating their intrinsic spatial and temporal variability. Consequently, many tools that use state-of-the-art data visualization and processing schemes to communicate complex environmental information are being adopted in a variety of fields (e.g., Demir et al., 2013; Gronewold et al., 2013; Kokkonen et al., 2010; Warne et al., 2013).

In this paper, we present a very fast, free and user-friendly visualization tool of predicted Lagrangian trajectories at the sea surface, named Alavai. This tool has been implemented off the Euro-region Galicia (Spain)-Northern Portugal Iberia as part of the Iberian Margin Ocean Observatory (RAIA, <http://www.marnaraia.org>). An automated system uploads ocean and meteorological forecasts to a public server. Alavai uses the latest available predictions to provide the end-user with a visualization of the predicted drift trajectory starting from any location in the Observatory area. Trajectories are usually displayed in less than one minute — depending on the internet connection —, facilitating a quick response strategy to combat hazardous substances at sea, or simply providing information to researchers about advection of biological material trapped near the surface. Moreover, the interface has been designed to be as intuitive as possible.

The application is based on a simple random walk model in the horizontal, with no vertical advection. In case of oil spill trajectories, a wind drag coefficient is also added to compute the

displacement. The Prestige accident in 2002 will be used here in hindcast mode to demonstrate Alavai capabilities.

Alavai is open source and has been built using the open source programming language Processing (<http://www.processing.org>), an extension of Java. Processing is a widely used environment for developing visually oriented applications with an emphasis on animation and providing users with instant feedback through interaction (Fry, 2007). Alavai has been coded to accept input from ROMS (Regional Ocean Modeling System: Haidvogel et al., 2000) which makes it easily adaptable to any region by any ocean modeling group with operational capability.

## **2. Material and Methods**

The Alavai plots forecast drift trajectories of both surface passive particles and oil spill, the last driven by the combined action of the sea surface dynamics and the direct dragging of the wind. Thus, the base data come from results of a hydrodynamic and a meteorological model executed in operational way.

### ***2.1 The operational configuration***

Forecast sea currents are obtained from an operational configuration of ROMS, a free surface primitive equation ocean model. The current configuration is maintained and executed by the Instituto Español de Oceanografía in the frame of the Iberian Margin Ocean Observatory. In this configuration, two grids are nested during online execution through the employ of the AGRIF version of the model (Penven et al., 2006). Data from the coarser (4 km horizontal resolution) 236 x 181 grid force the lateral open boundaries of the finer (~1.3 km) 312 x 167 embedded grid (right panel in Fig. 1). More details and some validations of this configuration can be found in Otero et al. (2013; 2009).

The ocean model is forced at the surface with meteorological data from a 12 km spatial and 1 h temporal resolution configuration of the Weather, Research and Forecasting (WRF) model, operationally run by the regional meteorological agency MeteoGalicia and disseminated through the RAI A Observatory.

Every day, and once the meteorological forecasts are available, these data are downloaded and interpolated to the grid of the ocean model configuration, which restarts from the output of the previous day. After the execution has finished, hourly ROMS standard outputs are interpolated from their original terrain following coordinates to levels of constant depth (to facilitate interpretation) and subsequently, converted to the NetCDF Climate and Forecast (CF) convention format (<http://cf-pcmdi.llnl.gov/>). Finally, these data are uploaded to a public data repository (THREDDS server, Thematic Realtime Environmental Distributed Data Services; <https://www.unidata.ucar.edu/software/thredds/current/tds/>) for dissemination purposes.

## 2.2 The Alavai Lagrangian model

Alavai displays drift pathways and simulates the motion of virtual particles by means of the advective transport induced by surface currents and wind, plus turbulent horizontal diffusion. The diffusivity velocity depends on the sea turbulent characteristics, simulated here as a Brownian motion of particles by means of a random walk process (e.g., Csanady, 1973; Visser, 1997). For an individual particle, the equation to solve is:

$$\frac{d\vec{x}_i}{dt} = \vec{u}_{sea}(x_i, t) + C_D \vec{u}_{wind}(x_i, t) + R(2K_h dt)^{1/2} \quad (1)$$

where  $x_i$  is the particle position,  $u_{sea}$  is the advective velocity of the sea surface at  $x_i$ ,  $u_{wind}$  is the 10 m height wind velocity at  $x_i$ ,  $C_D$  is the wind drag coefficient,  $K_h$  is the turbulent horizontal diffusion (set here to be  $1 \text{ m}^2 \text{ s}^{-1}$ ) and  $R$  is a random process; here, we use a standard normal distribution (e.g., North et al., 2006). The wind component is only used in the computation of oil spill trajectories, with  $C_D = 0.03$ . This value is dependent on wind speed and sea state (e.g., Abascal et al., 2009; Castanedo et al., 2006), although the value selected here has proven to produce good results during the simulation of the Prestige oil spill (e.g., Montero et al., 2003; Marta-Almeida et al., 2013). Sea surface currents are obtained from the child grid in the operational configuration of the ocean model (Fig. 1), whereas the wind speed fields are directly obtained from the outputs of the meteorological model.

Although the specific role of waves in the drift of oil slick becomes important in near-shore areas (e.g., Abascal et al., 2009), the present operational configuration does not compute waves, and therefore, the term has not been included in Equation 1. To not to lose focus on the scope of

this tool (to obtain a very fast and user-friendly visualization tool), effects of oil weathering have not been taken into account.

Much of the efficiency of Alavai comes from the fact that the two advective terms are computed in advance once the model forecasts are ready. The displacement by advection (in both horizontal directions) that a particle departing from the center of each grid cell will experience in one time step is stored as a series of NetCDF files. These intermediate files are uploaded on a public THREDDS server, from where any THREDDS-enabled data analysis and display tools like Alavai can efficiently access them. Each of these intermediate files is in effect a Lagrangian return map that gives the final position in the model grid of an array of particles released at the grid centers after a given time interval. Banas and Hickey (2005) used a static version of this method (and a single return map, not a time series) to produce an interactive visualization of the tidal circulation of Willapa Bay, Washington, USA

(<http://coast.ocean.washington.edu/willapa/tidemodel.html>; Banas et al. (2009) used a similar method to generate larval trajectories for a study of invasive green crab in Willapa Bay). Here, we have made a step forward and now we have a set of return maps, one for each time step (3600 s), but dynamically changing depending on the results of the operational model. In the present application, when the user locates the mouse cursor over the canvas, the software starts to plot the particle track from the correspondent pixel. The number of pixels in the canvas (in both directions) is higher than the number of model grid cells, which increases the resolution of the Lagrangian model. The software reads the displacement matrix in the intermediate file (lookup in the return-map) and interpolates data from the model grid to the current pixel. The software draws a first line linking the start and the end points (pixels in the canvas), after adding to the end location a random displacement (integration of the diffusive term in (1)). In iterative process, the software reads the displacement data for the next timesteps and repeats the steps, resulting in a complete drift track. One general advantage of the Lagrangian return map approach is that it separates the integration of three-dimensional model fields from the generation of the trajectory visualization. Thus, for other applications, one could choose a shorter advective timesteps or a more intensive integration scheme without any effect on the interactive tool.

The graphical interface of Alavai, adapted from the FlowWeaver project (<http://faculty.washington.edu/banasn/flowWeaver/>), is constructed not around a query-response model, but rather instantaneous feedback: trajectories are displayed starting from the current

mouse location and continuously regenerated as the user sweeps the mouse over the map. Exploration of the flow field is thus user-driven and extremely fast.

### 3. Results and discussion

Figure 1 shows the graphical interface of Alavai in combination with an illustration of the Prestige oil spill trajectory between the day of the sinking on November 19<sup>th</sup> 2002 and the day when part of the main spill stranded on the coast, on December 1<sup>st</sup>. The figure shows the location of the spill observed by planes and ships (obtained from Montero et al., 2003), and also the hindcast simulation of the trajectory once the spill entered through the left boundary of our domain included in Alavai. The main advantage of using this period in the testing is that there is information on the oil spill trajectories and on the oceanographic conditions during that period (e.g., Montero et al., 2003; Ruiz-Villarreal et al., 2006). Thus, we know that during the following days after the sinking the spill was moving in the same direction of the wind, speeding up or stopping as a function of the wind speed, and that during the night from November 30<sup>th</sup> to December 1<sup>st</sup>, the main spill arrived at the northern part of Cape Finisterre (43°N). Figure 1 shows the prediction of the trajectory starting on November 27<sup>th</sup> at midnight (the interface shows local time, +1 GMT), and the impact along the coast 97 h later, which matches the observed behavior of the main spill.

Observations at that time also showed that the spill dispersed during its route mainly due to the interaction with mesoscale activity (Ruiz-Villarreal et al., 2006) and rough sea state (Montero et al., 2003). Consequently, many patches were located off the Galician coast between 42.4°N and 43.0°N on November 30<sup>th</sup>. From December 1<sup>st</sup>, light northwesterly winds were dominant and some of those slicks impacted the National Park of the Atlantic Islands, located at the mouth of the *rías*, but no significant amount of oil entered in these estuaries. Northeasterly winds on December 3<sup>rd</sup> contributed to move the slicks further away from the coast. Figure 2 shows some snapshots of the Alavai tool during these dates, with drift trajectories starting at noon on December 1<sup>st</sup>. Three starting points are considered, with latitudinal separation of 0.15°. Alavai demonstrates that islands of the National Park were affected by the oil coming from northwesterly locations previous being taken away on December 3<sup>rd</sup>, a situation that would have not been reproduced by slicks departing from a southernmost location (right panel in Figure 2).



Previous figures show the ability of Alavai to visualize complex information in a very simple and intuitive way. However, the trust in the simulated drift trajectories mainly comes from the quality of the results provided by the model configuration and the value of  $C_D$ . The ability of the present model configuration to reproduce the high spatial and temporal coastal variability, that is under the strong influence of river plumes, has been also previously demonstrated with CTD, currentmeters and satellite-derived data (Otero et al., 2009, 2013) and also validated against a buoy track in April 2007 during a contingency exercise (Marta-Almeida et al., 2013). The selected  $C_D$  value has proven to properly simulate the drift of the Prestige oil spill, but this value should be tuned to simulate other substances (even other type of crude oil) or objects.

#### **4. Conclusions**

Alavai includes state-of-the art data visualization and processing schemes to visualize the ocean surface drift forecast in a free, fast and intuitive way. This allows a quick display of the estimated trajectories of surface-drifting substances in the coastal ocean to both planners and non-experts. Advances in computing power and software engineering are boosting the field of coastal ocean forecasting (e.g. Marta-Almeida et al. 2011, Penven et al. 2008) and nowadays state-of-the art numerical models with realistic forcing and detailed physics can be executed on a daily basis. Any validated ocean model configuration producing routine forecasts with high resolution CF-complaint outputs could be adapted to run inside Alavai with little effort. To help in this task, some preprocessing Matlab® scripts are also disseminated with the source code. The use of Java has facilitated the access to the NetCDF files in asynchronous way, in spite of using other language as Javascript will make this tool highly exportable. Future work should combine both requirements.

Examples in the Results section show hindcast simulations, in contrast with the nowcast and forecast mode that is made available through our web project. The end-user can play with a timeline that ranges from the previous day —to the current date— to the 3-day horizon prediction. Among the possible applications of Alavai in the Galician coast, where a strong shellfish culture activity takes place, we are currently using Alavai to display the possible drift of harmful algae populations to shellfish harvesting areas for early warning of harmful events.

## Acknowledgements

We acknowledge the predictive and modelling staff of the regional meteorological agency MeteoGalicia and the national agency AEMet who provided meteorological data. This visualization tool is part of the RAIA Coastal Observatory, financed by the Programa Operativo de Cooperación Transfronteriza España-Portugal POCTEP 2007-2013 (RAIA.co, 0520\_RAIA\_CO\_1\_E). Author contributions: P.O., N.B. and M.R.-V designed the research; P.O. and N.B. coded Alavai; P.O. and M.R.-V. performed the ROMS model configuration; P.O. analysed the data; M.R.-V. installed and configured the THREDDS server; all coauthors wrote and revised the paper. We are grateful to M. Cobas and L. García for their helpful comments about the model and the manuscript and G. González-Nuevo for his help with the THREDDS server. Many thanks to Dr. Rich Signell, Dr. Rob Hetland and two anonymous reviewers who contributed to improve the quality of the manuscript with their comments and suggestions. Part of the development was performed in relation to ASIMUTH project (EC FP7-SPACE-2010-1 grant agreement number 261860).

## References

- Abascal, A.J., Castanedo, S., Mendez, F.J., Medina, R., Losada, I.J., 2009. Calibration of a Lagrangian Transport Model Using Drifting Buoys Deployed during the Prestige Oil Spill. *J. Coast. Res.* 25(1) 80-90.
- Banas, N.S., Hickey, B.M., 2005. Mapping exchange and residence time in a model of Willapa Bay, Washington, a branching, macrotidal estuary. *J. Geophys. Res.-Oceans* 110(C11).
- Banas, N.S., McDonald, P.S., Armstrong, D.A., 2009. Green Crab Larval Retention in Willapa Bay, Washington: An Intensive Lagrangian Modeling Approach. *Estuaries and Coasts* 32(5) 893-905.
- Carr, S.D., Capet, X.J., McWilliams, J.C., Pennington, J.T., Chavez, F.P., 2008. The influence of diel vertical migration on zooplankton transport and recruitment in an upwelling region: estimates from a coupled behavioral-physical model. *Fish. Oceanogr.* 17(1) 1-15.
- Castanedo, S., Medina, R., Losada, I.J., Vidal, C., Mendez, F.J., Osorio, A., Juanes, J.A., Puente, A., 2006. The Prestige oil spill in Cantabria (Bay of Biscay). Part I: Operational forecasting system for quick response, risk assessment, and protection of natural resources. *J. Coast. Res.* 22(6) 1474-1489.

- Csanady, G.T., 1973. Turbulent diffusion in the environment. D. Reidel Publishing Company, Dordrecht.
- Daniel, P., Josse, P., Dandin, P., 2005. Further improvement of drift forecast at sea based on operational oceanography systems, Coastal Engineering VII: Modelling, Measurements, Engineering and Management of Seas and Coastal Regions, pp. 13-22.
- Daniel, P., Josse, P., Dandin, P., Lefevre, J.-M., Lery, G., Cabioch, F., Gouriou, V., 2004. Forecasting the Prestige Oil Spills, Proceedings of the Interspill 2004 conference: Trondheim, Norway.
- De Dominicis, M., Pinaridi, N., Zodiatis, G., 2013. MEDSLIK-II, a Lagrangian marine oil spill model for short-term forecasting - Part 1: Theory. Geoscientific Model Development Discussions 6 1949-1997.
- Demir, I., Krajewski, W.F., 2013. Towards an integrated Flood Information System: Centralized data access, analysis, and visualization. Environ. Model. Soft. 50, 77-84. doi: 10.1016/j.envsoft.2013.08.009
- Fry, B., 2007. Visualizing data. O'Reilly Media Inc., 384 pp.
- Gronewold, A.D., Clites, A.H., Smith, J.P., Hunter, T.S., 2013. A dynamic graphical interface for visualizing projected, measured, and reconstructed surface water elevations on the earth's largest lakes. Environ. Model. Soft. 49, 34-39. doi: 10.1016/j.envsoft.2013.07.003
- Hackett, B., Breivik, Ø., Wettre, C., 2006. Forecasting the drift of objects and substances in the ocean, In: Chassignet, E.P., Verron, J. (Eds.), Ocean weather forecasting, an integrated view of oceanography. Springer, pp. 507-524.
- Haidvogel, D.B., Arango, H.G., Hedstrom, K., Beckmann, A., Malanotte-Rizzoli, P., Shchepetkin, A.F., 2000. Model evaluation experiments in the North Atlantic Basin: simulations in nonlinear terrain-following coordinates. Dyn. Atmos. Ocean. 32(3-4) 239-281.
- Kokkonen, T., Ihaksi, T., Jolma, A., Kuikka, S., 2010. Dynamic mapping of nature values to support prioritization of coastal oil combating. Environ. Model. Soft. 25, 248-257. doi: 10.1016/j.envsoft.2009.07.017
- Marta-Almeida, M., Ruiz-Villarreal, M., Pereira, J., Otero, P., Cirano, M., Zhang, X., Hetland, R.D., 2013. Efficient tools for marine operational forecast and oil spill tracking. Mar. Pollut. Bull. 71(1-2), 139-151; <http://dx.doi.org/10.1016/j.marpolbul.2013.03.022>.
- Marta-Almeida, M., Ruiz-Villarreal, M., Otero, P., Cobas, M., Peliz, A., Nolasco, R., Cirano, M. and Pereira, J. 2011. OOF: a Python engine for automating regional and coastal ocean forecasts. Environ. Model. Soft. 26(5), 678-680.

- Montero, P., Blanco, J., Cabanas, J.M., Balseiro, C.F., Carracedo, P., Maneiro, J., Pazos, Y., Morono, A., Gómez, B., 2003. Oil spill monitoring and forecasting of the Prestige-Nassau accident, Arctic and marine oilspill program technical seminar. Environment Canada; 1999, pp. 1013-1030.
- North, E.W., Hood, R.R., Chao, S.Y., Sanford, L.P., 2006. Using a random displacement model to simulate turbulent particle motion in a baroclinic frontal zone: A new implementation scheme and model performance tests. *J. Mar. Sys.* 60(3-4) 365-380.
- Otero, P., Ruiz-Villarreal, M., Garcia-Garcia, L., Gonzalez-Nuevo, G., Manuel Cabanas, J., 2013. Coastal dynamics off Northwest Iberia during a stormy winter period. *Ocean Dyn.* 63(1) 115-129.
- Otero, P., Ruiz-Villarreal, M., Peliz, A., 2009. River plume fronts off NW Iberia from satellite observations and model data. *ICES J. Mar. Sci.* 66(9) 1853-1864.
- Penven, P., Debreu, L., Marchesiello, P., McWilliams, J.C., 2006. Evaluation and application of the ROMS 1-way embedding procedure to the central california upwelling system. *Ocean Model.* 12(1-2) 157-187.
- Penven, P., Marchesiello, P., Debreu, L., Lefvre, J., 2008. Software tools for pre- and post-processing of oceanic regional simulations. *Environ. Model. Soft.*, 23(5), 660-662.
- Ruiz-Villarreal, M., Gonzalez-Pola, C., Diaz del Rio, G., Lavin, A., Otero, P., Piedracoba, S., Cabanas, J.M., 2006. Oceanographic conditions in North and Northwest Iberia and their influence on the Prestige oil spill. *Mar. Pollut. Bull.* 53(5-7) 220-238.
- Visser, A.W., 1997. Using random walk models to simulate the vertical distribution of particles in a turbulent water column. *Mar. Ecol. Prog. Ser.* 158 275-281.
- Warne, D.J., Larsen, G., Young, J., Cox, M.E., 2013. Image-based flow visualisation (IBFV) to enhance interpretation of complex flow patterns within a shallow tidal barrier estuary. *Environ. Model. Soft.* 47, 64-73. doi: 10.1016/j.envsoft.2013.04.007




Figure 1. The predicted drift of the main spill after the Prestige accident is plotted starting from midnight on November 27<sup>th</sup> 2002. The impact on the coast is predicted 97 hours after the start of the drift. The random walk process is observed by plotting 10 different trajectories, all of them departing from the same location. To put this experiment into context, the location where the oil tanker collapsed and the overflight observations of the slicks are also shown. Gray arrows, which are not shown in the original display of the application, indicate points where rivers are imposed in the ocean model. On the right, the large grid of the ocean model configuration is shown together its embedded child grid. The user-friendly Alavai interface is formed by a time slider and only two buttons at the top right to select i) oil spill or ii) material trapped near the surface.




Figure 2. Snapshots of the Alavai tool for three virtual oil spills departing from different locations on December 1st 2002 at noon. Left and middle panels -in contrast with the right one- show the impact on the National Park of the Atlantic Islands for those particles departing from the northern-most virtual positions. This is one of the powerful characteristics of the tool: the user can visualize the drift starting at the same time from many locations just moving the mouse cursor over the screen.