

Antioxidant Molecular Brain Changes Parallel Adaptive Cardiovascular Response to Forced Running in Mice

Clara Bartra, Lars Andre Jager, Anna Alcarraz, Aline Meza-Ramos, Gemma Sangüesa, Rubén Corpas, Eduard Guasch, Montserrat Batlle and Coral Sanfeliu.

Table S1. List of TaqMan® Gene Expression Assays used for real time qPCR analysis

Gene Symbol	Assay ID	Gene Name
<i>Actb</i>	Mm02619580_g1	<i>Actin, beta</i>
<i>Bdnf</i>	Mm04230607_s1	<i>Brain Derived Neurotrophic Factor</i>
<i>Cat</i>	Mm00437992_m1	<i>Catalase</i>
<i>Creb1</i>	Mm00501607_m1	<i>CAMP Responsive Element Binding Protein 1</i>
<i>Gdnf</i>	Mm00599849_m1	<i>Glial Cell Derived Neurotrophic Factor</i>
<i>Nfe2l2</i>	Mm00477784_m1	<i>NFE2 Like BZIP Transcription Factor 2</i>
<i>Psmb5</i>	Mm07296970_g1	<i>Proteasome 20S Subunit Beta 5</i>
<i>Psmb6</i>	Mm01245590_g1	<i>Proteasome 20S Subunit Beta 6</i>
<i>Psmb7</i>	Mm01327044_m1	<i>Proteasome 20S Subunit Beta 7</i>
<i>Psmb8</i>	Mm01278980_g1	<i>Proteasome 20S Subunit Beta 8</i>
<i>PsmB9</i>	Mm00479004_m1	<i>Proteasome 20S Subunit Beta 9</i>
<i>Psmb10</i>	Mm00479052_g1	<i>Proteasome 20S Subunit Beta 10</i>
<i>Sirt1</i>	Mm00490758_m1	<i>Sirtuin 1</i>
<i>Sod2</i>	Mm01313000_m1	<i>Superoxide Dismutase 2</i>
<i>Ubc</i>	Mm01198158_m1	<i>Ubiquitin C</i>
<i>Vegfa</i>	Mm01281449_m1	<i>Vascular Endothelial Growth Factor</i>




Figure S1. Gene expression of neuroplasticity markers was not significantly increased by exercise training regimens. Genes for neurotrophic factors BDNF (a), GDNF (b) and VEGF (c); and for the signal transducers CREB1 (d) and SIRT1 (e). Experimental groups: SED (orange circles), sedentary;

MOD (green squares), moderate training; INT (blue triangles), high-intensity training. Values are mean \pm SEM ((a-c) SED $n = 11$, MOD $n = 11$, INT $n = 11$); (d) SED $n = 10$, MOD $n = 10$, INT $n = 10$; (e) SED $n = 10$, MOD $n = 11$, INT $n = 11$).




Figure S2. Gene expression of the catalytic 20S constitutive proteasome and 20S immunoproteasome was not significantly increased by exercise training regimens. Genes: *Psmb5* codifies for subunit $\beta 5$ with chymotrypsin-like activity (a); *Psmb6* codifies for subunit $\beta 1$ with caspase-like activity (b); *Psmb7* codifies for subunit $\beta 2$ with trypsin-like activity (c); *Psmb8* codifies for subunit $\beta 5i$ with chymotrypsin-like activity (d); *Psmb9* codifies for subunit $\beta 1i$ with caspase-like activity (e); *Psmb10* codifies for subunit $\beta 2i$ with trypsin-like activity (f); *Ubc* codifies for ubiquitin (g). Experimental groups: SED (orange circles), sedentary; MOD (green squares), moderate training; INT (blue triangles), high-intensity training. Values are mean \pm SEM ((a,b,d-f) SED $n = 10$, MOD $n = 11$, INT $n = 11$); (c) SED $n = 11$, MOD $n = 11$, INT $n = 10$; (g) SED $n = 11$, MOD $n = 11$, INT $n = 11$).