The historical origins of aridity and vegetation degradation in south-eastern Spain

José S. Carrión^{a *}, Santiago Fernández^a, Gonzalo Jiménez-Moreno^b, Séverine Fauquette^c, Graciela Gil-Romera^d, Penélope González-Sampériz^e, Clive Finlayson^f

^a *Departamento de Biología Vegetal, Facultad de Biología, Universidad de Murcia, 30100 Murcia, Spain*

^b *Departamento de Estratigrafía y Paleontología, Facultad de Ciencias, Universidad de Granada, Avenida Fuente Nueva S/N, 18002 Granada, Spain*

^c *Université de Montpellier 2, CNRS, Institut des Sciences de l'Évolution, CC 061, Université Montpellier II, Place Eugène Bataillon, 34095 Montpellier cedex 5, France*

^d *Institute of Geography and Earth Sciences, University of Wales, SY23 3DB Aberystwyth, United Kingdom*

^e *Instituto Pirenaico de Ecología-CSIC, Avenida Montañana 1005, 50059 Zaragoza, Spain*

^f *The Gibraltar Museum, 18-20 Bomb House Lane, P.O.Box 939, Gibraltar*

^{*}Corresponding author: J.S. Carrión: carrion@um.es, Phone and Fax: (34) 968 364995

Abstract

The complex relationships within modern landscapes cannot be understood without the benefit of retrospective studies. We review palaeoenvironmental data for southeastern Spain, a landscape vulnerable to desertification and with antiquity of human pressure on the landscape. A xerophytic component is discernible in the pollen diagrams of the southeastern peninsula ever since the Middle Miocene. During glacial stages of the Pleistocene, mountain grasslands and lowland steppes expanded, but tree vegetation, although episodically contracted, was ever present across the region, explaining part of the modern plant-species diversity. The magnitude of human impacts on vegetation during the Holocene has been highly variable, starting earlier (e.g. after c. 5000 cal yr BP) in low-elevation areas and river basins. Forest degradation of the mountains started rather late during the Argaric period, and reached its maximum during the Roman occupation. Over the last millennia, natural and/or human-set fires, combined with overgrazing, probably have pushed forests over a threshold leading to the spread of grassland,

1
2
3
4 thorny scrub, junipers, and nitrophilous communities. The high degree of xerophytization
5
6 observed today in southeastern Spain results from the long-term determinism of the mid-late
7
8 Holocene climate aridification, and the contingency of historical factors like fire events and
9
10 changes in prehistoric local economies involving resource exhaustion.
11
12
13
14

15 **Key words**

16
17 Palaeoecology, historical biogeography, Spain, Tertiary, Quaternary, Holocene
18
19
20
21

22 **Introduction**

23
24 This paper deals with the ecological history of a semi-desert region, the Murcian-Almerian
25
26 bioprovince, extending over 13,000 km² in the southeast of the Iberian Peninsula (Peinado et al.,
27
28 1992) (Fig.1). We review available evidence of past vegetation and climate-human-vegetation
29
30 interactions in the study area with the goal of establishing a temporal framework incorporating the
31
32 vegetation patterns and the natural processes observed today in an area that has witnessed one
33
34 of the highest rates of long-term land degradation within the European continent (Puigdefábregas
35
36 and Mendizábal, 1998). Specific issues to be addressed relate to (i) whether the arid belt with
37
38 xerothermic Mediterranean, and Saharo-Irano-Turanian elements in the Mediterranean coast of
39
40 southeastern Iberia is a recent feature or originates during or prior to the Quaternary, (ii) whether
41
42 the region served as glacial refugium for woody species during the Pleistocene, (iii) whether this
43
44 generally treeless region was forested any time during the Holocene, and, if so, which arboreal
45
46 species were involved, (iv) which factor(s) gave rise to the high degree of xerophytization
47
48 observed in the region today, and (v) whether the palaeo-record of biotic changes, fire
49
50 occurrence, and anthropogenic disturbance can shed light on the characteristics of modern
51
52 vegetation. This last issue includes the consideration of causal links with prehistoric patterns of
53
54 human settlement.
55
56
57
58
59
60
61
62
63
64
65

Present-day climate and vegetation

Although essentially Mediterranean, with summer drought and average annual temperatures of 16-17°C in the lowlands, the climate of the Murcian-Almerian bioprovince is characterised by greater aridity than other Mediterranean areas. Extreme summer temperatures are common, with daily maxima of up to 46 °C. The region is bounded by the Mediterranean Sea and surrounded by mountain ranges acting as a natural barrier to precipitation (IGME, 1999) (Fig. 1). Yearly rainfall ranges from 250 to 330 mm, though in mountainous areas it may exceed 400 mm and in some coastal areas it falls below 190 mm (Lázaro et al., 2001). Precipitation is irregular and often causes devastating floods of great erosive power, and evapotranspiration is greater than anywhere else in the Iberian Peninsula (Cantón et al., 2001).

Interestingly, while being one of the most arid regions on the European continent, it is among the richest floristically, perhaps due to the region's extensive bioclimatic gradients and geological diversity (Mota et al., 2004). More than half of the c. 7500 vascular plants found in Iberia (Devesa and Ortega, 2004) occur within the limited territory of this province, a majority of which are Mediterranean species (65%), including Iberian, Murcian-Almerian, Ibero-North African and Ibero-Irano-Turanian endemics (Mota et al., 2003). In the most arid environments, namely coastal mountains and littoral depressions, sclerophyllous brushwoods prevail, dominated by one or more of the following species: *Maytenus senegalensis* subsp. *europaea*, *Periploca angustifolia*, *Withania frutescens*, *Ziziphus lotus*, *Calicotome intermedia*, *Launaea arborescens*, and several species of *Genista*. Other abundant species include *Lycium intricatum*, *Rhamnus oleoides*, *Pistacia lentiscus*, *Olea europaea* var. *sylvestris*, *Stipa tenacissima*, *Ephedra fragilis*, as well as a diversity of Cistaceae, Lamiaceae, and Asteraceae species. Open forests with *Pinus halepensis* and *Quercus rotundifolia*, and *Quercus coccifera* scrub occur in the more continental depressions and less dry conditions of the mountainous hinterland. Saline depressions, as well as intermittent

watercourses, are colonised by *Artemisia*, Chenopodiaceae, *Nerium oleander*, *Tamarix*, *Limonium*, and Poaceae species.

Aridity was already present in the late Tertiary

Little is known about the Tertiary (65-1.8 Ma) evolution of Iberian floras. Available data suggest that aridity is a feature of the southeastern peninsula ever since the middle Miocene, about 16 million years ago. It is also possible that Mediterranean-type climates had appeared intermittently during the Miocene (23-5.3 Ma), as Tzedakis (2007) has recently suggested. Pollen spectra in records from Alborán A1 (36° 38' N 4° 13' E), Andalucía A1 (36° 35' N 2° 43' E) and Gor (37° 22' N 2° 59' W) (Fig. 1), include minor subdesertic elements since the Langhian (16-13.6 Ma) such as *Nitraria*, *Neurada*, *Prosopis*, *Lygeum*, *Ephedra*, Caesalpiniaceae, *Acacia* and *Calligonum*, while non-arboreal elements (Poaceae, Chenopodiaceae-Amaranthaceae, Plumbaginaceae, Caryophyllaceae) prevail in the pollen spectra (Jiménez-Moreno, 2005) (Fig. 2). These elements are but a part of a complex assemblage involving tropical and subtropical components (*Taxodium-Glyptostrobus*, *Bombax*, *Engelhardia*, Sapotaceae, *Myrica*, *Buxus bahamensis*, *Croton*, *Alcornea*, *Mussaenda*), mangrove species (*Avicennia*), and mesothermic (deciduous *Quercus*, *Carya*, and *Zelkova*), mid-altitude (*Cedrus*, *Tsuga*) and high-altitude (*Abies*, *Picea*) trees (Jiménez-Moreno and Suc, 2007). Physiographic heterogeneity must have been important at this time, as the rainfall requirements of these taxa vary greatly. Doubtless, the arid conditions were not widespread in the southeastern peninsula but there must have been sub-desertic areas because the *Neurada-Nitraria* assemblages are typical of the northern fringes of the Sahara today (Fauquette et al., 2006), and the Middle Miocene (c. 16-11 Ma) pollen sequences from the southeast contain a greater herbaceous component than northeastern records (Jiménez-Moreno and Suc, 2007).

The pollen sequence of the Andalucía G1 borehole (Fig.1) follows the erosional phase corresponding to the deep Mediterranean salinity crisis (Clauzon et al. 1990), and has been dated from c. 5.32 to 2.4 Ma (Suc et al. 1995). This sequence includes abundant herbs (Poaceae, Asteraceae, Chenopodiaceae-Amaranthaceae, *Artemisia*), deciduous *Quercus*, and subdesertic elements such as *Lygeum*, *Nitraria*, and *Calligonum*, indicating dry and hot conditions (Fauquette et al., 1998). These elements are, however, also present in the Tarragona E2 borehole, offshore northeastern Spain (Bessais and Cravatte, 1988) (Fig. 1), suggesting that an arid thermomediterranean belt with Saharo-Irano-Turanian elements continued to occur during the Pliocene along the Mediterranean coast of Iberia from Andalucía to southern Cataluña. Fauquette et al. (1999) quantified climatic conditions for this site through a pollen-derived method based on the mutual climatic range of plant taxa. Apart from the record of two major coolings that took place at 4.5 and 3.5 Ma (Suc and Zagwijn, 1983), annual temperatures remained equivalent to 6°C higher than today, and annual precipitation was eventually lower than today (if the Pliocene climate is calculated excluding deciduous *Quercus*, on the assumption that the species occupied mountainous areas adjacent to the study site (Fauquette et al., 1999)). The Pliocene at Andalucía G1 strongly contrasts with the site of Rio Maior in Portugal where Ericaceae dominates the non-arboreal pollen spectra and the subtropical component (*Cathaya*, *Engelhardia*, *Sequoia*, *Myrica*, *Taxodium*...) dominates the arboreal (Diniz, 1984). Using the former method of climatic estimation, their pollen data, and the plant functional types established by Prentice et al. (1992), Fauquette et al. (1999) reconstructed Pliocene biomes between 5.32 and 5 Ma for the West Mediterranean area. Hot desert, semi desert, xerophytic scrub, and warm grass/shrub biomes were reconstructed for southeastern Spain (Fig. 3).

Glacial refugium for tree taxa

Early to Mid-Pleistocene palaeobotanical records of the study region are lacking. If we take into account the biogeochemistry and ecomorphology of the large mammal assemblage, we can infer early Pleistocene landscapes characterised by the presence of shallow lakes with swampy marginal zones and extensive areas of savanna with tall grass and shrubs (Palmqvist et al., 2003). For the better documented Upper Pleistocene, the region can be broadly characterized by a diversity of landscapes including mountain grasslands and lowland steppes, *Pinus* woods at mid-altitudes, and the occurrence of a number of woody plants in the present-day meso and thermomediterranean belts along coastal shelves and intramontane valleys (Carrión et al., 2008). Glacial stages are characterized by the expansion of desert-like communities of Poaceae, *Artemisia*, Chenopodiaceae, Asteraceae, *Ephedra*, and Lamiaceae, among others (Finlayson and Carrión, 2007). However, mesothermophilous taxa occur along the coastal shelves such as in the Neanderthal sites of Cueva Pernerias (37° 31' N, 1° 25' W) (Carrión et al., 1995) and Sima de las Palomas (37° 47' N, 0° 53' W) (Carrión et al., 2003b) (Fig. 1). These sites have provided abundant pollen of *Pinus*, *Quercus* and Oleaceae, and continuous or frequent occurrence of broad-leaved trees (*Fraxinus*, *Alnus*, *Corylus*, *Juglans*, *Ulmus*, *Salix*) and thermophytes (*Myrtus*, *Erica arborea*, *Pistacia*, *Buxus*, *Periploca*, *Maytenus*, *Osyris*, *Withania*, *Lycium*, *Calicotome*, *Ephedra fragilis*). At San Rafael marshland there is continuous occurrence of evergreen and deciduous *Quercus* and *Olea* from c. 16000 to 10000 BP (Pantaleón-Cano et al., 2003). Anthracological studies on charcoal fragments, *Pinus* cone remains, and *Pinus* kernel shells in Nerja Cave, Málaga, document the occurrence of three species of *Pinus* between c. 24000 and 17500 yr BP: *Pinus pinea*, *Pinus nigra* and *Pinus halepensis* (Badal, 1998). Charcoal of *Pinus*, *Quercus*, *Juniperus*, *Cistus*, *Rhamnus-Phillyrea*, *Prunus*, and *Sorbus-Crataegus* have been found (Badal, 1998).

In general, the inland areas of south-eastern Spain fail to show continuous presence of mesothermophilous taxa during the last glacial stage. The pollen sequences from Beneito Cave

(38° 48' N, 0° 28' W) (Carrión and Munuera, 1997), Navarrés peatbog (39° 06' N, 0° 41' W) (Carrión and van Geel, 1999), and Villena peatbog (38°39' N, 0°52' W) (Yll et al., 2003) show episodic occurrences of *Quercus* and other woody taxa during OIS3, like Padul peatbog (37°N, 3° 40' W) (Pons and Reille, 1988) and Carihuela Cave (37° 26' N, 3° 25' W) (Carrión et al., 1999) during OIS4, OIS3 and OIS2. Lateglacial (after c. 15000 years BP) *Quercus* invasions (Pons and Reille, 1988; Fernández et al., 2007) suggest the proximity of refugia of evergreen and deciduous *Quercus*.

Refugia for woody taxa developed not only along the thermomediterranean belt in the coast, but also along intramontane valleys where sufficient precipitation may have provided enough moisture to support trees within a generally arid environment. Here, the lacustrine pollen site of Siles (Carrión, 2002) (38° 24' N, 2° 30' W), dating from upper pleniglacial times (c. 17030 yr BP), includes *Pinus pinaster*, deciduous *Quercus*, evergreen *Quercus*, Ericaceae, *Corylus*, *Betula*, and *Fraxinus*, in pollen percentages always above 2%, and also frequently with *Acer*, *Taxus*, *Arbutus*, *Buxus*, *Salix*, *Ulmus*, *Phillyrea*, *Pistacia*, and *Olea*.

Mid to late Holocene xerophytization

The Holocene is a time of great environmental variability in Mediterranean Spain. Several pollen sequences show an early-Holocene dominance by *Quercus* (Pons & Reille, 1988; González-Sampériz, 1998; Fernández et al., 2007), but in others *Pinus* continues to be dominant until the mid Holocene (Carrión & van Geel, 1999; Carrión et al. 2001b) or long thereafter (Stevenson, 2000; Franco et al., 2005). Overall, the period from c. 7500-5000 cal yr BP represents the mesophytic optimum, the xerophytic minimum, the period of lowest fire activity, and locally, a phase of relatively high lake levels (Carrión et al., 2001c, Carrión, 2002). During this period, the mountainous areas were forested with several species of *Pinus*, *Quercus faginea*, *Q. pyrenaica*, *Q. ilex*, *Q. suber*, broad-leaf trees (*Corylus*, *Alnus*, *Betula*, *Acer*, *Fraxinus*, *Ulmus*, *Juglans*,

1
2
3
4 *Arbutus*, *Salix*, *Castanea*) and other woody taxa (*Juniperus*, *Taxus*, *Olea*, *Phillyrea*, *Pistacia*,
5
6 *Buxus*, *Prunus*, *Ephedra fragilis*) (Carrión et al., 2007, 2008).
7

8
9 A relevant question is when did the arid southeast attain its present status? In addition,
10
11 what was the ultimate factor that provided the high degree of xerophytization observed today
12
13 (Calmel-Avila, 2000)? A comparative analysis of palaeobotanical data from the Murcian-Almerian
14
15 province and surroundings (Fig. 1) sheds light onto these questions. The pollen sequences of
16
17 Sierras de Gádor (2° 55'W, 36° 54'N, 1530 m a.s.l.) and Baza (37° 14'N, 2° 42'W, 1900 m a.s.l.)
18
19 are particularly useful in this respect.
20
21

22
23 Holocene ecological change in Sierra de Gádor has been constant, and at times, rapid
24
25 (Carrión et al., 2003a). After c. 3940 cal yr BP, frequent alternation of dominance of *Pinus* and
26
27 evergreen *Quercus* at the expense of deciduous *Quercus*, is observed. This change is preceded
28
29 by increases of microcharcoal particles at c. 4200 cal yr BP, suggesting enhanced fire (Fig. 4).
30
31 Fire events along the Gádor sequence are correlative of xerophyte pollen (*Lygeum*,
32
33 *Chenopodiaceae*, *Artemisia*, *Lamiaceae*, *Ephedra fragilis*), but the change towards sclerophyllous
34
35 vegetation is also contemporaneous with the Argaric settlement, which, according
36
37 palaeoanthracological analyses (Rodríguez-Ariza, 1992), would have intensified pastoralism,
38
39 forest destruction and matorralization. A *Poaceae* increase after c. 1760 cal yr BP is preceded by
40
41 rising percentages of *Sordariaceae*, *Polygonum aviculare*, *Riccia* and *Genisteae*, indicating heavy
42
43 grazing in the lake catchment (Carrión et al., 2003a). The first pollen records of *Vitis* are dated at
44
45 c. 1600 cal yr BP (Fig. 4). Pastoralism and agriculture are further supported by charcoal and seed
46
47 remains of *Olea*, *Vitis*, and heliophytic taxa in the excavation sites of the slopes (Rodríguez-Ariza,
48
49 1995; Buxó, 1997), and archaeological evidence suggests great demographic pressure during
50
51 Roman times (Cámlich and Martín, 1999). Sierra de Gádor today is a treeless mountain, but one
52
53 millennium ago, several taxa were found that are now extinct in the Sierra, such as *Taxus*
54
55 *baccata*, *Corylus avellana*, *Betula celtiberica*, *Alnus glutinosa*, and *Myrtus communis* (Carrión et
56
57
58
59
60
61
62
63
64
65

1
2
3
4 al., 2003a). Others, like *Arbutus unedo*, *Buxus sempervirens*, *B. balearica*, and *Acer granatense*,
5
6 have become extremely rare. In addition, historical records and toponimic information suggest
7
8 there was extensive brush and *Pinus* and *Quercus* forest cover in several mountain systems of
9
10 Almería only three centuries ago (García-Latorre and García-Latorre, 1996). These most recent
11
12 changes could have resulted from impact of lead extraction, beginning in 1822 (Grove and
13
14 Rackham, 2001). The current treeless condition of Sierra de Gádor is, therefore, a relatively
15
16 recent phenomenon. Under the long-term influence of increased dryness and burning, opening of
17
18 the landscape may have started patchily after c. 3940 cal yr BP, but the definitive loss of forest is
19
20 a human-induced case.
21
22

23
24 Like Gádor, the history of the vegetation in Sierra de Baza appears clearly influenced by
25
26 changes in local economy, with emphasis on the highly-populated Argaric period, in a context of
27
28 increased dryness over the last 5000 years (Carrión et al., 2007). The replacement of mesophytic
29
30 by more xeric Mediterranean vegetation around 3800 cal yr BP is preceded by greater fire activity
31
32 at c. 4100 cal yr BP. Grassland scrub and thorny matorral spread after c. 2560 cal yr BP (Fig. 4).
33
34 Sierra de Baza is today partially forested due to recent afforestations, but deforestation became
35
36 strong over the last two millennia, when anthropogenic disturbance (agriculture, mining,
37
38 pastoralism) reached its maximum.
39
40
41

42
43 Anthropogenic landscapes have their origins later in more continental territories, like the
44
45 Segura Mountains (Fig. 1). Yet, spatial complexity prevailed. *Plantago* and other indicators of
46
47 agriculture and arboriculture, and indicators of forest degradation, occur first in the pollen record
48
49 of the mesomediterranean Villaverde (c. 2200-1600 cal yr BP) (Fig. 4), then in Siles (c. 1400 cal
50
51 yr BP) and El Sabinar (c. 1353 cal yr BP), and finally in the high-elevation Cañada de la Cruz (c.
52
53 670 cal yr BP) (Carrión, 2002b) (Fig. 1). Archaeological data from the Neolithic to the Bronze Age
54
55 suggest that settlement was sparse in these mountains, and that agriculture was not important
56
57 before Roman times (Jordán, 1992). Documentary evidence suggests that only in the last
58
59
60
61
62
63
64
65

centuries has population growth and the improvement of agricultural technologies caused transformation of forests into croplands at low- and mid-altitudes, although many areas were left uncultivated or reverted to grazing until the present-day, and much of the forested territory was managed exclusively for timber (Sánchez-Gómez et al., 1995).

Important forest losses are detected earlier in thermomediterranean areas than in the mountains and most continental territories. The San Rafael marshland pollen record, in the littoral (36° 46' N, 2° 36' W), shows the development of shrublands (with expansion of communities of *Artemisia* and *Chenopodiaceae*) since the mid Holocene c. 5000 cal BP (Pantaleón-Cano et al., 2003) (Fig. 4). A similar sequence occurs at Roquetas (Pantaleón-Cano et al., 2003), Gata (Burjachs et al., 2000, Jalut et al., 2000), Antas (Pantaleón-Cano et al., 2003), and Caldereros (Fuentes et al., 2006) (Fig. 1). Geomorphological studies confirm a gradual erosive down cutting of many river catchments in the south-east from the Chalcolithic (Calmel-Avila, 2000). Anthracological studies also suggest that the anthropic impact in the thermomediterranean began earlier, since the Neolithic. This led to the progressive expansion of garrigues, depending on the locality, to the detriment of *Pinus* and arboreal *Quercus* (Badal et al., 1994, Rodríguez-Ariza, 2000). During and after the Chalcolithic, the well-developed riparian vegetation became progressively denuded in the lowland river valleys and their immediate surroundings (Rodríguez-Ariza, 1992).

Modern vegetation in southeastern Spain cannot be adequately understood in the absence of the palaeorecord. It is worth emphasizing that *Pinus* forests are natural in the region while evergreen *Quercus*-dominated formations are not systematically the pre-anthropogenic stage of vegetation (Carrión et al., 2004). Mixed conifer-*Quercus* forests including deciduous *Quercus* are a common feature in the middle Holocene (Fig. 4). Several species such as *Corylus avellana*, *Betula celtiberica*, *Acer granatense*, *Taxus baccata*, and *Arbutus unedo* are relics of the mid-Holocene mesic forests (Carrión et al., 2003a). The pollen sequences are also pertinent to

discern the ecological affinities of several species such as *Maytenus europaeus*, *Chamaerops humilis*, *Myrtus communis*, *Asparagus*, *Phillyrea* (cf. *P. media*), and *Buxus sempervirens/balearica*, which behaved as mesophytes by reaching a maximum during the mid Holocene (Carrión et al., 2003a). These taxa have declined since c. 4000 yr BP, and their variation does not correlate with that of the xerothermic scrub of *Ziziphus*, *Withania*, *Periploca*, and *Calicotome*, which have been phytosociologically related with (Peinado et al., 1992).

Conclusions

Aridity is a feature of southeastern Spain ever since the Middle Miocene, 16 million years ago, which is characterized by complex palynological assemblages including a subdesertic component with *Nitraria*, *Neurada*, *Prosopis*, *Lygeum*, *Ephedra*, *Caesalpinaceae*, *Acacia* and *Calligonum*. For the Pliocene (5.3-1.8 Ma), pollen records also reflect the occurrence of episodes with abundant herbs (Poaceae, Asteraceae, Chenopodiaceae-Amaranthaceae, *Artemisia*), and subdesert elements such as *Lygeum*, *Nitraria*, and *Calligonum* which indicate dry and hot conditions. During the Pleistocene (1.8-0.1 Ma), southeastern Iberia can be characterized by a diversity of landscapes including mountain grasslands and lowland steppes, *Pinus* woods in mid-altitudes and a number of woody plants in the present-day meso and thermomediterranean belts along coastal shelves and intramontane valleys. Forest cover was less extensive but ever present across the region during glacial stages, with the occurrence of *Pinus*, *Quercus*, *Juniperus*, *Abies*, *Corylus*, *Betula*, *Alnus*, *Ulmus*, *Salix*, *Castanea*, *Fraxinus*, *Juglans*, *Acer*, *Sorbus*, *Taxus*, *Olea*, *Pistacia*, *Phillyrea*, *Arbutus*, and *Myrtus*, among others.

Complicating the effects of climate on vegetation during the Holocene is the impact of humans, which has been highly variable over southeastern Spain, starting earlier in low-elevation areas and river basins. Forest degradation of the mountains started late during the Argaric period (4300-3600 cal yr BP), and reached its maximum during the Roman occupation, c. 2000 cal yr

BP. Over the last two millennia, natural and/or human-set fires, combined with overgrazing, would have pushed mountain forests over a threshold leading to the spread of grassland, thorny scrub, junipers, and nitrophilous communities. Specifically-orientated projects could be directed towards the investigation of whether ancient forests would be able to recover if anthropogenic pressures ceased, but the picture would remain speculative because demographic pressure is becoming greater over the last years. It can be concluded that the high degree of xerophytization observed today in southeastern Spain is not an unicausal process, but rather it is spatially heterogeneous, and resulting from a combination of anthropogenic disturbance, and resource exhaust under enhanced aridity.

Acknowledgements

This research has been funded by the projects CGL-2006-2956-BOS (Ministerio de Educación y Ciencia, Madrid) and Paleodiversitas Network (Fundación Séneca, Murcia). This paper is an ISEM contribution n°2008-***. This paper has benefited by changes suggested by R. Scott Anderson, Northern Arizona University, and an anonymous referee on an earlier draft.

References

- Badal, E., 1998. Interés económico del pino piñonero para los habitantes de la Cueva de Nerja. In: Sanchidrián, J.L., Simón, M.D. (Eds.), Las culturas del Pleistoceno Superior en Andalucía. Patronato de la Cueva de Nerja, Málaga, pp. 287-300.
- Badal, E., Bernabeu, J., Vernet, J.L., 1994. Vegetation changes and human action from the Neolithic to the Bronze Age (7000-4000 B.P.) in Alicante, Spain, based on charcoal analysis. *Vegetation History and Archaeobotany* 3, 155-166.

Bessais, E., Cravatte, J., 1988. Les écosystèmes végétaux pliocènes de Catalogne méridionale. Variations latitudinales dans le domaine nord-ouest méditerranéen. *Geobios* 21, 49-63.

Burjachs, F., Giralt, S., Roca J.R., Seret, G., Julià, R., 1997. Palinología holocénica y desertización en el Mediterráneo occidental. In: Ibáñez J.J., Valero, B.L., Machado, C. (Eds.), *El paisaje mediterráneo a través del espacio y del tiempo. Implicaciones en la desertificación*, Geoforma Editores, Logroño, pp. 379-394.

Burjachs, F., Febrero, A., Rodríguez-Ariza, M.O., Buxó, R., Araus, J.L., Julià, R. 2000. Holocene pollen sequences and carbon isotope discrimination of plant remains in Spain: evidence of a progressive increase in aridity. In: Balabanis, P., Peter, D., Ghazi, A., Tsogas, M. (Eds.), *Mediterranean desertification. Research Results and policy implications*. Directorate General Research, Brussels.

Buxó, R., 1997. *Arqueología de las plantas*. Crítica, Barcelona.

Calmel-Avila, M., 2000. Procesos hídricos holocenos en el Bajo Guadalentín (Murcia, SE España). *Cuaternario y Geomorfología* 14, 65-78.

Cámalich, M.D., Martín, D., 1999. El territorio almeriense desde los inicios de la producción hasta fines de la antigüedad. Un modelo: la Depresión de Vera y Cuenca del Río Almanzora. *Arqueología, Monografías*, Junta de Andalucía, Sevilla.

Cantón, Y., Domingo, F., Solé Benet, A., Puigdefábregas, J., 2001. Hydrological an erosion response of a badlands system in semiarid SE Spain. *Journal of Hydrology* 252, 65-84.

1
2
3
4
5
6 Carrión, J.S., 2002a. A taphonomic study of modern pollen assemblages from dung and surface
7
8 sediments in arid environments of Spain. *Review of Palaeobotany and Palynology* 120, 217-232.
9

10
11
12
13 Carrión, J.S., 2002b. Patterns and processes of Late Quaternary environmental change in a
14
15 montane region of southwestern Europe. *Quaternary Science Reviews* 21, 2047-2066.
16

17
18
19
20 Carrión, J.S., Andrade, A., Bennett, K.D., Navarro, C., Munuera, M., 2001b. Crossing forest
21
22 thresholds: inertia and collapse in a Holocene sequence from south-central Spain. *The Holocene*
23
24 11, 635-653.
25

26
27
28
29 Carrión, J.S., Dupré, M., Fumanal, M.P., Montes, R., 1995. A palaeoenvironmental study in semi-
30
31 arid southeastern Spain: the palynological and sedimentological sequence at Perneras Cave
32
33 (Lorca, Murcia). *Journal of Archaeological Science* 22, 355-367.
34
35

36
37
38 Carrión, J.S., Finlayson, C., Finlayson, G., Allué, E., López-Sáez, J.A., López-García, P.,
39
40 Fernández-Jiménez, S., Gil, G., Fuentes, N., González-Sampériz, P. 2008. A coastal reservoir of
41
42 biodiversity for Upper Pleistocene human populations. *Palaeoecological investigations in*
43
44 *Gorham's Cave (Gibraltar) in the context of the Iberian Peninsula. Quaternary Science Reviews*
45
46 (in press).
47
48
49

50
51
52 Carrión, J.S., Fuentes, N., González-Sampériz, P., Sánchez-Quirante, L., Finlayson, C.,
53
54 Fernández, S., Andrade, A. 2007. Holocene environmental change in a montane region of
55
56 southern Europe with a long history of human settlement. *Quaternary Science Reviews* 26, 1455-
57
58 1475.
59
60
61
62
63
64
65

1
2
3
4
5
6 Carrión, J.S., Munuera, M., 1997. Upper Pleistocene palaeoenvironmental change in eastern
7 Spain: new pollen analytical data from Cova Beneito (Alicante). *Palaeogeography,*
8
9 *Palaeoclimatology, Palaeoecology* 128, 287-299.
10
11
12
13
14

15 Carrión, J.S., Munuera, M., Dupré, M., Andrade, A., 2001c. Abrupt vegetation changes in the
16 Segura mountains of southern Spain throughout the Holocene. *Journal of Ecology* 89, 783-797.
17
18
19
20
21

22 Carrión, J.S., Munuera, M., Navarro, C., Burjachs, F., Dupré, M., Walker, M.J., 1999. The
23 palaeoecological potential of pollen records in caves: the case of Mediterranean Spain.
24
25 *Quaternary Science Reviews* 18, 1061-1073.
26
27
28
29
30

31 Carrión, J.S., Sánchez-Gómez, P., Mota, J.F., Yll, E.I., Chaín, C., 2003a. Fire and grazing are
32 contingent on the Holocene vegetation dynamics of Sierra de Gádor, southern Spain. *The*
33
34 *Holocene* 13, 839-849.
35
36
37
38
39

40 Carrión, J.S., van Geel, B., 1999. Fine-resolution Upper Weichselian and Holocene palynological
41 record from Navarrés (Valencia, Spain) and a discussion about factors of Mediterranean forest
42
43 succession. *Review of Palaeobotany and Palynology* 106, 209-236.
44
45
46
47
48

49 Carrión, J.S., Willis, K.J., Sánchez Gómez, P., 2004. Holocene forest history of the eastern
50 plateaux in the Segura Mountains (Murcia, southeastern Spain). *Review of Palaeobotany and*
51
52 *Palynology* 132, 219-236.
53
54
55
56
57
58
59
60
61
62
63
64
65

Carrión, J., Yll, E., Walker, M., Legaz, A., Chaíns, C., López, A., 2003b. Glacial refugia of temperate, Mediterranean and Ibero-North African flora in south-eastern Spain: new evidence from cave pollen at two Neandertal man sites. *Global Ecology and Biogeography* 12, 119-129.

Clauzon, G., Suc, J.P., Aguilar, J.P., Ambert, P., Capetta, H., Cravatte, J., Drivaliari, A., Domènech, R., Dubar, M., Leroy, S., Martinell, J., Michaux, J., Roiron, P., Rubino, J.L., Savoye, B., Vernet, J.L., 1990. Pliocene geodynamic and climatic evolutions in the French Mediterranean region. *Paleontology and Evolution, Special Memories* 2, 131-186.

Devesa, J.A., Ortega, A., 2004. *Especies vegetales protegidas en España: plantas vasculares. Naturaleza y Parques Nacionales, Serie Técnica*, Ministerio de Medio Ambiente, Madrid.

Fauquette, S., Quézel, P., Guiot, J., Suc, J.P., 1998. Signification bioclimatique de taxons-guides du Pliocène Méditerranéen. *Geobios* 31, 151-169.

Fauquette, S., Suc, J.P., Bertini, A., Popescu, S.M., Warny, S., Bachiri, N., Pérez Villa, M.J., Chikhi, H., Feddi, N., Subally, D., Clauzon, G., Ferrier, J., 2006. How much did climate force the Messinian salinity crisis ? Quantified climatic conditions from pollen records in the Mediterranean region. *Palaeogeography, Palaeoclimatology, Palaeoecology* 238, 281-301.

Fauquette, S., Suc, J.P., Guiot, J., Diniz, F., Feddi, N., Zheng, Z., Bessais, E., Drivaliari, A., 1999. Climate and biomes in the West Mediterranean area during the Pliocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 152, 15-36.

Fernández, S., Carrión, J.S., Fuentes, N., González-Sampériz, P., Gil, G., García-Martínez, M.S.,
Vega-Toscano, L.G., Riquelme, J.A., 2007. Palynology of Carihuela Cave, southern Spain:
completing the record. *Geobios* 40, 75-90.

Finlayson, C., Carrión, J. S., 2007. Rapid ecological turnover and its impact on Neanderthal and
other human populations. *Trends in Ecology and Evolution* 22, 213-222.

Franco, F., García-Antón, M., Maldonado, J., Morla, C., Sainz, H., 2005. Ancient pine forest on
inland dunes in the Spanish northern Meseta. *Quaternary Research* 63, 1-14.

Fuentes, N., García Martínez, M., González Sampériz, P., Fernández, S., Carrión, J.S., Ros, M.,
López Campuzano, M. & Medina, J., 2005. Degradación ecológica y cambio cultural durante los
últimos cuatro mil años en el sureste ibérico semiárido. *Anales de Biología* 27, 69-84.

García-Latorre, J., García-Latorre, J., 1996. Los bosques ignorados de Almería. Una
interpretación histórica y ecológica. In: Sánchez-Picón, A. (Ed.), *Historia y medio ambiente en el
territorio almeriense*, Almería: Servicio de Publicaciones, Universidad de Almería, pp. 99-126.

González-Sampériz, P., 1998. Estudio palinológico de la Cueva de En Pardo (Planes, Alicante).
Primeros resultados. *Cuaternario y Geomorfología* 12, 45-61.

Grove, A.T., Rackham, O., 2001. *The nature of Mediterranean Europe. An ecological history.*
Yale University Press, New Haven and London.

1
2
3
4 Harzhauser, M., Piller, W. 2007. Benchmark data of a changing sea. *Palaeogeography*,
5
6 *Palaeobiogeography and events in the Central Paratethys during the Miocene*. *Palaeogeography*,
7
8 *Palaeoclimatology, Palaeoecology* 253, 8-31.
9

10
11
12
13 IGME. 1999. Atlas del medio natural de la Región de Murcia. Consejería de Política Territorial y
14
15 Obras Públicas. Instituto Geológico y Minero, Madrid.
16
17

18
19
20 Jalut, G., Esteban, A., Bonnet, L., Gauquelin, T., Fontugne, M., 2000. Holocene climatic changes
21
22 in the western Mediterranean, from south-east France to south-east Spain. *Palaeogeography*,
23
24 *Palaeoclimatology, Palaeoecology* 160, 255-290.
25
26

27
28
29 Jiménez-Moreno, G. 2005. Utilización del análisis polínico para la reconstrucción de la
30
31 vegetación, clima y estimación de paleoaltitudes a lo largo del arco alpino europeo durante el
32
33 Mioceno (21-8 Ma). Thesis, University of Granada.
34
35

36
37
38 Jiménez-Moreno, G., Suc, J.-P., 2007. Middle Miocene latitudinal climatic gradient in western
39
40 Europe : evidence from pollen records. *Palaeogeography, Palaeoclimatology, Palaeoecology*
41
42 253: 224-241.
43
44

45
46
47 Jordán, J.F., 1992. Prospección arqueológica en la Comarca de Hellín-Tobarra. Metodología,
48
49 resultados y bibliografía. *Al-Basit* 31, 183-227.
50
51

52
53
54 Lázaro, R., Rodrigo, F.S., Gutiérrez, L., Domingo, F., Puigdefábregas, J., 2001. Análisis of a 30-
55
56 year rainfall record (1967-1997) in semi-arid Spain for implications on vegetation. *Journal of Arid*
57
58 *Environments* 48, 373-395.
59
60
61
62
63
64
65

1
2
3
4
5
6 Mota, J., Cabello, J., Cerrillo, M.I., Rodríguez-Tamayo, M.L., 2004. Los subdesiertos de Almería:
7 naturaleza de cine. Junta de Andalucía, Consejería de Medio Ambiente, Sevilla.
8
9

10
11
12
13 Mota, J., Cueto, M., Merlo, M.E., 2003. Flora amenazada de la provincia de Almería: una
14 perspectiva desde la biología de la conservación. Universidad de Almería, Secretariado de
15 Publicaciones, Almería.
16
17
18
19
20
21

22 Nogueras, P., Burjachs, F., Gallart, F., Puigdefàbregas, J., 2000. Recent gully erosion in the El
23 Cautivo badlands (Tabernas, SE Spain). Catena 40, 203-215.
24
25
26
27

28 Palmqvist, P., Gröcke, D.R., Arribas, A., Fariña, R.A., 2003. Paleoecological reconstruction of a
29 lower Pleistocene large mammal community using biogeochemical ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, Sr:Zn) and
30 ecomorphological approaches. Paleobiology 29, 205-229.
31
32
33
34
35
36
37

38 Pantaleón-Cano, J., Yll, E. I., Pérez-Obiol, R., Roure, J. M., 2003. Palynological evidence for
39 vegetational history in semi-arid areas of the western Mediterranean (Almería, Spain). The
40 Holocene 13, 109-119.
41
42
43
44
45
46

47 Peinado, M., Alcaraz, F., Martínez-Parras, J.M., 1992. Vegetation of southeastern Spain. J.
48 Cramer, Berlin.
49
50
51
52

53 Pons, A., Reille, M., 1988. The Holocene and Upper Pleistocene pollen record from Padul
54 (Granada, Spain): a new study. Palaeogeography, Palaeoclimatology, Palaeoecology 66, 243-
55 263.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 Prentice, I.C., Cramer, W., Harrison, S.P., Leemans, R., Monserud, R.A., Solomon, A.M., 1992. A
7
8 global biome model based on plant physiology and dominance, soil properties and climate.
9
10 Journal of Biogeography 19, 117-134.
11
12
13
14

15 Puigdefábregas, J., Mendizábal, T., 1998. Perspectives on desertification: western
16
17 Mediterranean. Journal of Arid Environments 39, 209-224.
18
19
20
21

22 Rivas-Martínez, S., Asensi, A., Díez-Garretas, B., Molero, J., 1997. Biogeographical synthesis of
23
24 Andalucía (southern Spain). Journal of Biogeography 24, 915-928.
25
26
27
28

29 Rodríguez-Ariza, M.O., 1992. Human plant relationships during the Copper and Bronze Ages in
30
31 the Baza and Guadix basins (Granada, Spain). Bulletin Société Botanique de la France 139, 451-
32
33 464.
34
35
36
37

38 Rodríguez-Ariza, M.O., 2000. El paisaje vegetal de la Depresión de Vera durante la Prehistoria
39
40 reciente. Una aproximación desde la antracología. Trabajos de Prehistoria 57, 145-156.
41
42
43
44
45

46 Sánchez-Gómez, P., Carrión, J.S., Jordán, J., Munuera, M., 1995. Aproximación a la historia
47
48 reciente de la flora y vegetación en las Sierras de Segura Orientales. Albasit 21, 87-111.
49
50
51
52
53

54 Stevenson, A.C., 2000. The Holocene forest history of the Montes Universales, Teruel, Spain.
55
56 The Holocene 10, 603-610.
57
58
59
60
61
62
63
64
65

Suc, J.P., Diniz, F., Leroy, S., Poumot, C., Bertini, A., Dupont, L., Clet, M., Bessais, E., Zheng, Z., Fauquette, S., Ferrier, J., 1995. Zanclean (Brunssumian) to early Piacenzian (early-middle Reuverian) climate from 4° to 54° north latitude (West Africa, West Europe and West Mediterranean areas). *Meded. Rijks Geol. Dienst* 52, 43-56.

Suc, J.P., Zagwijn, W.H., 1983. Plio-Pleistocene correlations between the northwestern Mediterranean region and northwestern Europe according to recent biostratigraphic and palaeoclimatic data. *Boreas* 12, 153-166.

Tzedakis, P.C., 2007. Seven ambiguities in the Mediterranean palaeoenvironmental narrative. *Quaternary Science Reviews* 26, 2042-2066.

Yll, E.I., Carrión, J.S., Pantaleón, J., Dupré, M., La Roca, N., Roure, J.M., Pérez Obiol, R., 2003. Palinología del Cuaternario reciente en la Laguna de Villena (Alicante). *Anales de Biología* 25, 65-72.

FIGURE LEGENDS

Figure 1. Palaeobotanical sites of south-eastern Spain discussed in this paper. Alborán A-1, Andalucía A-1, Andalucía G-1, Tarragona E2 and Gor (Jiménez-Moreno and Suc, 2007), Navarrés (Carrión and van Geel, 1999), Beneito (Carrión and Munuera, 1997), Villaverde (Carrión et al. 2001b), Siles (Carrión 2002a), Sabinar (Carrión et al., 2004), Cañada de la Cruz (Carrión et al., 2001a), Carihuela (Carrión et al., 1999, Fernández et al., 2007), Padul (Pons and Reille, 1988), Nerja (Badal, 1998), Gádor (Carrión et al., 2003a), Baza (Carrión et al., 2007), Las Palomas (Carrión et al., 2003b), Caldereros (Fuentes et al., 2005), Pernerias (Carrión et al., 1995),

Antas, San Rafael, Roquetas de Mar (Pantaleón-Cano et al., 2003), Gata (Jalut et al, 2000).

Asterisks refer to sites with evidences of tree pollen during glacial stages of the Pleistocene.

Figure 2. Synthetic pollen diagrams of the Middle Miocene sites Alborán-A1 and Andalucía-A1 in southeastern Spain. Subdesertic elements include *Nitraria*, *Calligonum*, *Lygeum*, *Neurada*, and *Prosopis*, among others. Mediterranean xerophytes include *Olea*, *Phillyrea*, *Ceratonia* and *Quercus ilex-coccifera*. Palaeogeography on palinspastic map of Harzhausser and Piller (2007). Redrawn from Jiménez-Moreno and Suc (2007).

Figure 3. Biomes of the Iberian Peninsula at 5.32-5 Ma using the typology established by Prentice et al. (1992), and climatic estimations. Redrawn from Fauquette et al. (1999).

Figure 4. Comparative chart of vegetational developments during the Holocene in four important pollen sequences from southeastern Spain, from the high-elevation Baza to the littoral San Rafael marshland. A climatically-induced mesophytic optimum is observed during the middle Holocene. Xerophytization starts from c. 5000-4500 cal yr BP, but it is modulated by changes in fire occurrence, and human activities. Deforested landscapes occur first in the thermomediterranean San Rafael.

Figure 1

[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)

Figure 3

[Click here to download high resolution image](#)

Figure 4
[Click here to download high resolution image](#)