

Vitamin B₁ in marine sediments: pore water concentration gradient drives benthic flux with potential biological implications

Danielle R. Monteverde^{1*}, Laura Gómez-Consarnau², Lynda Cutter², Lauren Chong¹, William Berelson¹ and Sergio A. Sañudo-Wilhelmy^{1,2}

¹ Department of Earth Sciences, University of Southern California, Los Angeles, CA, USA, ² Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA

OPEN ACCESS

Edited by:

Kurt O. Konhauser,
University of Alberta, Canada

Reviewed by:

Mustafa Yucel,
GEOMAR - Helmholtz Centre for
Ocean Research Kiel, Germany
Amy Michele Grunden,
North Carolina State University, USA
Ryan Paerl,
University of Copenhagen, Denmark

*Correspondence:

Danielle R. Monteverde,
Department of Earth Sciences,
Zumberge Hall of Science, University
of Southern California, 3651 Trousdale
Pkwy, Los Angeles, CA 90089-0740,
USA
dmonteve@usc.edu

Specialty section:

This article was submitted to
Microbiological Chemistry and
Geomicrobiology,
a section of the journal
Frontiers in Microbiology

Received: 02 February 2015

Paper pending published:
14 March 2015

Accepted: 22 April 2015

Published: 12 May 2015

Citation:

Monteverde DR, Gómez-Consarnau L, Cutter L, Chong L, Berelson W and Sañudo-Wilhelmy SA (2015) Vitamin B₁ in marine sediments: pore water concentration gradient drives benthic flux with potential biological implications. *Front. Microbiol.* 6:434. doi: 10.3389/fmicb.2015.00434

Vitamin B₁, or thiamin, can limit primary productivity in marine environments, however the major marine environmental sources of this essential coenzyme remain largely unknown. Vitamin B₁ can only be produced by organisms that possess its complete synthesis pathway, while other organisms meet their cellular B₁ quota by scavenging the coenzyme from exogenous sources. Due to high bacterial cell density and diversity, marine sediments could represent some of the highest concentrations of putative B₁ producers, yet these environments have received little attention as a possible source of B₁ to the overlying water column. Here we report the first dissolved pore water profiles of B₁ measured in cores collected in two consecutive years from Santa Monica Basin, CA. Vitamin B₁ concentrations were fairly consistent between the two years ranging from 30 pM up to 770 pM. A consistent maximum at ~5 cm sediment depth covaried with dissolved concentrations of iron. Pore water concentrations were higher than water column levels and represented some of the highest known environmental concentrations of B₁ measured to date, (over two times higher than maximum water column concentrations) suggesting increased rates of cellular production and release within the sediments. A one dimensional diffusion-transport model applied to the B₁ profile was used to estimate a diffusive benthic flux of ~0.7 nmol m⁻² d⁻¹. This is an estimated flux across the sediment-water interface in a deep sea basin; if similar magnitude B-vitamin fluxes occur in shallow coastal waters, benthic input could prove to be a significant B₁-source to the water column and may play an important role in supplying this organic growth factor to auxotrophic primary producers.

Keywords: vitamin B₁, thiamin, coenzyme, sediment, flux, auxotroph

Introduction

Vitamin B₁ (thiamin) is a soluble, biotically synthesized, heterocyclic sulfur, and nitrogen-containing catalyst required in trace amounts by all organisms (Jurgenson et al., 2009). It is primarily used as a coenzyme in forming and breaking C-C bonds and is required in central metabolic processes including the pentose-phosphate pathway and tricarboxylic acid cycle as well as in acetolactate synthase utilized in the synthesis of branched-chain amino acids

(Frank et al., 2007). This vitamin was originally identified as the molecule in rice husks which cures the human disease beriberi, caused by vitamin B₁ deficiency, a discovery which was recognized with the 1929 Nobel Prize in Physiology and Medicine (Eijkman, 1990). In the 1950s and 60s it was discovered that some species of marine phytoplankton are unable to synthesize B₁ *de novo* (B₁ auxotrophs) and instead must acquire the coenzyme from an exogenous source (Droop, 1957; Provasoli, 1958; Carlucci and Silbernagel, 1969). This included many of the major marine primary producers (Croft et al., 2006; Bertrand and Allen, 2012; Sañudo-Wilhelmy et al., 2014) as well as some ubiquitous marine bacteria (Giovannoni et al., 2005) and picoeukaryotic algae (Paerl et al., 2015). However, not every marine microbe (including both bacterioplankton and phytoplankton) requires an exogenous source of B₁; many microbes possess the full metabolic pathway needed to synthesize B₁ (Sañudo-Wilhelmy et al., 2014). Interestingly, B₁ auxotrophs do not appear to be related phylogenetically indicating that the loss of synthesis capability likely occurred multiple times (Helliwell et al., 2013). Additionally, it was discovered that B₁ synthesizers (or B₁ prototrophs) are able to self-regulate physiological concentrations within the cell via the use of a riboswitch (Croft et al., 2007). Recent work has revealed an additional layer of complexity regarding B₁ proto- and auxotroph dynamics, in that some species may only possess part of the synthesis pathway and can scavenge B₁ and/or its precursor moieties (4-amino-5-hydroxymethyl-2-methylpyrimidine or 4-methyl-5-β-hydroxyethylthiazole) in order to obtain the complete and active form of this vitamin (Jurgenson et al., 2009). Such organisms include climatologically relevant eukaryotic species such as *Emiliani huxleyi* (McRose et al., 2014) as well as environmentally abundant bacteria in the SAR11 clade (Carini et al., 2014). Field studies investigating dissolved B-vitamins in marine systems have shown that phytoplankton species succession and biomass production are influenced by the availability of vitamins B₁₂ (cobalamin) and B₁ (Sañudo-Wilhelmy et al., 2006; Panzeca et al., 2009; Koch et al., 2011; Bertrand et al., 2012). Additionally, it has been found that large regions of the ocean appear to be depleted in B₁ as well as other B-vitamins (Sañudo-Wilhelmy et al., 2012). Despite this, the sources of this organic growth factor have not been clearly identified and research in the area has mainly focused on B-vitamin production within the water column (e.g., Koch et al., 2012).

Marine sediments pose a potentially significant source for B₁ since sediments contain some of the highest cellular densities and diversity of any environment on Earth (where cellular abundance can reach as high as 10⁹ cells/cm³; Kallmeyer et al., 2012). Pioneering work in the 1950s and 1960s revealed that marine sediments may serve as a source of some B-vitamins including vitamin B₁ (Burkholder and Burkholder, 1958; Burkholder and Lewis, 1968). Based on our survey of required synthesis genes in whole genome sequenced sediment isolates (see Supplementary Material Table S1), marine sediments include many potential B₁ prototrophs. However, the majority of sediment microbes have remained uncultured (Eilers et al., 2000), and the dynamics of B₁ production and extracellular release remain largely unexplored. Thus, measuring dissolved B₁ in sediment pore waters is

essential to determine if the marine sediment community as a whole serves as a source of this critically required vitamin. In comparison to the many decades of study on trace metal and inorganic nutrient requirements (e.g., Fe and NO₃⁻), vitamins have received substantially less attention as a limiting agent to productivity. This is due in part to difficulties encountered measuring a labile molecule found in trace amounts (femto to pico molar concentrations) via the classic bioassay techniques or with the more recently developed liquid chromatography-mass spectrometry (LC/MS) techniques which can provide compound-specific information (Carlucci and Silbernagel, 1966; Okbamichael and Sañudo-Wilhelmy, 2005). As a result, the existing published environmental measurements of dissolved B₁ are almost entirely focused on the water column with little attention given to marine sediments and their pore waters. As such, we pose the following targeted research questions: (1) What are the sediment pore water B₁ concentrations? (2) Is there a flux of B₁ from sediments to the overlying water? (3) How relevant are those concentrations and fluxes to biological communities in the water column?

To address these questions, this study presents the first dissolved B₁ concentration profiles in marine pore waters, collected from the California Borderlands in Santa Monica Basin (SMB), CA from two sampling years (2011 and 2012). Pore water concentrations were compared to water column concentrations collected at the same station (Sañudo-Wilhelmy et al., 2012). Finally a simple diffusion-transport model was applied to the B₁ pore water concentrations in order to establish the first diffusive benthic flux estimates of B₁ from marine sediments.

Materials and Methods

Study Site

SMB lies ~10 miles offshore from Los Angeles within the California Continental Borderlands region. The basin is steep-walled with a flat-bottom covering an area of ~1800 km² with a basin floor at ~910 m and a sill at ~740 m isolating subsill waters from mixing with nearby basins; flushing events are estimated to occur every 1–8 years (Hammond et al., 1990; Berelson, 1991; Hickey, 1991; Berelson and Stott, 2003). Bottom waters and surface sediments are nearly but never completely anoxic (<10 μM oxygen) yet oxygen is undetectable within the first few millimeters of the sediment column (Shaw et al., 1990; Berelson et al., 2005). As a result of the low oxygen concentrations in bottom waters, the sediments are laminated with no evidence of infauna and minimal bioturbation indicating little to no advective mixing of pore waters (Jahnke, 1990; Christensen et al., 1994; Berelson et al., 2005; Tems et al., 2015). Previous studies indicate a ~5 cm-deep ferruginous/manganous zone defined by maximum concentrations of dissolved iron, and manganese (Jahnke, 1990; McManus et al., 1998; Prokopenko et al., 2011). Beneath this lies a zone with decreasing dissolved iron and manganese concentrations (Jahnke, 1990; McManus et al., 1998; Burdige and Komada, 2011). Flushing events, minor bioturbation, and changes in surface primary productivity may cause seasonal changes in shallow sediment (~0–5 cm) geochemical zonation by introducing increased

concentrations of oxygen, nitrate, and/or particulate organic carbon (Berelson, 1991). Multiple studies have investigated SMB sediment accumulation rates and found that roughly 9–11% of surface water primary productivity is exported to the basin floor resulting in consistent hemipelagic-sourced sediments accumulating at $\sim 16.0 \pm 3 \text{ mg cm}^{-2} \text{ y}^{-1}$ (Huh et al., 1990; Christensen et al., 1994; Berelson and Stott, 2003). In nearby San Pedro Basin particle flux was found to be seasonal and SMB likely experiences similar seasonality in sediment input (Collins et al., 2011). Of the organic carbon that reaches the sediment floor, $\sim 40\%$ is buried and preserved while the rest is remineralized and escapes to the water column (Jahnke, 1990). SMB's sediments are characterized as a silty-clay with $\sim 10\%$ calcium carbonate content and $\sim 4\text{--}6\%$ organic carbon (Craven and Jahnke, 1992; Gorsline, 1992). Sediments follow a typical porosity profile starting around a porosity of 0.98 which exponentially decreases with depth to values of ~ 0.85 at 8 cm depth (Berelson et al., 2005; Komada et al., 2013).

Core Collection

Sediment cores were collected from SMB ($33^\circ 48.76' \text{ N}$, $118^\circ 46.60' \text{ W}$; **Figure 1**) far enough away from basin walls and on a small regional high to avoid turbidite sampling. Cruises occurred in January 2011 and March 2012, just prior/during the expected maximum particle flux but before any spring flushing. Cores of 25–45 cm length were collected with an Ocean Instruments (MC 400) multicorer (Barnett et al., 1984) containing 9.5 cm diameter core liners. Upon retrieval, cores were inspected for a well-preserved sediment-water interface, minimal overlying water turbidity, and a lack of bubbles in the sediment in order to minimize collection artifacts. All cores were stored on board ship in an ice bath protected from light until transport to the laboratory cold room for sampling $\sim 9 \text{ h}$ after retrieval. Cores were sampled at depths of 1, 3, 5, 7, 9, 11, 15, 20, 25, and 35 cm in 2011 and 1.5, 3.5, 5.5, 7.5, 11.5,

15.5, 19.5, 25.5, 31.5, and 39.5 cm in 2012 using Rhizon soil samplers (Rhizosphere Research Products) fitted with $0.2 \mu\text{m}$ pore size filters. Rhizons were inserted into pre-drilled holes in the core liner and pore water was collected on cm-scale resolution using plastic syringes (Norm-ject) which had been acid-cleaned and methanol-rinsed. Sample volume ranged from 5 to 30 mL. Samples were then passed through an acid-washed $0.2 \mu\text{m}$ polypropylene capsule filter and stored frozen in acid-cleaned and methanol-rinsed high-density polyethylene (HDPE) amber bottles until analysis. Samples were protected from light as much as possible throughout sample processing. The 2012 dissolved iron samples were collected and filtered in the same way as the vitamin samples. Samples were stored in trace metal-cleaned HDPE bottles and acidified with Optima grade hydrochloric acid to a $\text{pH} < 2$ following standard trace clean techniques.

Analytical Methods

Vitamin B₁ was measured according to the technique described previously (Sañudo-Wilhelmy et al., 2012). The technique involves a solid-phase extraction onto a C₁₈ resin at $\text{pH} 6.5$ and 2.0 followed by elution with methanol, drying, and quantification using high-performance liquid chromatography/tandem triple quadrupole mass spectrometer (LC/MS) with an electrospray ionization interface. Reagent grade thiamin hydrochloride ($\geq 99\%$) was obtained from Sigma-Aldrich and used as an external standard. Samples were triple injected into the LC/MS to confirm instrument stability. Because sample volume is so low replicate sample splits were not performed. Method analytical blanks were measured with Milli-Q water subjected to the same preconcentration and quantification steps resulting in 0 pM in 2011 and 3.7 pM in 2012, which was subtracted from all of the sample measurements. The detection limits of the technique were 11 pM in 2011 and 0.53 pM in 2012 defined as three times the standard deviation of the procedural blank for 2012 and three times the standard deviation of the y-intercept of the calibration curve divided by the slope following the method outlined by Snyder et al. (2010) since the procedural blank was equal to zero in 2011. The improvement in detection limits between years resulted from an optimization of the method by increasing the sample injection volume from 50 to $100 \mu\text{L}$. Dissolved iron concentrations were quantified by ICP-MS using external calibration curves and an internal indium standard.

Benthic Flux Model

We applied a one-dimensional diffusion-transport model following Fick's First Law to the B₁ pore water concentration profile. The data (from 0 to 9 cm) was fit with a polynomial function:

$$C = C_0 + m_1x + m_2x^2$$

Where x is depth (cm), m_1 and m_2 are fitting parameters, and C_0 is the concentration at the sediment water interface (SWI). The diffusive flux of B₁ across the SWI was determined using Fick's First Law, applied to the derivative of the polynomial function fit evaluated at $x = 0$:

$$J = -\varphi^3 D_o \left(\frac{dc}{dx} \right)$$




FIGURE 1 | Santa Monica Basin station location ($33^\circ 48.76' \text{ N}$, $118^\circ 46.60' \text{ W}$). This figure was generated using Ocean Data View (Schlitzer, R. Ocean Data View, <http://odv.awi.de>, 2015).

Where ϕ represents sediment porosity at the SWI and D_0 is the molecular diffusion coefficient, and $\frac{dc}{dx}$ is the slope at the SWI. The D_0 of citrate ($3.22 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$) was used due to similarities in composition and molecular weight to B₁. The model ignores advection as is standard in similar sedimentation rate environments lacking bioturbation (e.g., Hammond et al., 1996).

Results

Pore Water Profiles

Vitamin B₁ concentrations in sediment pore water showed a consistent depth-profile shape in both sampling years (Figure 2). Concentrations were higher than water column values in most sampling depths of 2012 and all depths of 2011. Pore water

concentrations ranged from 330 to 770 pM in 2011 and 30–480 pM in 2012 as compared to water column concentrations of 30–280 pM previously reported by Sañudo-Wilhelmy et al. (2012) for the same station location (Supplementary Material Table S3). Additionally, the water column concentrations consistently increased with depth such that the deepest water column sample (890 m) had the highest B₁ concentration, ~280 pM. In both pore water profiles, vitamin B₁ exhibited consistent maximum concentrations at ~5 cm sediment depth and subsequently decreased with depth in both sampling years. The maximum concentrations of B₁ at ~5 cm sediment depth coincided with a maximum of dissolved iron (Figure 2; Supplementary Material Table S4).

Modeled Flux

The well-defined convex-upward B₁ profile in 2011 allowed a one dimensional diffusion-transport model, based on Fick's First Law, to be applied to the pore water concentrations (see Supplementary Figure S1). A simple quadratic curve fit evaluated the inflection point to be at ~9 cm. This model was then applied to the five pore water data points within the top 9 cm of sediment and the bottom water concentration was fixed at 280 pM based on the deepest water column value. The model produced a statistically significant model fit with a high chi-squared value ($\text{chisq} = 160$). The concentration gradient was evaluated at the SWI and a potential B₁ flux of $0.7 \text{ nmol m}^{-2} \text{ d}^{-1}$ was calculated out of the sediment. The less smooth shape of the 2012 profile (Figure 2) especially just below the sediment water interface, due to possible disturbance during transport and sampling or simply spatial variability, did not allow a good model fit for this sampling year. This is not uncommon for pore water sampling in deep marine sediments, for example of 8 total cores collected for DOC analysis in the same basin only 4 showed profiles consistent enough to allow a model fit (Komada et al., 2013).


Potential Algal Growth Yield

The calculated vitamin B₁ flux described above is for the exchange across the SWI at depths of 900 m in a sedimentary basin, yet this represents the first flux estimate for any marine setting. Assuming our calculated flux is representative of similar fluxes occurring in shallow water environments, and that B₁ degradation is minimal, a mass balance was applied to estimate the hypothetical growth response of such a sediment flux on a B₁-limited surface ocean phytoplankton community. Using the estimated sediment flux ($0.7 \text{ nmol m}^{-2} \text{ d}^{-1}$), a series of experimental phytoplankton cell growth yields ranging from 2.2×10^{-8} to $3.6 \times 10^{-8} \text{ pmol B}_1 \text{ cell}^{-1}$ (Paerl et al., 2015), and a photic zone of ~20 m (Small et al., 1989), we estimated that this magnitude sediment flux could support an algal growth yield ranging from 9.8×10^5 to $1.6 \times 10^6 \text{ cells L}^{-1} \text{ d}^{-1}$ (Supplementary Material Table S2).

Discussion

Vitamin B₁ Pore Water Profiles

The pore water depth profiles of B₁ from two sampling years showed high maximum concentrations compared to previous



field measurement as well as a consistent profile shape, especially considering the vitamin is present in such trace concentrations. This is in contrast to upper water column concentrations of B₁ which can vary widely on fairly short time scales (months to days; e.g., Gobler et al., 2007; Koch et al., 2012) and do not necessarily show consistent profile shape (Sañudo-Wilhelmy et al., 2012). This is likely due to the stratified nature of deep marine sediments which result in predictable geochemical zones that are not as susceptible to mixing or large diurnal shifts in bacterio-phytoplankton activity which likely affect vitamin production and uptake in the surface ocean (Sañudo-Wilhelmy et al., 2014). The consistent B₁ pore water concentration profile shape points to the existence of a stable mechanism for B₁ release to the dissolved phase. Furthermore, the maximum dissolved pore water concentrations for B₁ (770 pM) in 2011 were among the highest concentrations of any previously published values (see Table 1) almost 2 times higher than maximum water column concentrations. In fact the only other measurement that is within the same range was a single pore water value measured via HPLC (Okbamichael and Sañudo-Wilhelmy, 2005), supporting the hypothesis that sediments may represent universally elevated concentrations. Of the other water column measurements (Table 1), we note that some of the highest measurements are either found in shallow embayments likely affected by high benthic fluxes (Okbamichael and Sañudo-Wilhelmy, 2005) or anoxic marine basins such as SMB (Sañudo-Wilhelmy et al., 2012). These high concentrations suggest that vitamin production within the pore waters could be an important vitamin source for both the sediments and the water column. Additionally, of the 56 sediment bacteria and archaea surveyed in our genomic review, 74% of them were B₁ prototrophs with all of the genes necessary to synthesize the vitamin *de novo* (see Supplementary Material Table S1).

The shape of the B₁ pore water profile showed similarity to our dissolved iron profile as both showed peaks at ~5 cm. Manganese, which has been measured in the same basin in other studies, also shows a coincident peak at ~5 cm depth (Jahnke, 1990). This implies that the largest B₁ production was occurring within a geochemical zone of iron and manganese reduction as defined by the classic redox cascade of terminal electron

acceptors (Figure 2; Froelich et al., 1979). We are unaware of any biological mechanism linking B₁ to metal reduction, however many iron and manganese reducers are B₁ prototrophs (see Supplementary Material Table S1). Future culture experiments on sediment isolates from this sediment zone may help to explain why the elevated pore water concentrations occurred at this depth.

Previous measurements of dissolved organic carbon (DOC) in this same basin also showed elevated concentrations starting around ~5 cm (Komada et al., 2013). As an organic molecule, B₁ is part of the DOC pool, albeit a very small proportion (pM versus mM concentrations). Thus, processes driving changes in pore water DOC may also contribute to the profile shape of B₁, namely organic carbon remineralization. B₁ is a required cofactor for many important C-C breaking decarboxylase enzymes (Sañudo-Wilhelmy et al., 2014) which may serve as a possible mechanism linking the dissolved concentrations of DOC to B₁. Future environmental sampling and culture experiments targeted at carbon remineralization and B₁ production will be needed to confirm the validity of this proposed connection.

Vitamin B₁ Benthic Flux

The diffusive flux of 0.7 nmol m⁻² d⁻¹ out of the sediment represents the first such estimate ever made and therefore we lack a good comparison in order to judge the magnitude or significance of this flux. However, the algal growth yield calculation resulted in rates of cellular production (see Supplementary Material Table S2) some of which fall within the range for a phytoplankton bloom (Anderson et al., 2002). Certain caveats go along with these calculations, the most important being that we are explicitly not implying that the flux measured in SMB is reaching the surface waters. Instead, we assume that the calculated sediment flux may be representative of similar fluxes in more shallow environments as has been hypothesized in other studies (Okbamichael and Sañudo-Wilhelmy, 2005). Such a shallow water environment where a B₁ sediment flux would be particularly relevant would include shallow embayments, marshes, lagoons, or other environments that experience significant mixing and/or deep seasonal upwelling in order to allow transport of B₁-rich bottom water to surface

TABLE 1 | Environmental marine measurements of vitamin B₁.

Study area	Concentration range (pM)	References
SMB pore water	30–770	This study
Marine pore water from Flax Pond, NY	750	Okbamichael and Sañudo-Wilhelmy, 2005
Shallow Embayments in Peconic River and Stony Brook Harbor, NY	230–310	Okbamichael and Sañudo-Wilhelmy, 2005
California-Baja Pacific Margin	<0.81–314	Sañudo-Wilhelmy et al., 2012
Western Tropical North Atlantic, Amazon River Plume	<0.81–230	Barada et al., 2013
Long Island Sound, NY	<10–220	Vishniac and Riley, 1961
Peconic River, NY	12–190	Gobler et al., 2007
Quantuck Bay, NY	7–169	Koch et al., 2013
Old Fort Pond, NY	0.1–112	Gobler et al., 2007; Koch et al., 2012
Long Island Sound, NY	<0.10–99	Koch et al., 2012
Scripps Institute of Oceanography Pier	20–40	Carlucci and Silbernagel, 1966

waters. Furthermore, if the hypothesis that B₁ is linked to DOC proves correct, we would expect that shallow sediments, which are generally more organic rich and host higher bacterial abundances, likely produce significantly higher B₁ fluxes, and therefore the hypothetical growth yield can be considered a conservative estimate. Another assumption is that the surface algal community could be vitamin B₁-limited, which is possible given that 20% of genomic surveyed algae and 27% of cultured phytoplankton are B₁ auxotrophs (Croft et al., 2006; Sañudo-Wilhelmy et al., 2014), including many harmful algal bloom species (Tang et al., 2010). Of course additional variables would affect the initiation and response of the microbial community to the addition of B₁ including potential degradation prior to biologic uptake, competitive auxotrophic consumption of B₁, and additional release or transport during bloom die off (Sañudo-Wilhelmy et al., 2014). Despite these unknowns, our model suggests that benthic fluxes of B₁ occur at physiologically relevant rates and could impact surface primary production under a vitamin-limited regime, which appears to exist in many regions of today's oceans (Bertrand and Allen, 2012; Sañudo-Wilhelmy et al., 2012).

Summary and Future Directions

Here we presented the first deep-sea sediment pore water profiles of the universally required vitamin B₁. Our data showed a stable profile, hinting at a yet-to-be-determined link to a fundamental metabolic sediment cycle. Additionally we showed that sediments might serve as a source of B₁ to the water column. Future studies are needed in order to constrain spatial and temporal variability of sediment B₁ fluxes. While we cannot provide an unequivocal link between B₁ and microbial producers or

consumers, these pore water profiles serve as a starting point to formulate future hypotheses. Interesting avenues for future studies include whether B₁ can be limiting to sediment microbes in a similar way to their demonstrated limitation to surface water organisms despite the high pore water concentrations. Recent studies support the idea that B₁ has the potential to act as an ectocrine intermediate and perhaps a limiting nutrient based on a recent finding of auxotrophy in the widespread and highly abundant phylum chloroflexi (Rodionova et al., 2014), the members of which represent a significant abundance of bacteria in some shallow and deep sediments based on genomic studies (e.g., Blazejak and Schippers, 2010; Jorgensen et al., 2012). Such hypotheses, when coupled with *in situ* prokaryotic diversity techniques, and physiological studies using bacterial isolates from the different biogeochemical zones, will elucidate the role of vitamin B₁ on community function and composition both within the sediment and as a source to the water column.

Acknowledgments

Financial support for this research has been provided by the National Science Foundation (OCE 1435666; OCE 0934073; and ANT 1029878). The authors would like to acknowledge Doug Hammond, Nick Rollins, and Jotautas Baronas for their help in sample collection and analysis.

Supplementary Material

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2015.00434/abstract>

References

- Anderson, D. M., Glibert, P. M., and Burkholder, J. M. (2002). Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* 25, 704–726. doi: 10.1007/bf02804901
- Barada, L. P., Cutter, L., Montoya, J. P., Webb, E. A., Capone, D. G., and Sañudo-Wilhelmy, S. A. (2013). The distribution of thiamin and pyridoxine in the western tropical North Atlantic Amazon River plume. *Front. Microbiol.* 4:25. doi: 10.3389/fmicb.2013.00025
- Barnett, P. R. O., Watson, J., and Connelly, D. (1984). A multiple corer for taking virtually undisturbed samples from shelf, bathyl and abyssal sediments. *Oceanologica Acta* 7, 399–408.
- Berelson, W. M. (1991). The flushing of two deep-sea basins, Southern California Borderland. *Limnol. Oceanogr.* 36, 1150–1166.
- Berelson, W. M., Prokopenko, M., Sansone, F. J., Graham, A. W., McManus, J., and Bernhard, J. M. (2005). Anaerobic diagenesis of silica and carbon in continental margin sediments: discrete zones of TCO₂ production. *Geochim. Cosmochim. Acta* 69, 4611–4629. doi: 10.1016/j.gca.2005.05.011
- Berelson, W. M., and Stott, L. D. (2003). Productivity and organic carbon rain to the California margin seafloor: modern and paleoceanographic perspectives. *Paleoceanography* 18, 2.1–2.15. doi: 10.1029/2001pa000672
- Bertrand, E. M., and Allen, A. E. (2012). Influence of vitamin B Auxotrophy on nitrogen metabolism in eukaryotic phytoplankton. *Front. Microbiol.* 3:375. doi: 10.3389/fmicb.2012.00375
- Bertrand, E. M., Allen, A. E., Dupont, C. L., Norden-Krichmar, T. M., Bai, J., Valas, R. E., et al. (2012). Influence of cobalamin scarcity on diatom molecular physiology and identification of a cobalamin acquisition protein. *Proc. Natl. Acad. Sci. U.S.A.* 109, E1762–E1771. doi: 10.1073/pnas.1201731109
- Blazejak, A., and Schippers, A. (2010). High abundance of JS-1– and Chloroflexi–related Bacteria in deeply buried marine sediments revealed by quantitative, real-time PCR. *FEMS Microbiol. Ecol.* 72, 198–207. doi: 10.1111/j.1574-6941.2010.00838.x
- Burdige, D. J., and Komada, T. (2011). Anaerobic oxidation of methane and the stoichiometry of remineralization processes in continental margin sediments. *Limnol. Oceanogr.* 56, 1781–1796. doi: 10.4319/lo.2011.56.5.1781
- Burkholder, P. R., and Burkholder, L. M. (1958). Studies on B vitamins in relation to productivity of the Bahia Fosforescente. *P. R. Bull. Mar. Sci. Gulf Caribb.* 8, 201–223.
- Burkholder, P. R., and Lewis, S. (1968). Some patterns of B vitamin requirements among neritic marine bacteria. *Can. J. Microbiol.* 14, 537–543.
- Carini, P., Campbell, E. O., Morré, J., Sañudo-Wilhelmy, S. A., Thrash, J. C., Bennett, S. E., et al. (2014). Discovery of a SAR11 growth requirement for thiamin's pyrimidine precursor and its distribution in the Sargasso Sea. *ISME J.* 8, 1727–1738. doi: 10.1038/ismej.2014.61
- Carlucci, A. F., and Silbernagel, S. B. (1966). Bioassay of seawater. II. Methods for determination of concentrations of dissolved vitamin B₁ in seawater. *Can. J. Microbiol.* 12, 1079.
- Carlucci, A. F., and Silbernagel, S. B. (1969). Effect of vitamin concentrations on growth and development of vitamin-requiring algae. *J. Phycol.* 5, 64–67. doi: 10.1111/j.1529-8817.1969.tb02578.x

- Christensen, C. J., Gorsline, D. S., Hammond, D. E., and Lund, S. P. (1994). Nonannual lamination and expansion of anoxic basin-floor conditions in Santa-Monica Basin, California Borderland, over the past 4 centuries. *Mar. Geol.* 116, 399–418. doi: 10.1016/0025-3227(94)90054-x
- Collins, L. E., Berelson, W., Hammond, D. E., Knapp, A., Schwartz, R., and Capone, D. (2011). Particle fluxes in San Pedro Basin, California: a four-year record of sedimentation and physical forcing. *Deep Sea Res. Part I* 58, 898–914. doi: 10.1016/j.dsr.2011.06.008
- Craven, D. B., and Jahnke, R. A. (1992). Microbial utilization and turnover of organic-carbon in Santa-Monica Basin sediments. *Prog. Oceanogr.* 30, 313–333. doi: 10.1016/0079-6611(92)90017-t
- Croft, M. T., Moulin, M., Webb, M. E., and Smith, A. G. (2007). Thiamine biosynthesis in algae is regulated by riboswitches. *Proc. Natl. Acad. Sci. U.S.A.* 104, 20770–20775. doi: 10.1073/pnas.0705786105
- Croft, M. T., Warren, M. J., and Smith, A. G. (2006). Algae need their vitamins. *Eukaryotic Cell* 5, 1175–1183. doi: 10.1128/ec.00097-06
- Droop, M. R. (1957). Auxotrophy and organic compounds in the nutrition of marine phytoplankton. *J. Gen. Microbiol.* 16, 286–293.
- Eijkman, C. (1990). Anti-neuritis vitamin and beriberi. Nobel prize paper. 1929. *Ned. Tijdschr. Geneesk.* 134, 1654–1657.
- Eilers, H., Pernthaler, J., Glockner, F. O., and Amann, R. (2000). Culturability and *in situ* abundance of pelagic bacteria from the North Sea. *Appl. Environ. Microbiol.* 66, 3044–3051. doi: 10.1128/aem.66.7.3044-3051.2000
- Frank, R., Leeper, F., and Luisi, B. (2007). Structure, mechanism and catalytic duality of thiamine-dependent enzymes. *Cell. Mol. Life Sci.* 64, 892–905. doi: 10.1007/s00018-007-6423-5
- Froelich, P. N., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D., et al. (1979). Early oxidation of organic-matter in pelagic sediments of the eastern equatorial Atlantic—suboxic diagenesis. *Geochim. Cosmochim. Acta* 43, 1075–1090. doi: 10.1016/0016-7037(79)90095-4
- Giovannoni, S. J., Tripp, H. J., Givan, S., Podar, M., Vergin, K. L., Baptista, D., et al. (2005). Genome streamlining in a cosmopolitan oceanic bacterium. *Science* 309, 1242–1245. doi: 10.1126/science.1114057
- Gobler, C. J., Norman, C., Panzeca, C., Taylor, G. T., and Sañudo-Wilhelmy, S. A. (2007). Effect of B-vitamins (B-1, B-12) and inorganic nutrients on algal bloom dynamics in a coastal ecosystem. *Aquat. Microb. Ecol.* 49, 181–194. doi: 10.3354/ame01132
- Gorsline, D. S. (1992). The geologic setting of Santa-Monica and San-Pedro Basins, California Continental Borderland. *Prog. Oceanogr.* 30, 1–36. doi: 10.1016/0079-6611(92)90008-n
- Hammond, D. E., Marton, R. A., Berelson, W. M., and Ku, T. L. (1990). Ra-228 distribution and mixing in San-Nicolas and San-Pedro Basins, Southern California Borderland. *J. Geophys. Res. Oceans* 95, 3321–3335. doi: 10.1029/JC095iC03p03321
- Hammond, D. E., McManus, J., Berelson, W. M., Kilgore, T. E., and Pope, R. H. (1996). Early diagenesis of organic material in equatorial Pacific sediments: stoichiometry and kinetics. *Deep Sea Res. Part II* 43, 1365–1412. doi: 10.1016/0967-0645(96)00027-6
- Helliwell, K. E., Wheeler, G. L., and Smith, A. G. (2013). Widespread decay of vitamin-related pathways: coincidence or consequence? *Trends Genet.* 29, 469–478. doi: 10.1016/j.tig.2013.03.003
- Hickey, B. M. (1991). Variability in two deep coastal basins (Santa-Monica and San-Pedro) off Southern California. *J. Geophys. Res. Oceans* 96, 16689–16708. doi: 10.1029/91jc01375
- Huh, C. A., Small, L. F., Niemi, S., Finney, B. P., Hickey, B. M., Kachel, N. B., et al. (1990). Sedimentation dynamics in the Santa-Monica San-Pedro Basin off Los Angeles: radiochemical, sediment trap and transmissometer studies. *Cont. Shelf Res.* 10, 137–164. doi: 10.1016/0278-4343(90)90027-j
- Jahnke, R. A. (1990). Early diagenesis and recycling of biogenic debris at the sea-floor, Santa-Monica Basin, California. *J. Mar. Res.* 48, 413–436. doi: 10.1357/002224090784988773
- Jorgensen, S. L., Hannisdal, B., Lanzen, A., Baumberger, T., Flesland, K., Fonseca, R., et al. (2012). Correlating microbial community profiles with geochemical data in highly stratified sediments from the Arctic Mid-Ocean Ridge. *Proc. Natl. Acad. Sci. U.S.A.* 109, E2846–E2855. doi: 10.1073/pnas.1207574109
- Jurgenson, C. T., Begley, T. P., and Ealick, S. E. (2009). The structural and biochemical foundations of thiamin biosynthesis. *Annu. Rev. Biochem.* 78, 569–603. doi: 10.1146/annurev.biochem.78.072407.102340
- Kallmeyer, J., Pockalny, R., Adhikari, R. R., Smith, D. C., and D'hondt, S. (2012). Global distribution of microbial abundance and biomass in subseafloor sediment. *Proc. Natl. Acad. Sci. U.S.A.* 109, 16213–16216. doi: 10.1073/pnas.1203849109
- Koch, F., Hattenrath-Lehmann, T. K., Goleski, J. A., Sañudo-Wilhelmy, S. A., Fisher, N. S., and Gobler, C. J. (2012). Vitamin B₁ and B₁₂ uptake and cycling by plankton communities in coastal ecosystems. *Front. Microbiol.* 3:363. doi: 10.3389/fmicb.2012.00363
- Koch, F., Marcoval, M. A., Panzeca, C., Bruland, K. W., Sañudo-Wilhelmy, S. A., and Gobler, C. J. (2011). The effect of vitamin B(12) on phytoplankton growth and community structure in the Gulf of Alaska. *Limnol. Oceanogr.* 56, 1023–1034. doi: 10.4319/lo.2011.56.3.1023
- Koch, F., Sañudo-Wilhelmy, S. A., Fisher, N. S., and Gobler, C. J. (2013). Effect of vitamins B₁ and B₁₂ on bloom dynamics of the harmful brown tide alga, *Aureococcus anophagefferens* (Pelagophyceae). *Limnol. Oceanogr.* 58, 1761–1774. doi: 10.4319/lo.2013.58.5.1761
- Komada, T., Burdige, D. J., Crispo, S. M., Druffel, E. R. M., Griffin, S., Johnson, L., et al. (2013). Dissolved organic carbon dynamics in anaerobic sediments of the Santa Monica Basin. *Geochim. Cosmochim. Acta* 110, 253–273. doi: 10.1016/j.gca.2013.02.017
- McManus, J., Berelson, W. M., Klinkhammer, G. P., Johnson, K. S., Coale, K. H., Anderson, R. F., et al. (1998). Geochemistry of barium in marine sediments: implications for its use as a paleoproxy. *Geochim. Cosmochim. Acta* 62, 3453–3473. doi: 10.1016/s0016-7037(98)00248-8
- McRose, D., Guo, J., Monier, A., Sudek, S., Wilken, S., Yan, S., et al. (2014). Alternatives to vitamin B₁ uptake revealed with discovery of riboswitches in multiple marine eukaryotic lineages. *ISME J.* 8, 2517–2529. doi: 10.1038/ismej.2014.146
- Okbami, M., and Sañudo-Wilhelmy, S. A. (2005). Direct determination of vitamin B-1 in seawater by solid-phase extraction and high-performance liquid chromatography quantification. *Limnol. Oceanogr. Methods* 3, 241–246. doi: 10.4319/lom.2005.3.241
- Paerl, R. W., Bertrand, E. M., Allen, A. E., Palenik, B., and Azam, F. (2015). Vitamin B₁ ecophysiology of marine picoeukaryotic algae: strain-specific differences and a new role for bacteria in vitamin cycling. *Limnol. Oceanogr.* 60, 215–228. doi: 10.1002/lno.10009
- Panzeca, C., Beck, A. J., Tovar-Sanchez, A., Segovia-Zavala, J., Taylor, G. T., Gobler, C. J., et al. (2009). Distributions of dissolved vitamin B(12) and Co in coastal and open-ocean environments. *Estuar. Coast. Shelf Sci.* 85, 223–230. doi: 10.1016/j.ecss.2009.08.016
- Prokopenko, M. G., Sigman, D. M., Berelson, W. M., Hammond, D. E., Barnett, B., Chong, L., et al. (2011). Denitrification in anoxic sediments supported by biological nitrate transport. *Geochim. Cosmochim. Acta* 75, 7180–7199. doi: 10.1016/j.gca.2011.09.023
- Provasoli, L. (1958). Nutrition and ecology of protozoa and algae. *Annu. Rev. Microbiol.* 12, 279–308. doi: 10.1146/annurev.mi.12.100158.001431
- Rodionova, I. A., Li, X., Plymale, A. E., Motamedchaboki, K., Konopka, A. E., Romine, M. F., et al. (2014). Genomic distribution of B-vitamin auxotrophy and uptake transporters in environmental bacteria from the Chloroflexi phylum. *Environ. Microbiol. Rep.* 7, 204–210. doi: 10.1111/1758-2229.12227
- Sañudo-Wilhelmy, S. A., Cutter, L. S., Durazo, R., Smail, E. A., Gómez-Consarnau, L., Webb, E. A., et al. (2012). Multiple B-vitamin depletion in large areas of the coastal ocean. *Proc. Natl. Acad. Sci. U.S.A.* 109, 14041–14045. doi: 10.1073/pnas.1208755109
- Sañudo-Wilhelmy, S. A., Gobler, C. J., Okbami, M., and Taylor, G. T. (2006). Regulation of phytoplankton dynamics by vitamin B-12. *Geophys. Res. Lett.* 33, 1–4. doi: 10.1029/2005gl025046
- Sañudo-Wilhelmy, S. A., Gómez-Consarnau, L., Suffridge, C., and Webb, E. A. (2014). The role of B Vitamins in marine biogeochemistry. *Ann. Rev. Mar. Sci.* 6, 14.11–14.29. doi: 10.1146/annurev-marine-120710-100912
- Shaw, T. J., Gieskes, J. M., and Jahnke, R. A. (1990). Early diagenesis in differing depositional environments—The response of transition-metals in pore water. *Geochim. Cosmochim. Acta* 54, 1233–1246. doi: 10.1016/0016-7037(90)90149-f
- Small, L. F., Landry, M. R., Eppley, R. W., Azam, F., and Carlucci, A. F. (1989). Role of plankton in the carbon and nitrogen budgets of Santa-Monica-Basin, California. *Mar. Ecol. Prog. Ser.* 56, 57–74. doi: 10.3354/meps056057
- Snyder, L. R., Kirkland, J. J., and Dolan, J. W. (2010). “Qualitative and quantitative analysis,” in *Introduction to Modern Liquid Chromatography* (John Wiley & Sons, Inc.), 499–530.

- Tang, Y. Z., Koch, F., and Gobler, C. J. (2010). Most harmful algal bloom species are vitamin B-1 and B-12 auxotrophs. *Proc. Natl. Acad. Sci. U.S.A.* 107, 20756–20761. doi: 10.1073/pnas.1009566107
- Tems, C. E., Berelson, W. M., and Prokopenko, M. G. (2015). Particulate $\delta^{15}\text{N}$ in laminated marine sediments as a proxy for mixing between the California Undercurrent and the California Current: a proof of concept. *Geophys. Res. Lett.* 42, 419–427. doi: 10.1002/2014GL061993
- Vishniac, H. S., and Riley, G. A. (1961). Cobalamin and thiamine in Long Island Sound: patterns of distribution and ecological significance. *Limnol. Oceanogr.* 6, 36–41.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Monteverde, Gómez-Consarnau, Cutter, Chong, Berelson and Sañudo-Wilhelmy. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.