1 Pyrethroid use-malaria control and individual applications by households for other
2 pests and home garden use.

3
4 Maria Luisa Feo^a, Ethel Eljarrat^{a*}, Maria N. Manaca^{a,b,c}, Carlota Dobaño^{b,c,d}, Damia Barcelo^a, Jordi
5 Sunyer^{d,e}, Pedro L. Alonso^{b,c,d}, Clara Menendez^{b,c,d} and Joan O. Grimalt^a

6
7 ^aInstitute of Environmental Assessment and Water Research (IDÆA-CSIC). Jordi Girona, 18.
8 08034-Barcelona. Catalonia. Spain.

9 ^bCentro de Investigação em Saúde da Manhiça (CISM). Maputo. Mozambique

10 ^cBarcelona Centre for International Health Research (CRESIB). Hospital Clínic. Universitat de
11 Barcelona. Rosselló 132, 4a. 08036-Barcelona. Catalonia. Spain.

12 ^dCiber Epidemiología y Salud Pública, Spain

13 ^eCentre for Research in Environmental Epidemiology (CREAL). Doctor Aiguader, 88. 08003-
14 Barcelona. Catalonia. Spain

15
16
17 **Abstract**

18 Presence of pyrethroid insecticides in human breast milk and in thatch wall material of dwellings
19 from Southern Africa subtropical area (Manhiça, Mozambique) was investigated to assess potential
20 pyrethroid route of human exposure. Human breast milk samples were collected during 2002 when
21 pyrethroids were widely used as insecticides for mosquito bed nets in Mozambique for malaria
22 control. The median concentration value of total pyrethroids ranged between 87 and 1200 ng/g lw,
23 with λ -cyhalothrin being the most predominant pyrethroid in human breast milk contributing for
24 35% of the total amount. Moreover, and for the first time, an isomer-specific enrichment was found
25 in human breast milk, showing a selective isomeric accumulation or metabolism in the human body.

26 Based on the calculated pyrethroid concentrations in human breast milk, the daily ingestion rate of
27 pyrethroid was estimated. The nursing infant dietary intake ranged from 0.67 to 9.0 μg (kg of body
28 weight)⁻¹ day⁻¹. In addition, thatch materials collected after the reintegration of dichlorodiphenyl-
29 trichloroethene (DDT) as insecticide residual spraying (IRS) in Mozambique, showed the presence
30 of pyrethroids with concentration values ranging between 6.9 and 700 ng/g dw. In thatch material as
31 well as in human breast milk, pyrethroid contamination was mainly attributed to the agriculture
32 usage of this insecticide knowing that agriculture represent the 80% of the economy in
33 Mozambique. However, a possible usage of this insecticide as IRS in Mozambique cannot be
34 excluded despite their low efficiency for malaria control. The continued use of these compounds
35 (both for agricultural and malaria prevention) and the ingestion rates calculated from the breast milk
36 concentrations indicate that these insecticides cannot be overlooked for the assessment of the
37 lactation risks of breastfeeding infants from the Manhiça region.

38

39

40 **Introduction**

41 In 2004, an estimated 350-500 million people contracted malaria and 0.85 million died (91% in
42 Africa, 85% of them children under 5 years) ([WHO 2007a](#), [WHO and UNICEF 2005](#)). The World
43 Health Organization Pesticide Evaluation Scheme (WHOPES) ([WHO 2008](#)) supports the use of
44 recommended insecticides for malaria control based on the evaluation of human and environmental
45 safety conditions ([WHO 2006](#)). In tropical Africa, these insecticides were used for treatment of
46 mosquito nets (ITNs) ([Kapp, 2004](#)) and indoor residual spraying (IRS) on walls and roofs to kill the
47 mosquitoes that land and rest there ([Montgomery et al., 2010](#)). In the last years increases of
48 international funding for malaria control allowed to protecting larger numbers of people in sub-
49 Saharan Africa, from 13 million in 2005 to 75 million in 2009 by IRS as well as 66% of the 765
50 million at risk by use of 254 million ITNs between 2008 and 2010 (World Malaria Report, WHO
51 2010). The implementation of dichlorodiphenyltrichloroethene (DDT) and pyrethroids for IRS

constitute one of the major interventions for reduction and interruption of malaria transmission by vector control in all epidemiological settings (World Malaria Report, WHO 2010).

In Mozambique DDT was introduced in 1946 for agriculture and health programs. The IRS program with DDT broke down in the late 1970s due to the civil war. After this event (1993), the National Malaria Control Program (MNMCP) decided to restart IRS with pyrethroids in suburban areas of most provincial capitals. However, *Anopheles funestus*, one of the main mosquito vectors became resistant to this group of insecticides (Hargreaves et al., 2000; Sereda and Meinhardt, 2005). Thus, in 2000 carbamates (bendiocarb) were used in the rural areas of Maputo province within a coordinated effort for protection of the population of the Lubombo region (Mozambique, Swaziland and South Africa; Mabaso et al., 2004) while the use of pyrethroids was continued for mosquito nets. By end 2005, DDT was reintroduced for IRS following the WHO recommendations for areas of potential human life loss as consequence of unstable malaria transmission and epidemics (WHO, 2006),

Pyrethroids are synthesized derivatives of pyrethrins, which are natural insecticides produced by certain species of chrysanthemum (*Chrysanthemum cinerariaefolium*). Even though effects to humans are still unclear, the US Environmental Protection Agency (EPA) has classified some of them (cypermethrin, permethrin and bifenthrin) as possible human carcinogens (Cox, 1996). Pyrethroids are persistent compounds with high hydrophobicity ($\log K_{ow} = 5.7-7.6$) and low water solubility (a few $\mu\text{g L}^{-1}$) (Laskowski, DA., 2002). Despite these properties there is evidence of human pyrethroid metabolism and urine excretion of these compounds (ATSDR, 2003).

The accumulation of some pyrethroids in human milk has been considered in a limited number of studies (Bouwman et al., 2006; Sereda et al., 2009; Zehringer and Herrmann, 2001) showing appreciable pyrethroid levels in breast milk together with DDT. In some individuals, pyrethroid levels were higher than DDT levels suggesting domestic and home garden use of the former, while the presence of DDT was attributed to activities for control of malaria vectors. Except for DDT, safety of insecticide residues in breast milk has not been considered during the WHOPES

78 evaluation and very little is known on the effect of these chemicals to infants. This issue is
79 important because milk is the best sole nutrient source for infants, particularly in Africa.

80 In the present study assessment of pyrethroid exposure in a rural area located in the south of
81 Mozambique (Manhiça district) is undertaken. The study encompasses a comprehensive
82 examination of the compounds belonging to the pyrethroid group, e.g. bifenthrin, λ -cyhalothrin,
83 permethrin, cyfluthrin, cypermethrin, esfenvalerate, fenvalerate, fenpropathrin, deltamethrin,
84 tetramethrin, phenothrin and resmethrin, including the isomeric composition of some of these
85 compounds. Human milk was analysed as body burden estimate. Moreover, pyrethroid content in
86 walls (thatch material) of dwellings was also determined for assessment of potential human
87 exposure. To the best of our knowledge this is the first time in which this combined human-
88 environmental approach is addressed.

89

90

91 **2. Material and Method**

92 *2.1. Study area*

93 Manhiça district is a rural area located in the Northern of Maputo province in Mozambique. The
94 climate is subtropical characterized by a warm and rainy season between November-April and a dry
95 and cold season during the rest of the year.

96

97 *2.2. Samples*

98 Mature breast milk samples were collected in 2002 (n = 22) in the context of studies conducted at
99 the Centro de Investigação em Saúde da Manhiça (CISM). The research protocol was approved by
100 the ethic committees of Mozambique and Hospital Clinic in Barcelona. All women signed an
101 informed consent before they were enrolled in the study. Samples were stored in sterile polyester
102 containers at -80°C at CISM and at -20°C in IDÆA-CSIC until analysis, which was performed in
103 this institute.

104 Thatch samples ($n = 14$) covering surfaces of about 10 cm^2 were collected in-door during
105 2006-2007 and introduced in sterile polyester bags (Kapak Corporation, Minneapolis, USA) which
106 were closed with a heat sealer and stored at -20°C .

107 Thatch was elaborated from *Typha* plants (particular *Typha latifolia*).

108

109

110 2.3. Standards and reagents

111 All certified pyrethroid standards were obtained from Dr. Ehrenstorfer (Augsburg, Germany). They
112 encompassed a standard mixture of seven pyrethroids, cyfluthrin, cypermethrin, deltamethrin,
113 fenvalerate, permethrin, phenothrin and tetramethrin and single analytical standards of bifenthrin, λ -
114 cyhalothrin, esfenvalerate, fenpropathrin and resmethrin. d_6 -trans-permethrin and d_6 -trans-
115 cypermethrin were used as surrogate standard. Hexane, dichloromethane and acetonitrile were
116 obtained from Sigma Aldrich (St. Louis, MO, USA). The solvents used in this study were all
117 pesticide grade.

118 The standard solutions were prepared in ethyl acetate. In order to check the linearity of the
119 method two calibration curves were prepared at five different concentrations ranging between 0.08
120 and 2.5 ng mL^{-1} (first curve) and between 5 and 45 ng mL^{-1} (second curve). These calibration lines
121 contained d_6 -trans-permethrin and d_6 -trans-cypermethrin at 45 ng mL^{-1} and 22 ng mL^{-1} ,
122 respectively.

123

124 2.4. Sample preparation

125 Thatch material (0.3 g) and breast milk (0.1 g dry weight) were placed in 40 mL glass-centrifuge
126 tubes. They were fortified with d_6 -trans-permethrin (4.5 ng) and d_6 -trans-cypermethrin (2.5 ng) as
127 surrogate standards. The samples were stirred and extracted by sonication with 20 ml of
128 hexane:dichloromethane (2:1) in a Raypa, UCI-200 bath for 15 min. Then, the samples were
129 centrifuged at 3500 rpm for 5 min. The organic phase remained at the top of the conical tube and

130 was entirely transferred to a vial and evaporated under a nitrogen stream. This extraction step was
131 repeated two additional times and all the solvent residues were collected together.

132 Thatch material extracts were cleaned up by elution through Florisil cartridges (2g/15 ml).
133 Each cartridge was conditioned with 15 mL of ethyl acetate:dichloromethane (2:1). The sample was
134 loaded onto the cartridges and the pyrethroids were eluted with 25 mL of ethyl acetate. The eluate
135 was evaporated under a nitrogen stream and re-dissolved with 100 µL ethyl acetate for GC-NCI-
136 MS-MS analysis (Feo et al., 2010).

137 The breast milk extracts were cleaned up by elution through C18 cartridges (2g/15ml) coupled
138 to basic alumina (5g/ 25ml) and conditioned with 25 ml of acetonitrile. Then the sample was eluted
139 with 30 ml of acetonitrile. The acetonitrile extract was evaporated under a nitrogen stream and the
140 residue was dissolved in 100 µl of ethyl acetate for GC-NCI-MS-MS analysis.

141

142 2.5. GC-NCI-MS-MS operating conditions

143 GC-MS-MS analysis was performed in NCI mode on Agilent Technologies 7890A GC system
144 coupled to 7000A GC/MS Triple Quad. A DB-5MS capillary column (15m x 0.25mm i.d., 0.1 µm
145 film thickness) containing 5% phenyl methyl siloxane was used with helium as carrier gas at
146 constant flow of 1 ml min⁻¹. The temperature program was from 100°C (held for 1 min) to 230°C at
147 15°C min⁻¹, then from 230 to 310°C (held for 2 min) at 10°C min⁻¹, using the splitless injection
148 mode during 0.8 min. Inject volume was 3 µl. The inlet temperature was set at 275°C and ion
149 source temperature at 250°C. Ammonia was used as reagent gas at 2.04 x 10⁻⁴ torr. More details on
150 MS-MS condition and selected transitions were reported elsewhere (Feo et al., 2011).

151

152 2.6. Lipid content

153 Total milk lipid content was determined by crematocrit method (Mayans et al., 1994). However,
154 due to the low breast milk volume available, lipid content was not calculated in all the collected
155 samples, thus a median value was used for the calculation of pyrethroid concentrations.

156

157 2.7. *Quality control*

158 Recovery tests were carried out by addition of each pyrethroid to a thatch sample at concentrations
159 of 16 ng/g dry weight (dw) and to a breast milk sample at concentrations of 100 ng/g lipid weight
160 (lw) (Table 1). These samples were previously analyzed in order to determine pyrethroid presence
161 before spiking. Five replicates were prepared for evaluation of the reproducibility of the method.
162 Recovery values were higher than 77% in straw and ranged between 48 and 91% in breast milk
163 with relative standard deviation values lower than 3-20% (n=5; Table 1). Method detection limits
164 (MLODs) defined as the minimum amount of analyte which produces a peak with a signal-to-noise
165 ratio equal to 3 were determined for each single pyrethroid isomers by estimating the relative
166 isomer abundance of the relative peak areas. They ranged between 0.10 to 75 pg/g dw and 3.1 to
167 1100 pg/g lw for straw and breast milk, respectively (Table 1). Limits of quantification, defined as
168 the minimum amount of analyte that produces a peak with a signal-to noise ratio equal to 10, ranged
169 between 0.33 to 230 pg/g and between 8.3 to 3600 pg/g lw for straw and breast milk, respectively.

170

171 2.8. *Estimation of Nursing Infant Dietary Intake*

172 In order to evaluate the magnitude of exposure to pyrethroids by infants, the estimated daily intake
173 (EDI) was calculated as $EDI_i = C_i F Mb$ where EDI_i is the estimated intake [micrograms per
174 kilograms of body weight (bw) per day], C_i is the median concentration of pyrethroid in milk
175 samples (micrograms per grams of lipid weight), F is the lipid content in milk samples (grams of
176 lipid per 100 g of milk) and Mb is the daily consumption of milk (grams per kilograms of body
177 weight per day). The infant's average milk consumption (Mb), 175 g of milk (kg bw)⁻¹ day⁻¹, was
178 calculated from US EPA recommendations (US EPA, 2002) by assuming an average daily intake
179 breast milk rate of 702 mL of milk per day (723 g of milk per day) and a 1-month-old infant body
180 weight of 4.14 kg. The mean value of lipid content in analyzed samples was used for F estimation,
181 with a value of 4.4 g of lipid per 100 g of milk

182

183

184 3. Results and Discussion

185 3.1. Pyrethroid levels in human breast milk

186 Basic statistics of pyrethroid levels found in breast milk from Manhiça mothers are reported in
187 Table 2. λ -Cyhalothrin and permethrin were detected in all samples followed by
188 esfenvalerate/fenvalerate (found in 21 samples), cypermethrin (found in 20 samples) and
189 tetramethrin and bifenthrin (found in 19 samples) while cyfluthrin was only detected in 9 samples.
190 Phenothrin, resmethrin and deltamethrin were not found in any milk sample. The concentration
191 ranges were 1.1-36, 16-440, 10-230, 11-220, 3.3-160, 9.7-200 and 6.7-230 pg/g lw for bifenthrin, λ -
192 cyhalothrin, permethrin, cyfluthrin, cypermethrin, esfenvalerate/fenvalerate and tetramethrin,
193 respectively. The median values estimated from case-wise data were 110 ng/g lw for λ -cyalothrin,
194 70 ng/g lw for tetramethrin, 60 ng/g lw for cyfluthrin, 55 ng/g lw for permethrin and 42 ng/g lw for
195 esfenvalerate/fenvalerate. Total pyrethroid concentration ranged between 87 and 1200 ng/g lw.
196 Figure 1 shows single pyrethroid contribution (%) to the total concentration. In this case, a pair-
197 wise statistical approach was used: the most predominant pyrethroid was λ -cyhalothrin (35%)
198 followed by permethrin (21%) cypermethrin, esfenvalerate/fenvalerate and tetramethrin (14%).

199 The literature on pyrethroid levels in human breast milk samples is very scarce. Our results
200 can be compared to those found in human breast milk in Basle (Switzerland) during 1998/99
201 (Zehringer and Herrmann, 2001). In that study, 13 pyrethroids were analyzed with median
202 concentration values ranging between 15 and 31 ng/g lw. In our study median concentration values
203 of the 10 detected pyrethroid ranged between 87 and 1200 ng/g lw showing higher levels than those
204 found in Basle. This was probably due to the different use of these insecticides. In Basle,
205 pyrethroids were used only for agricultural and, in minor part, urban (e.g. pet sprays) applications
206 (Zehringer and Herrmann, 2001).

207 More recently, Sereda et al., 2009 found high permethrin levels up to 1.2 µg/g lw (mean
208 value) which occurred together with cypermethrin and cyfluthrin at lower concentration in human
209 breast samples collected from northern KwaZulu Natal, South Africa (Sereda et al., 2009). The
210 authors associated the pyrethroid contamination to home garden and indoor use. In the same region,
211 during 2006 Bouwman et al. found permethrin, cyfluthrin, cypermethrin and deltamethrin at
212 concentrations of 14.5, 42, 4.2 and 8.4 µg/l, respectively (Σ pyrethroid concentration of 31.5 µg/l)
213 which the authors associated to agriculture (Bouwman et al., 2006). These levels are higher than
214 those found in our study. According to the estimated fat content of 4% (Bouwman et al., 2006)
215 total pyrethroids ranged between 110 and 1050 with mean concentrations of 790 ng/g lw.
216 Pyrethroid concentrations found in Manhiça mothers are similar to those found in South Africa
217 (Bouwman et al., 2006; Sereda et al., 2009).

218

219 3.2. Pyrethroid levels in thatch materials.

220 Basic statistics of pyrethroid levels found in thatch material are reported in Table 2. Cypermethrin
221 was detected in 13 samples followed by λ-cyhalothrin and tetramethrin (detected in 12 samples).
222 Cyfluthrin and esfenvalerate/fenvalerate were found in 11 samples. Permethrin was found in 9
223 samples while deltamethrin, the pyrethroid used during the IRS program together with λ-cyhalothrin,
224 were found in 8 samples. Bifenthrin, phenothrin and resmethrin were also found in a few samples.
225 The concentration ranges were 0.45-7.7, 0.45-510, 0.45-695, 0.75-150, 0.50-210, 1.2-18, 2.9-30,
226 0.18-2.3, 0.52-3.1 and 0.05-0.76 ng/g dw for bifenthrin, λ-cyhalothrin, permethrin, cyfluthrin,
227 cypermethrin, esfenvalerate/fenvalerate, deltamethrin, tetramethrin, phenothrin and resmethrin,
228 respectively. The median values estimated from pairwise data were 7.4 ng/g dw for deltamethrin,
229 4.5 ng/g dw for cyfluthrin, 3.5 ng/g dw for λ-cyhalothrin, 3.2 ng/g dw for cypermethrin, 2.8 ng/g dw
230 for permethrin and esfenvalerate/fenvalerate. Total pyrethroid concentration ranged between 7.0
231 and 700 ng/g dw. In Figure 1 the most predominant pyrethroid was cypermethrin (contribution of
232 37% of the total amount) followed by cyfluthrin (25%) and λ-cyhalothrin (19%).

233

234 3.3. *Exposure and bioaccumulation of pyrethroids*

235 Pyrethroids are generally used in agriculture. This activity constitutes 80% of total income in
236 Mozambique, which traditionally was related to cotton and now also encompasses maize, soybeans
237 and rice (Arlindo and Keyser, 2007). In Manhiça people are mostly subsistence farmers and
238 workers in an agricultural cooperative that grows sugarcane, bananas and rice and also operate a
239 large sugarcane-processing factory (Alonso et al., 2002).

240 Pyrethroids have also been used for IRS being applied indoors and under the outside rafters
241 of dwellings. These insecticide compounds have also been used for ITN contributing to a reduction
242 of malaria transmission risk (Lindsay and Gibson 1998, Takken 2002). The use of λ -cyhalothrin and
243 deltamethrin for IRS was common (USAID, 2008) till the reintroduction of DDT in 2005 (WHO,
244 2006). In particular, deltamethrin was used for malaria control since 1970, when it was an
245 impregnating agent in bed nets or curtains and later it was used for IRS in spite of its marked
246 excitorepellency, which in some situations may be an advantage as it reduces human-vector contact
247 (USAID, 2008). Deltamethrin was found in thatch samples collected during 2006-2007 but it was
248 not detected in any human breast milk sample collected in 2002. Conversely, λ -cyhalothrin was
249 found in both thatch and breast milk samples. The concentration of this compound was higher
250 (median value 110 ng/g lw) than that of the other pyrethroids identified.

251 The replacement period of thatch from the dwellings is generally four years. Taken into
252 account that thatch samples were collected indoor and that the half lives of pyrethroid range
253 between 11.5 and 425 days in aerobic and anaerobic soils and between 1.83 and 619 days in water
254 (Oros et al., 2005 and Laskowski et al., 2002), we can attribute the presence of pyrethroid
255 contamination in the dwelling thatch samples analyzed during 2006-2007 to agricultural
256 applications but also to their use for IRS.

257 Human exposure to pyrethroids, including λ -cyhalothrin, can result from various routes,
258 including dermal uptake, inhalation (dust and gas phase) and ingestion of food containing residues

259 of this group of insecticides (Bouwman and Kylin, 2009). The use of these compounds in
260 agriculture and for IRS may therefore explain their occurrence in the milk samples collected in
261 2002.

262 Results on pyrethroid levels found in human breast milk demonstrate that mothers exposed to
263 insecticide contamination accumulate pyrethroids that could be transfer to infants via breast milk.
264 Some studies reported that pyrethroids are metabolized by humans: the chrysanthemic acid ester is
265 usually cleaved via esterase or mixed function oxidase activity and the resulting alcohol moieties
266 are converted to their corresponding acids. It is reported that these metabolites are partly conjugated
267 to glucoronide and both the conjugates and free acids are excreted in urine (ATSDR, 2003).
268 However, and based on our results, bioaccumulation of pyrethroid in women is evident and it seems
269 to differ depending on the pyrethroid. Figure 1 shows the percentage contribution of each detected
270 pyrethroid in thatch material and human breast milk. The distribution patterns are different which
271 may indicate changes in pyrethroid use through time or the combination of different pyrethroid
272 sources, e.g. domestic and agricultural applications, in breast milk.

273 Pyrethroid molecules typically contain 2-3 asymmetric carbon atoms (chiral centers), making
274 them a family of pesticides with high chirality. Figure 2 shows the relative contributions of the two
275 isomers of permethrin and esfenvalerate/fenvalerate found in a commercial technical mixture
276 (standard), as well as in thatch material and human breast milk. For permethrin, the abundance in
277 the commercial technical mixture is 84% and 16%, for isomer I and II respectively. In thatch
278 material, an abundance of 69% and 31% was found, showing a roughly similar distribution.
279 However, for human breast milk samples, the percentage contribution of both isomers is very
280 similar, with 52% and 48%, respectively. The observed enrichment in isomer II may reflect higher
281 bioaccumulation potential of this compound or, conversely, a higher degree of human
282 metabolization of isomer I. To the best of our knowledge this selective enrichment in isomer
283 composition is described here for the first time. Analysis of milk samples in forthcoming studies are
284 needed for a better understanding of the processes leading to this preferential accumulation.

285 In the case of esfenvalerate/fenvalerate, the abundance in the commercial technical mixture is
286 of 60% and 40%, for isomer I and II respectively. These percentages were similar to those found in
287 thatch material and also in human breast milk samples, with 60% and 40% and 56% and 44% for
288 isomer I and II in thatch material and human breast milk, respectively. In this case, no differential
289 isomeric behavior was observed.

290

291 *3.4. Nursing Infant Estimated Dietary Intake*

292 EDI values calculated from the breastmilk concentrations (Table 2) were 0.12, 0.28, 1.5, 1.7, 1.75,
293 1.8 and 3.4 $\mu\text{g/kg bw}$ and per day for cypermethrin, bifenthrin, esfenvalerate/fenvalerate, cyfluthrin,
294 permethrin, tetramethrin and cyhalothrin, respectively (Table 3). These values were compared to the
295 recommended acceptable daily intake (ADI) values reported by FAO and WHO 2005 for bifenthrin
296 ($4 \mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$), cyfluthrin ($20 \mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$), cypermethrin ($20 \mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$),
297 deltamethrin ($10 \mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$), λ -cyhalothrin ($5 \mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$) and permethrin ($50 \mu\text{g}$
298 ($\text{kg bw)}^{-1} \text{ day}^{-1}$) (FAO and WHO 2005 and FAO 2005). In general, the rates estimated from the
299 mothers in Manhiça are lower than these recommended levels but the levels of λ -cyhalothrin in
300 some mothers, the pyrethroid used during IRS program in Mozambique, were close to the ADI
301 WHO-recommended value ($5 \mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$).

302 Further comparison of these breast milk pyrethroid concentrations can be obtained by
303 comparison to the maximum residue levels (MRLs) after unit transformation in mg/kg of milk
304 (FAO and WHO, 2005). These MRLs are proposed for food commodities or animal feeds. As
305 shown in Table 4, the median concentrations of the six pyrethroids found in milk samples above
306 limit of detection correspond to concentrations that are about six times lower than MRL values at
307 the most. The mothers exhibiting highest concentrations have also values below these MRL
308 reference levels.

309

310

311 4. Conclusions

312 Pyrethroids were found in human breast milk despite the discontinuation in the use of these
313 compounds for IRS. Their occurrence may reflect an influence from the insecticide impregnated
314 bed nets, agricultural sources or use for IRS in some cases despite their known low efficiency for
315 malaria control. The presence of these compounds in breast milk confirms their bioaccumulation
316 potential in humans. Some pyrethroid compounds are accumulated with isomeric discrimination.
317 The concentrations of some of these insecticides found in some mothers, namely λ - cyhalothrin
318 used for IRS in Mozambique, involves EDI values close to the upper limits recommended by FAO.
319 The presence of pyrethroids in thatch from dwellings evidences that these insecticides are still used
320 for IRS. The observed occurrence of pyrethroids in dwellings despite the preferential use of other
321 insecticides for IRS and the concentrations of some of these compounds found in human breastmilk
322 evidence that these compounds need to be considered in the evaluation of infant risks associated to
323 lactation in areas where insecticides are used for elimination of malaria vectors.

324

325

326 ACKNOWLEDGEMENTS

327 We thank all the families for their participation in the study and the staff of the Manhiça Health
328 Research Center for their support during data and sample collection, particularly Jahit Sacarlal,
329 Raquel Gonzalez, Inacio Mandomando, for procurement of the straw and breast milk samples used
330 in this study. MNM is funded by a PhD Scholarship from Fundació Marfà. CD is supported by the
331 Spanish Ministry of Science and Innovation (MICINN; RYC-2008-02631). Funding was received
332 from MICINN (INMA G03/176, Consolider Ingenio GRACCIE, CSD2007-00067), CSIC (PIF06-
333 053) and ArcRisk EU Project (FP7-ENV-2008-1-226534). The Centro de Investigaçao em Saúde da
334 Manhiça receives core support from the Spanish Agency for International Cooperation and
335 Development (AECID).

336

337 **Figure Caption:**

338

339 **Figure 1.** Percentage contribution of each pyrethroid to total contamination estimated in straw
340 material and human breast milk collected from Manhiça (Mozambique).

341

342 **Figure 2.** Abundance (%) of the two isomers of permethrin and esfenvalerate/fenvalerate in straw
343 material and human breast milk collected from Manhiça (Mozambique).

344

345 **References:**

- 346 Alonso P, Saute F, Aponte JJ, Gomez-Olive FX, Nhacolo A, Thompson R, et al. Manhiça DSS,
347 Mozambique In: INDEPTH, editor. Population and Health in Developing countries. Ottawa:
348 International Development Research Centre (IDRC); 2002. p. 189–95.
- 349 Arlindo P., Keyser J.C. Background paper for the competitive commercial agriculture in Africa
350 study (CCAA). Mozambique country case study. December 10, 2007.
- 351 ATSDR. Toxicological profile for pyrethrins and pyrethroids, Agency for toxic substances and
352 disease registry, Atlanta, 2003.
- 353 Bouwan H, Sereda B, Meinhardt HM. Simultaneous presence of DDT and pyrethroid residue in
354 human breast milk from a malaria endemic area in South Africa. Environ. Pollut 2006, 144:
355 902-917.
- 356 Bouwman H., Kylin H. Malaria control insecticides residues in breast milk: the need to consider
357 infant health risks. Environ health perspect 2009, 117, 10: 1477-1480.
- 358 Cox, C. Herbicide factsheet: glufosinate. Pestic. Reform 1996; 16 : 15-19.
- 359 FAO 2005. Codex Alimentarius. < www.codexalimentarius.net>.
- 360 FAO and WHO, 2005. Joint meeting of the panel of experts on pesticide residues.
361 www.who.int/ipcs/publications/jmpr/en/.
- 362 Feo ML, Eljarrat E, Barceló D. Presence of pyrethroid pesticides in water and sediments of Ebro
363 River Delta. J Hydrol 2010, 393: 156-162.
- 364 Feo ML, Eljarrat E, Barceló D. A rapid and sensitive analytical method for the determination of
365 fourteen pyrethroids in water samples. J Chromatogr A 2010, 1227: 2248-2253.
- 366 Feo M.L., Eljarrat E., Barceló D. Performance of gas chromatography/tandem mass spectrometry
367 in the analysis of pyrethroid insecticides in environmental and food samples. Rapid Commun
368 Mass Spec 2011, 25: 869-876.

369 Hargeaves K, Koekemoer LL, Brooke BD, Hint RH, Mthembu J, Coetzee M. *Anopheles funestus*
 370 resistant to pyrethroid insecticides in South Africa. *Medical and Veterinary Entomology*
 371 2000, 14: 181-189.

372 Kapp C. Hazard or help? *The Lancet* 2004, 364: 1113-1114.

373 Laskowski, DA. Physical and chemical properties of pyrethroids. *Rev. Environ. Contam. Toxicol.*
 374 2002, 174: 49-170.

375 Lindsay SW and Gibson ME. Bednets revised old idea new angle. *Parasitology Today* 1998, 4:
 376 270-272.

377 LSDI. Spatial Development Initiative. Annual Report 2006

378 Mabaso MLH, Sharp B, Lengeler C. Historical review of malarial control in southern Africa with
 379 emphasis on the use of indoor residual house-spraying. *Tropical Medicine and International*
 380 *Health* 2004, 9: 846-856.

381 Manaca MN, Grimalt JO, Gari M, Sacarlal J, Sunyer J, Gonzalez R, Dobaño C, Menendez C,
 382 Alonso PL. Assessment of exposure to DDT and metabolites after indoor residual spraying
 383 through the analysis of straw material from rural African dwellings. *Environ Sci Pollut Res*,
 384 submitted.

385 Mayans E, Martell M. Estimacion del valor calorico de leche materna mediante la tecnica del
 386 crematocrito. *Red Med Uruguay* 1994, 10: 160-164.

387 Montgomery CM, Munguambe K, Pool R. Group-based citizenship in the acceptance of indoor
 388 residual spray (IRS) for malaria control in Monzabique. *Soc Sci Med.* 2010, 70: 1648-55.

389 Oros, DR, Werner I. Pyrethroid insecticides: Analysis of Use Patterns, Distributions, Potential toxicity, and
 390 Fate in the Sacramento-San Joaquin Delta and Central Valley. White paper for the Interagency
 391 Ecological Program. 2005 SFEI Contribution 415. San Francisco Estuary Institute, Oakland,
 392 California.

393 Sereda BL, Meinhardt HR. Contamination of the Water Environment in Malaria Endemic Areas of
394 KwaZulu-Natal, South Africa by DDT and Its Metabolites. Bull. Environ. Contam. Toxicol.
395 2005, 75: 538–545.

396 Sereda B, Bouwan H, Kylin H. Comparing water, bovine milk and indoor residual spraying as
397 possible sources of DDT and pyrethroid residues in breast milk. J. Toxicol Environ Health A
398 2009, 72: 842-851.

399 Takken W. Do insecticide treated bednets have an effect on malaria vectors? Trop Med Intern
400 Health. 2002. 7,12: 1022-1030.

401 US Environmental Protection Agency. Child-specific exposure factors handbook, National Center
402 for Environmental Assessment: Washington, DC, 2002, 2 and 11: 20.

403 WHO and UNICEF 2005. World Malaria Report 2005. WHO/HTM/MAL/2005.1102. Geneva:
404 World Health Organization and United Nations Children's Fund.

405 WHO 2006. Indoor Residual Spraying: Use of indoor residual spraying for Scaling up Global
406 Malaria Control and Elimination. WHO/HTM/MAL/2006:1112. Geneva: World Health
407 Organization.

408 WHO. 2007a. Malaria Elimination: A field manual for low and moderate endemic countries.
409 Geneva: World Health Organization.

410 WHO 2008. WHO Pesticide Evaluation Scheme. Available: <http://www.who.int/whopes/en/>.

411 World Malaria Report. WHO. 2010.
412 http://www.who.int/malaria/world_malaria_report_2010/en/index.html2.

413 USAID from the American people. United States Agency International Development. IVM PEA
414 Supplemental Environmental Assessment: Indoor residual spraying using registered carbamates,
415 pyrethroids and DDT in Mozambique. January 30, 2008.

416 Zehring M, Herrmann A. Analysis of polychlorinated biphenyls, pyrethroid insecticides and
417 fragrance in human milk using a laminar cup liner in the GC injector. Eur Food Res Technol
418 2001, 212: 247-251.

419

420

421

Table 1. Analytical quality parameters of pyrethroid methodologies applied to straw material and breast milk samples.

	Straw material					Breast Milk				
	Blank (ng/g dw)	Recovery (%)	RSD (%)	MLODs ^a (pg/g dw)	MLOQs ^a (pg/g dw)	Blank (ng/g lw)	Recovery (%)	RSD (%)	MLODs ^a (pg/g lw)	MLOQs ^a (pg/g lw)
Bifenthrin	1.2	86	5	1.7	5.1	0.82	70	20	32	97
λ-Cyhalothrin	2.2	78	5	1.1	3.3	9.05	82	20	3.6	11
Permethrin	3.8	82	8	11; 9.5	33; 28.5	24	80	4	1100; 1100	3200; 3200
Cyfluthrin	1.1	85	20	2.7; 2.5; 0.10	8.1; 7.6; 0.30	nd	60	20	160; 160; 3.1	480; 480; 9.2
Cypermethrin	1.6	90	19	2.0; 4.5; 3.7	6.0; 13.5; 11	3.55	72	6	140; 140; 140	430; 430; 430
Es/fenvalerate	2.8	98	20	1.0; 0.90	3.0; 2.7	nd	70	7	63; 63	190; 190
Deltamethrin	3.0	104	20	13	39	nd	48	8	280	8.3
Tetramethrin	0.75	77	3	0.41; 0.75	1.2; 2.25	17	86	20	45	140
Phenothrin	0.84	60	20	67; 69	200; 210	nd	50	20	1100; 1100	3200; 3600
Resmethrin	Nd	87	20	69; 75	210; 230	nd	91	16	800; 780	2400; 2300
Total	17					54				

^aMLODs and MLOQs were estimated for each isomer of a specific pyrethroid.

Table 2. Basic statistics for single pyrethroid and Σ pyrethroid concentrations in straw materials (ng/g dw) and human breast milk (ng/g lw) of Mozambique.

	Bifenthrin	λ -Cyhalothrin	Permethrin	Cyfluthrin	Cypermethrin	Es/Fenvalerate	Deltamethrin	Tetramethrin	Phenothrin	Resmethrin	Σ PYR
<i>Straw</i>											
Mean	2.9	59	80	17	24	5.4	11	0.68	1.4	0.43	162
Median	2.6	3.5	2.8	4.5	3.2	2.8	7.4	0.54	1.1	0.47	55
Max	7.7	510	695	150	210	18	30	2.3	3.1	0.76	700
Min	0.45	0.45	0.45	0.75	0.50	1.2	2.9	0.18	0.52	0.05	7.0
SD	2.6	150	230	45	60	5.2	9.1	0.59	1.0	0.36	220
<i>Milk</i>											
Mean	6.5	140	79	80	54	55	nd	80	nd	nd	425
Median	4.0	110	55	60	34	42	nd	70	nd	nd	370
Max	36	440	230	220	160	200	nd	230	nd	nd	1200
Min	1.1	16	10	11	3.3	9.7	nd	6.7	nd	nd	87
SD	8.0	120	62	65	50	44	nd	64	nd	nd	265

Table 3. Estimated daily intake (EDI) of pyrethroids (expressed as $\mu\text{g (kg bw)}^{-1} \text{ day}^{-1}$) calculated from the breastmilk concentrations of these compounds in Manhiça (Mozambique).

	Bifenthrin	λ -Cyhalothrin	Permethrin	Cyfluthrin	Cypermethrin	Es/Fenvalerate	Tetramethrin	ΣPYR
Mean	0.05	1.05	0.61	0.61	0.42	0.42	0.62	3.3
Median	0.03	0.85	0.42	0.46	0.03	0.33	0.54	2.8
Max	0.28	3.4	1.75	1.7	0.12	1.5	1.8	9.0
Min	0.01	0.13	0.08	0.09	0.00	0.07	0.05	0.67

Table 4. Median values of pyrethroids concentration in milk samples expressed as mg/kg compared to MRL values reported by WHO FAO 2005.

	Median concentrations in milk samples (mg/kg)	MRL values (mg/kg) FAO WHO 2005.
Cyfluthrin	0.0019	0.04
Deltamethrin	nd	0.05
Cypermethrin	0.0009	0.05
Fenvalerate	0.0128	0.1
Permethrin	0.0059	0.1

Figure 1.

Figure 2.