Submitted to *Environmental Research*

An evaluation of the sexual differences in the accumulation of organochlorine compounds in children at birth and at the age of four years

**Joan O. Grimalt^a, Daniel Carrizo^a, Mercè Garí^a, Laia Font-Ribera^a, Nuria Ribas-Fito^b,
Maties Torrent^c, and Jordi Sunyer^b**

^aDepartment of Environmental Chemistry, Institute of Environmental Assessment and Water Research (IDÆA-CSIC), Jordi Girona, 18, 08034-Barcelona, Catalonia, Spain

^bCentre for Research in Environmental Epidemiology (CREAL), Barcelona, Catalonia, Spain

^cÀrea de Salut de Menorca, IB-SALUT, Menorca, Spain.

Address correspondence to J.O. Grimalt, Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (IDÆA-CSIC), Jordi Girona, 18, 08034-Barcelona, Catalonia, Spain. Telephone: 34934006122. Fax: 34932045904. E-mail: jgoqam@cid.csic.es

19 **Abstract**

20

21 This study of the body burden and serum concentrations, of organochlorine compounds (OCs),
22 represents a general population in a cohort from Menorca Island (birth 1997-1998) of children at
23 birth and at four years of age; the study has shown that the concentrations of hexachlorobenzene
24 (HCB), 4,4'-DDE, 4,4'-DDT, polychlorobiphenyl (PCB) congeners #153, #138 and #180 and total
25 PCBs in sera collected at four years exhibit much higher values in breastfed children than in those
26 fed with formula, e.g. HCB 0.48 vs 0.21 ng/ml, β -HCH 0.32 vs 0.24 ng/ml, total DDTs 2.2 vs 0.57
27 ng/ml, total PCBs 1.4 vs 0.52 ng/ml, respectively. Comparison of gender differences in four years
28 old children shows higher concentrations of all examined OCs in females than in males with the
29 exception of HCB and PeCB in breastfed children which are higher in males than in females, e.g. β -
30 HCH 0.34 vs 0.28 ng/ml, total DDTs 2.6 vs 1.7 ng/ml, total PCBs 1.6 vs 1.0 ng/ml, respectively, for
31 breastfed children and e.g. β -HCH 0.23 vs 0.19 ng/ml, total DDTs 0.59 vs 0.48 ng/ml, total PCBs
32 0.58 vs 0.45 ng/ml, respectively, for formula fed children. Gender comparison of the body burden
33 between children fed with breast milk or formula also shows higher concentrations in females than
34 males, e.g. β -HCH 0.47 vs 0.35 μ g, total DDTs 3.0 vs 1.8 μ g, total PCBs 1.9 vs 1.2 μ g,
35 respectively, for breastfed children, and β -HCH 0.39 vs 0.17 μ g, total DDTs 0.48 vs 0.27 μ g, total
36 PCBs 0.66 vs 0.55 μ g, respectively, for formula fed children. The results may suggest a higher
37 capacity in female children for the retention of OC incorporated through breast feeding. However,
38 these results should be taken with caution because the differences of the gender averages have low
39 statistically significance when evaluated with the Student test.

40

41 Key words: Organochlorine compounds, persistent organic pollutants, breastfeeding,
42 polychlorobiphenyls, 4,4'-DDE, pollutants in cord serum, background concentrations of
43 organochlorine compounds in children sera, newborns

Informe justificativo del Coordinador Científico de la Red

TO WHOM IT MAY CORRESPOND

Barcelona, 16th October 2006

The manuscript entitled **An evaluation of the sexual differences in the accumulation of organochlorine compounds in children at birth and at the age of four years** by Carrizo et. al. to be submitted to *Environmental Research* has been sponsored by the Spanish Research Network "Infancia y Medio Ambiente" (G03/176) that is funded by the Spanish Health Funding Agency: Instituto de Salud Carlos III.

This study was approved by the ethics committee of the Institut Municipal d'Investigació Mèdica and all mothers provided a signed informed consent.

Yours sincerely,

Jordi Sunyer
Coordinator of the INMA Network
ISCIII (G03/176)
jsunyer@imim.es

1. Introduction

4,4'-DDE and polychlorobiphenyls (PCBs) are incorporated into children *in utero* and through diet, namely breastfeeding (Huisman et al. 1995; Karmaus et al. 2001; Rhainds et al. 1999; Ribas-Fito et al. 2003; 2005; Carrizo et al., 2006). The lipophilic properties and high stability to chemical degradation of these compounds favours their accumulation in human fat. These properties are those characteristics of persistent organic pollutants (POPs) and define a group that encompasses those mentioned above and others such as hexachlorobenzene (HCB), hexachlorocyclohexanes (HCHs), 4,4'-DDT, involving up to twelve types of organochlorine compounds (OCs) that have been banned by the Stockholm agreement (Stockholm Convention on Persistent Organic Pollutants, 2005).

One key aspect needed for the assessment of the risks of these compounds into human health is the understanding of the processes leading to their accumulation into humans, namely in the first period of growth. Several studies have documented the significance of intake through breastmilk in children that is a major determinant of their body burden, even after several years of discontinuation of this feeding mode (Lanting et al., 1998; Jacobson et al., 1989; Karmaus et al., 2001; Carrizo et al., 2006). Other studies have also shown the influence of environmental exposure in relation to some specific activities (e.g. Ribas-Fito et al., 2005; Sala et al., 2001).

In contrast, little attention has been paid to the importance of the own characteristics of the exposed individuals in OC intake. In this respect, sex is one of the most obvious aspects to consider. Accordingly, the present paper reports a study devoted to assessing the gender differences in the accumulation of these compounds in newborns and children at the age of four years. This goal has been addressed by considering whether children were fed with breast milk or formula; since, as mentioned above, these two feeding practices are major determinants of the accumulation of these compounds in the early stages of life.

The population selected for this study is from Menorca, one of the Balearic Islands in the northwest Mediterranean Sea. The island does not have factories producing OCs. DDT was used for agriculture in the past. Thus, the participating children were exposed to background POP levels and can be taken as examples of the regular exposure to these pollutants in western countries. A general population birth cohort was set up in 1997 within the Asthma Multicenter Infants Cohort study (AMICS) (Polk et al. 2004). The cohort recruited all women presenting for antenatal care over 12 months starting in mid 1997. 482 children were subsequently enrolled and 470 (97.5%) provided complete outcome data up to the fourth year visit. Among these 410 (85%) had OCs measured in cord serum and 285 (59%) in serum collected at four years.

2. Materials and Methods

Materials

Standards of tetrabromobenzene (TBB), pentachlorobenzene (PeCB), HCB, α -, β -, γ -, and δ -HCH, PCBs, 4,4'-DDT and 4,4'-DDE were purchased from Dr. Ehrenstoffer (Augsburg, Germany). Analytical grade concentrated sulphuric acid (conc.-H₂SO₄), iso-octane and *n*-hexane were all purchased from Merck (Darmstadt, Germany).

Sample extraction and clean up

Samples were obtained in the context of the procedures defined by the INMA Project (Fernandez et al., 2007). Serum samples (0.5 mL) were introduced into 10 mL centrifuge tubes and TBB and PCB 209 were added as recovery standards. Two mL of conc.-H₂SO₄ and 3 mL of *n*-hexane were added, mixed in a vortex (*ca.* 1500 rpm, 30 seconds) and then centrifuged (*ca.* 1500 rpm, 10 minutes). The supernatant *n*-hexane layer was aspirated into a second centrifuge tube using

136 a Pasteur pipette. Further, *n*-hexane (2 mL) was added to the first tube containing the H₂SO₄ /
137 serum, stirred (vortex *ca.* 1500 rpm, 30 s) and then centrifuged (*ca.* 1500 rpm, 10 min). This last
138 step was repeated, yielding a combined extract of 7 mL of *n*-hexane, to which 2 mL conc-H₂SO₄
139 was added, the sample mixed (vortex mixer, *ca.* 1500 rpm, 90 s), centrifuged as before, and the
140 supernatant transferred to a conical bottomed, graduated tube. The combined extracts were then
141 reduced to near dryness under a gentle stream of nitrogen and an injection standard (PCB 142 in
142 isooctane; 10 µL) was added. Then, the sample was quantitatively transferred to gas
143 chromatographic vials using four 25 µL rinses of isooctane. If an emulsion was formed at any stage
144 of the extraction, 10 – 15 drops of MilliQ water were added before sample centrifugation.

145

146 *Composition of organochlorine compounds*

147 Selection of OC for analysis was based on a literature search of commonly-studied
148 contaminants and included representatives of industrial and agrochemical products. Four HCH
149 isomers were analyzed (α -, β -, γ - and δ -). The α -, γ - and δ -isomers were only found above
150 quantification limit in less than 5% of total samples. Therefore, these compounds were not included
151 in the database of the study. Among the large family of PCB congeners, those often referred to as
152 the “ICES 7” (from International Council for the Exploration of the Sea) were selected: PCBs #28,
153 #52, #101, #118, #138, #153 and #180. These congeners are frequently found at high
154 concentrations in humans and wildlife. In all cases, when the concentrations were below the limit of
155 detection or quantification (Table 2), the value of 0 was introduced. Substitution of these values by
156 half of the limit of quantification (Table 2) did not change the results or the significance of the
157 statistical tests nor the correlation analyses discussed below.

158

159 *Instrumental determinations*

160 A gas chromatograph with electron capture detection (Hewlett Packard 6890N GC-ECD)
161 was used to quantify PeCB and HCB, PCB congeners #28, #52, #101, #118, #138, #153, #180,
162 *p,p'*-DDT and *p,p'*-DDE. α -, β -, γ -, and δ -HCH were quantified with GC-MS (HP 5973 MSD) in
163 negative chemical ionisation mode using ammonia as reagent gas (1.0 mL/min). In both instruments
164 samples were injected (2 μ L) in splitless mode onto a 60 m, DB-5 column with a retention gap
165 (both from J&W and Agilent). Helium was the carrier gas (1.5 mL/min). The temperature program
166 was: 90°C 2 min, 20°C/min to 140°C, 4°C/min to 200°C held for 13 min, 4°C/min to 310°C held
167 for 10 min.

168 In both instruments, quantification was performed by external standards using PCB 142
169 injection standard to correct for volume. Recoveries of TBB and PCB 209 (mean $\pm$ standard error =
170 102 ± 4.2 and 93 ± 1.8 , respectively) were used to correct results. Limits of detection (LOD) and
171 quantification (LOQ) were calculated from blanks (LOD = mean of all blanks plus three times the
172 standard deviation, LOQ = mean plus five times the standard deviation) or from instrumental LOD
173 using diluted standards if the compound was absent from the blanks.

174 This method performed satisfactorily in repeated international intercalibration exercises
175 within the Arctic Monitoring and Assessment Programme (AMAP 2004).

176 This study was approved by the ethics committee of the Institut Municipal d'Investigació
177 Mèdica of Barcelona and all mothers provided a signed informed consent.

178

179 *Statistical analyses*

180 Means and standard deviations of the OC concentrations or body burden in the diverse
181 groups of children, e.g. breastfeeding and formula fed, males and females, were used for
182 comparison of their differences with were evaluated according to the t test.

183

184

185 3. Results

186

187 *Characteristics of the population under study*

188 Sex, feeding practices, and duration of lactation of children in the population studied are
189 shown in Table 1. 83% of children were breastfed. Duration of lactation encompassed from very
190 short periods (2 months or less) to large periods (more than one year). No significant selection
191 biases between the group of participants at birth (n = 410) and 4 years later (n = 285) are observed.
192 No significant differences in feeding practices, either maternal or formula, or duration of lactation
193 are found between the two gender groups of this population.

194 Over the whole population included in the study, 4,4'-DDE is the most abundant OC, in
195 both cord serum and serum collected at 4 years (average 1.6 ng/ml in both cases; Table 2). The
196 lower concentrations of 4,4'-DDT (0.18 ng/ml and 0.073 ng/ml, respectively) than 4,4'-DDE likely
197 reflect that the whole mixture of DDT metabolites correspond to past exposure of this pesticide
198 because a substantial amount of the 4,4'-DDT initially introduced into the environment has already
199 been transformed into 4,4'-DDE (Wedemeyer, 1967; Aguilar, 1984).

200 HCB is the second major OC involving average values of 0.75 ng/ml and 0.42 ng/ml in cord
201 and four years sera, respectively (Table 2). Total concentrations of the ICES 7 PCB congeners are
202 0.70 ng/ml and 1 ng/ml, respectively. The PCB distributions are dominated by congener #153 in
203 both types of samples. β -HCH is found at concentrations of 0.21 ng/ml and 0.28 ng/ml in cord and
204 four years sera, respectively. PeBC is the OC found in lowest concentration, 0.081 ng/ml and 0.023
205 ng/ml, in serum collected at birth and at four years, respectively, among those included in the
206 present study.

207

208 *Gender differences*

A summary of the gender differences in the populations of newborns and 4 year-old children is represented in Fig. 1. The results are in agreement with those discussed above and summarized in Table 2. However, small gender differences can be observed. Thus, in newborns slightly higher concentrations of HCB and 4,4'-DDE are observed in male whereas PCBs are found in higher concentrations in female (Fig. 1). In contrast, in the examined population of children at the age of 4 years, 4,4'-DDE, PCBs and β -HCH are present in higher concentrations in females than in males.

Influence of breastfeeding

The average concentrations of HCB, 4,4'-DDE, 4,4'-DDT, PCB congeners #153, #138 and #180 and total PCBs in serum collected at four years exhibit significant higher values in breastfed children than in those fed with formula (Table 3). The degree of statistical probability of the dissimilar averages is very high ($p < 0.0001$) for HCB and most hydrophobic compounds. Therefore, breastfeeding is a key factor for the presence of these OC in children at age four even after 3.5-2.3 years that this practice was discontinued. A full discussion of the maternal aspects such as age, body mass index (BMI) and period of lactation that are correlated to OC body burden in children at four years of age is given elsewhere (Carrizo et al., 2006). None of those was statistically significantly correlated with the gender differences examined in the present study.

The gender averages of OCs in the population of four years old children fed with breastmilk show higher concentrations in female than male for all OCs except PeCB and HCB (Table 3). Formula fed children also show higher concentrations in female than male but the gender differences are higher in breastfed than formula fed children (Table 3). These gender differences are statistically significant in the case of 4,4'-DDT among breastfed children ($p < 0.1$; Table 3). Neither BMI nor age of the mother were statistically significantly correlated with the gender differences examined in the present study.

233 Calculation of the concentration differences in serum OC at four years and at birth shows a
234 general increase for breastfed children (Table 4). This increase is observed for all OCs with the
235 exception of HCB and a small decrease found in the case of PeCB, 4,4'-DDT and PCB-180. In
236 contrast, formula fed children show small variations and those involving more than 0.1 ng/ml,
237 HCB, 4,4'-DDE, 4,4'-DDT, PCB-180 and total PCBs, exhibited decreases (Table 4). The
238 differences between breast and formula fed children are statistically significant for HCB, 4,4'-DDE,
239 4,4'-DDT and most of the hydrophobic PCBs.

240 Comparison of the concentration changes in breastfed children after grouping by gender
241 shows, for all OCs except PeCB and HCB, higher concentration increases, or lower concentration
242 decreases, in female than male (Table 4). The differences are statistically significant in the case of
243 4,4'-DDE ($p < 0.1$; Table 4). Gender comparison of the concentration changes in formula fed
244 children also shows, for all OCs except HCB, PCB-28 and PCB-52, higher increases, or lower
245 decreases, in female than male children.

246 Straightforward comparison of the concentrations at birth and at four years overlooks the
247 changes in blood volume due to children growth. Calculation of OC body burdens allows the
248 inclusion of this aspect in the calculations.

249

250 *Body burden*

251 Transformation of the measured concentrations at birth and at the age of four years into
252 body burden using the estimated total amount of serum from weight, *ca.* 0.075 L/kg, allows the
253 calculation of the increase in OC amounts in each individual between these two periods (Table 5).
254 The average values show that there is an OC increase between the two ages for all compounds.
255 Except in the case of PeCB, the magnitude of the increase is higher among breastfed than formula
256 fed children which is consistent with the above mentioned concentration differences. The OC body
257 burden differences are statistically significant for HCB and all the most hydrophobic OC (Table 5).

258 Examination of the gender averages of this body burden data shows again a higher intake in female
259 than male children. The differences are about the same as those previously discussed on
260 concentrations.

261

262

263 **4. Discussion**

264

265 *OC concentrations in children from Menorca and other populations*

266 The OC concentrations found in the children from the cohort of Menorca correspond to
267 background values of general population. The observed average concentrations of total PCB in cord
268 serum, 0.68 ng/ml, are lower than those found in studies from Norway (3.0 ng/ml, Skaare et al.
269 1988), USA (2.5 ng/ml, Schwartz et al. 1983), Faroe Islands (1.1 ng/ml, Grandjean et al. 1997) and
270 Germany (0.96 ng/ml or 1.4 ng/ml, Lackmann et al. 1996). On the other hand, they are higher than
271 those reported in the Netherlands (0.38 ng/ml, Huisman et al. 1995), Canada (0.50 ng/ml, Rhainds
272 et al. 1999) or Catalonia (0.36 ng/ml, Sala et al. 2001). Comparison of these data must be done with
273 caution because different PCB congeners were used for the calculation of total PCBs in each study.
274 However, the reported figures may vary by a factor of two at the most. In this cohort the PCB
275 congeners of higher volatility were also considered for quantification which is not usually
276 performed in other studies.

277 The concentrations of HCB in cord serum from these Menorcan children, 0.75 ng/ml, are
278 lower than those reported in Norway (1.0 ng/ml, Skaare et al, 1988), Germany between 1984-85
279 (2.0 ng/ml, Lackmann et al. 1996) or Catalonia (1.2 ng/ml, Sala et al., 2001) but higher than the
280 average concentrations found in Germany between 1994-95 (0.61 ng/ml, Lackmann et al. 1996) or
281 Canada (0.04 ng/ml, Rhainds et al. 1999). The concentrations of 4,4'-DDE, 1.6 ng/ml, are lower
282 than those reported in Norway (3.0 ng/ml, Skaare et al. 1988) and higher than the average

measurements of Canada (0.4 ng/ml, Rhainds et al. 1999) or Catalonia (0.83 ng/ml, Sala et al. 2001). Previous studies have outlined the capacity of these compounds to cross the placenta leading to *in utero* exposure (DeKoning and Karmaus, 2000).

OCs in male and female children

The cord serum OC concentrations show higher concentrations of 4,4'-DDE and HCB in male than female newborns, about the same concentrations of PeCB and β -HCH in both genders and higher PCB concentrations in female than male (Fig. 1) but the differences are not statistically significant ($p < 0.1$). These differences change when comparing the same children at age 4. Now, the concentrations of all OC except PeCB and HCB are higher in female than male breastfed children but again the differences are only statistically significant for 4,4'-DDT ($p < 0.1$; Table 3). These differences are not related with different BMI of the two gender groups since both of them have the same average value, 16.1 kg/m² (standard deviation 1.4 kg/m²). In formula fed children all OC concentrations are higher in female than male (Table 3). Examination of the BMI of both gender groups show higher average values in male, 17.3 kg/m² (standard deviation 2.5 kg/m²), than female, 16.3 kg/m² (standard deviation 2.6 kg/m²). The differences are not statistically significant but, in any case, the higher BMI values are found in male children (the group in which lower OC concentrations were found). Thus, the OC differences in formula fed children cannot be attributed to BMI differences. In a previous study, on children at 7 years of ages from the Federal State of Hessen (Germany), the concentrations of 4,4'-DDE, β -HCH and HCB in both genders were practically the same but total PCBs were significantly higher in male than female ($p < 0.001$) (Karmaus et al. 2001). Higher levels of PCBs were also found in male than female children, at age four, from Michigan (USA) but the differences were small (Jacobson et al. 1989).

Breastfeeding

308 As shown in Table 3, breast feeding is a major way of incorporation of OCs into children of
309 the population of Menorca, which is in agreement with previous observations for some OCs in other
310 studies, e.g. 4,4'-DDE (Karmaus et al. 2001), 4,4'-DDT (Jacobson et al. 1989), β -HCH (Karmaus et
311 al. 2001), HCB (Karmaus et al. 2001) and PCB mixtures encompassing diverse congeners with
312 more than five chlorine substituents (Jacobson et al. 1989; Lanting et al. 1998; Karmaus et al.
313 2001). A full account of the maternal characteristics influencing the intake of OCs through
314 breastfeeding is given in Carrizo et al. (2006).

315 Examination of the gender differences between breast and formula feeders shows higher
316 contrasts in breast fed female children. Thus, all OC exhibit higher concentrations in female than
317 male children (Table 3). The dependence between breastfeeding and gender differentiation is also
318 illustrated in Fig. 2 where the average composition of the main OC groups is represented after
319 grouping by gender and feeding practices. As shown in this figure higher duration of breastfeeding
320 involves higher gender differentiation of the average values but the coefficients of variation are also
321 larger. These differences suggest that female children could retain more efficiently OC incorporated
322 through breastmilk. In this context, the parity differences between the two groups have also to be
323 considered. For breastfed children they are 1.1 (standard deviation 0.74) and 0.8 (standard deviation
324 0.74) for males and females, respectively, and for formula fed children they are 1.2 (standard
325 deviation 0.97) and 1.1 (standard deviation 0.76), respectively. These differences are small and not
326 statistically significant.

327 Further insight into this retention capacity can be evaluated from the comparison of the
328 concentration changes of these OC in children between birth and at the age of four years (Table 4).
329 These data show a general concentration increase in breastfed female children (all OC except
330 PeCB, HCB, 4,4'-DDT and PCB-180). In male children there is also a concentration increase but
331 with more exceptions (all OC except PeCB, HCB, PCB-28, 4,4'-DDE, 4,4'-DDT and PCB-180).
332 On the other hand, all OC except PeCB and HCB exhibit higher concentration increases (or lower

concentration decreases) in breastfed female than male children (Table 4). Formula fed children show a predominance of OC concentration decreases in both female and male children (Table 4).

Changes in body burden of OCs between birth and four years

Examination of the OC intakes between four years and birth in each individual by subtraction of the body burden of these compounds at four years and at birth confirms that breastfeeding is a major way of incorporation of these compounds in the first years of children growth (Table 5). Higher gender differentiation involving higher accumulation in female children is observed for all PCBs. These observations give more ground to a possible higher retention capacity of female children in the retention of OC incorporated through breastfeeding. Nevertheless, the high standard deviations resulting from these body burden calculations give rise to lack of statistical significance to the observed gender average differences.

Acknowledgements

M.C. Alvaro is thanked for her useful advice for the interpretation of results. This research was supported by the Instituto de Salud Carlos III. Red de Grupos INMA (G03/176) and projects PI041666, PI051987, ArcRisk (FP7-ENV-2008-1-226534) and GRACCIE Consolider Ingenio 2010 Project (CSD2007-00067).

References

Aguilar, A., 1984. Relationship of DDE/sigma DDT in marine mammals to the chronology of DDT input into the ecosystem. Can. J. Fish. Aquat. Sci. 41, 840-844.

- 359 AMAP, 2004. Arctic Monitoring and Assessment Programme. Oslo:AMAP Secretariat. Available:
360 <http://www.amap.no> [accessed 22 August 2009]
- 361 Carrizo, D., Grimalt, J.O., Ribas-Fito, N., Sunyer, J., Torrent, M., 2006. Physical-chemical and
362 maternal determinants of the accumulation of organochlorine compounds in four-year-old
363 children. Environ. Sci. Technol. 40, 1420-1426.
- 364 DeKoning, E.P., Karmaus, W., 2000. PCB exposure *in utero* and via breastmilk. A review. J. Expo.
365 Anal. Environ. 10, 285-293.
- 366 Fernandez, M.F., Sunyer, J., Grimalt, J.O., Rebagliato, M., Ballester, F., Ibarluzea, J., Ribas-Fito,
367 N., Tardon, A., Fernandez-Patier, R., Torrent, M., Olea, N., 2007. The Spanish Environment
368 and Childhood Research Network (INMA study). Int. J. Hyg. Env. Health 210, 491-493.
- 369 Grandjean, P., Weihe, P., White, R.F., Debes, F., Araki, S., Yokoyama, K., Marata, K., Sorensen,
370 N., Dahl, R., Jorgensen, P.J., 1997. Cognitive deficit in 7-year-old children with prenatal
371 exposure to methylmercury. Neurotoxicol. Teratol. 19, 417-428.
- 372 Huisman, M., Koopman-Esseboom, C., Fidler, V., Hadders-Algra, M., van der Paauw, CG.,
373 Tuinstra, L.G.M.T., Weisglas-Kuperus, N., Sauer, P.J.J., Touwen, B.C.L., Boersma, E.R.,
374 1995. Perinatal exposure to polychlorinated biphenyls and dioxins and its effect on neonatal
375 neurological development. Early Hum. Dev. 41, 111-127.
- 376 Jacobson, J.L., Humphrey, H.E.B., Jacobson, S.W., Schantz, S.L., Mullin, M.D., Welch, R., 1989.
377 Determinants of polychlorinated biphenyls (PCBs), polybrominated biphenyls (PBBs), and
378 dichlorodiphenyl trichloroethane (DDT) levels in the sera of young children. Am. J. Pub.
379 Health 79, 1401-1404.
- 380 Karmaus, W., deKoning, E.P., Kruse, H., Witten, J., Osius, N., 2001. Early childhood determinants
381 of organochlorine concentrations in school-aged children. Pediatric Res. 50, 331-336.

- 382 Lanting, C.I., Patandin, S., Fidler, V., Weisglas-Kuperus, N., Sauer, P.J.J., Boersma, E.R., Touwen,
383 B.C., 1998. Neurological condition in 42-month-old children in relation to pre- and
384 postnatal exposure to polychlorinated biphenyls and dioxins. *Early Hum. Dev.* 50, 283-292.
- 385 Lackmann, G.M., Goën, T., Töllner, U., Schaller, K.H., Angerer, J., 1996. PCBs and HCB in serum
386 of full-term German neonates. *Lancet* 348, 1035.
- 387 Polk, S., Sunyer, J., Munoz-Ortiz, L., Barnes, M., Torrent, M., Figueroa, C., Harris, J., Vall, O.,
388 Anto, J.M., Cullinan, P., 2004. A prospective study of Fel d1 and Der p1 exposure in
389 infancy and childhood wheezing. *Am. J. Respir. Crit. Care Med.* 170, 273-8.
- 390 Rhainds, M., Levallois, P., Ayotte, P., 1999. Lead, mercury, and organochlorine compound levels
391 in cord blood in Québec, Canada. *Arch. Environ. Health* 54, 40-47.
- 392 Ribas-Fito, N., Cardo, E., Sala, M., de Muga, E., Mazon, C., Verdu, A., Kogevinas, M., Grimalt,
393 J.O., Sunyer, J., 2003. Breastfeeding, exposure to organochlorine compounds, and
394 neurodevelopment in infants. *Pediatrics* 111, e580-e585
- 395 Ribas-Fito, N., Grimalt, J.O., Marco, E., Sala, M., Mazon, C., Sunyer, J., 2005. Breastfeeding and
396 concentrations of HCB and p,p'-DDE at the age of 1 year. *Environ. Res.* 98, 8-13.
- 397 Sala, M., Ribas-Fito, N., Cardo, E., deMuga, M.E., Marco, E. Mazon, C., Verdu, A., Grimalt, J.O.,
398 Sunyer, J., 2001. Levels of hexachlorobenzene and other organochlorine compounds in cord
399 blood: exposure across placenta. *Chemosphere* 43, 895-901.
- 400 Schwartz, P.M., Jacobson, S.W., Fein, G., Jacobson, J.L., Price, H.A., 1983. Lake Michigan fish
401 consumption as a source of polychlorinated biphenyls in human cord serum, maternal serum,
402 and milk. *Am. J. Pub. Health* 73, 283-296.
- 403 Skaare, J.U., Tuveng, J.M., Sande, H.A., 1988. Organochlorine pesticides and polychlorinated
404 biphenyls in maternal adipose tissue, blood, milk, and cord blood from mothers and their
405 infants living in Norway. *Arch. Environ. Contam. Toxicol.* 17, 55-63.

406 Stockholm Convention on Persistent Organic Pollutants, 2005. Available: <http://www.pops.int>
407 (accessed 22 August 2009).

408 Wedemeyer, G., 1967. Dechlorination of 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane by
409 *Aerobacter aerogenes*. I Metabolic products. Appl. Microbiol. 15, 569-574.

FIGURE CAPTIONS

Fig. 1. Average concentrations of organochlorine compounds in cord blood and sera collected at four years of female and male children. The intervals represent standard error ($p < 0.05$).

Fig. 2. Average concentrations and coefficients of variation of diverse gender and feeding practice groups of children included in the cohort of Menorca. BF: breastfeeding. Long BF > 3 months, short BF ≤ 3 months. F: female. M: male. As shown in the plots, the OC concentrations at four years of age are strongly influenced by breastfeeding and its duration at the early stages of life. Female accumulate larger concentrations of OC except HCB but the differences with respect to male are not statistically significant.