**Spatiotemporal patterns of seed dispersal in a wind-dispersed Mediterranean tree
(*Acer opalus* subsp. *granatense*): implications for regeneration**

LORENA GÓMEZ-APARICIO*, JOSE M. GÓMEZ and REGINO ZAMORA

Grupo de Ecología Terrestre, Dpto. Ecología, Facultad de Ciencias.

Universidad de Granada. E-18071 Granada, Spain.

* Correspondence: Lorena Gómez Aparicio (fax: 845 677-5976; email: gomezl@ecostudies.org).

Present address: Institute of Ecosystem Studies, 65 Sharon Turnpike, P.O. Box AB, Millbrook, New York, 12545-0129 USA

Summary

Seed dispersal can severely limit the quantity of plant recruits and their spatial distribution. However, our understanding of the role of dispersal in regeneration dynamics is limited by the lack of knowledge of seed deposition patterns in space and time. In this paper, we analyse the spatiotemporal variability of seed dispersal patterns in the Mediterranean maple, *Acer opalus* subsp. *granatense*, by monitoring seed rain along two years at a broad spatial scale (2 mountain ranges, 2 populations per range, 4 microhabitats per population). We quantified seed limitation and its components (source and dispersal limitation), and explored dispersal limitation in space by analysing dispersal distances, seed aggregation, and microhabitat seed distribution. *Acer opalus* subsp. *granatense* was strongly seed-limited throughout the gradients explored, being always dispersal limitation much higher than source limitation. The distribution of seeds with distance from adult individuals was leptokurtic and right-skewed in all populations, being both kurtosis and skewness higher the year of the highest seed production. Dispersal distances were shorter than expected by random in the four populations, which suggests distance-limited dispersal. Dispersal patterns were highly aggregated and showed a preferential direction around adults. At the microhabitat scale, most seeds accumulated under adult maples. However, there were no more seeds under trees and shrubs other than maple than in open interspaces, implying that established vegetation does not disrupt patterns of seed deposition by physically trapping seeds. When compared with patterns of seedling establishment, limited dispersal ability and inter-annual spatial concordance in seed rain patterns suggest that several potentially safe sites for recruitment have a very low probability of receiving seeds in most maple populations. These findings are especially relevant for rare species such as *Acer opalus* subsp. *granatense*, and illustrate how dispersal studies are not only crucial for our understanding of plant population dynamics but also to provide conservation directions.

1 Introduction

2 Seed dispersal plays a key role in the organization and dynamics of plant communities (Harper 1977,
3 Chambers and Macmahon 1994, Venable and Brown 1993, Schupp and Fuentes 1995). The spatial
4 pattern of dispersal will not only determine the distribution of a species, but also the environmental
5 variability encountered by its seedlings and saplings and thus, ultimately, the probability of
6 incorporation of new adults into the population (Gómez et al. 2004). Despite their relevance, the
7 patterns of seed deposition were for a long time scarcely included in dispersal studies under the
8 general assumption that seed dispersal does not limit recruitment. However, in 1995, Schupp and
9 Fuentes explicitly highlighted the relevance of the spatial relation of dispersed seeds both to seed
10 sources and among patches in the landscape. Since then, dispersal patterns have been analysed in
11 several demographic studies as crucial for understanding plant regeneration dynamics (e.g. Jordano
12 and Herrera 1995, Rey and Alcántara 2000, Gómez 2003, Hampe 2004).

13 Seed dispersal can limit regeneration if seeds are dispersed only to a fraction of suitable sites
14 for recruitment (Prymack and Miao 1992, Hurtt and Pacala 1995, Tilman 1997, Zobel et al. 2000). This
15 phenomenon, called seed limitation, can be due to a limited number of seeds available (source
16 limitation) or to low dispersal capacity (dispersal limitation; Clark et al. 1998). In addition, dispersal
17 limitation can be caused when most seeds are distributed at short distances from the parent tree
18 (limitation in distance), and/or in a spatially aggregated manner (limitation in space; Jordano and
19 Godoy 2002). Seed shadows of zoochorous species seem to be more aggregated than those of wind-
20 dispersed species, mainly due to non-random movements of dispersers (Schupp et al. 2002, Jordano
21 & Godoy 2002, Gómez 2003). However, anemochorous species can also show highly heterogeneous
22 seed rain for reasons such as the aggregation of seed sources in space (Muller-Landau et al. 2002) or
23 the occurrence of prevailing wind directions (Stewart et al. 1998, Bullock and Clarke 2000).

24 Microhabitat patchiness can also play a major role in generating heterogeneous patterns of seed
25 arrival, although it has frequently been ignored especially for wind-dispersed species. Given the
26 capacity of the vegetation to act as barriers to primary dispersal (Nathan et al. 2002), more seeds
27 should be expected to reach covered microhabitats than open spaces (Bullock and Moy 2004), which
28 in turn could have relevant demographic effects (e.g. seed-seedling conflicts, Schupp 1995) if
29 subsequent recruitment phases are also microhabitat-dependent (Nathan and Muller-Landau 2000).

30 Dispersal limitation can undergo substantial inter-annual variation (Houle 1998), mainly as a

consequence of differences in seed production. It has been shown that in years of high production maximum dispersal distances increase due to a greater probability for at least one seed landing farther than usual (Dirzo and Domínguez 1986) or because of a trade-off between seed size and number, resulting in the production of lighter seeds with greater flying capacity (Clark et al. 1998). In addition, seed aggregation can also be influenced by annual reproduction patterns, spatial homogeneity increasing with the number of reproductive individuals (Houle 1998).

Overall, due to the strong spatiotemporal variability in seed dispersal patterns, to know the extent to which they can limit regeneration requires the analysis of dispersal processes in different locations and years. However, such analyses have rarely been made (Clark et al. 1999, Nathan and Muller-Landau 2000, Hampe 2004). The main goal of this study is to analyse the spatiotemporal variability of seed dispersal patterns in the Mediterranean wind-dispersed tree, *Acer opalus* subsp. *granatense* (Boiss.) Font Quer & Rothm. We monitored abundance and spatial distribution of seeds over two years (2001 and 2002) and across a broad spatial scale (2 mountain ranges, 2 populations per range, 4 microhabitats per population). Specifically, our objectives were to: (i) quantify seed limitation and its components (source and dispersal limitation) (ii) explore dispersal limitation with distance and in space, and (iii) determine the spatiotemporal congruence of seed dispersal patterns and its implications for natural regeneration at the local scale. We consider that understanding the relative importance of source and dispersal limitations as well as the spatial patterns of seed dispersal relative to favourable microsites is essential to assess conservation status and planning restoration efforts for threatened or rare species such as *Acer opalus* subsp. *granatense*.

Material and methods

STUDY SPECIES

Mediterranean maple, *Acer opalus* subsp. *granatense* is an Iberian-Mauritanian endemic deciduous tree catalogued as Vulnerable by the IUCN (2000) and included in the Red List of Threatened Vascular Plants of Andalusia (Blanca et al. 2000). Maple has undergone reductions in number and size of populations in recent decades due to human disturbance (e.g. tree felling, deforestation, overgrazing; Blanca et al. 1998). Currently its distribution is composed of many small patches scattered throughout medium and high mountains (1100-2000 m a.s.l.) in the SE Iberian Peninsula, Balearic Islands and North Morocco (López-González 1994). Maple is a small tree, adults rarely surpassing 7-8

meters in height. Individuals of this species are usually intermingled with Mediterranean evergreens such as *Pinus sylvestris* L., *Pinus nigra* Arnold, or *Quercus ilex* L. Populations are sexually polymorphic, composed of 50% of protogynous trees, the other 50% comprising males and protandrous (Gabriela Gleiser, unpublished data). Since seed production in protandrous individuals is very low, about 50% of the maple adults can be considered to contribute to the seed pool in each population (seeding adults, hereafter). The indehiscent fruits are composed of two samaras with convergent wings, each weighting around 20-50 mg. Parthenocarpic development of fruits is rather common (de Jong 1976), especially in years of low seed production (Gómez-Aparicio unpublished data). Dispersal occurs between September and December.

STUDY AREAS

The study was conducted in two mountain ranges of SE Spain, Sierra Nevada (SN) and Sierra de Baza (SB), situated some 80 km apart. The climate in the zone is continental Mediterranean, with cold winters and hot, dry summers. Rainfall averages 846.5 ± 55.7 mm year⁻¹ in SN and 527.4 ± 40.9 mm year⁻¹ in SB (mean 1991-2002), with most of the precipitation falling in autumn and spring. Two maple populations were chosen per mountain range, one intermingled within a *Pinus sylvestris* forest (forest population [F], hereafter) and another on a stony slope (stony-slope population [SL], hereafter). Density of adult maples in these populations ranged between 17-111 individuals/m², mean adult height varied between 3.5-7.1 m, and mean seed mass varied between 20 and 27 mg (Table 1). We defined four microhabitats per population: (i) "Maple", under the canopy of adult maples; (ii) "Canopy", under the canopy of heterospecific adult trees and shrubs >1.5 m in height (1-2 species per population); (iii) "Shrub", under the canopy of shrubs <1.5 m in height (1-3 species per population); and (iv) "Open", in open interspaces between woody vegetation. Tree and shrub species were chosen according to their availability. The Canopy microhabitat included *Pinus sylvestris* L. and *Taxus baccata* L. in F-SN, *Amelanchier ovalis* Medik in SL-SN, and *P. sylvestris* in F-SB and SL-SB. The Shrub microhabitat included *Ononis aragonensis* Asso., *Juniperus communis* L. and *Berberis hispanica* Boiss. & Reuter in F-SN, *O. aragonensis* in SL-SN, *Prunus ramburii* Boiss. and *Crataegus monogyna* Jacq. in F-SB, and *J. communis* and *B. hispanica* in SL-SB.

SEED RAIN SAMPLING

Seed rain of *A. opalus* subsp. *granatense* was quantified during two consecutive years (2001 and 2002) by placing seed traps (0.045 m² aluminium trays) in a 1-ha study area per population. Traps were located randomly (and consequently without any reference adult), implying that sampling effort was not equitably distributed among distance ranges but, as strongly recommended by Willson (1993), proportional to their abundance in each population (median distance to the nearest seeding adult [Q25%-Q75%]: 8.5 m [4.8-12.7] in F-SN; 6.7 m [4.1-10.6] in SL-SN; 6.0 m [3.9-10.3] in F-SB; and 8.0 m [5.1-11.9] in SL-SB). Traps were placed at 30 points per microhabitat and population, each point containing 2 adjacent traps. In Canopy and Shrub microhabitats, sampling points were replicated for each species. The total number of traps was 420 in F-SN, 240 in SL-SN, 300 in F-SB and 300 in SL-SB (n=1260 traps). The distance distribution of traps did not differ between the Canopy, Shrub and Open microhabitats ($p>0.05$ in all cases, Kolmogorov-Smirnoff tests), implying that the different microhabitats were similarly distributed in relation to adult maples. Traps were covered with 1.3-cm diameter mesh in order to avoid post-dispersal predation. This mesh size has been shown to effectively restrict predators in seed predation experiments conducted in the four study sites (Gómez-Aparicio 2004). Traps were sampled fortnightly during the dispersal period (September-December), and their content transported to the laboratory, where seeds were individually weighed.

Since the number of seeds found in a trap depends both on its position in relation to the sources and on the amount of seed production, we calculated a seed availability index (SAI) for each trap (see Russell and Schupp 1998 for a similar approach). By using this index as a covariate, we can discern the role of the microhabitat in determining spatial patterns of dispersal regardless of its location. Thus, for each trap, we recorded the distance and orientation to the five nearest maples. Seed production for each adult was quantified in a semi-quantitative scale from 0 (no production) to 5 (very high production). SAI was calculated for each trap as:

$$SAI = \sum_{i=1}^5 \frac{p_i}{d_i}$$

where p is the seed production for each of the 5 nearest adult maples, and d its distance to the trap. This index is considered to represent the true relevance of being near maples, since factor 1 (distance to the nearest adult) explained 70% of the variability in the index, factor 2, 3, 4 and 5 explaining 19%,

7%, 3%, and 1%, respectively (averages for the four populations).

DATA ANALYSIS

Analysis of seed limitation and its components: source limitation and dispersal limitation

For each population, we quantified seed limitation and its components (source limitation and dispersal limitation) at the 0.09 m² spatial scale (pair of seed traps per sampling point). *Seed limitation* is defined as the proportion of empty traps over the time period: *source limitation* as the expected proportion of empty traps if seeds were uniformly distributed across space (uniform distribution defined stochastically as a random distribution); and *dispersal limitation* as the proportion of sampling points that received seeds in relation to sampling points that would receive them if seeds were randomly distributed (Clark et al. 1998, Nathan and Muller-Landau 2000). By using this approach, we assessed seed limitation in maple populations by describing spatial variation in natural seed rain patterns, an alternative to experimental manipulations. Therefore, seed limitation calculated in this way refers to “fundamental” and not “realized” seed limitation, assuming optimal conditions for seedling emergence. However, this method provides accurate information concerning the role of dispersal patterns in regeneration (Dalling et al. 2002, Muller-Landau et al. 2002), and enables the monitoring of a wider area, which is especially useful for studies covering several populations, as in the present study. Limitation measurements were calculated for every year (2001 and 2002), at the 1-year scale (average of both years) and at the 2-year scale (considering the total fallen seeds in both years). Limitation values range from 0 (no limitation) to 1 (maximum limitation).

Analysis of dispersal limitation in distance and space

The distribution of seed density with distance to the nearest seeding maple was described by 6 parameters: kurtosis, skewness, mean, median, mode and maximum. For each study year, differences among populations in the 6 parameters were analysed using Welch t-tests (Zar 1996). To explore any effect of distance on dispersal limitation, the density of seeds collected in the different distance classes during the 2 years of the study was compared to the density expected in each class if seeds were dispersed at random, the latter being approximated as the density of seed traps per distance interval. Since the nearest reproductive individual was assumed as the seed source, dispersal distances reported in this study are not “real” but “conservative” distances of dispersal, a

proportion of the seeds probably coming from a more distant adult.

Aggregation of seeds in space was analysed using Morisita's index of dispersion (Morisita 1959), calculated for each population at the sampling point (i.e. pair of traps) level. Deviations from random distributions were analysed by χ^2 tests with $q-1$ degrees of freedom (Greig-Smith 1983). We also determined the directionality of seeds around adult maples by using circular statistics (Oriana 2.0, Kovach Computing Services). Since the angular distribution of seeds depends partially on the position of traps, we first explored whether trap location follows a regular distribution in the circular space (or von Mises distribution) by using Rayleigh tests (Batschelet 1981). Values were not significant for any population ($p>0.05$ in the four cases), showing that traps were uniformly distributed around maple adults and that therefore sampling effort was not biased towards any specific direction around adults. Consequently, we also applied Rayleigh tests to analyse whether seeds were distributed uniformly around maple adults. Finally, Watson's F tests (Batschelet 1981) were used to compare the distribution of the seeds around maple adults in the two years of the study.

The role of the microhabitat in determining spatial patterns of dispersal was analysed by using Generalized Lineal Mixed Models (Proc MIXED, SAS 2002), using the maximum likelihood as estimation method (Stokes et al. 1995). For each population, total seed number recorded per sampling point was adjusted to a Poisson distribution, using Log as link function. Microhabitat was introduced as a fixed factor and year as a random. The seed availability index (SAI) was introduced as a continuous predictor. Differences among microhabitats were explored with *post-hoc* Bonferroni tests. To control for the experiment wise type I error produced by multiple comparisons, we adjusted the probabilities of error to $\alpha=0.05$ by using the sequential Bonferroni technique (Rice 1989).

Results

ANALYSIS OF SEED LIMITATION AND ITS COMPONENTS: SOURCE LIMITATION AND DISPERSAL LIMITATION

The four populations registered high values of seed limitation (0.63-1) in the two years of study (Table 2). Seed limitation decreased from 2001 to 2002 in all populations, matching an increase in seed density. However, this inter-annual difference affected mostly the shorter distance classes, as seed limitation in all cases was ≥ 0.90 at 6-8 meters from the nearest seeding maple (Fig. 1). Seed limitation was lower when calculated at the 2-year scale (0.60-0.80) than at the 1-year scale (0.74-0.90). The

magnitude of source limitation in all populations was much lower than the magnitude of dispersal limitation, especially at the 2-year scale, when seed limitation was due almost entirely to dispersal limitation (Table 2).

ANALYSIS OF DISPERSAL LIMITATION IN DISTANCE AND SPACE

The distribution of seeds with distance to the nearest seeding tree was leptokurtic and right-skewed in all populations, both leptokurtosis and skewness increasing in the second year of study (Fig. 2).

Although there were differences among populations and years, in all cases the mean ranged between 2-4 m, the median was less than 4 m, the mode was ~ 2 m, and the maximum observed distances ranged 7-12.5 m. The observed distribution of seeds per distance class significantly differed from the distribution expected if seeds were dispersed at random ($p < 0.05$ in the four populations, Kolmogorov-Smirnov tests), the observed median distances being in all cases much shorter than expected (3.5 vs 8.5 m in F-SN; 2.1 vs 6.7 m in SL-SN; 2.5 vs 6.0 m in F-SB; and 3.2 vs 8.0 m in SL-SB). Seed mass was not related to dispersal distance in any population ($p > 0.05$ in all cases, Spearman rank correlations).

Seeds showed a significant spatial aggregation in all populations and years (Table 3), which was generally higher in 2001. The density of seeds around adult maple trees differed significantly from uniformity in 6 of the 7 combinations of population and year explored ($p < 0.05$, Rayleigh tests; Table 3). There was no inter-annual difference in seed distribution around parent maples in any of the populations ($p > 0.05$, Watson tests).

Microhabitat identity did significantly affect seed density in all populations, even after introducing the seed availability index (SAI) as a continuous predictor (Table 4). In all populations and years, seed deposition was much higher under Maple than in any other microhabitat (67% of the seeds trapped). Shrub and Open showed similar values in all cases, whereas Canopy presented the lowest seed density (Fig. 3). The inter-annual variation in seed density was not the same for all microhabitats, as indicated by significant Microhabitat x Year interactions in the four populations (Table 4). Thus, although seed density was higher in 2002 than in 2001 in all microhabitats, the increase was comparatively higher under Maple and generally lower under Canopy.

Discussion

SEED LIMITATION, SOURCE LIMITATION AND DISPERSAL LIMITATION

Acer opalus subsp. *granatense* was strongly seed-limited throughout the spatiotemporal gradient explored. Seed limitation at the 1-year scale ranged 0.74-0.90 in all populations, despite their differences in two key characteristics influencing the values of the index, adult height (which indirectly influences seed production) and abundance (Muller-Landau *et al.* 2002). When compared with other populations throughout the range of the species in the Iberian Peninsula, our four populations covered the whole range of possibilities for both adult size and density (Gómez-Aparicio 2004). Therefore, seed limitation at the 1-year scale could be considered as a general pattern in maple populations across broad spatial scales. In addition, we did not detect any strong reduction in seed limitation calculated at the 2-year scale. Thus, although our temporal scale was not long enough to rule out possible limitation changes over the long term (see Muller-Landau *et al.* 2002), our results suggest that maple populations in Mediterranean mountains undergo high seed limitation at least over the short term, as reported for other tree species in tropical (Harms *et al.* 2000, Muller-Landau *et al.* 2002) and temperate forests (Clark *et al.* 1998).

The relative contribution of the two components of seed limitation (source and dispersal limitation, *sensu* Clark *et al.* 1998) showed a high spatiotemporal congruence, with source limitation being consistently much lower than dispersal limitation. Source limitation decreased in 2002 due to a higher seed production. Consequently, when calculated at the 2-year scale, seed limitation was almost entirely due to dispersal limitation, as in other *Acer* species (Clark *et al.* 1998).

DISPERSAL LIMITATION IN DISTANCE AND SPACE

Dispersal distances were shorter than expected according to a random distribution in the four populations, which suggests the existence of limitation in distance even at high densities of conspecific trees (>100 individuals/ha). In fact, 50% of the seeds dispersed were located less than 4 m from the nearest seeding tree, and no seeds were found beyond 13 m. Moreover, maximum dispersal distances did not differ in our two years, despite the variation in seed production and the fact that an increase in seed number is an important parameter for enlarging seed shadows (Dirzo and Domínguez 1986, Tanaka *et al.* 1998). Two main factors that generally affect dispersal in anemochorous species could constrain the dispersal distance of the species, source height and seed

mass (Augspurger and Franson 1987, Thiede and Augspurger 1996, Nathan et al. 2001). The small size of maple trees restricts the horizontal reach of seeds (Cousens and Rawlinson 2001, Nathan et al. 2002). In fact, differences of adult height in our populations likely influenced dispersal distances, since the two populations with the tallest maples (forest population in Sierra Nevada and stony-slope population in Sierra de Baza) presented maximum dispersal distances as much as 5 m longer (i.e., 30-40% of the overall dispersal distance) than the two populations including shorter adults. On the other hand, the mass of maple seeds ranged between 20 and 40 mg, being much heavier than seeds of most co-occurring wind-dispersed trees such as *Pinus nigra* or *Pinus sylvestris* (e.g. 5-12 mg for *P. sylvestris* in Castro 1999). Moreover, the range of mass variation of maple seeds did not appear to be big enough to influence dispersal distances, since lighter seeds did not travel farther (see also Augspurger and Franson 1987).

Morisita indices showed aggregated dispersal patterns in all four maple populations. This high heterogeneity in seed shadow may be a consequence of a large fraction of seeds falling near parent trees (Venable and Brown 1993), but also implies that those that travelled farther were deposited in clumps. The formation of such clumps could be influenced by directional dispersal related to local wind regimes, because most seed concentrated in specific orientations in relation to adults. Moreover, more than 50% of the traps that received seeds in 2001 also did it in 2002, suggesting a high degree of inter-annual concordance in seed rain patterns ($p < 0.05$ in the four populations, Kendall correlations), a result that in turn has being rarely reported for wind-dispersed species (Houle 1998, Nathan et al. 2000, Jones et al. 2005).

Seed deposition patterns were significantly affected by microhabitat. However, this influence was restricted basically to Maple, which consistently showed much higher seed density than any other microhabitat. Contrary to our expectations of higher seed arrival to covered microhabitats (Canopy and Shrubs) than to the Open microhabitat, the magnitude of seed dispersal did not differ among the three microhabitats. These findings agree with those reported by Russell & Schupp (1998) for *Cercocarpus ledifolius*, and Castro et al. (1999) for *Pinus sylvestris*, but differ with several reports that reached the opposite conclusion (Fuentes et al. 1984, McEvoy and Cox 1987, Thiede and Augspurger 1996, Bullock and Moy 2004). Therefore, seed trapping seems to be highly dependent on the species and system considered.

IMPLICATIONS FOR REGENERATION

A direct consequence of limited dispersal ability of maple seeds is that they are unable to explore larger fractions of the landscape. Dispersal kernels were leptokurtic and right-skewed, especially in the year of highest seed production, seed limitation being above 0.9 just at around 7 m from the nearest seeding adult maple in all populations and years. Maple early recruitment has been reported to be highly dependent on the microhabitat, seedling and sapling survival being much higher under pre-established woody vegetation than in open interspaces (Gómez-Aparicio *et al.* 2005). Thus, the fact that the 50% of the seed traps randomly located under trees and shrubs were beyond 7 m from any adult maple implies that several suitable microhabitats for regeneration have low probabilities of receiving seeds. Furthermore, this pattern seems to be temporally stable, since the 2002 increase in seed availability decreased seed limitation but only at short distances (<6-8 meters). This finding suggests that, in the long term, seed limitation could be expected to reach a minimum once seeds have saturated the space near maples whereas many safe sites remain empty because they are too far away (Schupp *et al.* 2002).

The fact that seed deposition was highly aggregated and spatially concordant over the years implies that even near maples seeds will land in the same microsites every year, reducing the number of sites colonized. Secondary seed dispersal could attenuate this aggregation by redistributing seeds, but it does not appear to substantially alter primary dispersal patterns, since emerging seedlings are highly aggregated around adults (Gómez-Aparicio 2004). Aggregated patterns of seed deposition could cascade through later demographic stages giving rise to intra-specific competition (Levin *et al.* 1984, Augspurger and Kitajima 1992, Hurtt and Pacala 1995), not only between individuals of the same cohort, but also of different cohorts.

The results presented here show that the variation in maple seed deposition patterns did not preclude seed and dispersal limitation throughout the spatiotemporal scope of the study. This is especially relevant for rare species such as *Acer opalus* subsp. *granatense*, since seed limitation slows down rates of abundance change (Muller-Landau *et al.* 2002), presumably hampering the recovery of its past distribution. In these scenarios, protection of remaining reproductive individuals and the introduction of propagules in distant, high-quality, unoccupied microhabitats could counterbalance the negative consequences of restricted dispersal patterns for maple regeneration. This way, seed dispersal analyses become not only crucial for understanding plant population

dynamics, but also for the conservation and restoration of natural populations of species that suffer from seed limitation (Primack and Miao 1992, Eriksson and Ehrlén 1992, Turnbull *et al.* 2000, Bullock *et al.* 2002, Makana and Thomas 2004).

Acknowledgements

We thank the Consejería de Medio Ambiente, Junta de Andalucía, for permitting the field work in Sierra Nevada and Sierra de Baza. Sergio de Haro provided invaluable help in the field. Corinna Riginos and an anonymous reviewer made constructive comments on the manuscript. David Nesbitt improved the English version. Financial support was provided by a PFPU-MECD grant to L.G.A., the Andalusian government grant (PAI) rnm-220, and projects REN2001-4552-E and HETEROMED REN2002-04041-CO2-01/GLO from the Spanish Ministry of Science and Technology (MCYT).

References

- Augsburger, C. K. and Franson, S. E. 1987. Wind dispersal of artificial fruits varying in mass, area, and morphology. - *Ecology* 68: 27-42.
- Augsburger, C. K. and Kitajima, K. 1992. Experimental studies of seedling recruitment from contrasting seed distributions. - *Ecology* 73: 1270-1284.
- Batschelet, E. 1981. Circular statistics in biology. - Academic Press, New York.
- Blanca, G. *et al.* 1998. Threatened vascular flora of Sierra Nevada (Southern Spain). - *Biol. Conserv.* 85: 269-285.
- Blanca, G. *et al.* 2000. Libro Rojo de Flora Amenazada de Andalucía. - Consejería de Medio Ambiente, Junta de Andalucía, Sevilla.
- Bullock, J. M. and Clarke R. T. 2000. Long distance seed dispersal by wind: measuring and modeling the tail of the curve. - *Oecologia* 124: 506-521.
- Bullock, J. M. and Moy, I. L. 2004. Plant as seed traps: inter-specific interference with dispersal. - *Acta Oecol.* 25: 35-41.
- Bullock, J. M. *et al.* 2002. Plant dispersal and colonization processes at local and landscape scales. In: Bullock, J. M., Kenward, R. E. and Hails, R. (eds.), *Dispersal Ecology*. -Blackwell, pp. 173-194.
- Castro, J. 1999. Seed mass *versus* seedling performance in Scots pine: a maternally dependent trait. - *New Phytol.* 144: 153-161.

- 1 Castro, J. et al. 1999. Seed predation and dispersal in relict Scots pine forests in southern Spain. -
2 Plant Ecology 145: 115-123.
- 3 Chambers, J. C. and MacMahon, J. A. 1994. A day in the life of a seed: movements and fates of
4 seeds and their implications for natural and managed systems. - Annu. Rev. Ecol. Syst. 25: 263-
5 292.
- 6 Clark, J. S. et al. 1998. Stages and spatial scales of recruitment in southern Appalachian forests. -
7 Ecol. Monogr. 68: 213-235.
- 8 Clark, J. S. et al. 1999. Interpreting recruitment limitation in forests. - Am. J. Bot. 86: 1-16.
- 9 Cousens, R. D. and Rawlinson, A. A. 2001. When will plant morphology affect the shape of a seed
10 dispersal kernel? - J. Theor. Biol. 211: 229-238.
- 11 Dalling, J. W. et al. 2002. Role of dispersal in the recruitment limitation of neotropical pioneer species.
12 - J. Ecol. 90: 714-727.
- 13 de Jong, P. C. 1976. Flowering and Sex Expression in *Acer* L. A biosystematic study. - Mededelingen
14 Landbouwhogeschool, Wageningen. The Netherlands.
- 15 Dirzo, R. and Domínguez, C. A. 1986. Seed shadows, seed predation and the advantages of
16 dispersal. - In: Estrada, A. and Fleming, T. (eds.), Frugivores and Seed Dispersal. Dr. W. Junk,
17 Dordrecht, the Netherlands, pp. 237-250.
- 18 Eriksson, O. and Ehrlén, J. 1992. Seed and microsite limitation of recruitment in plant populations. -
19 Oecologia 91: 360-364.
- 20 Fuentes, E. R. et al. 1984. Shrub clumps of the Chilean matorral vegetation: structure and possible
21 maintenance mechanisms. - Oecologia 62: 405-411.
- 22 Gómez, J. M. 2003. Spatial patterns in long-distance dispersal of *Quercus ilex* acorns by jays in a
23 heterogeneous landscape. - Ecography 26: 573-584.
- 24 Gómez, J. M. et al. 2004. Differences between structural and functional environmental heterogeneity
25 caused by seed dispersal. - Funct. Ecol. 18: 787-792.
- 26 Gómez-Aparicio, L. 2004. *Papel de la heterogeneidad en la regeneración del Acer opalus subsp.*
27 *granatense en la montaña mediterránea: implicaciones para la gestión y conservación de sus*
28 *poblaciones*. PhD thesis, University of Granada, Granada.
- 29 Gómez-Aparicio, L. et al. 2005. Microhabitats shift rank in suitability for seedling establishment
30 depending on habitat type and climate. - J. Ecol. 93: 1194-1202.

- 1 Greig-Smith, P. 1983. Quantitative Plant Ecology. 3rd Ed. - University of California Press, Berkeley,
- 2 CA.
- 3 Hampe, A. 2004. Extensive hydrochory uncouples spatiotemporal patterns of seedfall and seedling
- 4 recruitment in a “bird-dispersed” riparian tree. - J. Ecol. 92: 797-807.
- 5 Harms, K. E. et al. 2000. Pervasive density- dependent recruitment enhances seedling diversity in a
- 6 tropical forest. - Nature 404: 493-495.
- 7 Harper, J. L. 1977. Population Biology of Plants. - Academic Press, London.
- 8 Houle, G. 1998. Seed dispersal and seedling recruitment of *Betula alleghaniensis*: spatial
- 9 inconsistency in time. - Ecology 79: 807-818.
- 10 Hurtt, G. C. and Pacala, S. W. 1995. The consequences of recruitment limitation: reconciling chance,
- 11 history and competitive differences between plants. - J. Theor. Ecol. 176: 1-12.
- 12 IUCN. 2000. *IUCN Red List Categories*. IUCN, Gland.
- 13 Jones, F. A. et al. 2005. A genetic evaluation of seed dispersal in the neotropical tree *Jacaranda*
- 14 *copaia* (Bignoniaceae). - Am. Nat. 166: 543-555.
- 15 Jordano, P. and Herrera, C. M. 1995. Shuffling the offspring: uncoupling and spatial discordance of
- 16 multiple stages in vertebrate seed dispersal. - Ecoscience 2: 230-237.
- 17 Jordano, P. and Godoy, J. A. 2002. Frugivore-generated seed shadows: a landscape view of
- 18 demographic and genetic effects. - In: Levey, D. J., Silva, W. R. and Galetti, M. (eds.), Seed
- 19 dispersal and Frugivory: Ecology, Evolution and Conservation. CAB International, pp. 305-321.
- 20 Levin, S. A. et al. 1984. Dispersal in patchy environments. - Theor. Pop. Biol. 26: 165-191.
- 21 López-González, G. 1994. La guía de Incafo de los árboles y arbustos de la Península Ibérica. -
- 22 Editorial Incafo, Madrid.
- 23 Makana, J. -R. and Thomas, S. C. 2004. Dispersal limits natural recruitment of African mahoganies. -
- 24 Oikos 106: 67-72.
- 25 McEvoy, P. B. and Cox, C. S. 1987. Wind dispersal distances in dimorphic achenes of ragwort,
- 26 *Senecio jacobaea*. - Ecology 68: 2006-2015.
- 27 Morisita, M. 1959. Measuring of the dispersion of individuals and analysis of the distributional patterns.
- 28 - Mem. Fac. Sci. Kyushu Univ. 2: 215-235.

- 1 Muller-Landau, H. C. et al. 2002. Assessing recruitment limitation: concepts, methods and case-
2 studies from a tropical forest. - In: Levey, D. J., Silva, W. R. and Galetti, M. (eds.), Seed Dispersal
3 and Frugivory: Ecology, Evolution and Conservation. CAB International, pp. 35-54.
- 4 Nathan, R. and Muller-Landau, H. C. 2000. Spatial patterns of seed dispersal, their determinants and
5 consequences for recruitment. - Trends Ecol. Evol. 15: 278-285.
- 6 Nathan, R. et al. 2000. Spatiotemporal variation in seed dispersal and recruitment near and far from
7 *Pinus halepensis* trees. - Ecology 81: 2156-2169.
- 8 Nathan, R. et al. 2001. Field validation and sensitivity analysis of a mechanistic model for tree seed
9 dispersal by wind. - Ecology 82: 374-388.
- 10 Nathan, R. et al. 2002. Mechanistic models for tree seed dispersal by wind in dense forests and open
11 landscapes. - In: Levey, D. J., Silva, W. R. and Galetti, M. (eds.), Seed Dispersal and Frugivory:
12 Ecology, Evolution and Conservation. CAB International, pp. 69-82.
- 13 Primack, R. B. and Miao, S. L. 1992. Dispersal can limit local plant distribution. - Conserv. Biol. 6: 513-
14 519.
- 15 Rey, P. J. and Alcántara, J. M. 2000. Recruitment dynamics of a fleshy-fruited plant (*Olea europaea*):
16 connecting patterns of seed dispersal to seedling establishment. - J. Ecol. 88: 622-633.
- 17 Rice, W. R. 1989. Analyzing tables of statistical tests. - Evolution 43: 223-225.
- 18 Russell, S. K. and Schupp, E. W. 1998. Effects of microhabitat patchiness on patterns of seed
19 dispersal and seed predation of *Cercocarpus ledifolius* (Rosaceae). - Oikos 81: 434-443.
- 20 SAS. 2002. *SAS/STAT user's guide*. Version 8.2. SAS Institute, Cary, NC.
- 21 Schupp, E. W. 1995. Seed-seedling conflicts, habitat choice, and patterns of plant recruitment. -
22 American Journal of Botany 82: 399-409.
- 23 Schupp, E. W. and Fuentes, M. 1995. Spatial patterns of seed dispersal and the unification of plant
24 population ecology. - Ecoscience 2: 267-275.
- 25 Schupp, E. W. et al. 2002. Dissemination limitation and the origin and maintenance of species-rich
26 tropical forests. - In: Levey, D. J., Silva, W. R. and Galetti, M. (eds.), Seed Dispersal and Frugivory:
27 Ecology, Evolution and Conservation. CAB International, pp. 19-34.
- 28 Stewart, J. D. et al. 1998. Dispersal of white spruce in mature aspen stands. - Can. J. Bot. 76: 181-
29 188.
- 30 Stokes, M. E. et al. 1995. Categorical data analysis using the SAS System. - SAS Institute, Cary, NC.

- 1 Tanaka, H. et al. 1998. A mechanistic approach for evaluation the role of wind dispersal in tree
2 population dynamics. - J. Sustainable Forestry 6: 155-174.
- 3 Thiede, D. A. and Augspurger, C. K. 1996. Intraspecific variation in seed dispersion of *Lepidium*
4 *campestre* (Brassicaceae). - Am. J. Bot. 83: 856-866.
- 5 Tilman, D. 1997. Community invasibility, recruitment limitation, and grassland biodiversity. - Ecology
6 78: 81-92.
- 7 Turnbull, L. A. et al. 2000. Are plants seed-limited? A review of seed sowing experiments. - Oikos 88:
8 225-238.
- 9 Venable, D. L. and Brown, J. S. 1993. The population-dynamic functions of seed dispersal. - Vegetatio
10 107/108: 31-55.
- 11 Willson, M. F. 1993. Dispersal mode, seed shadows, and colonization patterns. - Vegetatio 107/108:
12 261-280.
- 13 Zar, J. H. 1996. Biostatistical Analysis. 3rd Ed. - Prentice Hall, Englewood Cliffs, N.J.
- 14 Zobel, M. et al. 2000. Is small-scale species richness limited by seed availability or microsite
15 availability? - Ecology 81: 3274-3282.

Table 1 Summary of the main characteristics of the four study populations of *Acer opalus* subsp. *granatense*. Adult density and height were sampled using 25x10 m transects (n=10) haphazardly distributed in a representative area of each population 1-ha in size. Seed mass was calculated from seeds collected in traps during Fall 2002. Mean $\pm$ 1SE [minimum-maximum] are shown.

	Site			
	Sierra Nevada		Sierra de Baza	
	Forest	Stony-slope	Forest	Stony-slope
<i>Location (UTM)</i>	30SVG5905	30SVG5904	30SWG1438	30SWG1433
<i>Altitude (m a.s.l.)</i>	1850	1920	1850	2000
<i>Orientation</i>	NW	NW	NW	NE
<i>Slope (°)</i>	40	40	30	35
<i>Adult density (n°/ha)</i>	17.2 $\pm$ 6.5	80.2 $\pm$ 12.2	111.5 $\pm$ 18.8	61.1 $\pm$ 10.9
<i>Adult height (m)</i>	7.1 $\pm$ 0.3	4.5 $\pm$ 0.2	3.5 $\pm$ 0.2	5.5 $\pm$ 0.3
	[6.0-8.0]	[2.5-6.0]	[2.0-4.5]	[3.0-8.0]
<i>Seed mass (mg)</i>	22.1 $\pm$ 0.4	20.2 $\pm$ 0.9	21.7 $\pm$ 1.1	26.9 $\pm$ 0.6
	[20.4-35.8]	[19.1-44.8]	[18.9-35.3]	[23.9-39.6]

Table 2 Seed density (seeds/m²; mean $\pm$ 1SE) and limitation measurements at the sampling point spatial scale (0.09 m²) for the four study populations of *Acer opalus* subsp. *granatense*. Values were calculated for each year independently, as well as at the 1-year (mean $\pm$ 1SE of the 2 years) and 2-year (sum of the 2 years) scale. Limitation values were between 0 (no limitation) and 1 (maximum limitation).

Population	Measure	Time scale			
		2001	2002	1-year	2-year
Sierra Nevada					
Forest	Seed density	0	34.18 ± 10.95	17.1 ± 5.51	34.07±10.92
	Seed limitation	1	0.80	0.90 ± 0.10	0.80
	Source limitation	1	0.04	0.52 ± 0.48	0.04
	Dispersal limitation	‡	0.80	‡	0.80
Stony-slope	Seed density	8.11 ± 4.42	22.42 ± 6.81	15.2 ± 4.12	30.46±9.23
	Seed limitation	0.91	0.69	0.80 ± 0.11	0.67
	Source limitation	0.48	0.13	0.30 ± 0.17	0.06
	Dispersal limitation	0.82	0.64	0.73 ± 0.09	0.65
Sierra de Baza					
Forest	Seed density	11.32 ± 4.21	15.57 ± 3.82	13.4 ± 2.85	28.59±7.99
	Seed limitation	0.79	0.71	0.75 ± 0.04	0.64
	Source limitation	0.33	0.27	0.30 ± 0.03	0.01
	Dispersal limitation	0.68	0.63	0.65 ± 0.02	0.63
Stony-slope	Seed density	10.35 ± 4.93	33.36 ± 8.33	21.8 ± 4.94	43.63±11.45
	Seed limitation	0.85	0.63	0.74 ± 0.11	0.60
	Source limitation	0.39	0.05	0.22 ± 0.17	0.02
	Dispersal limitation	0.74	0.60	0.67 ± 0.07	0.59

‡/Note: Since no seeds were produced in the forest population of Sierra Nevada in 2001, seed and source limitation were given the maximum value (1), and thereby dispersal limitation could not be calculated.

Table 3 Description of seed aggregation patterns in the two study years. Values of the Morisita's index of dispersion significantly higher than 0 indicate aggregated patterns, whereas values non-significantly different from 0 indicate a regular distribution. Mean angle (in degrees) represents the prevailing dispersal direction around adult maples. Concentration represents the deviation of the seed distribution around adult maples from a uniform distribution in the circular space (or von Mises distribution), and was evaluated with Rayleigh tests.

Year	Population	Morisita index	Angular distribution	
			Mean angle (± 1 SE)	Concentration
2001	Sierra Nevada			
	Stony-slope	35.51****	216.25 $\pm$ 4.28	5.12****
	Sierra de Baza			
	Forest	18.10****	359.92 $\pm$ 10.01	1.05****
	Stony-slope	34.73****	274.82 $\pm$ 20.38	1.02*
2002	Sierra Nevada			
	Forest	22.17****	327.13 $\pm$ 3.90	1.58****
	Stony-slope	11.69****	223.39 $\pm$ 20.95	0.59*
	Sierra de Baza			
	Forest	10.74****	54.56 $\pm$ 18.12	0.51**
	Stony-slope	9.99****	226.91 $\pm$ 80.18	0.08

**** $P < 0.0001$, *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$

Table 4 Summary of the Generalized Lineal Mixed Model analysing differences among microhabitats in the four populations and during the two years of study. SAI (seed availability index) was introduced as a continuous predictor.

Population	Factor	df	L-R χ^2	P
Sierra Nevada				
Forest	Microhabitat	3	21.38	<0.0001
	Year	1	96.25	<0.0001
	Microhabitat x Year	3	20.09	0.0002
	SAI	1	1177.64	<0.0001
Stony-slope	Microhabitat	3	23.98	<0.0001
	Year	1	6.34	0.012
	Microhabitat x Year	3	8.74	0.033
	SAI	1	325.53	<0.0001
Sierra de Baza				
Forest	Microhabitat	3	11.45	0.009
	Year	1	0.99	0.326
	Microhabitat x Year	3	9.56	0.023
	SAI	1	314.26	<0.0001
Stony-slope	Microhabitat	3	17.47	0.0005
	Year	1	19.99	<0.0001
	Microhabitat x Year	3	37.98	<0.0001
	SAI	1	556.20	<0.0001

Figure captions

Fig. 1 Variation in seed limitation with distance (in meters) to the canopy edge of the nearest seeding maple for each population and year of study. Seed limitation is calculated as the proportion of sampling points not receiving seeds in the time period. Sampling points were classified in 18 distance classes (20 for the forest population in Sierra Nevada) of 1 meter. The grey line indicates the position of the value 0.9 of seed limitation. F, forest population; SL, stony-slope population; SN, Sierra Nevada; SB, Sierra de Baza.

Fig. 2 Average seed density (mean $\pm$ 1SE) as a function of distance (in meters) from the canopy edge of the nearest seeding adult in a) 2001 and b) 2002. For each population and year, characteristics of the shape and central tendencies of the dispersal distributions are given (K, kurtosis; S, skewness; Mn, mean; Md, mode, Md, median, Mx, maxima). Different letters show statistical differences among populations (within the same year) after Welch t-tests. No seeds were produced in the forest population of Sierra Nevada in 2001. Letter codes same as Fig. 1.

Fig. 3 Differences between microhabitats in seed density (mean $\pm$ 1SE) in the two years of study. Different letters show significant differences among microhabitats within populations at $\alpha < 0.05$ (after Bonferroni correction). Letter codes same as Fig. 1. No seeds were produced in the forest population of Sierra Nevada in 2001.

Fig. 1

Fig. 2

Fig. 3