EXTENDED PARENTAL CARE IN THE FRESHWATER SHRIMP GENUS
DUGASTELLA BOUVIER, 1912 (DECAPODA, ATYIDAE, PARATYINAE)

BY

D. HUGUET^{1,4}), J. E. GARCÍA MUÑOZ²), J. E. GARCÍA RASO²) and J. A. CUESTA^{3,5})

¹) Département Milieux et Peuplements aquatiques, Muséum national d'Histoire
naturelle, 61 rue Buffon, CP 53, F-75253 Paris cedex 05, France

²) Departamento de Biología Animal, Facultad de Ciencias, Universidad de Málaga,
Campus de Teatinos s/n, E-29071 Malaga, Spain

³) Instituto de Ciencias Marinas de Andalucía (CSIC), Avenida República Saharaui, 2,
11519 Puerto Real, Cadiz, Spain

⁴) e-mail: denise.hughet9@orange.fr

⁵) e-mail: jose.cuesta@icman.csic.es

Parental care and extended parental care are known behaviours in decapod crustaceans (see Thiel, 2000, 2003). While parental care before hatching, which involves incubation of the eggs (carrying, cleaning, and aeration) by females, is a common behaviour in the majority of decapod taxa (with the exception of Dendrobranchiata), extended parental care is a rare behaviour in decapods, restricted only to some species of brachyuran crabs (Diesel, 1989, 1992, 1997; Diesel & Horst, 1995; Ng & Tan, 1995), hermit crabs (Hazlett, 1983; López Greco et al. 2004; Calado et al., 2006), astacid crayfishes (Johnston & Fiegel, 1997; Vogt & Tolley, 2004), and caridean shrimps (Duffy, 1996). In the majority of these cases, the parental female presents different types of parental activity towards the offspring after hatching, like habitat control, elimination of predators, or sharing meals. Also, there are some cases in which larvae develop transient structures that allow them to remain fixed to the pleopods of the female abdomen, like in astacids (see Vogt & Tolley, 2004) and several species of the genus Sclerocrangon Sars, 1883 (see Makarov, 1968; Lacoursière-Roussel & Sainte-Marie, 2009). However, the most infrequent cases of extended parental care in decapods are those where females carry their juvenile offspring under their abdomen after the hatched larvae develop there (cf. abdominal brood pouches, or incubation chamber). These cases have only been reported in one species of Brachyura, Tunicotheres moseri (Rathbun, 1918) (see Bolaños et al., 2004), and one caridean, the kangaroo shrimp Dugastella valentina (Ferrer Galdiano, 1924) (see Cuesta et al. 2006). Taking into account this last case, and considering that there are only two species in the genus Dugastella Bouvier, 1912, the objective of the present study was to corroborate whether the other species, D. marocana Bouvier, 1912, also presents the same behaviour. These two species have a restricted distribution, D. valentina occurs in the Gulf of Valencia (southern Europe) and D. marocana is found in some localities of Morocco (northwestern Africa) (see fig. 1).

Females of Dugastella marocana with few, large eggs have been observed by the first author in 1975 but those observations were not published at that time (lecitotrophic eggs about 1.0 mm in length, 0.60 mm width; number of eggs varying from 16 to 24). D. marocana was only known from its topotypic population of Settât (Morocco) (see fig. 1). After its description by Bouvier (1912) there have no more data been reported on its distribution, biology, or ecology until recently. For this reason, the objective of the present work was to collect ovigerous females of this species and study larval development. Different field sampling campaigns were carried out in Morocco, with unequal success. Those samplings in Settât show that, due to the transformation of this area through urbanization, the species is not longer present there. The last specimens, which were collected in the vicinity of Settât (at about 35 km distance) by the first author, in June 1975 at three stations, represent the last available samples from the proximity of the topotypic locality.

The locality data of the material collected by the first author are as follows: (1) in the Kihane River next to Aïn-Belmesk, not very far from the Oum er Rbia River; (2) in a separate spring (resurgence) next to Aïn-Belmesk; (3) in the As Sila River, N.W. of Aïn-Belmesk (see fig. 1). This region was facing a great drought during the last years, so a recent expedition (20 May 2007) to the same localities did not find any shrimps. Some specimens were collected together with individuals of Atyaephyra desmarestii (Millet, 1831) near the Oum er Rbia River (at Bzou, N.E. of Marrakech, Morocco) by J.E. Garcia Muñoz and J.E. Garcia Raso on 24 October 2007 (Garcia Muñoz et al., 2009).

According to mitochondrial gene data (genes 16S and COI) Dugastella marocana is a valid species, well separated from D. valentina (cf. Garcia Muñoz et al., 2009). Unfortunately, there were no ovigerous females in the sample from Bzou, but a female was observed with larvae (zoea II) in its incubation chamber, demonstrating that in D.

marocana, as in D. valentina, females bore larvae in their abdominal brood pouches. An extended parental care behaviour is thus presented here as a common feature of the species in the genus Dugastella. The same observation was made earlier by the first author, who has found females with larvae attached in the population captured in the spring next to Aïn-Belmesk.

Later, a revision of the material deposited at the Muséum national d'Histoire naturelle, Paris, allowed us to locate among the material collected by M. Pallary in 1914 in the spring of Settât, topotypic locality, a female specimen of D. marocana with larvae (zoea I and II) inside its incubation chamber. This confirms the breeding mechanism observed in the specimen collected in the proximity of the Oum er Rbia River (at Bzou) and in that from the spring next to Aïn-Belmesk.

Species of Atyidae have been described with a wide range of types of larval development, ranging from direct development to long, planktonic larval development (10-12 zoeal stages) (see Walsh, 1993). Until now, however, the two species of the genus Dugastella are the only ones showing this peculiar extended parental care. Probably re-examination of ovigerous females of other atyid species, i.e., those with reported direct development, might reveal other cases of similar extended parental care, since without detailed observation of the abdominal enclosure, one cannot confirm the presence of larval stages instead of eggs. A female releasing juveniles could constitute evidence for direct development, without larval stages at all, or actually could be showing a true case of extended parental care.

ACKNOWLEDGEMENTS

The first author is very grateful to the Major of Settat who had authorized to collect shrimp samples in the area of this town in 1975. The other authors would like to express their gratitude to Mohamed Ghamizi (Muséum d'Histoire Naturelle de Marrakech, Université Cadi Ayyad) for his help during sample collecting in Morocco. Collections in Morocco by JEGM and JEGR were possible thanks to fundings provided by “CSIC Proyectos Intramurales Especiales” 2006 3 01 238 to JAC. We express our thanks to Bernard Sainte-Marie for calling our attention to the cases of extended parental care in Sclerocrangon. Thanks are also due to Christoph D. Schubart and Martin Thiel for their valuable revision of a first version of this manuscript, two anonymous referees for their comments and corrections and Carel von Vaupel Klein for his editorial effort. The first author thanks Danielle Defaye and Danièle Guinot (Muséum national d'Histoire naturelle, Département Milieux et Peuplements aquatiques) for their encouragement and help.

REFERENCES

- BOLAÑOS, J., J. A. CUESTA, G. HERNÁNDEZ, J. HERNÁNDEZ & D. L. FELDER, 2004. Abbreviated larval development of Tunicotheres moseri (Rathbun, 1918) (Decapoda: Pinnotheridae), a rare case of parental care among brachyuran crabs. *Scientia Marina*, **68**: 373-384.
- BOUVIER, E.L., 1912. Dugastella marocana, crevette primitive nouvelle de la famille des Atyidés. *Comptes Rendus de l'Academie des Sciences Paris*, **155**: 993-998.
- CALADO, R., N. NOGUEIRA & A. dos SANTOS, 2006. Extended parental care in a hermit crab of the genus Calcinus (Anomura: Diogenidae). *Journal of the Marine Biological Association of United Kingdom*, **86**: 121-123.

- CUESTA, J. A., E. PALACIOS-THEIL, P. DRAKE & A. RODRÍGUEZ, 2006. A new rare case of parental care in decapods. *Crustaceana*, **79**: 1401-1405.
- DIESEL, R., 1989. Parental care in an unusual environment: Metopaulias depressus (Decapoda: Grapsidae), a crab that lives in epiphytic bromeliads. *Animal Behaviour*, **38**: 561-575.
- , 1992. Managing the offspring environment: brood care in the bromeliad crab, Metopaulias depressus. *Behavioral Ecology and Sociobiology*, **30**: 125-134.
- , 1997. Maternal control of calcium concentration in the larval nursery of the bromeliad crab, Metopaulias depressus (Grapsidae). *Proceeding of the Royal Society of London, (B)* **264**: 1403-1406.
- DIESEL, R. & D. HORST, 1995. Breeding in a snail shell: ecology and biology of the Jamaican montane crab Sesarma jarvisi (Decapoda: Grapsidae). *Journal of Crustacean Biology*, **15**: 179-195.
- DUFFY, J. E., 1996. Eusociality in a coral-reef shrimp. *Nature, London*, **381**: 512-514.
- GARCÍA MUÑOZ, J. E., A. RODRÍGUEZ, J. E. GARCÍA RASO & J. A. CUESTA, 2009. Genetic evidence for cryptic speciation in the freshwater shrimp genus Atyaephyra de Brito Capello (Crustacea, Decapoda, Atyidae). *Zootaxa*, **2025**: 32-42.
- HAZLETT, B. A., 1983. Parental behavior in decapod crustacean. In: S. Rebach & D. W. Dunham (eds.), *Studies in adaptation - the behavior of higher Crustacea*: 171-193. (John Wiley and Sons, Inc., New York).
- JOHNSTON, C. E. & C. FIEGEL, 1997. Microhabitat parameters and life-history characteristics of Fallicambarus gordonii Fitzpatrick, a crayfish associated with pitcher-plant bogs in southern Mississippi. *Journal of Crustacean Biology*, **17**: 687-691.

- LACOURSIERE-ROUSSEL, A. & B. SAINTE-MARIE, 2009. Sexual system and female spawning frequency in the sculptured shrimp Sclerocrangon boreas (Decapoda: Caridea: Crangonidae). *Journal of Crustacean Biology*, **29**: 192-200.
- LÓPEZ GRECO, L. S., V. VIAU, M. LAVOLPE, G. BOND-BUCKUP & E. M. RODRIGUEZ, 2004. Juvenile hatching and maternal care in Aegla uruguayana (Anomura, Aeglidae). *Journal of Crustacean Biology*, **24**: 309-313.
- MAKAROV, R., 1968. On the larval development of the genus Sclerocrangon G. O. Sars (Caridea, Crangonidae). *Crustaceana Supplement* **2**: 27-37.
- NG, P. K. L. & C. G. S. TAN, 1995. Geosesarma notophorum sp. nov. (Decapoda, Brachyura, Grapsidae, Sesarminae), a terrestrial crab from Sumatra, with novel brooding behaviour. *Crustaceana*, **68**: 390-395.
- THIEL, M., 2000. Extended parental care behavior in crustaceans - a comparative overview. In: J.C.V.V. Klein & F.R. Schram (eds.), *The Biodiversity Crisis and Crustacea. Crustacean Issues*. **12**: 211-226. (A.A. Balkema, Rotterdam).
- , 2003. Extended parental care in crustaceans- an update. *Revista Chilena de Historia Natural*, **76**: 205-218.
- VOGT, G. & L. TOLLEY, 2004. Brood care in freshwater crayfish and relationship with the offspring's sensory deficiencies. *Journal of Morphology*, **262**: 566-582.
- WALSH, C. J., 1993. Larval development of Paratya australiensis Kemp, 1917 (Decapoda: Caridea: Atyidae), reared in the laboratory, with comparisons of fecundity and egg and larval size between estuarine and riverine environments. *Journal of Crustacean Biology*, **13**: 456-480.

CAPTION

Fig. 1. Geographic distribution of: 1, Dugastella marocana Bouvier, 1912; and, 2, D. valentina (Ferrer Galdiano, 1924) according to literature reports and present collection data. a, Settât, topotypic locality of D. marocana; b, Almenara ponds, topotypic locality of D. valentina; w, Kihane River; x, spring next to Aïn-Belmesk; y, As Sila River; z, Oum er Rbia River.