1 Dynamics of liver GH/IGF axis and selected stress markers in juvenile gilthead sea
2 bream (*Sparus aurata*) exposed to acute confinement. Differential stress response of
3 growth hormone receptors

4
5
6 Alfonso Saera-Vila^a, Josep Alvar Calduch-Giner^a, Patrick Prunet^b and Jaume Pérez-
7 Sánchez^a

8 ^a Nutrition and Fish Growth Endocrinology, Institute of Aquaculture Torre de la Sal
9 (CSIC), 12595 Ribera de Cabanes, Castellón, Spain

10 ^b INRA-Physiology, Rennes, France

11
12
13 Corresponding author: Jaume Pérez-Sánchez

14 E-mail address: jperez@iats.csic.es

15 Phone: +34 964319500

16 Fax: +34 964319509

Abstract

The time courses of liver GH/IGF axis and selected stress-markers were analyzed in juvenile gilthead sea bream (*Sparus aurata*) sampled at zero time and at fixed intervals (1.5, 3, 6, 24, 72 and 120 h) after acute confinement (120 kg/m³). Fish remained unfed throughout the course of the confinement study, and the fasting-induced increases in plasma growth hormone (GH) levels were partially masked by the GH-stress inhibitory tone. Hepatic mRNA levels of growth hormone receptor-I (GHR-I) were not significantly altered by confinement, but a persistent 2-fold decrease in GHR-II transcripts was found at 24 and 120 h. A consistent decrease in circulating levels of insulin-like growth factor-I (IGF-I) was also found through most of the experimental period, and the down-regulated expression of GHR-II was positively correlated with changes in hepatic IGF-I and IGF-II transcripts. This stress-specific response was concurrent with plasma increases in cortisol and glucose levels, reflecting the cortisol peak (60-70 ng/ml) the intensity and duration of the stressor when data found in the literature were compared. Adaptive responses against oxidative damage were also found, and a rapid enhanced expression was reported in the liver tissue for mitochondrial heat-shock proteins (glucose regulated protein 75). At the same time, the down-regulated expression of proinflammatory cytokines (tumour necrosis factor- α) and detoxifying enzymes (cytochrome P450 1A1) might dictate the hepatic depletion of potential sources of reactive oxygen species. These results provide suitable evidence for a functional partitioning of hepatic GHRs under states of reduced IGF production and changing cellular environment resulting from acute confinement.

Key words: cortisol, glucose regulated protein 75, tumour necrosis factor- α , cytochrome P450 1A1.

1. Introduction

Over the last decade, growth hormone receptors (GHRs) have been cloned and sequenced in more than forty fish species covering almost all fish lineages (Calduch-Giner et al., 2001; Lee et al., 2001; Benedet et al., 2005). Initially, these receptors were clustered in two clades encompassing GHRs of salmonid (GHR-II) and non-salmonid fish (GHR-I), although this observation led to the suggestion that both receptors are retained in the same fish through teleost radiation and evolution. This hypothesis is supported by the coexistence of duplicated GHRs in rainbow trout and several Mediterranean perciform fish, such as gilthead sea bream, common dentex and European sea bass (Saera-Vila et al., 2005; Bermejo-Nogales et al., 2007). Likewise, Jiao *et al.* (2006) demonstrated the occurrence of two GHRs in black sea bream, Southern catfish and Nile tilapia. More recently, duplicated GHRs have been demonstrated in Mozambique tilapia (Pierce et al., 2007), orange-spotted grouper (Li et al., 2007) and Atlantic halibut (Hildahl et al., 2007; Hildahl et al., 2008), which indicates that duplication and divergence of fish GHRs might have taken place in an early fish ancestor.

Nevertheless, receptor specificity remains unclear and GHR-I of masu salmon binds somatolactin (SL) and growth hormone (GH), but the binding affinity is higher for SL than GH and it was then named SL receptor by Fukada et al. (2005). The orthologous medaka gene might also mediate SL signalling (Fukamachi et al., 2005), but the switch and diversification of GH/SL receptors are highly probable among modern and primitive fish lineages. Thus, GHRs of Japanese eel bind specifically to GH (Ozaki et al., 2006), whereas the GHR might be a promiscuous receptor for GH and SL in the archaic sturgeon and lungfish (Fukamachi and Meyer, 2007). This promiscuity is

not unusual in the Class I cytokine receptor superfamily, although each ligand/receptor interaction may result in unique signalling outcomes (Denley et al., 2005) that may differ among fish species. Indirect evidence for this also exists in gilthead sea bream, as transcriptional studies indicate that insulin-like growth factors (IGFs) in growth (liver, skeletal muscle) and immune (head kidney) relevant tissues are positively correlated with GHR-I rather than GHR-II in parasite-challenges (Sitjà-Bobadilla *et al.*, 2008) and different growth models (Benedito-Palos et al., 2007; Saera-Vila et al., 2007). Thus, overall data suggests that GHR-I has evolved in perciform fish, and particularly in gilthead sea bream, as a true orthologous GHR.

Computational analyses also evidence a different promoter organization of GHRs in gilthead sea bream (Saera-Vila et al., 2007). It is noteworthy that surrounding the transcription start site of GHR-II, but not in GHR-I, there are several consensus elements for redox and stress regulatory elements (activating proteins, APs, 1 and 4, and cAMP-responsive elements, CRE), which would contribute to delineate a differential and stress-specific regulation of fish GHRs. Confinement exposure is a common stressor in aquaculture practice that impacts negatively on the immune system, reproduction and growth performance (Pickering 1993; Van Weerd and Komen, 1998). Thus, this experimental condition was chosen to analyze in juvenile fish the functional partitioning of hepatic GHRs after acute stress confinement. The study includes data on GHR transcripts, hepatic IGF production and plasma levels of GH. Circulating levels of cortisol and glucose were measured as conventional stress markers to assess the reliability of stress response triggered under our experimental conditions. Since mitochondrial heat-shock proteins (glucose regulated protein 75, GRP75), proinflammatory cytokines (tumour necrosis factor- α , TNF- α) and detoxifying enzymes (cytochrome P450 1A1, CYP1A1) are also stress-sensitive markers in sparid fish

(Bermejo-Nogales et al., 2007; 2008), these genes were chosen to assess how protein misfolding and potential sources of reactive oxygen species (ROS) were regulated in concert with the liver GH/IGF axis.

2. Material and methods

2.1. Fish rearing and sampling

Juvenile gilthead sea bream (*Sparus aurata*) of 110-130 g final body weight were reared from July to December in a seawater re-circulatory system (500-l tanks) equipped with physical/biological filters, and a heat-unit system that maintained water temperature above 18-19 °C. Voluntary feed intake was near to maintenance ration at the time of the experiment (December), and it was stopped one day before confinement exposure to avoid result disturbances due to differences in feed intake between control (undisturbed fish) and stressed fish. Batches of 10 fish were transferred from 500-l tanks (9-10 kg/m³) to cylinder net baskets of 10-l volume (117-123 Kg/m³), each one suspended in 90-l tanks with a seawater flow of 10-l/min to avoid water deterioration (oxygen > 5 ppm; unionised ammonia < 0.02 mg/l). These fish served as zero time and stressed fish at specific sampling times (1.5, 3, 6, 24, 72 and 120 h) after confinement exposure. Additional 500-l tanks (one per each sampling time) were used as control fish donors. No mortality was registered in control and stressed fish over the course of the confinement period.

At each sampling time, eight fish from control and confinement tanks were netted into a bucket containing 0.1 g/l of 3-aminobenzoic acid ethyl ester (MS-222;

Sigma, Saint Louis, MO, USA). Blood was taken from caudal vessels (in less than 2 min for all fish), centrifuged at 3000 g for 20 min at 4 °C, and plasma samples were frozen and stored at -30 °C until hormone and metabolite analyses were performed. Prior to tissue collection, fish were killed by cervical section and the liver was extracted, immediately frozen in liquid nitrogen, and stored at -80 °C awaiting RNA isolation. All procedures were carried out according to national (Consejo Superior de Investigaciones Científicas, Institute of Aquaculture Torre de la Sal Review Board) and the current EU legislation on the handling of experimental animals.

2.2. Hormone and metabolite assays

Plasma cortisol levels were assayed using an enzyme immunoassay kit (Diagnostic Systems Laboratories, Webster, TX, USA) based on the competition between unlabelled cortisol and cortisol-horseradish peroxidase for a fixed number of antibody-binding sites. Tetramethylbenzidine was used as a chromogen solution with sensitivity (90% of binding) of 1 ng/ml. Plasma glucose levels were measured by the glucose oxidase method (Thermo Electron, Louisville, CO, USA).

Plasma GH levels were determined by a homologous gilthead sea bream radioimmunoassay (RIA) as reported elsewhere (Martínez-Barberá et al., 1995). The sensitivity and midrange (ED₅₀) of the assay were 0.15 and 1.8 ng/ml, respectively. Plasma IGFs were extracted by acid-ethanol cryoprecipitation (Shimizu et al., 2000), and the concentration of IGF-I was measured by means of a generic fish IGF-I RIA validated for Mediterranean perciform fish (Vega-Rubín de Celis et al., 2004). The assay is based on the use of red sea bream (*Pagrus major*) IGF-I (GroPep, Adelaide, Australia) as tracer and standard, and anti-barramundi (*Lates calcarifer*) IGF-I serum

(GroPep) (1:8000) as a first antibody. The sensitivity and midrange of the assay were 0.05 and 0.7–0.8 ng/ml, respectively.

2.3. RNA extraction and RT procedure

Total RNA extraction from target tissues was performed with the ABI PRISM™ 6100 Nucleic Acid PrepStation (Applied Biosystems, Foster City, CA, USA). Briefly, tissue samples were homogenized at a ratio of 25 mg/ml with a guanidine-detergent lysis reagent. The reaction mixture was treated with proteinase K, and RNA purification was achieved by passing the tissue lysate (0.4 - 0.5 ml) through a purification tray containing an application-specific membrane. Wash solutions containing DNase were applied, and total RNA was eluted into a 96-well PCR plate. The RNA yield was 30-50 µg with absorbance measures ($A_{260/280}$) of 1.9-2.1. Reverse transcription (RT) with random decamers was performed with the High-Capacity cDNA Archive Kit (Applied Biosystems). For this purpose, 500 ng total RNA were reverse transcribed into a final volume of 100 µl. RT reactions were incubated for 10 min at 25 °C and 2 h at 37 °C. Negative control reactions were run without reverse transcriptase.

2.4. Real-time PCR

Measurements of hepatic transcripts (GRP75, TNF- α , CYP1A1, GHR-I, GHR-II, IGF-I and IGF-II) were taken using an iCycler IQ Real-time Detection System (Bio-Rad, Hercules, CA, USA) as described elsewhere (Bermejo-Nogales et al., 2007). Briefly, diluted RT reactions were used for PCR reactions in 25-µl volume. Each PCR-well contained a SYBR Green Master Mix (Bio-Rad), and specific primers at a final

concentration of 0.3 – 0.9 μ M were used to obtain amplicons of 75 – 169 bp in length (Table 1).

β -actin was used as the housekeeping gene, and the efficiency of PCR reactions for the target and the reference gene varied between 87% and 99%, respectively. The dynamic range of standard curves (serial dilutions of RT-PCR reactions) spanned five orders of magnitude, and the amount of product in a particular sample was determined by interpolation of the cycle threshold (Ct) value. The specificity of reaction was verified by analysis of melting curves and by electrophoresis and sequencing of PCR amplified products. Reactions were performed in triplicate and the fluorescence data acquired during the extension phase were normalized to β -actin by the delta-delta method (Livak and Schmittgen, 2001). No changes in β -actin expression were found in response to confinement exposure.

2.5. Statistical analysis

The time course of circulating levels of hormones and metabolites was analyzed by one-way analysis of variance (ANOVA) followed by Student-Newman-Keuls. At each sampling time, stress-specific changes in circulating parameters and hepatic transcripts were analyzed by Student t-test. Pearson Product Moment correlations was used for correlation analyses of circulating GH and IGF-I, and hepatic transcripts of IGFs and GHRs. SPSS for Windows Version 14.0.1 (SPSS Inc) was used as statistical software.

3. Results

3.1. Circulating and hepatic stress markers

Plasma cortisol titres (< 3 ng/ml) in control fish did not change significantly over the course of the study (Fig. 1A). In stressed fish, plasma cortisol levels peaked up to 60-70 ng/ml after 1.5 h of confinement exposure with a recovery of control values at the 6 h sampling time. After 24 h of confinement exposure plasma cortisol levels peaked again, with a recovery of control values at the last sampling time (120 h).

Plasma glucose levels in control fish did not vary over the course of the confinement study (Fig. 1B). In stressed fish, the plasma glucose concentration was 2-fold increased at the 3 h sampling time, with a progressive recovery of control values during the subsequent 6-120 h sampling times (Fig. 1B).

Hepatic transcripts of GRP75 were significantly up-regulated (2 or 3-fold increase) over the course of the 6-72 h sampling period, with a recovery of control values at the last sampling time (120 h) (Fig. 2A). After 6 h of confinement exposure, TNF- α (Fig. 2B) and CYP1A1 (Fig. 2C) transcripts were not significantly altered, but a 2 or 5-fold decrease was found during the subsequent 24-120 h sampling times.

3.2. Liver GH/IGF axis

All fish over the course of the confinement period remained unfed, and GH levels in control fish increased significantly with time (Fig. 3A). This fasting-induced GH rise was partially masked by confinement exposure, and plasma GH levels in stressed fish were 2-fold lower than in control fish at the 72 and 120 h sampling times.

Circulating IGF-I concentration was also lowered by confinement exposure, and stressed fish showed a significant reduction in plasma levels of IGF-I over the course of the 6-72 h period (Fig. 3B), although there was not a significant correlation with GH levels. The maximum decrease (3-fold) was found at the 24 h sampling time, with a recovery of control values at the last sampling (120 h).

After 6 h of confinement exposure, hepatic IGF-I (Fig. 4A) and IGF-II (Fig. 4B) expression was not altered by confinement exposure, but a 2-fold decrease in both IGF transcripts was evidenced at the 24 h sampling time. Later on, the trend was towards recovery of the control values, but a significant reduction was still found at the 120 h sampling time. Regarding the hepatic expression of GHRs, we failed to demonstrate any significant change in the expression of GHR-I (Fig. 5A). In contrast, a 2-fold reduction in GHR-II transcripts was found over the course of the 24-120 h sampling times (Fig. 5B). This stress-specific response was positively correlated with changes in transcripts encoding for IGF-I ($r=0.605$, $P < 0.01$) and IGF-II ($r=0.514$, $P < 0.05$).

4. Discussion

Gilthead sea bream is an important aquaculture species in the Mediterranean region, and the stress response of cortisol, glucocorticoid receptors and energy metabolites has been examined in fish subjected to different stressors (Tort et al., 1996; Arends et al., 1999; Acerete et al., 2008). There is also now experimental evidence for an inhibitory effect of handling and crowding stress upon circulating GH levels (Rotllant et al., 2000; 2001), but the effects of acute and chronic stressors on hepatic IGF production and expression of duplicated fish GHRs have not been evaluated simultaneously. This study is, therefore, the first report in non salmonid fish considering the time course of selected stress markers and key components of the liver GH/IGF axis in a changing cellular environment resulting from acute confinement.

Cortisol and glucose are currently used as primary/secondary stress biomarkers in a wide variety of fish species (Barton and Iwama, 1991; Wendelaar Bonga, 1997). In the present study, cortisol titres in control fish enter into the ideal fish levels (>5 ng/ml) (Pickering and Pottinger, 1989), and the stress-associated increases in plasma cortisol and glucose levels followed the expected response in this fish species (Ortuno et al., 2001; Sangiao-Alvarellos et al., 2005). Moreover, the cortisol peak (60-70 ng/ml) did not deviate from gilthead sea bream literature data when plasma titres were related to stocking density: (i) 15-20 ng/ml at $26-30 \text{ kg/m}^3$ (Rotllant et al., 2000; Barton et al., 2005), (ii) 30 ng/ml at 70 kg/m^3 (Sangiao-Alvarellos et al., 2005) and (iii) 178 ng/ml at 200 kg/m^3 (Rotllant et al., 2001). This finding supports a close relationship between cortisol response and the intensity/duration of confinement, although we found a bimodal cortisol rise as reported elsewhere by Arends et al. (1999). These authors considered that the first cortisol peak is due to a rapid activation of the hypothalamic-

257 pituitary-interrenal axis (HPI) leading to exhaustion and/or negative feedback regulation
258 of this axis. The second cortisol peak might be ACTH-independent and, in our case, it
259 was associated to a wide variety of tissue repair and remodelling processes that were
260 mostly mediated by the stress response of the endoplasmic reticulum (Calduch-Giner et
261 al., 2008).

262 The transition from normal to stressful conditions is also accompanied by a
263 robust up-regulation of heat shock proteins (Hsp), which damps the cytotoxicity caused
264 by misfolded and denaturated proteins (Anckar and Sistonen, 2007). These proteins thus
265 have housekeeping functions, which makes the mitochondrial GRP75 and its yeast
266 homologue (SSC1p) life-essential (Craig et al., 1989; Kaul et al., 2007). Few studies
267 have examined the role of GRP75 in fish, but it is now recognized that hepatic GRP75
268 is up-regulated in zebrafish toxicogenomic models (Lam et al., 2006). The up-regulated
269 expression of hepatic GRP75 has also been demonstrated in a previous gilthead sea
270 bream study with fish exposed for short (24 h) and long (3 weeks/pair fed study) periods
271 of time to high (120 kg/m³) and mild (50 kg/m³) loading densities, respectively
272 (Bermejo-Nogales et al., 2008). This enhanced expression protects mitochondria against
273 oxidative damage, and not surprisingly the hepatic GRP75 showed in the present study
274 a rapid and robust up-regulation, which stopped after 5 days of acute confinement
275 exposure. This loss of responsiveness can be understood as the reestablishment of a new
276 redox homeostasis that was encompassed by a down-regulated expression of hepatic
277 TNF α and CYP1A1 genes. The former is a pro-inflammatory cytokine that increases
278 leukocyte and mitochondrial ROS production (Yang et al., 2007), and experimental data
279 support a reduced or enhanced expression of TNF α and TNF decoy receptors in stressed
280 rats and fish, respectively (Connor et al., 2005; Momoda et al., 2007). Likewise, hepatic
281 CYP1A1 is a major hepatic metabolizing enzyme that transforms endogenous

substrates, procarcinogens and pollutants into less- or non-toxic metabolites (van der Oost et al., 2003). These processes are, however, potential sources of ROS, and CYP1A1 expression is generally repressed by oxidative stress (Barouki and Morel, 2001).

The modulation of ROS production and antioxidant defences by the endocrine system is also a well-documented phenomenon (Haddad et al., 2002). Thus, in long-lived dwarf mice, the reduced signalling of GH and IGF-I contributes to maintain an appropriate cellular redox state (Holzenberger et al., 2003; Bartke and Brown-Borg, 2004). Conversely, animals over expressing GH combat oxidative stress less efficiently than normal and dwarf mice (Brown-Borg et al., 1999; Brown-Borg and Rakoczy, 2000). In this way, it is not surprising the stress-induced reduction in plasma GH levels. This stress response after confinement exposure has been reported in a wide variety of fish species, including gilthead sea bream (Rotllant et al., 2000; 2001), tilapia (Auperin et al., 1997), rainbow trout (Pickering et al., 1991) and Atlantic salmon (Wilkinson et al., 2006). However, the link between somatotrophic and HPI axes is not clear, since the *in vivo* cortisol and GH negative correlation disagrees with the *in vitro* GH stimulatory action of cortisol (Nishioka et al., 1985; Wendelaar Bonga, 1997; Yada et al., 2005). This apparent inconsistency is also found in mammals, and short-term glucocorticoid treatments stimulate GH secretion whereas long-term treatments exert an inhibitory action (Casanueva et al., 1990; Guiustina et al., 1992; Miell et al., 1991). The present study was conducted under a regimen of natural photoperiod, and the shortened light-dark cycle at the time of the study produced the known winter decreases in cortisol and GH titres (Mingarro et al., 2002). This scenario of seasonal hyposomatotropism and hypocortisolism makes difficult any additional enhancement of the inhibitory GH-tonus, but the achieved results indicate that confinement exposure partially blunted the fasting-

induced increase in plasma GH levels. This elevated plasma GH levels during fasting and malnutrition presumably reflects the insensitivity of liver to GH action, and thereby a reduced negative feedback effect of IGF-I on pituitary GH release (Pérez-Sánchez et al., 1992; 1995). Indeed, more recent studies in gilthead sea bream (Saera-Vila et al., 2005) and hybrid striped bass (Picha et al., 2008) indicate that this fasting state of GH resistance is mediated at long-term by a down-regulated expression of GHR-I and II.

Time series analyses of IGFs evidenced a lag-time among changes in hepatic IGF transcripts and circulating levels of protein. However, the hepatic tissue is the primary source of circulating IGF-I and its depletion by stress and nutritional disorders reflects, in this and previous gilthead sea bream studies (Gómez-Requeni et al., 2004; Benedito-Palos et al., 2007), the down-regulated expression of hepatic IGF-I. In rainbow trout and Atlantic salmon, handling and confinement also reduce circulating levels of IGF-II (Wilkinson et al., 2006). Thus far, there is no information in gilthead sea bream on the effects of aquaculture stressors on circulating levels of IGF-II. Nevertheless, the results presented here evidenced a robust down-regulation of hepatic IGF-II, which might reflect reduced circulating levels of IGF-II. Experimental evidence in catfish (Small et al., 2006), tilapia (Kajimura et al., 2003) and sunshine bass (Davis and Peterson, 2006) also indicate that confinement and cortisol treatment reduce IGF activity and sensitivity to GH. Therefore, the mechanisms by which stressful conditions inhibit growth are apparently conserved throughout evolution, but more complete information on the sequence of events that adjusts growth to each particular condition requires species-specific studies. Thus, in gilthead sea bream, the reduced hepatic IGF-I production can be compensated at the local tissue level (skeletal muscle) by the enhanced expression of IGF-II (Benedito-Palos et al., 2007; Saera-Vila et al., 2007). In

support of this, growth differences between families of channel catfish have been related to differences in muscle IGF-II expression (Peterson et al., 2004).

In gilthead sea bream, local compensatory mechanisms of growth can also be mediated by GHRs, and the muscle expression of GHR-II is increased by the large dietary replacement of fish oil with vegetable oils regardless of that found for IGFs (Benedito-Palos et al., 2007). Both in gilthead sea bream (Saera-Vila et al., 2005) and hybrid striped bass (Picha et al., 2008), the muscle expression of GHR-II is also up-regulated by fasting, which may serve to repair and preserve tissue functions as reported for rat GHRs in atrophied muscle fibers (Casse et al., 2003). In the same way, we found in the present study that hepatic transcripts of GHR-II were down-regulated after acute confinement exposure, whereas no significant changes were reported for GHR-I. Since glucocorticoids inhibit the expression of GHRs in mammals (Beauloye et al., 1999; Gabrielsson et al., 1995) and GHR-I exhibit a higher affinity for SL in salmonids (Fukada et al., 2005), it may be argued that GHR-II is the true orthologous of mammalian GHR whereas GHR-I might be mostly evolved as a SL receptor. However, SL has not a growth-promoting action in gilthead sea bream (Pérez-Sánchez et al., 2002; Vega-Rubín de Celis et al., 2003), and nice positive correlations between GHR-I and IGF transcripts have been reported in different experimental models of this marine fish (Benedito-Palos et al., 2007; Saera-Vila et al., 2007; Sitjà-Bobadilla *et al.*, 2008). Moreover, GHRs of Japanese eel bind specifically to GH (Ozaki et al., 2006) and Jiao et al. (2006) did not found in black sea bream any response to SL in GHR transfected cells. If so, there is not an easy pattern of transcription and function of GHRs in non-salmonid fish, although it is reasonable to imagine that GHR-II has a dual function preserving cell survival and limiting at the same time growth and hepatic IGF production under life-stress conditions.

Prior to the present study, there is not a time course study addressing the differential stress regulation of GHRs in fish, although several lines of evidence support a complex cross talk between glucocorticoids and fish GHRs. Thus, cortisol injection in black sea bream enhances the expression of hepatic GHR-I without significant effects on GHR-II (Jiao et al., 2006). Pierce *et al.* (2005) reported an increased expression of GHR-II in primary cultures of salmon hepatocytes exposed to dexamethasone, and Small *et al.* (2006) demonstrated a reduced expression of GHR-II in catfish fed with cortisol for 4 weeks. More recently, Uchida et al. (2009) reported that handling and confinement exposure enhanced in tilapia the hepatic expression of GHR-II. To date there is no an easy explanation for these apparent contradictory findings, although it is suspected that the glucocorticoid response is a dynamic process having both stimulatory and inhibitory effects upon the somatotrophic axis, which may reflect physiological and pharmacological responses and/or differences between short- and long-term treatments as found in mammals (Vottero et al., 2003). In particular for gilthead sea bream, we found in the present study that the stress-specific response of GHR-II was encompassed by adaptive changes in ROS production and mitochondrial chaperones. Furthermore, a previous computational study (Saera-Vila et al., 2007) recognized a subset of stress-sequence elements (CRE and APs) surrounding the transcription start site of GHR-II. These sequence elements can activate and/or inhibit the expression of target genes depending on promoter organization and cellular redox context (Abate et al., 1991; Hai and Hartman, 2001; Glahder et al., 2003). Taking all these findings together, there is now increasing evidence for a functional partitioning of GHRs in gilthead sea bream, but some overlapping and redundancy may occur, and the possibility of GHR-I/GHR-II heterodimers cannot be excluded.

In summary, stress-mediated changes in primary and secondary stress markers (cortisol, glucose, GRP75, TNF- α , CYP1A1) evidenced a robust stress response with the establishment of a new cellular homeostasis in fish exposed to acute confinement. In this changing cellular environment, the time course of the liver GH/IGF axis revealed delayed stimulatory effects of fasting on plasma GH titres, a reduced hepatic IGF production and a differential expression of the two types of gilthead sea bream GHRs. This can represent an interesting field of research for the exploration of the different stress-susceptibilities of fish species and strains to aquaculture stressors.

5. Acknowledgments

This work was funded by EU (contract no. SSP98-CT-2004-513692; Combined genetic and functional genomic approaches for stress and disease resistance markers assisted selection in fish and shellfish, AQUAFIRST) and Spanish (Ingenio-2010 Programme; Improvement of Aquaculture Production by the use of biotechnological tools, AQUAGENOMICS) projects. AS-V was recipient of a Spanish PhD fellowship from the Diputación Provincial de Castellón. The authors are grateful to M.A. González for the excellent technical assistance in molecular analysis.

6. References

- Abate, C., Luk, D., and Curran, T., 1991. Transcriptional regulation by Fos and Jun In vitro - interaction among multiple activator and regulatory domains. *Mol. Cell. Biol.* 11, 3624-3632.
- Acerete, L., Balasch, J.C., Castellana, B., Redruelo, B., Roher, N., Canario, A.V., Planas, J.V., MacKenzie, S., and Tort, L., 2008. Cloning of glucocorticoid receptor (GR) in gilthead seabream (*Sparus aurata*) - Differential expression of GR and immune relevant genes in gilthead seabream after an immune challenge. *Comp. Biochem. Physiol. B. Biol.* 148, 32-43.
- Anckar, J. and Sistonen, L., 2007. Heat shock factor 1 as a coordinator of stress and developmental pathways. *Adv. Exp. Med. Biol.* 594, 78-88.
- Arends, R.J., Mancera, J.M., Muñoz, J.L., Wendelaar Bonga, S.E., and Flik, G., 1999. The stress response of the gilthead sea bream (*Sparus aurata* L.) to air exposure and confinement. *J Endocrinol* 163, 149-157.
- Auperin, B., Baroiller, J.F., Ricordel, M.J., Fostier, A., and Prunet, P., 1997. Effect of confinement stress on circulating levels of growth hormone and two prolactins in freshwater-adapted tilapia (*Oreochromis niloticus*). *Gen. Comp. Endocrinol.* 108, 35-44.
- Barouki, R. and Morel, Y., 2001. Repression of cytochrome P450 1A1 gene expression by oxidative stress: mechanisms and biological implications. *Biochem. Pharmacol.* 61, 511-516.
- Bartke, A. and Brown-Borg, H., 2004. Life extension in the dwarf mouse. *Curr Top Dev Biol* 63, 189-225.
- Barton, B.A. and Iwama, G.K., 1991. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases* 1, 3-26.
- Barton, B.A., Ribas, L., Acerete, L., and Tort, L., 2005. Effects of chronic confinement on physiological responses of juvenile gilthead sea bream, *Sparus aurata* L., to acute handling. *Aquac. Res.* 36, 172-179.
- Benedet, S., Johansson, V., Sweeney, G., Galay-Burgos, M., and Björnsson, B., 2005. Cloning of two Atlantic salmon growth hormone receptor isoforms and *in vitro* ligand-binding response. *Fish Physiol. Biochem.* 31, 315-329.
- Benedito-Palos, L., Saera-Vila, A., Calduch-Giner, J.A., Kaushik, S., and Pérez-Sánchez, J., 2007. Combined replacement of fish meal and oil in practical diets for fast growing juveniles of gilthead sea bream (*Sparus aurata* L.): networking of systemic and local components of GH/IGF axis. *Aquaculture* 267, 199-212.
- Bermejo-Nogales, A., Benedito-Palos, L., Saera-Vila, A., Calduch-Giner, J.A., Sitjà-Bobadilla, A., and Pérez-Sánchez, J., 2008. Confinement exposure induces glucose regulated protein 75 (GRP75/mortalin/mtHsp70/PBP74/HSPA9B) in the

440 hepatic tissue of gilthead sea bream (*Sparus aurata* L.). Comp. Biochem.
441 Physiol. B 149, 428-438.

442 Bermejo-Nogales, A., Saera-Vila, A., Calduch-Giner, J.A., Navarro, J.C., Sitjà-
443 Bobadilla, A., and Pérez-Sánchez, J., 2007. Differential metabolic and gene
444 expression profile of juvenile common dentex (*Dentex dentex* L.) and gilthead
445 sea bream (*Sparus aurata* L.) in relation to redox homeostasis. Aquaculture 267,
446 213-224.

447 Brown-Borg, H., Bode, A., and Bartke, A., 1999. Antioxidative mechanisms and plasma
448 growth hormone levels. Endocrine 11, 41-48.

449 Brown-Borg, H.M. and Rakoczy, S.G., 2000. Catalase expression in delayed and
450 premature aging mouse models. Exp. Gerontol. 35, 199-212.

451 Calduch-Giner, J.A., Saera-Vila, A., Cairns, M., Davey, G., Prunet, P., and Pérez-
452 Sánchez, J., 2008. Time series analyses of sea bream (*Sparus aurata* L.) stress
453 response after confinement exposure. Comp. Biochem. Physiol. A 151, S41.

454 Calduch-Giner, J.A., Duval, H., Chesnel, F., Boeuf, G., Pérez-Sánchez, J., and Boujard,
455 D., 2001. Fish growth hormone receptor: molecular characterization of two
456 membrane-anchored forms. Endocrinology 142, 3269-3273.

457 Casanueva, F.F., Burguera, B.A.R.T., Muruais, C.O.V.A., and Dieguez, C.A.R.L.,
458 1990. Acute administration of corticoids: a new and peculiar stimulus of growth
459 hormone secretion in man. J Clin Endocrinol Metab 70, 234-237

460 Connor, T.J., Brewer, C., Kelly, J.P., and Harkin, A., 2005. Acute stress suppresses pro-
461 inflammatory cytokines TNF- α and IL-1 β independent of a catecholamine-
462 driven increase in IL-10 production. J. Neuroimmunol. 159, 119-128.

463 Craig, E.A., Kramer, J., Shilling, J., Werner-Washburne, M., Holmes, S., Kasic-
464 Smithers, J., and Nicolet, C.M., 1989. SSC1, an essential member of the yeast
465 HSP70 multigene family, encodes a mitochondrial protein. Mol. Cell. Biol. 9,
466 3000-3008.

467 Davis, K.B. and Peterson, B.C., 2006. The effect of temperature, stress, and cortisol on
468 plasma IGF-I and IGFBPs in sunshine bass. Gen. Comp. Endocrinol. 149, 219-
469 225.

470 Denley, A., Cosgrove, L.J., Booker, G.W., Wallace, J.C., and Forbes, B.E., 2005.
471 Molecular interactions of the IGF system. Cytokine Growth Factor Rev. 16,
472 421-439.

473 Fukada, H., Ozaki, Y., Pierce, A.L., Adachi, S., Yamauchi, K., Hara, A., Swanson, P.,
474 and Dickhoff, W.W., 2005. Identification of the salmon somatolactin receptor, a
475 new member of the cytokine receptor family. Endocrinology 146, 2354-2361.

476 Fukamachi, S. and Meyer, A., 2007. Evolution of receptors for growth hormone and
477 somatolactin in fish and land vertebrates: Lessons from the Lungfish and
478 Sturgeon orthologues. J. Mol. Evol. 65, 359-372.

- 479 Fukamachi, S., Yada, T., and Mitani, H., 2005. Medaka receptors for somatolactin and
480 growth hormone: phylogenetic paradox among fish growth hormone receptors.
481 Genetics 171, 1875-1883.
- 482 Giustina, A. and Wehrenberg, W.B., 1992. The role of glucocorticoids in the regulation
483 of growth hormone secretion Mechanisms and clinical significance. Trends
484 Endocrinol. Metab. 3, 306-311.
- 485 Glahder, J.A., Hansen, C.N., Vinther, J., Madsen, B.S., and Norrild, B., 2003. A
486 promoter within the E6 ORF of human papillomavirus type 16 contributes to the
487 expression of the E7 oncoprotein from a monocistronic mRNA. J. Gen. Virol.
488 84, 3429-3441.
- 489 Gómez-Requeni, P., Mingarro, M., Calduch-Giner, J.A., Médale, F., Martin, S.A.M.,
490 Houlihan, D.F., Kaushik, S., and Pérez-Sánchez, J., 2004. Protein growth
491 performance, amino acid utilisation and somatotropic axis responsiveness to fish
492 meal replacement by plant protein sources in gilthead sea bream (*Sparus*
493 *aurata*). Aquaculture 232, 493-510.
- 494 Haddad, J.J., Saadé, N.E., and Safieh-Garabedian, B., 2002. Cytokines and neuro-
495 immune-endocrine interactions: a role for the hypothalamic-pituitary-adrenal
496 revolving axis. J. Neuroimmunol. 133, 1-19.
- 497 Hai, T. and Hartman, M.G., 2001. The molecular biology and nomenclature of the
498 activating transcription factor/cAMP responsive element binding family of
499 transcription factors: activating transcription factor proteins and homeostasis.
500 Gene 273, 1-11.
- 501 Hildahl, J., Power, D., Björnsson, B.T., and Einarsson, I.E., 2008. Involvement of
502 growth hormone-insulin-like growth factor I system in cranial remodeling
503 during halibut metamorphosis as indicated by tissue- and stage-specific receptor
504 gene expression and the presence of growth hormone receptor protein. Cell
505 Tissue Res. 320, 211-225.
- 506 Hildahl, J., Sweeney, G., Galay-Burgos, M., Einarsson, I.E., and Björnsson, B.T.,
507 2007. Cloning of Atlantic halibut growth hormone receptor genes and
508 quantitative gene expression during metamorphosis. Gen. Comp. Endocrinol.
509 151, 143-152.
- 510 Holzenberger, M., Dupont, J., Ducos, B., Leneuve, P., Gélöën, A., Even, P.C., Cervera,
511 P., and Le Bouc, Y., 2003. IGF-1 receptor regulates lifespan and resistance to
512 oxidative stress in mice. Nature 421, 182-187.
- 513 Jiao, B., Huang, X., Chan, C.B., Zhang, L., Wang, D., and Cheng, C.H.K., 2006. The
514 co-existence of two growth hormone receptors in teleost fish and their
515 differential signal transduction, tissue distribution and hormonal regulation of
516 expression in seabream. J Mol Endocrinol 36, 23-40.
- 517 Kajimura, S., Hirano, T., Visitacion, N., Moriyama, S., Aida, K., and Grau, E.G., 2003.
518 Dual mode of cortisol action on GH/IGF-I/IGF binding proteins in the tilapia,
519 *Oreochromis mossambicus*. J Endocrinol 178, 91-99.

520 Kaul, S.C., Deocaris, C.C., and Wadhwa, R., 2007. Three faces of mortalin: A
521 housekeeper, guardian and killer. *Exp. Gerontol.* 42, 263-274.

522 Lam, S.H., Winata, C.L., Tong, Y., Korzh, S., Lim, W.S., Korzh, V., Spitsbergen, J.,
523 Mathavan, S., Miller, L.D., Liu, E.T., and Gong, Z., 2006. Transcriptome
524 kinetics of arsenic-induced adaptive response in zebrafish liver. *Physiol.*
525 *Genomics* 27, 351-361.

526 Lee, L.T.O., Nong, G., Chan, Y.H., Tse, D.L.Y., and Cheng, C.H.K., 2001. Molecular
527 cloning of a teleost growth hormone receptor and its functional interaction with
528 human growth hormone. *Gene* 270, 121-129.

529 Li, Y., Liu, X., Zhang, Y., Zhu, P., and Lin, H., 2007. Molecular cloning,
530 characterization and distribution of two types of growth hormone receptor in
531 orange-spotted grouper (*Epinephelus coioides*). *Gen. Comp. Endocrinol.* 152,
532 111-122.

533 Livak, K.J. and Schmittgen, T.D., 2001. Analysis of relative gene expression data using
534 real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* 25, 402-408.

535 Martínez-Barberá, J.P., Pendón, C., Martí-Palanca, H., Calduch-Giner, J.A., Rodríguez,
536 R.B., Valdivia, M.M., and Pérez-Sánchez, J., 1995. The use of recombinant
537 gilthead sea bream (*Sparus aurata*) growth hormone for radioiodination and
538 standard preparation in radioimmunoassay. *Comp. Biochem. Physiol. A* 110,
539 335-340.

540 Miell, J.P., Corder, R., Pralong, F.P., and Gaillard, R.C., 1991. Effects of
541 dexamethasone on GHRH, arginine and dopaminergic stimulated GH
542 stimulation and total plasma IGF-I levels in normal male volunteers. *J Clin*
543 *Endocrinol Metab* 72, 675-681.

544 Mingarro, M., Vega-Rubín de Celis, S., Astola, A., Pendón, C., Martínez Valdivia, M.,
545 and Pérez-Sánchez, J., 2002. Endocrine mediators of seasonal growth in gilthead
546 sea bream (*Sparus aurata*): the growth hormone and somatolactin paradigm.
547 *Gen. comp. endocrinol.* 128, 102-111.

548 Momoda, T.S., Schwindt, A.R., Feist, G.W., Gerwick, L., Bayne, C.J., and Schreck,
549 C.B., 2007. Gene expression in the liver of rainbow trout, *Oncorhynchus mykiss*,
550 during the stress response. *Comp. Biochem. Physiol. D* 2, 303-315.

551 Nishioka, R.S., Grau, E.G., and Bern, H.A., 1985. In vitro release of growth hormone
552 from the pituitary gland of tilapia, *Oreochromis mossambicus*. *Gen. comp.*
553 *endocrinol* 60, 90-94

554 Ortuno, J., Esteban, M.A., and Meseguer, J., 2001. Effects of short-term crowding stress
555 on the gilthead seabream (*Sparus aurata* L.) innate immune response. *Fish*
556 *Shellfish Immunol.* 11, 187-197.

557 Ozaki, Y., Fukada, H., Kazeto, Y., Adachi, S., Hara, A., and Yamauchi, K., 2006.
558 Molecular cloning and characterization of growth hormone receptor and its
559 homologue in the Japanese eel (*Anguilla japonica*). *Comp. Biochem. Physiol. B*
560 143, 422-431.

- 561 Pérez-Sánchez, J., Martí-Palanca, H., and Kaushik, S.J., 1995. Ration size and protein
562 intake affect circulating growth hormone concentration, hepatic growth hormone
563 binding and plasma insulin-like growth factor-I immunoreactivity in a marine
564 teleost, the gilthead sea bream (*Sparus aurata*). J. Nutr. 125, 546-552.
- 565 Pérez-Sánchez, J., Weil, C., and Le Bail, P.-Y., 1992. Effects of human insulin-like
566 growth factor-I on release of growth hormone by rainbow trout (*Oncorhynchus*
567 *mykiss*) pituitary cells. J. Exp. Zool. 262, 287-290.
- 568 Pérez-Sánchez, J., Caldach-Giner, J.A., Mingarro, M., Vega-Rubín de Celis, S., Gómez-
569 Requeni, P., Saera-Vila, A., Astola, A., & Valdivia, M.M., 2002 Overview of
570 Fish Growth Hormone Family. New Insights in Genomic Organization and
571 Heterogeneity of Growth Hormone Receptors. Fish Physiol. Biochem. 27 243-
572 258.
- 573 Picha, M.E., Turano, M.J., Tipsmark, C.K., and Borski, R.J., 2008. Regulation of
574 endocrine and paracrine sources of IGFs and GH receptor during compensatory
575 growth in hybrid striped bass (*Morone chrysops* × *Morone saxatilis*). J.
576 Endocrinol. 199, 81-94.
- 577 Pickering, A. and Pottinger, T., 1989. Stress responses and disease resistance in
578 salmonid fish: Effects of chronic elevation of plasma cortisol. Fish Physiol.
579 Biochem. 7, 253-258.
- 580 Pickering, A.D., 1993. Growth and stress in fish production. Aquaculture 111, 51-63.
- 581 Pickering, A.D., Pottinger, T.G., Sumpter, J.P., Carragher, J.F., and Le Bail, P.Y., 1991.
582 Effects of acute and chronic stress on the levels of circulating growth hormone
583 in the rainbow trout, *Oncorhynchus mykiss*. Gen. Comp. Endocrinol. 83, 86-93.
- 584 Pierce, A.L., Fukada, H., and Dickhoff, W.W., 2005. Metabolic hormones modulate the
585 effect of growth hormone (GH) on insulin-like growth factor-I (IGF-I) mRNA
586 level in primary culture of salmon hepatocytes. J. Endocrinol. 184, 341-349.
- 587 Pierce, A.L., Fox, B.K., Davis, L.K., Visitacion, N., Kitahashi, T., Hirano, T., and Grau,
588 E.G., 2007. Prolactin receptor, growth hormone receptor, and putative
589 somatolactin receptor in Mozambique tilapia: tissue specific expression and
590 differential regulation by salinity and fasting. Gen. Comp. Endocrinol. 154, 31-
591 40.
- 592 Ross, R.J.M. and Chew, S.L., 1995. Acquired growth hormone resistance. Eur J
593 Endocrinol 132, 655-660.
- 594 Rotllant, J., Balm, P.H.M., Pérez-Sánchez, J., Wendelaar-Bonga, S.E., and Tort, L.,
595 2001. Pituitary and interrenal function in gilthead sea bream (*Sparus aurata* L.,
596 Teleostei) after handling and confinement stress. Gen. Comp. Endocrinol. 121,
597 333-342.
- 598 Rotllant, J., Balm, P.H.M., Ruane, N.M., Pérez-Sánchez, J., Wendelaar-Bonga, S.E.,
599 and Tort, L., 2000. Pituitary proopiomelanocortin-derived peptides and
600 hypothalamus-pituitary-interrenal axis activity in gilthead sea bream (*Sparus*
601 *aurata*) during prolonged crowding stress: differential regulation of

602 adrenocorticotropin hormone and α -melanocyte-stimulating hormone release by
603 corticotropin-releasing hormone and thyrotropin-releasing hormone. Gen.
604 Comp. Endocrinol. 119, 152-163.

605 Saera-Vila, A., Calduch-Giner, J.A., and Pérez-Sánchez, J., 2005. Duplication of growth
606 hormone receptor (GHR) in fish genome: gene organization and transcriptional
607 regulation of GHR type I and II in gilthead sea bream (*Sparus aurata*). Gen.
608 Comp. Endocrinol. 142, 193-203.

609 Saera-Vila, A., Calduch-Giner, J.A., and Pérez-Sánchez, J., 2007. Co-expression of
610 IGFs and GH receptors (GHRs) in gilthead sea bream (*Sparus aurata* L.):
611 sequence analysis of the GHR-flanking region. J Endocrinol 194, 361-372.

612 Sangiao-Alvarellos, S., Guzmán, J.M., Láiz-Carrión, R., Míguez, J.M., Martín del Río,
613 M.P., Mancera, J.M., and Soengas, J.L., 2005. Interactive effects of high
614 stocking density and food deprivation on carbohydrate metabolism in several
615 tissues of gilthead sea bream *Sparus auratus*. J. Exp. Zool. A 303, 761-775.

616 Shimizu, M., Swanson, P., Fukada, H., Hara, A., and Dickhoff, W.W., 2000.
617 Comparison of extraction methods and assay validation for salmon insulin-like
618 growth factor-I using commercially available components. Gen. Comp.
619 Endocrinol. 119, 26-36.

620 Sitjà-Bobadilla, A., Calduch-Giner, J., Saera-Vila, A., Palenzuela, O., Álvarez-Pellitero,
621 P., and Pérez-Sánchez, J., 2008. Chronic exposure to the parasite *Enteromyxum*
622 *leei* (Myxozoa: Myxosporea) modulates the immune response and the
623 expression of growth, redox and immune relevant genes in gilthead sea bream,
624 *Sparus aurata* L. Fish Shellfish Immunol. 24, 610-619.

625 Small, B.C., Murdock, C.A., Waldbieser, G.C., and Peterson, B.C., 2006. Reduction in
626 channel catfish hepatic growth hormone receptor expression in response to food
627 deprivation and exogenous cortisol. Domest. Anim. Endocrinol. 31, 340-356.

628 Tort, L., Sunyer, J.O., Gómez, E., and Molinero, A., 1996. Crowding stress induces
629 changes in serum haemolytic and agglutinating activity in the gilthead sea bream
630 *Sparus aurata*. Vet. Immunol. Immunopathol. 51, 179-188.

631 Uchida, K., Moriyama, S., Breves, J.P., Fox, B.K., Pierce, A.L., Borski, R.J., Hirano, T.
632 and Gordon Grau, E., 2009. cDNA cloning and isolation of somatolactin in
633 Mozambique tilapia and effects of seawater acclimation, confinement stress, and
634 fasting on its pituitary expression. Gen. Comp. Endocrinol. 161, 162-170.

635 van der Oost, R., Beyer, J., and Vermeulen, N.P.E., 2003. Fish bioaccumulation and
636 biomarkers in environmental risk assessment: a review. Environ. Toxicol.
637 Pharmacol. 13, 57-149.

638 Van Weerd, J.H. and Komen, J., 1998. The effects of chronic stress on growth in fish: a
639 critical appraisal. Comp. Biochem. Physiol. A 120, 107-112.

640 Vega-Rubín de Celis, S., Gómez-Requeni, P., and Pérez-Sánchez, J., 2004. Production
641 and characterization of recombinantly derived peptides and antibodies for
642 accurate determinations of somatolactin, growth hormone and insulin-like

643 growth factor-I in European sea bass (*Dicentrarchus labrax*). Gen. Comp
644 Endocrinol 139, 266-277.

645 Vega-Rubín de Celis, S., Rojas, P., Gómez-Requeni, P., Albalat, A., Gutiérrez, J.,
646 Médale, F., Kaushik, S.J., Navarro, I., Pérez-Sánchez, J., 2003. Nutritional
647 assessment of somatolactin function in gilthead sea bream (*Sparus aurata*):
648 concurrent changes in somatotrophic axis and pancreatic hormones. Comp.
649 Biochem. Physiol. A 138, 533-542.

650 Vijayan, M.M. and Moon, T.W., 1992. Acute handling stress alters hepatic glycogen
651 metabolism in food-deprived rainbow trout (*Oncorhynchus mykiss*). Can. J. Fish.
652 Aquat. Sci. 49, 2260-2262.

653 Weil, C., Carré, F., Blaise, O., Breton, B., and Le Bail, P.Y., 1999. Differential effect of
654 insulin-like growth factor I on in vitro gonadotropin (I and II) and growth
655 hormone secretions in rainbow trout (*Oncorhynchus mykiss*) at different stages
656 of the reproductive cycle. Endocrinology 140, 2054-2062.

657 Wendelaar Bonga, S.E., 1997. The stress response in fish. Physiol. Rev. 77, 591-625.

658 Wilkinson, R.J., Porter, M., Woolcott, H., Longland, R., and Carragher, J.F., 2006.
659 Effects of aquaculture related stressors and nutritional restriction on circulating
660 growth factors (GH, IGF-I and IGF-II) in Atlantic salmon and rainbow trout.
661 Comp. Biochem. Physiol. A 145, 214-224.

662 Yada, T., Muto, K., Azuma, T., Hyodo, S., and Schreck, C.B., 2005. Cortisol stimulates
663 growth hormone gene expression in rainbow trout leucocytes in vitro. Gen.
664 comp. endocrinol. 142, 248-255.

665 Yang, D., Elner, S.G., Bian, Z.-M., Till, G.O., Petty, H.R., and Elner, V.M., 2007. Pro-
666 inflammatory cytokines increase reactive oxygen species through mitochondria
667 and NADPH oxidase in cultured RPE cells. Exp. Eye Res. 85, 462-472.
668

669
670

Table 1. Gilthead sea bream primer sequences used for real-time PCR.

Gene	accession number	primer sequence										position
GRP-75	DQ524993	F	TCC	GGT	GTG	GAT	CTG	ACC	AAA	GAC		358-381
		R	TGT	TTA	GGC	CCA	GAA	GCA	TCC	ATG		500-477
TNF α	AJ413189	F	CAG	GCG	TCG	TTC	AGA	GTC	TC			587-606
		R	CTG	TGG	CTG	AGA	GGT	GTG	TG			663-644
CYP 1A1	AF011223	F	GCA	TCA	ACG	ACC	GCT	TCA	ACG	C		903-924
		R	CCT	ACA	ACC	TTC	TCA	TCC	GAC	ATC	TGG	1071-1045
GHR-I	AF438176	F	ACC	TGT	CAG	CCA	CCA	CAT	GA			1275-1294
		R	TCG	TGC	AGA	TCT	GGG	TCG	TA			1373-1354
GHR-II	AY573601	F	GAG	TGA	ACC	CGG	CCT	GAC	AG			1690-1709
		R	GCG	GTG	GTA	TCT	GAT	TCA	TGG	T		1764-1743
IGF-I	AY996779	F	TGT	CTA	GCG	CTC	TTT	CCT	TTC	A		112-133
		R	AGA	GGG	TGT	GGC	TAC	AGG	AGA	TAC		195-172
IGF-II	AY996778	F	TGG	GAT	CGT	AGA	GGA	GTG	TTG	T		406-427
		R	CTG	TAG	AGA	GGT	GGC	CGA	CA			514-495
β -actin	X89920	F	TCC	TGC	GGA	ATC	CAT	GAG	A			811-829
		R	GAC	GTC	GCA	CTT	CAT	GAT	GCT			861-841

671
672

LEGENDS

Figure 1. Time course of plasma levels of cortisol (A) and glucose (B) in control (○) and stressed (●) fish. Data are the mean $\pm$ SEM (n = 6-8). Different letters indicate statistically significant changes over the course of the experiment in stressed fish (ANOVA, $P < 0.05$). Statistically significant differences between stressed and control fish were analyzed at each sampling time by means of Student t-test (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Figure 2. Box-Whisker plots representing the time course of the relative gene expression of GRP75 (A), TNF- α (B) and CYP1A1 (C) in stressed fish. Data in control fish were used as arbitrary reference values at each sampling time in the normalization procedure (values > 1 or < 1 indicate increase or decrease respect to reference values). The lower boundary of the box indicates the 25th percentile and the upper boundary of the box indicates the 75th percentile. Whiskers above and below the box indicate the 90th and 10th percentiles. Continuous line inside the box is the median; non-continuous line inside the box is the mean. Statistically significant differences respect to the control group were analysed by means of Student t-test (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Figure 3. Time course of plasma levels of GH (A) and IGF-I (B) of control (white) and stressed (black) gilthead sea bream. Data are the mean $\pm$ SEM (n = 6-8). Different letters indicate statistically significant changes over the course of the experiment in control (regular font) and stressed fish (italic font, ANOVA, $P < 0.05$). Statistically significant differences between stressed and control fish were analyzed at each sampling time by means of Student t-test (* $P < 0.05$, *** $P < 0.001$).

Figure 4. Box-Whisker plots representing the time course of the relative gene expression of IGF-I (A), and IGF-II (B) in stressed fish. Data in control fish were used as arbitrary reference values at each sampling time in the normalization procedure (values > 1 or < 1 indicate increase or decrease with respect to reference values). The lower boundary of the box indicates the 25th percentile and the upper boundary of the box indicates the 75th percentile. Whiskers above and below the box indicate the 90th and 10th percentiles. Continuous line inside the box is the median; non-continuous line inside the box is the mean. Statistically significant differences respect to the control group were analysed by means of Student t-test (* $P<0.05$, ** $P<0.01$).

Figure 5. Box-Whisker plots representing the time course of the relative gene expression of GHR-I (A), and GHR-II (B) in stressed fish. Data in control fish were used as arbitrary reference values at each sampling time in the normalization procedure (values > 1 or < 1 indicate increase or decrease with respect to reference values). The lower boundary of the box indicates the 25th percentile and the upper boundary of the box indicates the 75th percentile. Whiskers above and below the box indicate the 90th and 10th percentiles. Continuous line inside the box is the median; non-continuous line inside the box is the mean. Statistically significant differences respect to the control group were analysed by means of Student t-test (* $P<0.05$).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5