CHARACTERIZATION OF A NITROREDUCTASE WITH SELECTIVE
NITROREDUCTION PROPERTIES IN THE FOOD- AND INTESTINAL-
OCCURRING LACTIC ACID BACTERIUM *LACTOBACILLUS PLANTARUM* WCFS1

H. Guillén, J. Curiel, J.M. Landete, R. Muñoz, T. Herraiz*

Instituto de Fermentaciones Industriales, CSIC. Juan de la Cierva 3, 28006,
Madrid. Spain. Phone: 34915644853. Fax: 34915644853. E-mail:
therraiz@ifi.csic.es

Running title: A selective nitroreductase in *Lactobacillus plantarum*

* Corresponding author. T. Herraiz. E-mail: therraiz@ifi.csic.es

SUMMARY

Nitroreductases reduce nitroaromatic compounds and other oxidants in living organisms having interesting implications in environmental and human health. A putative nitrobenzoate reductase encoding gene (lp_0050) was recently annotated in the completed DNA sequence of lactic acid bacterium *Lactobacillus plantarum* WCFS1 strain. In this research, this *L. plantarum* gene was cloned, expressed and the corresponding protein (PnbA) biochemically characterized. This *L. plantarum* PnbA reductase is a 216 amino acid residues FMN-flavoprotein which exhibits a 23 % identity with *Pseudomonas putida* and *Ralstonia eutropha* nitroreductases, and less than 11 % with those from enterobacteria such as *E. cloacae*. This reductase also showed 32-43% identity (65-72 % similarity) to predicted PnbA proteins from other lactic acid bacteria. It utilized a wide range of electron acceptors including dichlorophenolindophenol (DCPIP), nitroblue tetrazolium (NBT), ferricyanide and quinones (menadione, benzoquinone), but not pyridinium cations (paraquat and N-methyl- β -carbolinium), and it was inhibited by dicoumarol and diphenyliodonium. HPLC-MS and spectroscopic data showed that it specifically catalyzed the reduction of 4-nitroaromatic group to the corresponding hydroxylamino in the presence of NAD(P)H. Kinetics parameters (V_{max} and K_m) showed a much higher efficiency to reduce 2,4-dinitrobenzoate than 4-nitrobenzoate. It was chemoselective for reduction of 4-nitrobenzoates being unable to reduce other nitroaromatics. Then, *L. plantarum* PnbA reductase and perhaps similar reductases in other lactic bacteria might be more specific than other microbial nitroreductases which reduce a wider range of nitroaromatic compounds. The physiological and functional role of nitroreductases remain unknown, however their presence in

lactic acid bacteria widely occurring in foods and the human intestinal tract should be of further interest.

INTRODUCTION.

Lactic acid bacteria are common in foods and the mammalian intestine. Among them, *Lactobacillus plantarum* is a versatile lactic acid bacterium encountered in a range of environmental niches that has a proven ability to survive and can colonize the intestinal tract of human and other mammals (12). The complete genome sequence of the *L. plantarum* WCFS1 has become recently available (24), making this bacteria a suitable model to explore the molecular mechanisms underlying the targeted intestinal properties of this species (12). From its complete genome sequence analysis, a sequence (lp_0050) putatively encoding a protein annotated as *p*-nitrobenzoate reductase (PnbA) was found, although it has not been further produced and/or characterized. Nitroreductases usually catalyze the reduction of nitrocompounds that are cytotoxic and mutagenic (2, 38). Despite their toxicity, some bacteria have been found to harbor enzymes for the metabolism of nitroaromatics, and they have been considered of great potential application in environmental bioremediation, biocatalysis, and as chemotherapeutic agents following the bioactivation of prodrugs (4, 45, 47, 52, 54, 55, 57).

Nitroreductases belong to two main classes: oxygen-insensitive (type I) and oxygen-sensitive (type II) (25). The former abounds in bacteria and fungus (11), and the latter in mammalian systems. Oxygen-sensitive nitroreductases usually catalyze one-electron reduction yielding a nitro anion radical (42) that reacts with oxygen to form superoxide radical regenerating the nitro group. In

this “futile cycle” pyridine nucleotides are oxidized without net reduction of the nitro group. In absence of oxygen further reduction of the nitroaromatic may yield nitroso, hydroxylamine and amino derivatives (42). Oxygen-insensitive nitroreductases are a family of bacterial FMN-containing enzymes that catalyzes the reduction of nitro compounds producing nitroso, hydroxylamino and/or amino derivatives (47, 49, 52). These nitroreductases have been studied in a number of bacteria including enterobacteria (e.g. *E. coli*, *S. typhimurium*, *E. cloacae*) , *Pseudomonas*, and *Rhodobacter capsulatus* (5, 6, 19, 41, 44, 46, 53, 58), and exhibit varying substrate specificity being able to reduce a wide range of nitroaromatics such as nitrophenols, nitrobenzenes and nitrobenzoates. So far, the physiological and functional role of microbial nitroreductases remain unknown (47), but it may include: a possible role in quinone (or oxidants) reduction and detoxification (46), oxidative stress (32, 33), bioluminescence (28), cobalamin synthesis (43), or even a function involving specific degradation pathways of nitrocompounds (20, 45, 47).

A large number of nitroaromatics are present in the environment because of their use in manufacturing processes, as antimicrobial agents or generated as by-products of combustion processes. Biodegradation of these compounds by using selected microorganisms and nitroreductases can be of great interest in biodegradation of environmental pollutants (7, 45, 47). Nitroaromatic compounds have also attracted a considerable health concern since their metabolization through reductive pathways may lead to formation of potent genotoxic and/or mutagenic metabolites (30, 34, 38). Indeed, nitro compounds are able to generate reactive nitrogen oxide species which readily react with biological molecules (50). Although the metabolic pathways of these

compounds are complex, nitroreductases appear to have a central role in their bioactivation (11, 14). Thus, enzymatic reduction by nitroreductases gives rise to reactive intermediates that can undergo nucleophilic additions with DNA and other macromolecules, suggesting a possible mechanism for their cytotoxicity (2).

Lactic acid bacteria occurring in mammalian intestinal microbiota and fermented foods might metabolize nitroaromatics having possible environmental, health and toxicological implications (12, 44). In this regard, the goal of this work was to study the biochemical and functional characteristics of a novel Pnb reductase from the lactic acid bacterium, *Lactobacillus plantarum* WCFS1. The gene encoding the PnbA reductase from this bacterium was cloned and the recombinant protein overexpressed in *Escherichia coli*. This functional nitroreductase was a FMN flavoenzyme able to reduce several electron acceptors, including various quinones but not pyridinium cations, and it was selective for the reduction of specific 4-nitroaromatic compounds, exhibiting a potent reduction for 4-nitrobenzoate and 2,4-dinitrobenzoate to 4-hydroxylamine metabolites. Then, this nitroreductase presents in lactic acid bacteria that occur both in foods and intestinal human microflora might play a role in the metabolism and reduction of nitroaromatics and oxidants.

MATERIAL AND METHODS

Chemicals and reagents. FMN, NADPH, NADH dicoumarol, diphenyliodonium chloride (DPI), dichlorophenolindophenol (DCPIP), ferricyanide, nitroblue tetrazolium (NBT), cytochrome C, menadione (2-methyl-1,4-naphthoquinone), 1,4-benzoquinone, p-nitrophenyl- β (D)-glucopyranoside and 1,1'-dimethyl-4,4'-

bipyridinium dichloride hydrate, 4-nitrobenzoic acid, 3-nitrobenzoic acid, 2-nitrobenzoic acid, 2,4-dinitrophenol, 5-nitroindazole and 2-amino-4-nitrophenol were obtained from Sigma-Aldrich, 2,4-dinitrobenzoic acid and nitrobenzene were from Fluka, 3,5-dinitrosalicylic acid from Merck, and picric acid from Probus. *N*-methyl- β -carbolinium iodide (2-methyl-9*H*-pyrido-(3,4-*b*)-indole) (*i.e.* 2-methylnorharmanium and 2-methylharmanium) were synthesized from norharman and harman with methyl iodide in acetone (16, 17). Several nitroderivatives such as *N*-(2-nitrophenyl)-*L*-proline, *N*-(2-nitrophenyl)pipecolic acid, *N*-(2,4-dinitrophenyl)glycine, *N*-(2-nitrophenyl)glycine were obtained as previously (9). *p*-Hydroxylaminobenzoic acid was synthesized from 4-nitrobenzoic acid by reduction with Zn in presence of ClNH_4 , (3). For that, 17 g of 4-nitrobenzoic acid were dissolved in water and sodium hydroxide. After addition of 20 g of ammonium chloride, the mixture was cooled and 15 g of zinc dust was added gradually and stirring. After one-half hour stirring the zinc sludge was filtered off. To the filtrate hydrochloric acid was added slowly and amorphous flakes which separated while the reaction was still neutral, were removed by filtration. Upon addition of more hydrochloric acid to the filtrate, white needles separated of *p*-hydroxylaminobenzoic acid that was characterized by spectral data. This compound was used for quantification using the corresponding calibration curves.

Bacterial strains, plasmids, and enzymes. The *L. plantarum* strain that has been completely sequenced, *L. plantarum* WCFS1 (NCIMB 8826), was used in this study. *E. coli* DH5 α was used for all DNA manipulations. *E. coli* JM109 (DE3) was used for expression in pURI3 vector (10). *L. plantarum* strain was

grown in MRS medium at 30 °C without shaking. *E. coli* strains were cultured in Luria-Bertani (LB) medium at 37 °C and 200 rpm. When required, ampicillin or chloramphenicol were added to the medium at a concentration of 100 or 34 µg/mL, respectively. Chromosomal DNA, plasmid purification, and transformation of *E. coli* were carried out as described elsewhere (48).

Expression and purification of *L. plantarum* PnbA. The gene coding for the putative *p*-nitrobenzoate reductase, *pnbA* (lp_0050) from *L. plantarum* WCFS1 has been cloned and overexpressed in pURI3 vector to avoid the enzyme and ligation steps during the cloning (10). Expression vector pURI3 was constructed based on the commercial expression vector pT7-7 (USB) but containing a leader sequence with a six-histidine affinity tag. The *pnbA* gene was PCR amplified with Hot-start Turbo *Pfu* DNA polymerase by using the primers 321 (5'- CATCATGGTGACGATGACGATAAGatggaacaattaaagcgattcaca) and 322 (5'- AAGCTTAGTTAGCTATTATGCGTAttagtgataatatgcaaaacttgcg) (the nucleotides pairing the expression vector sequence are indicated in italics, and the nucleotides pairing the *pnbA* gene sequence are written in lowercase letters). The 648 pb purified PCR product was inserted into the pURI3 vector by using a restriction enzyme- and ligation-free cloning strategy described previously (10). *E. coli* DH5α cells were transformed, recombinant plasmids were isolated and those containing the correct insert were identified by restriction-enzyme analysis, verified by DNA sequencing, and then transformed into *E. coli* JM109(DE3) (pLysS) cells for expression.

Cells carrying the recombinant plasmid, pURI3-pnbA, were grown at 37 °C in Luria-Bertani media containing ampicillin (100 µg/ml) and chloramphenicol

(34 µg/ml), until they reach an optical density at 600 nm of 0.4 and induced by adding IPTG (0.4 mM final concentration). After induction, the cells were grown at 22 °C during 20 h and collected by centrifugation. Cells were resuspended in 20 mM Tris-HCl, pH 8.0, containing 0.1 M NaCl. Crude extracts were prepared by French press lysis of the cell suspension (three times at 1100 psi). The insoluble fraction of the lysate was removed by centrifugation at 47,000g for 30 min at 4 °C.

The supernatant was filtered through a 0.45 µm filter and applied to a His-Trap-FF crude chelating affinity column (Amersham Biosciences) equilibrated with 20 mM Tris-HCl, pH 8.0, 100 mM NaCl containing imidazole, to improve the interaction specificity in the affinity chromatography step. The bound enzyme was eluted by applying a stepwise gradient of imidazole concentration, from 20 mM Tris-HCl, pH 8.0, 100 mM NaCl containing 10 mM imidazole to the same buffer but containing 500 mM imidazole. Fractions containing the His₆-tagged protein were pooled and dialysed overnight at 4 °C against 10 mM Tris-HCl, pH 8.0, containing 10 mM NaCl in a membrane (3,500 cutoff). The purity of the enzyme was determined by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) in Tris-glycine buffer. Protein concentration was measured according to the method of Bradford using a protein assay kit (Bio-Rad) with bovine serum albumin as standard.

Reductase assays with electron acceptors. Activity of PnbA reductase was monitored using several electron acceptors as substrates as follows: a) Reduction of dichlorophenolindophenol (DCPIP) was assayed in a reaction mixture containing 100 mM phosphate buffer (pH 7.4), 500 µM NADPH, 200 µM DCPIP

and PnbA reductase (1-5 $\mu\text{g/ml}$ protein) in a final volume of 0.5 ml. The reaction was initiated by the addition of substrate and the rate of DCPIP reduction with time was determined ($\lambda = 600 \text{ nm}$; $\epsilon = 21.5 \text{ mM}^{-1} \text{ cm}^{-1}$). Reaction rate was also calculated in absence of enzyme or NADPH as controls, and in presence of diphenyliodonium (0-10 mM) or dicoumarol (0-500 μM) used as inhibitors. Reduction of DCPIP was also used to determinate the activity in presence of NADH (500 μM) replacing NADPH as electron donor and to study the optimal temperature of reduction after incubating the enzyme for 20 min at different temperatures; b) Reduction of cytochrome C in presence or absence of menadione as an electron acceptor was measured using a reaction mixture containing 90 mM Tris-HCl buffer (pH 7.4), 500 μM NADPH, menadione (0 or 20 μM), 0.96 mg/ml of cytochrome C and reductase (1-5 $\mu\text{g/ml}$ protein) in a final volume of 0.5 ml. The activity of reductase was started by addition of NADPH and the reduction rate of cytochrome C at 550 nm was measured ($\epsilon = 21 \text{ mM}^{-1} \text{ cm}^{-1}$); c) Reduction of ferricyanide was measured by using a mixture containing 90 mM phosphate buffer (pH 7.4), 500 μM NADPH, 1mM potassium ferricyanide ($\text{K}_3[\text{Fe}(\text{CN})_6]$) and reductase enzyme (1-5 $\mu\text{g/ml}$ protein). Reduction of ferricyanide was measured at 420 nm ($\epsilon = 1020 \text{ M}^{-1} \text{ cm}^{-1}$). This assay was also used for determination of optimal pH using buffers adjusted at pHs between 2 and 9; d) Reduction of nitroblue tetrazolium (NBT) was monitored in a reaction mixture containing 80 mM phosphate buffer (pH 7.4), 500 μM NADPH, 200 μM NBT and enzyme (1-5 $\mu\text{g/m}$ protein) in a final volume of 0.5 ml. The reaction was started with the addition of substrate and monitored with time at 580 nm ($\epsilon = 7 \text{ mM}^{-1} \text{ cm}^{-1}$).

Reduction of nitroaromatics, quinones and pyridinium cations by *L. plantarum* PnbA reductase.

The activity of recombinant PnbA reductase from *L. plantarum* was assayed with different substrates, including a number of nitrocompounds, pyridinium cations and quinones by two different methods: HPLC-DAD-mass spectrometry and by following the spectrophotometric NADPH depletion. Thus, reaction mixtures with a final volume of 0.5 ml containing reductase enzyme (usually from 0.25 to 5 μ g/ml protein) , NADPH (500 μ M) and the corresponding substrate (from 0 to 1 mM) in 100 mM buffer phosphate pH 7.4 were mixed and incubated at room temperature for 3 min; the reaction stopped with the addition of 10% 2N HCl, and subsequently injected into HPLC and HPLC-MS to quantify and identify metabolites of *L. plantarum* reductase and to calculate the reaction rates. 4-Nitrobenzoic acid, 2,4-dinitrobenzoic acid, 2-nitrobenzoic acid, 4-hydroxylaminobenzoic acid, 2,4-dinitrophenol and 3-nitrobenzoic acid were tested as substrates for PnbA reductase. The corresponding controls without reductase and/or NADPH were also carried out. Kinetic values of K_m and V_{max} were determined by non-linear regression analysis fitting to Michaelis-Menten curves of reaction rates of *p*-hydroxylamino products as a function of the concentration of nitroaromatic substrates from 0 to 1mM. Enzyme reaction rate was calculated as μ moles of hydroxylamino product/min mg of protein. On the other hand, the activity of *L. plantarum* PnbA reductase was also measured by monitoring the decrease in absorbance of NADPH at 340 nm in a spectrophotometer for two 2-6 min in presence of various nitrocompounds or quinones as substrates. For that, reaction mixtures (0.5 ml) containing reductase enzyme (from 1 to 5 μ g/ml), NADPH (300 μ M) and substrate (50 or 100 μ M) in 100 mM phosphate buffer (pH 7.4) were added to quartz cuvettes. The reaction was

started by the addition of substrates and the reduction rate was calculated as the μ moles of NADPH oxidized/min mg of protein.

Reversed phase HPLC chromatographic analysis. The analysis of enzymatic reaction media was performed by RP-HPLC with UV diode array and fluorescence detection using a HPLC 1050 (Hewlett Packard) with a 1100 diode array detector (DAD) and a 1046A-fluorescence detector. A 150 mm x 3.9 mm i.d., 4 μ m, Nova-pak C18 column (Waters, Milford, MA, USA) was used for chromatographic separation. Chromatographic conditions were: 50 mM ammonium phosphate buffer (pH 3) (buffer A) and 20% of A in acetonitrile (buffer B). The gradient was programmed from 0% (100% A) to 32% B in 12 min, and 90% B at 15 min. The flow rate was 1 ml/min, the column temperature was 40 °C and the injection volume was 20 μ l. Absorbance detection was set at 280 nm (analysis of 4-hydroxylaminobenzoic acid) and 243 nm (analysis of 2-nitro-4-hydroxylaminobenzoic acid). Identification of compounds was performed by UV spectra, mass spectrometry and coinjection with synthetic standards. For quantification of metabolites, a calibration curve of synthesized 4-hydroxylaminobenzoic acid was constructed (0 to 50 μ M) whereas for quantification of 2-nitro-4-hydroxylaminobenzoic acid, the response factor at 243 nm of a calibration curve of 2,4-dinitrobenzoic acid was used because both substrate and product exhibited similar spectra.

Identification by HPLC-ESI-mass spectrometry. To carry out mass spectrometric identification, reactions media of *L. plantarum* PnbA reductase, nitroaromatics and NADPH obtained as above were analyzed on a 150 X 2.1 mm i.d. Zorbax SB-C18, 5 μ m, column (Agilent Technologies) by using a series 1100

275 HPLC-MSD (Hewlett-Packard) (electrospray-positive and negative ion mode).
276 Eluent A: acetic acid (0.5%); B: 20 % of A in acetonitrile. Gradient: 100 % A at 0
277 min and 60 % B at 20 min and 100 % B at 25 min, flow rate 0.25 ml/min; T: 40 °C;
278 mass range 50-500 *amu*, flow gas 11 l/min and T^a 330 °C, and cone voltage 55 V.
279 Pressure of nebulizador 40 psi and capilar voltage 4000 V.

280

281 **Spectrophotometric studies.** UV-VIS spectra were recorded with a Beckman
282 spectrophotometer. Samples with the purified reductase from *L. plantarum* were
283 placed in quartz cuvettes and spectra were determined in anaerobic conditions,
284 before and after the addition of an excess of sodium dithionite (Na₂S₂O₃). The
285 same apparatus was used for the determination of enzyme activities based on the
286 decrease of NADPH (340 nm) in presence of substrate at room temperature as
287 indicated above.

288

289 **RESULTS**

290 **Cloning, expression and purification of *L. plantarum* PnbA.** The deduced
291 product of *L. plantarum pnbA* is a protein of 216 amino acid residues, 23.9 kDa,
292 and pI of 6.65. This protein is included in the nitroreductase Pfam family
293 (PF00881) that contains proteins involved in the reduction of nitrogen
294 compounds (<http://pfam.sanger.ac.uk/family?acc=PF00881>). BLAST databases
295 searches of the translated *L. plantarum* DNA sequence showed high-scoring
296 similarities (32-43% identity) with putative *p*-nitrobenzoate reductases (PnbA)
297 from the complete sequenced lactic acid bacteria (data not shown), suggesting
298 that these enzymes may exhibit similar functional properties in different lactic
299 acid bacteria. The predicted sequence of the *L. plantarum* PnbA was aligned

with selected nitrobenzoate reductases which biochemical activity has been previously reported (Figure 1). From them, *L. plantarum* PnbA showed the highest overall identity (23%) with that of *Pseudomonas putida* and *Ralstonia eutropha*. *Enterobacter cloacae* PnbA showed only an 11% identity to *L. plantarum* protein. Other nitroreductases from enterobacteria such as *E. coli* and *S. typhimurium* also gave identities lower than 10 %. In contrast, identities among nitroreductases from enterobacteria are higher than 80 % (59). As deduced from Figure 1, all these proteins showed some of the residues conserved in this family, belonging to the characteristic nitroreductase motif. A 2-nitroreductase enzyme from *Pseudomonas fluorescens* (20) not included in the alignment was only 3-7% identical to the other nitroreductases, and it did not present the conserved residues described for this family. In conclusion, on the basis of its amino acid sequence, it could be assumed that PnbA (lp_0050) from *L. plantarum* WCFS1 is a member of the nitroreductase family.

Functional expression of *L. plantarum* PnbA. To confirm that the *pnbA* (lp_0050) gene from *L. plantarum* WCFS1 encodes a functional *p*-nitrobenzoate reductase (PnbA), we expressed this gene in *E. coli* under the control of the T7 RNA polymerase-inducible $\Phi 10$ promoter. Cell extracts were used to detect the presence of hyperproduced proteins by SDS-PAGE analysis. Control cells containing the pURI3 vector plasmid alone did not show expression over the time course analyzed, whereas expression of an additional protein was apparent with cells harboring pURI3-pnbA (Figure 2). The molecular mass of the recombinant protein corresponded to that inferred from the nucleotide sequence (23.9 KDa). As the gene was cloned containing a purification poly-His tag, PnbA was purified on a His-Trap- FF crude chelating column and eluted

with a stepwise gradient of imidazole. Highly purified PnbA protein was obtained from pURI3-pnbA (Figure 2) and fractions showing activity and usually eluting at 125 mM imidazole had yellow colour. The presence of the protein band in the gel of the elution fractions, correlated well with the occurrence of reductase activity using DCPIP (dichlorophenolindophenol) as electron acceptor in presence of NAD(P)H. The presence of the Histag had no apparent effect on catalytic activity. Thus, we could conclude that the *pnbA* (lp_0050) gene encoded a functional reductase in *L. plantarum* WCFS1. The eluted *L. plantarum* nitroreductase was dialyzed to eliminate the imidazole and the purified protein was used for its biochemical characterization.

The purified PnbA reductase from *L. plantarum* was yellow indicating the presence of a bound cofactor and displayed UV-visible absorption spectra characteristic of flavin-containing enzymes (Figure 3). Thus, spectra of the protein showed two bands (maxima at 366 and 446 nm) resembling an oxidized FMN cofactor which is usually present in bacterial nitroreductase whereas this profile quickly disappeared after reduction of the flavoenzyme with an excess of sodium dithionite as seen by the lost of absorbance at 446 nm and as expected for FMN enzymes. The enzyme appeared to lost part of the activity and colour with time suggesting that the coenzyme might not be tightly bound to the reductase (53). A further incubation of the protein with FMN appeared to partly recover activity (not shown).

Catalytic properties of PnbA reductase from *L. plantarum*. The catalytic properties of PnbA reductase from *L. plantarum* were studied with a number of electron acceptors in presence of NAD(P)H as electron donor (Table 1).

Nitroreductase from *L. plantarum* reduced DCPIP (absorbance at 600 nm) in presence of NADPH. It reduced cytochrome C in presence of menadione, whereas in the absence of menadione as an electron carrier and acceptor it was a poor reductant (7.5 times lower) of cytochrome C, suggesting that this enzyme was not a cytochrome P450-like reductase. Moreover, *L. plantarum* PnbA reductase also reduced nitroblue tetrazolium (NBT) to give formazan and was able to give electrons to the single electron acceptor ferricyanide. Therefore, various types of substrates were electron acceptors of *L. plantarum* PnbA reductase. As determined for the reduction of DCPIP, the enzyme could use both NADPH and NADH as electron donors although using NADPH as reductant was 1.8 times more efficient than NADH (500 μ M). The optimal pH was measured in the range of 2-9 and the best reduction rates (reduction of ferricyanide) was achieved in the interval pH 5 and 7 (highest value around pH 6). The optimal temperature (reduction of DCPIP) was 25 °C and the activity highly decreased after incubating the enzyme at temperatures higher than 37 °C for 20 min. Inhibition studies were carried out with dicoumarol and diphenyliodonium, that are inhibitors of quinone reductases and flavoproteins. Both compounds inhibited in a dose-dependent manner the PnbA reductase of *L. plantarum*. Dicoumarol inhibited reduction of DCPIP with a IC₅₀ of 0.15 mM whereas diphenyliodonium (DPI) inhibited it with IC₅₀ of 5.5 mM. Inhibition by DPI was expected as it is an inhibitor of flavoproteins (8), whereas inhibition by dicoumarol appears to need higher concentrations than those reported for NADPH:quinone reductases (18).

As shown in Table 1, several electron acceptors of reductases were good substrates of *L. plantarum* PnbA reductase in presence of NAD(P)H.

Subsequently, a number of nitroaromatics, quinones and naturally occurring and toxic pyridinium cations were studied as substrates and electron acceptors of this nitroreductase. The reduction rate was performed by monitoring the decrease of NADPH in presence of the corresponding substrate (Table 2). The pyridinium cations, paraquat and 2-methyl- β -carbolinium cations, were not reduced significantly by PnbA reductase. However, the quinones, menadione and benzoquinone, were good substrates of this enzyme with benzoquinone showing five times more activity than menadione. Among a range of nitroaromatic compounds containing nitro groups in different positions in the benzenic ring, including nitrophenols, nitrobenzene and nitrobenzoic acids, only two nitro compounds: 4-nitrobenzoate and 2,4-dinitrobenzoate were substrates of this enzyme as showed by an increased removal of NADPH (340 nm) when compared with a control in absence of the nitroaromatic substrate. The activity for reduction of 2,4-dinitrobenzoic acid was 12-fold higher than for 4-nitrobenzoate, suggesting that the former was a better substrate for *L. plantarum* PnbA reductase. From these results only specific nitro compounds could be reduced in presence of NADPH by PnbA reductase.

Identification of metabolites in the reaction mixtures by HPLC-MS. In order to characterize the reduction products of nitrocompounds, enzymatic reaction media containing *p*-nitrobenzoic acid, NADPH and *L. plantarum* PnbA reductase were studied by HPLC-DAD and MS (electrospray) (Figure 4). *p*-Hydroxylaminobenzoic acid was identified as a major metabolite in the reaction media containing *p*-nitrobenzoic acid as shown by HPLC-MS, with ions at m/z 154 ($M+H$)⁺, 137 ($M+1-17$) (positive-ESI) and m/z 152 ($M-H$)⁻ (negative-ESI)

(Figure 4a). This metabolite coeluted with a synthetic standard of *p*-hydroxylaminobenzoic acid and afforded similar absorption spectra (*max.* at 278 nm). In the reaction mixtures, a peak of *N*-nitrosobenzoic acid (negative-ESI) at *m/z* 150 (M-H)⁻ also appeared, suggesting that it could be an intermediate in the reaction. In addition, traces of *p*-aminobenzoic acid eluting after the corresponding *p*-hydroxylamine compound (*m/z* at 136 (M-H)⁻ and 138 (M+H)⁺ in negative and positive ion mode) were detected. This amine product was not increased in the reaction media suggesting that it was not a final product of PnbA reductase, and it might have appeared from chemical reduction. Indeed, no reduction was observed when synthetic *p*-hydroxylaminobenzoic acid was used as substrate of PnbA reductase in presence of NADPH. On the other hand, 2,4-dinitrobenzoic acid was also reduced in the reaction media by *L. plantarum* PnbA reductase and a major metabolite was identified by HPLC-MS (Figure 4b) as *p*-hydroxylamino-2-nitrobenzoic acid in ESI-positive at *m/z* 199 (M+H)⁺ and 181, and ESI-negative at *m/z* 197 (M-H)⁻ and 395 (2M-H)⁻. The two substrates in Figure 4 provided ESI-negative at *m/z* 166 (M-H)⁻, 122 (M-45)⁻ (4-nitrobenzoic acid), and *m/z* at 211 (M-H)⁻, 167 (M-45)⁻ and 423 (2M-1)⁻ (2,4-dinitrobenzoic acid).

In agreement with previous spectrophotometric results (Table 2), no reduction was observed by HPLC(DAD)-MS of 2,4-dinitrophenol, 2-nitrobenzoic or 3-nitrobenzoic acid, suggesting that PnbA reductase was selective for some specific nitrocompounds as electron acceptors and with regard to position of the nitro substituent in the benzenic ring. The reduction of nitro in *para* position respective of the carboxylic group was required since 2- or 3-nitrobenzoic acid were not reduced by *L. plantarum* PnbA reductase. As mentioned above 4-

hydroxylaminebenzoic acid (100 μ M) was not reduced to the corresponding amine by this reductase in presence of NADPH showing that the hydroxylamine was a final product of the catalytic reaction.

Enzyme reduction kinetics of PnbA reductase from *L. plantarum*. The reduction of 4-nitrobenzoic acid in presence of NADPH to give the corresponding hydroxylamine was rapid and linear in the first 10 min of incubation. Then, 3 min were subsequently used for kinetic studies by HPLC-DAD in order to calculate reaction rates and kinetic parameters. Figure 5 shows the Michaelis-Menten curves with the experimental reaction rates as a function of the concentration of substrate both for 4-nitrobenzoic acid and 2,4-dinitrobenzoic acid that gave 4-hydroxylaminobenzoic acid and 4-hydroxylamino-2-nitrobenzoic acid, respectively. Table 3 gives the kinetic values of $V_{\max}$, K_m and the ratio $V_{\max}/K_m$. The calculated $V_{\max}$ was 2.3 times higher for reduction of 2,4-dinitrobenzoic acid than for 4-nitrobenzoic acid, whereas the K_m was much lower for 2,4-dinitrobenzoic acid (32-fold) than for 4-nitrobenzoate. This suggests a much higher affinity of *L. plantarum* reductase for 2,4-dinitrobenzoate than for 4-nitrobenzoate. Thus, $V_{\max}/K_m$ ratio was much higher for 2,4-dinitrobenzoate, indicating a higher suitability of this specific nitrocompound for reduction by PnbA reductase. On the other hand, the reaction rate to give the *p*-hydroxylamine derivative from 4-nitrobenzoic acid was similar in aerobic and anaerobic conditions (not shown), suggesting that this reductase was not highly affected by the oxygen as expected for an oxygen-insensitive nitroreductase. By following the elimination of NADPH (absorbance) and the corresponding formation of hydroxylamine by HPLC, the

stoichiometry calculated was about 2 moles of NADPH consumed per mole of hydroxylamine.

DISCUSSION

L. plantarum is a lactic acid bacteria species encountered in a wide range of environmental niches and fermented foods, and with ability to survive gastric transit and colonize the intestinal tract of human and other mammals (11, 12). In this research, the gen (Ip_0050) encoding a functional nitrobenzoate reductase (Pnb A) from *L. plantarum* WCFS1 was cloned and expressed. This gen had been previously annotated as a PnbA reductase during its complete DNA sequencing (24). In addition, similar proteins annotated as PnbA reductases were found in the complete genomes of several lactic acid bacteria belonging to the *Lactobacillus* genera (*L. sakei*, *L. gasseri*, *L. reuteri*, *L. rhamnosus*, *L. jhonsonii* and *L. delbrueckii*) as well as *Lactococcus* (*L. lactis*) and *Pediococcus* (*P. pensosaceus*) genera (11, 35). However, none of these putative reductases had been previously cloned and/or biochemically characterized. The amino acid sequence of the PnbA reductase from *L. plantarum* showed a 23 % identity to known nitroreductases from *P. putida*, and *R. eutropha*, but less than 11 % to nitroreductases from enterobacteria such as *E. cloacae*, *E. coli*, and *S. typhimurium*. A recombinant plasmid containing the *pnbA* gen was cloned into *E. coli* and the encoded protein showed to be a catalytically active *L. plantarum* PnbA reductase, resembling kinetic properties and inhibitor susceptibility of flavin reductases. This enzyme was a FMN-flavoprotein (yellow colour) that used both NADPH and NADH as electron donors, and it was inhibited by dicoumarol and DPI. It could use a broad spectrum of electron acceptors such

as DCPIP, NBT, ferricyanide and quinones suggesting that its active site could accommodate molecules of varying size and structure as substrates. In contrast, it was more selective regarding the reduction of nitrocompounds, with only 4-nitrobenzoates being reduced which suggests a higher selectivity for two electron reduction of nitroaromatic groups. Moreover, *L. plantarum* PnbA reductase was not able to reduce potentially toxic pyridinium cations such as the herbicide paraquat and the naturally-occurring 2-methyl- β -carbolinium alkaloids. These compounds might generate free radicals and superoxide anion if reduced contributing to their toxicity (15-17, 31).

As occur with other characterized nitroreductases, the natural substrates of *L. plantarum* nitroreductase and its physiological and/or metabolic role in lactic bacteria are unknown (47). Since NAD(P)H is used as an electron donor, it has been speculated that nitroreductases could link pyrimidine nucleotide oxidation to the reduction of substrates to maintain the steady supply of oxidized pyrimidine nucleotides. *L. plantarum* PnbA reductase has the ability to utilize a variety of electron acceptors such as quinones and other substrates; thus, it could remove oxidants which otherwise would be reduced to a free-radical state and form intermediate species reacting with DNA and proteins. This raises the possibility that nitroreductases could somehow protect cells against oxidative stress and the toxic effects caused by exposure to quinones and/or related cellular oxidants (21, 32, 46). On the other hand, nitroreductases could function as specific enzymes involved in degradation pathways of nitrocompounds (47). In this regard, biodegradation of these compounds by using selective nitroreductases and microorganisms are of great interest for removal of environmental pollutants (45, 47). On the other hand, these

nitroreductases might be also of interest in the therapeutic action of antimicrobial and anticancer prodrugs (4, 13, 55).

Living organisms can utilize nitroaromatic compounds which are metabolized by nitroreduction catalyzed by two types of nitroreductases: oxygen-sensitive and oxygen-insensitive (25, 57). Oxygen-sensitive such as cytochrome P-450 oxidoreductase may catalyze the one-electron reduction of nitro to the anion free radical which is reoxidized to the parent compounds by O₂ in a futile redox cycle producing superoxide anion. Oxygen-insensitive such as NAD(P)H-quinone oxidoreductase and nitroreductases of enteric bacteria (*E. coli*, *E. cloacae*, *Salmonella*) catalyze the two-electron reduction of the nitro group to nitroso, hydroxylamine and/or amine derivatives (26, 39). The functional PnbA from *L. plantarum* belongs to this latter family of nitroreductases and reduced *p*-nitrobenzoates (4-nitrobenzoates and 2,4-dinitrobenzoate) initially to 4-nitroso compounds which were rapidly converted to hydroxylamino derivatives as final products (Figure 6). The enzyme was not able to reduce the hydroxylamine further to *p*-aminobenzoate. This behaviour agrees well with other microbial nitroreductases (39) that afford nitrosoaromatic as a limiting step that quickly react to give hydroxylamine products (29, 49).

Nitroheterocyclic compounds constitute a wide range of chemicals whose potent biological activity has significant human health and environmental implications. Many nitrocompounds are well-known as potent mutagens and carcinogens (1, 37, 38). The biological activity of these compounds is linked to their reductive metabolism catalyzed by nitroreductases to hydroxylamine, nitroso and electrophilic species (30, 34, 51). Bacterial nitroreductases such as that one of *L. plantarum* perform two-electron reduction of nitroaromatic

compounds to hydroxylamine derivatives (6, 19, 36, 56). They are distributed among different microorganisms suggesting a certain relevance within microbial metabolic activities (11, 47). Nevertheless, certain aspects of their catalytic process such as substrate specificity and reduction mechanism are not well understood. Thus, nitroreductases from *E. cloacae* (6), *S. typhimurium* (53), *E. coli* (22, 46, 58), *Ralstonia eutropha* (49), *Rhodobacter capsulatus* (40, 41), *Pseudomonas* (7, 19, 27) are able to reduce a wide range of nitroaromatics including nitroalkylbenzenes, nitrophenols, nitrobenzoates, nitrofurazone, and *p*-nitroacetophenone, among others. Then, they appear to be relaxed on substrate specificity and can reduce nitrocompounds and other electron acceptors such as quinones (23, 36, 46). In this regard, the PnbA reductase from *L. plantarum* was selective for reduction of nitrocompounds, and only reduced those with a 4-nitrobenzoate moiety such as 4-nitrobenzoate and 2,4-dinitrobenzoate. The efficiency ($V_{\max}/K_m$) for reduction of 2,4-dinitrobenzoate was much higher than for 4-nitrobenzoate, whereas it was unable to reduce other nitrobenzoates or nitrophenols. This might suggest that this PnbA reductase, and perhaps other nitroreductases annotated in lactic acid bacteria (11, 35), may conform a class of more selective 4-nitrobenzoate nitroreductases. Moreover, this reductase might have similar functional properties in different lactic acid bacteria as it shows a high-scoring similarity (32-43% identity) to other putative *p*-nitrobenzoate reductases from lactic bacteria.

In conclusion, we have cloned, expressed and characterized a novel *p*-nitrobenzoate (PnbA) reductase from *L. plantarum* WCFS1 that was previously annotated on its complete genome sequence. Similar proteins seem to be

present in several other lactic acid bacteria. The PnbA reductase of *L. plantarum* utilized a wide range of electron acceptors including quinones, DCPIP, NBT and ferricyanide but it was not able to reduce pyridinium cations such as paraquat and 2-methyl- β -carbolinium substances. However, it was selective for nitro compounds reducing only 4-nitrobenzoates (4-nitrobenzoate and 2,4-dinitrobenzoate) among a number of nitrocompounds assayed. This PnbA nitroreductase may be distinct to other previously described bacterial nitroreductases because of its selectivity for 4-nitrobenzoates. This different selectivity could be due to that *L. plantarum* PnbA reductase exhibits a 23 % identity to nitroreductases from *Pseudomonas putida* and *R. eutropha* and even was less identical (< 11 %) to enterobacteria PnbA (e.g. *E. cloacae*, *E. coli* and *S. typhimurium*). This report is the first characterization of a nitroreductase in lactic acid bacteria that widely occur in foods, the environment and human intestinal tract, and thereby these results might be of further interest.

ACKNOWLEDGEMENTS

The authors thank the MCINN (Spanish Government) for the support of this research through the grants AGL2006-02414, AGL2009-09917, and AGL2008-01052. H. Guillén and J. A. Curiel thank CSIC and MICINN for a JAE predoctoral and a FPI predoctoral fellowships, respectively.

REFERENCES

1. Andre, V., C. Boissart, F. Sichel, P. Gauduchon, J. Y. LeTalaer, J. C. Lancelot, C. Mercier, S. Chemtob, E. Raoult, and A. Tallec. 1997. Mutagenicity of nitro- and amino-substituted carbazoles in *Salmonella*

577 *typhimurium*.3. Methylated derivatives of 9*H*-carbazole. Mutat. Res. Genet.
578 Toxicol. Environ. Mutagen. **389**:247-260.

579 2. **Bartel, L. C., M. M. de Mecca, and J. A. Castro.** 2009. Nitroreductive
580 metabolic activation of some carcinogenic nitro heterocyclic food
581 contaminants in rat mammary tissue cellular fractions. Food Chem.
582 Toxicol. **47**:140-144.

583 3. **Bauer, H., and S. M. Rosenthal.** 1944. 4-
584 hydroxylaminobenzenesulfonamide, its acetyl derivatives and diazotization
585 reaction. J. Am. Chem. Soc. **66**:611-614.

586 4. **Benouchan, M., F. Do Nascimento, G. Y. Perret, and B. M. Colombo.**
587 2006. Delivery of the bacterial nitroreductase gene into endothelial cells
588 prolongs the survival of tumour-bearing mice by bystander mechanisms.
589 Int. J. Oncol. **28**:457-462.

590 5. **Blasco, R., and F. Castillo.** 1993. Characterization of a nitrophenol
591 reductase from the phototrophic bacterium *Rhodobacter capsulatus* E1f1.
592 Appl. Environ. Microbiol. **59**:1774-1778.

593 6. **Bryant, C., and M. Deluca.** 1991. Purification and characterization of an
594 oxygen-insensitive NAD(P)H nitroreductase from *Enterobacter cloacae*. J.
595 Biol. Chem. **266**:4119-4125.

596 7. **Caballero, A., J. J. Lazaro, J. L. Ramos, and A. Esteve-Nuñez.** 2005.
597 PnrA, a new nitroreductase family enzyme in the TNT-degrading strain
598 *Pseudomonas putida* JLR11. Environ. Microbiol. **7**:1211-1219.

599 8. **Chakraborty, S., and V. Massey.** 2002. Reaction of reduced flavins and
600 flavoproteins with diphenyliodonium chloride. J. Biol. Chem. **277**:41507-
601 41516.

9. **Chicharro, R., S. de Castro, J. L. Reino, and V. J. Aran.** 2003. Synthesis of tri- and tetracyclic condensed quinoxalin-2-ones fused across the C-3-N-4 bond. *Eur. J. Org. Chem.* 2314-2326.
10. **de las Rivas, B., J. A. Curiel, J. M. Mancheno, and R. Muñoz.** 2007. Expression vectors for enzyme restriction and ligation-independent cloning for producing recombinant his-fusion proteins. *Biotechnol. Prog.* **23**:680-686.
11. **de Oliveira, I. M., J. A. P. Henriques, and D. Bonatto.** 2007. In silico identification of a new group of specific bacterial and fungal nitroreductases-like proteins. *Biochem. Biophys. Res. Commun.* **355**:919-925.
12. **de Vries, M. C., E. E. Vaughan, M. Kleerebezem, and W. M. de Vos.** 2006. *Lactobacillus plantarum* survival, functional and potential probiotic properties in the human intestinal tract. *Int. Dairy J.* **16**:1018-1028.
13. **Emptage, C. D., R. J. Knox, M. J. Danson, and D. W. Hough.** 2009. Nitroreductase from *Bacillus licheniformis*: A stable enzyme for prodrug activation. *Biochem. Pharmacol.* **77**:21-29.
14. **Goodwin, A., D. Kersulyte, G. Sisson, S. van Zanten, D. E. Berg, and P. S. Hoffman.** 1998. Metronidazole resistance in *Helicobacter pylori* is due to null mutations in a gene (rdxA) that encodes an oxygen-insensitive NADPH nitroreductase. *Mol. Microbiol.* **28**:383-393.
15. **Herraiz, T., H. Guillen, and V. J. Aran.** 2008. Oxidative metabolism of the bioactive and naturally occurring β -carboline alkaloids, norharman and harman, by human cytochrome P450 Enzymes. *Chem. Res. Toxicol.* **21**:2172-2180.

- 627 16. **Herraiz, T., H. Guillén, V. J. Arán, J. R. Idle, and F. J. Gonzalez.** 2006.
628 Comparative aromatic hydroxylation and *N*-demethylation of MPTP
629 neurotoxin and its analogs, *N*-methylated β -carboline and isoquinoline
630 alkaloids, by human cytochrome P450 2D6. *Toxicol. Appl. Pharmacol.*
631 **216**:387-398.
- 632 17. **Herraiz, T., H. Guillen, and J. Galisteo.** 2007. *N*-methyltetrahydro- β -
633 carboline analogs of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)
634 neurotoxin are oxidized to neurotoxic β -carbolinium cations by heme
635 peroxidases. *Biochem. Biophys. Res. Commun.* **356**:118-123.
- 636 18. **Hodnick, W. F., and A. C. Sartorelli.** 1997. Measurement of dicumarol-
637 sensitive NADPH:(menadione cytochrome c) oxidoreductase activity
638 results in an artifactual assay of DT-diaphorase in cell sonicates. *Anal.*
639 *Biochem.* **252**:165-168.
- 640 19. **Hughes, M. A., and P. A. Williams.** 2001. Cloning and characterization of
641 the *pnb* genes, encoding enzymes for 4-nitrobenzoate catabolism in
642 *Pseudomonas putida* TW3. *J. Bacteriol.* **183**:1225-1232.
- 643 20. **Iwaki, H., T. Muraki, S. Ishihara, Y. Hasegawa, K. N. Rankin, T. Sulea,**
644 **J. Boyd, and P. C. K. Lau.** 2007. Characterization of a pseudomonad 2-
645 nitrobenzoate nitroreductase and its catabolic pathway-associated 2-
646 hydroxylaminobenzoate mutase and a chemoreceptor involved in 2-
647 nitrobenzoate chemotaxis. *J. Bacteriol.* **189**:3502-3514.
- 648 21. **Jaiswal, A. K.** 2000. Characterization and partial purification of
649 microsomal NAD(P)H : quinone oxidoreductases. *Arch. Biochem. Biophys.*
650 **375**:62-68.

22. **Kadiyala, V., L. J. Nadeau, and J. C. Spain.** 2003. Construction of *Escherichia coli* strains for conversion of nitroacetophenones to ortho-aminophenols. *Appl. Environ. Microbiol.* **69**:6520-6526.
23. **Kitts, C. L., C. E. Green, R. A. Otley, M. A. Alvarez, and P. J. Unkefer.** 2000. Type I nitroreductases in soil enterobacteria reduce TNT (2,4,6-trinitrotoluene) and RDX (hexahydro-1,3,5-trinitro-1,3,5-triazine). *Can. J. Microbiol.* **46**:278-282.
24. **Kleerebezem, M., J. Boekhorst, R. van Kranenburg, D. Molenaar, O. P. Kuipers, R. Leer, R. Tarchini, S. A. Peters, H. M. Sandbrink, M. Fiers, W. Stiekema, R. M. K. Lankhorst, P. A. Bron, S. M. Hoffer, M. N. N. Groot, R. Kerkhoven, M. de Vries, B. Ursing, W. M. de Vos, and R. J. Siezen.** 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. U. S. A.* **100**:1990-1995.
25. **Kobori, T., H. Sasaki, W. C. Lee, S. Zenno, K. Saigo, M. E. P. Murphy, and M. Tanokura.** 2001. Structure and site-directed mutagenesis of a flavoprotein from *Escherichia coli* that reduces nitrocompounds. Alteration of pyridine nucleotide binding by a single amino acid substitution. *J. Biol. Chem.* **276**:2816-2823.
26. **Koder, R. L., C. A. Haynes, M. E. Rodgers, D. W. Rodgers, and A. F. Miller.** 2002. Flavin thermodynamics explain the oxygen insensitivity of enteric nitroreductases. *Biochemistry* **41**:14197-14205.
27. **Lee, B. U., S. C. Park, Y. S. Cho, H. Y. Kahng, and K. H. Oh.** 2008. Expression and characterization of the TNT nitroreductase of *Pseudomonas* sp HK-6 in *Escherichia coli*. *Curr. Microbiol.* **56**:386-390.

- 675 28. **Lei, B. F., M. Y. Liu, S. Q. Huang, and S. C. Tu.** 1994. *Vibrio harveyi*
676 NADPH-flavin oxidoreductase: cloning, sequencing and overexpression of
677 the gene and purification and characterization of the cloned enzyme. J.
678 Bacteriol. **176**:3552-3558.
- 679 29. **Leskovac, V., J. Svircevic, S. Trivic, M. Popovic, and M. Radulovic.**
680 1989. Reduction of aryl nitroso-compounds by pyridine and flavin
681 coenzymes. Int. J. Biochem. **21**:825-834.
- 682 30. **Letelier, M. E., P. Iquierdo, L. Godoy, A. M. Lepe, and M. Faundez.**
683 2004. Liver microsomal biotransformation of nitro-aryl drugs: Mechanism
684 for potential oxidative stress induction. J. Appl. Toxicol. **24**:519-525.
- 685 31. **Liochev, S. I., A. Hausladen, W. F. Beyer, and I. Fridovich.** 1994.
686 NADPH-ferredoxin oxidoreductase acts as a paraquat diaphorase and is a
687 member of the soxRS regulon. Proc. Natl. Acad. Sci. U. S. A. **91**:1328-
688 1331.
- 689 32. **Liochev, S. I., A. Hausladen, and I. Fridovich.** 1999. Nitroreductase A is
690 regulated as a member of the soxRS regulon of *Escherichia coli*. Proc.
691 Natl. Acad. Sci. U. S. A. **96**:3537-3539.
- 692 33. **Liu, G. F., J. T. Zhou, R. F. Jin, M. Zhou, J. Wang, H. Lu, and Y. Y. Qu.**
693 2008. Enhancing survival of *Escherichia coli* by expression of
694 azoreductase AZR possessing quinone reductase activity. Appl. Microbiol.
695 Biotechnol. **80**:409-416.
- 696 34. **Maeda, T., R. Nakamura, K. Kadokami, and H. I. Ogawa.** 2007.
697 Relationship between mutagenicity and reactivity or biodegradability for
698 nitroaromatic compounds. Environ. Toxicol. Chem. **26**:237-241.

- 699 35. **Makarova, K., A. Slesarev, Y. Wolf, A. Sorokin, B. Mirkin, E. Koonin, A.**
700 **Pavlov, N. Pavlova, V. Karamychev, N. Polouchine, V. Shakhova, I.**
701 **Grigoriev, Y. Lou, D. Rohksar, S. Lucas, K. Huang, D. M. Goodstein, T.**
702 **Hawkins, V. Plengvidhya, D. Welker, J. Hughes, Y. Goh, A. Benson, K.**
703 **Baldwin, J. H. Lee, I. Diaz-Muniz, B. Dosti, V. Smeianov, W. Wechter,**
704 **R. Barabote, G. Lorca, E. Altermann, R. Barrangou, B. Ganesan, Y.**
705 **Xie, H. Rawsthorne, D. Tamir, C. Parker, F. Breidt, J. Broadbent, R.**
706 **Hutkins, D. O'Sullivan, J. Steele, G. Unlu, M. Saier, T. Klaenhammer,**
707 **P. Richardson, S. Kozyavkin, B. Weimer, and D. Mills. 2006.**
708 Comparative genomics of the lactic acid bacteria. *Proc. Natl. Acad. Sci. U.*
709 *S. A.* **103**:15611-15616.
- 710 36. **Nivinskas, H., S. Staskeviciene, J. Sarlauskas, R. L. Koder, A. F.**
711 **Miller, and N. Cenas. 2002. Two-electron reduction of quinones by**
712 *Enterobacter cloacae* NAD(P)H : nitroreductase: quantitative structure-
713 activity relationships. *Arch. Biochem. Biophys.* **403**:249-258.
- 714 37. **Ostergaard, T. G., L. H. Hansen, M. L. Binderup, A. Norman, and S. J.**
715 **Sorensen. 2007. The cda GenoTox assay: A new and sensitive method**
716 **for detection of environmental genotoxins, including nitroarenes and**
717 **aromatic amines. Mutat. Res. Genet. Toxicol. Environ. Mutagen.** **631**:77-
718 **84.**
- 719 38. **Padda, R. S., C. Y. Wang, J. B. Hughes, R. Kutty, and G. N. Bennett.**
720 **2003. Mutagenicity of nitroaromatic degradation compounds. Environ.**
721 **Toxicol. Chem.** **22**:2293-2297.
- 722 39. **Peres, C. M., R. Russ, H. Lenke, and S. N. Agathos. 2001.**
723 **Biodegradation of 4-nitrobenzoate, 4-aminobenzoate and their mixtures:**

new strains, unusual metabolites and insights into pathway regulation.
FEMS Microbiol. Ecol. **37**:151-159.

40. **Perez-Reinado, E., R. Blasco, F. Castillo, C. Moreno-Vivian, and M. D. Roldan.** 2005. Regulation and characterization of two nitroreductase genes, *nprA* and *nprB*, of *Rhodobacter capsulatus*. Appl. Environ. Microbiol. **71**:7643-7649.

41. **Perez-Reinado, E., M. D. Roldan, F. Castillo, and C. Moreno-Vivian.** 2008. The NprA nitroreductase required for 2,4-dinitrophenol reduction in *Rhodobacter capsulatus* is a dihydropteridine reductase. Environ. Microbiol. **10**:3174-3183.

42. **Peterson, F. J., R. P. Mason, J. Hovsepian, and J. L. Holtzman.** 1979. Oxygen-sensitive and oxygen-insensitive nitroreduction by *Escherichia coli* and rat hepatic microsomes. J. Biol. Chem. **254**:4009-4014.

43. **Pollich, M., and G. Klug.** 1995. Identification and sequence analysis of genes involved in late steps of cobalamin (Vitamin B-12) synthesis in *Rhodobacter capsulatus*. J. Bacteriol. **177**:4481-4487.

44. **Rafii, F., W. Franklin, R. H. Heflich, and C. E. Cerniglia.** 1991. Reduction of nitroaromatic compounds by anaerobic bacteria isolated from the human gastrointestinal tract. Appl. Environ. Microbiol. **57**:962-968.

45. **Ramos, J. L., M. M. Gonzalez-Perez, A. Caballero, and P. van Dillewijn.** 2005. Bioremediation of polynitrated aromatic compounds: plants and microbes put up a fight. Curr. Opin. Biotechnol. **16**:275-281.

46. **Rau, J., and A. Stolz.** 2003. Oxygen-insensitive nitroreductases NfsA and NfsB of *Escherichia coli* function under anaerobic conditions as lawsone-dependent azo reductases. Appl. Environ. Microbiol. **69**:3448-3455.

- 749 47. **Roldan, M., E. Perez-Reinado, F. Castillo, and C. Moreno-Vivian.** 2008.
750 Reduction of polynitroaromatic compounds: the bacterial nitroreductases.
751 FEMS Microbiol. Rev. **32**:474-500.
- 752 48. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. Molecular cloning: A
753 laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold
754 Spring Harbor, New York, NY.
- 755 49. **Schenzle, A., H. Lenke, J. C. Spain, and H. J. Knackmuss.** 1999.
756 Chemoselective nitro group reduction and reductive dechlorination initiate
757 degradation of 2-chloro-5-nitrophenol by *Ralstonia eutropha* JMP134.
758 Appl. Environ. Microbiol. **65**:2317-2323.
- 759 50. **Sies, H., and H. Degroot.** 1992. Role of reactive oxygen species in cell
760 toxicity. Toxicol. Lett. **64-5**:547-551.
- 761 51. **Silvers, K. J., L. H. Couch, E. A. Rorke, and P. C. Howard.** 1997. Role
762 of nitroreductases but not cytochromes P450 in the metabolic activation of
763 1-nitropyrene in the HepG2 human hepatoblastoma cell line. Biochem.
764 Pharmacol. **54**:927-936.
- 765 52. **Spain, J. C.** 1995. Biodegradation of nitroaromatic compounds. Annu.
766 Rev. Microbiol. **49**:523-555.
- 767 53. **Watanabe, M., T. Nishino, K. Takio, T. Sofuni, and T. Nohmi.** 1998.
768 Purification and characterization of wild-type and mutant "Classical"
769 nitroreductases of *Salmonella typhimurium* L33R mutation greatly
770 diminishes binding of FMN to the nitroreductase of *S. typhimurium*. J. Biol.
771 Chem. **273**:23922-23928.
- 772 54. **White, C. L., T. Menghistu, K. R. Twigger, P. F. Searle, S. A. Bhide, R.**
773 **G. Vile, A. A. Melcher, H. S. Pandha, and K. J. Harrington.** 2008.

Escherichia coli nitroreductase plus CB1954 enhances the effect of radiotherapy in vitro and in vivo. Gene Ther. **15**:424-433.

55. **Wilkinson, S. R., M. C. Taylor, D. Horn, J. M. Kelly, and I. Cheeseman.** 2008. A mechanism for cross-resistance to nifurtimox and benznidazole in trypanosomes. Proc. Natl. Acad. Sci. U. S. A. **105**:5022-5027.

56. **Yabannavar, A. V., and G. J. Zylstra.** 1995. Cloning and characterization of the genes for p-nitrobenzoate degradation from *Pseudomonas pickettii* YH105. Appl. Environ. Microbiol. **61**:4284-4290.

57. **Ye, J., A. Singh, and O. P. Ward.** 2004. Biodegradation of nitroaromatics and other nitrogen-containing xenobiotics. World J. Microbiol. Biotechnol. **20**:117-135.

58. **Zenno, S., H. Koike, A. N. Kumar, R. Jayaraman, M. Tanokura, and K. Saigo.** 1996. Biochemical characterization of NfsA, the *Escherichia coli* major nitroreductase exhibiting a high amino acid sequence homology to Frp, a *Vibrio harveyi* flavin oxidoreductase. J. Bacteriol. **178**:4508-4514.

59. **Zenno, S., H. Koike, M. Tanokura, and K. Saigo.** 1996. Gene cloning, purification, and characterization of NfsB, a minor oxygen-insensitive nitroreductase from *Escherichia coli*, similar in biochemical properties to FRase I, the major flavin reductase in *Vibrio fischeri*. J. Biochem. (Tokyo) **120**:736-744.

Table 1. Reduction rates (μmol of substrate/min mg PnbA protein) of several electron acceptors by PnbA reductase from *Lactobacillus plantarum* WCFS1.

Electron acceptor	v ($\mu\text{mol}/\text{min mg protein}$) ¹
DCPIP	6.6 ± 0.065
Cytochrome C	0.2 ± 0.03
Cytochrome C/menadione	1.51 ± 0.05
Ferricyanide	158 ± 1.4
NBT	1.9 ± 0.09

¹ $v \pm \text{SD}$. Assays were carried out as indicated in experimental and performed at least in triplicate.

Table 2. Activity of PnbA reductase from *L. plantarum* WCFS1 strain with different nitrocompounds, pyridinium cations and quinones as substrates.

Substrate	v ($\mu\text{mol NADPH}/\text{min mg prot}$) ¹
1,4-benzoquinone	12.3 ± 3.0
Menadione	2.26 ± 0.24
4-nitrobenzoic acid	2.92 ± 0.8
2,4-dinitrobenzoic acid	36.1 ± 6.1
2-nitrobenzoic acid	ND
3-nitrobenzoic acid	ND
2-hydroxy-3,5-dinitrobenzoic acid	ND
2,4-dinitrophenol	ND
2,4,6-trinitrophenol (picric acid)	ND
2-amino-4-nitrophenol	ND
Nitrobenzene	ND
5-nitroindazole	ND
4-nitrophenyl- β -D-glucopiranoside	ND
N-(2-nitrophenyl)proline	ND
N-(2-nitrophenyl)pipecolic acid	ND
N-(2,4-dinitrophenyl)glycine	ND
N-(2-nitrophenyl)glycine	ND
1,1-dimethyl-1,4-bypyridyl	ND
2-methyl-norharman	ND
2-methyl-harman	ND

¹The enzymatic activity ($v \pm \text{SD}$) was measured as indicated in Experimental section in presence of enzyme (1-5 $\mu\text{g}/\text{ml}$, 300 μM NADPH, and 100 μM substrate with the exception of menadione and 2,4-dinitrophenol that were assayed at 50 μM to avoid signal saturation. The rate was measured as the decrease of NADPH (340 nm) in the first 1-2 min reaction at room temperature and the NADPH reduction in absence of substrate was subtracted. ND, not detectable activity.

814 Table 3. Kinetic parameters ($V_{\max}$ and K_m) of the reduction of 4-nitrobenzoic
 815 acid and 2,4-dinitrobenzoic acid by *L. plantarum* PnbA reductase.
 816

Substrate	K_m (μM) $\pm$ SE	$V_{\max} \pm$ SE ($\mu\text{mol}/\text{min mg protein}$)	$V_{\max}/K_m$ (l min^{-1} mg prot^{-1})
4-nitrobenzoic acid	536 ± 102	15.1 ± 1.6	0.028
2,4-dinitrobenzoic acid	16.69 ± 6.4	35.5 ± 1.99	2.13

FIGURE CAPTIONS

Figure 1. Comparison of PnbA reductase from *L. plantarum* WCFS1 (LPL) (accession CAD62742) and well-known PnbA reductase from *P. putida* (PPU) (AAG01540), *R. euthropha* (REU) (YP_297809), and *E. cloacae* (ECO) (Q01234). The ClustalW program was used to compare sequences. Conserved residues of the nitroreductase family (Pfam 00881) are indicated (●). Residues that are identical (*), conserved (:), or semiconserved (.) in all sequences of the alignment are also indicated. Dashes represent gaps introduced to maximize similarities.

Figure 2. SDS-PAGE analysis showing the PnbA expression and PnbA purification from *L. plantarum* WCFS1 strain. Lanes: 1, soluble extract from *E. coli* JM109 (pURI3) cells; 2, extract from *E. coli* JM109 (pURI3-PnbA); 3, extract from *E. coli* JM109 (pURI3-PnbA) not retained on the affinity His-Trap FF column; 4, purified PnbA reductase protein retained on the affinity column and eluted on 125 mM imidazole buffer. SDS-polyacrylamide gels were stained with Coomassie blue. The position of some molecular mass markers (Bio-Rad) are indicated on the left.

Figure 3. Absorption spectra of the purified flavoprotein PnbA from *L. plantarum* WCFS1 (a), and loss of the FMN bands following reduction with an excess of sodium dithionite (b).

Figure 4. HPLC-MS (ESI) analysis of the reaction media of 4-nitrobenzoic acid (280 nm) (a), and 2,4-dinitrobenzoic acid (243 nm) (b) reduction with *L. plantarum* PnbA reductase. Electrospray spectra of in negative ion mode of the corresponding hydroxylamine products are given. Reaction media in 100 mM phosphate buffer pH 7.4, contained substrate (100 μ M) NADPH (500 μ M), PnbA reductase (5 μ g/ml) and were incubated at room temperature and reaction stopped as indicated in Experimental.

Figure 5. Michaelis-Menten curves of the enzymatic reduction of 4-nitrobenzoic acid (a) and 2,4-dinitrobenzoic acid (b) by PnbA reductase of *L. plantarum*.

Reaction media contained 4-nitrobenzoic acid or 2,4-dinitrobenzoic acid in 100 mM phosphate buffer pH 7.4, NADPH (500 μ M), and PnbA reductase (0.25 μ g/ml for 2,4-dinitrobenzoic acid and 2 μ g/ml for 4-nitrobenzoic acid). Reaction was performed at room temperature for 3 min. Velocity (v) is given as μ mol of corresponding hydroxylamine product RNHOH/min mg of PnbA reductase.

Figure 6. Selective reduction of 4-nitrobenzoic acid compounds to nitroso and hydroxylamino derivatives by PnbA reductase from the lactic acid bacterium *L. plantarum* WCFS1.

862

863

PPU MALLTDDFDFAVVASRRRAVRAFLP--TPISRKLISEIIDRIARLAPSNSNTQPWSIHVLTGEP 59
 REU -----MKVSQAVESRKSVRGFLP-NPIDPDTIRRVLAAAARAPSGGNLQPWHIHVVGGEA 54
 LPL -----METIKAIHTRHSVRAFKD-DPIEPQLLTMIVTDAQQTPSWGN SQPWQVYIATGHA 54
 ECL -----MDIISVALKRHSTKAFDASKKLTAEEAEKIKTLLQYSPSSTNSQPWHFIVASTE- 54
 . . .*::::.* : . : :** * *** . : .

PPU KQALSALLGIAHNDPSADP---LAHL PDDLARKYRERQEKWGELFYGLHQIDKCDDAGRA 116
 REU MDRLMDIMRQRVTEAPGGEEREYDIYPRELVS PYRDRRFEVGEALYRSLGIPREDKQRR 114
 LPL LTNIKQHYATAAEQGI AEDADLAKVHRGDFS AFASQNMGHVVGTFRFPVIDSDP-----T 108
 ECL -EGKARVAKSAAGTYVFNERKMLDASHVVVFC AKTAMDDAWLERVVDQE EADGRFNTPEA 113
 . .

PPU RVSGLNFDFFGAPVGLIFTIDS NLKKYSWLDYGLFLQTLMLTARSRLSTCPQVSFARFQ 176
 REU AQFANNFAFFGAPLALFCSVDRRMGPPQWSD LGMYLQTVMLLLREEGLDSCAQECWAIYP 174
 LPL TYWDSRANLYRAPAIAYLVLDANPN SWSIYDLGAFSQTLM LAATARGVQSVPSYELVKYP 168
 ECL KAA NHKGRTYFAD MHRVDLKD DD--QWMAKQVYLVN VGNFLLGVGAMGLDAVP IEGFD--A 169
 . : * * : ..:* *::: .

PPU SLIKDFLHLLDDS-QEIVCGMSLGYADENATVNSLRMPREMAQGFTHFMGFDR 227
 REU QTIGGFSLPPE-RMLFTGMAIGYEDPEAPANQLRAPRAPLEAFTEFMGI-- 223
 LPL AIVRNALGMPAN-KVVAMGIALGYEKPEKPLNQFRTDRVATPQVLHIIN--- 216
 ECL AILDEEFGLEKKGFTSLVVVPVGHHSVEDFNATLPKSRLPLSTIVTEC---- 217
 : : : : :*:

Figure

Figure 3

Figure 5

Figure 6