1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

**Interaction of the attenuated recombinant rIHNV-Gvhsv GFP
virus with macrophages from rainbow trout (*Oncorhynchus
mykiss*)**

Alejandro Romero¹, Sonia Dios¹, Michel Bremont², Antonio Figueras¹, Beatriz
Novoa^{1*}

- 1. Instituto de Investigaciones Marinas, CSIC. Eduardo Cabello 6, 36208
Vigo, Spain.
- 2. INRA, Unité de Virologie et Immunologie Moléculaires, 78352 Jouy-en-
Josas, France.

*Corresponding author:
Tel: 0034 986 21 44 63
Fax: 0034 986 29 27 62
E-mail: virus@iim.csic.es

Abstract

One of the most important threats to the salmonid aquaculture industry is infection caused by novirhabdoviruses such as infectious haematopoietic necrosis virus (IHNV) or viral haemorrhagic septicaemia virus (VHSV). Using reverse genetics, an avirulent recombinant rIHNV-Gvhsv GFP strain was generated, which was able to replicate as effectively as wild type IHNV in a fish cell line and in macrophages. Although this recombinant virus induced protective responses against IHNV and VHSV, the response did not involve the production of antibodies or modulate the expression of some antiviral genes. To determine the immune mechanisms underlying the protection conferred by the rIHNV-Gvhsv GFP virus, different immune parameters (NO production, respiratory burst activity and the induction of apoptosis) were assessed in the macrophage population. The results obtained in the present work may indicate that the Nv protein could be important in the modulation of NO and ROS production. rIHNV-Gvhsv GFP did not appear to have a clear effect on nitric oxide production or apoptosis. However, an increased respiratory burst activity (with levels induced by the recombinant virus significantly higher than the levels induced by the wild type virus), suggests a stimulation of the macrophage population, which could be related to the protection against virulent viruses.

Keywords: rIHNV-Gvhsv GFP; IHNV; Rhabdovirus; Macrophages; TEM; ROS; Apoptosis; Trout

1. Introduction

Vaccination has an important role in large-scale commercial fish farming and has been a key reason for the success of salmonid fish cultivation. This culture has been highly affected by infections caused by the infectious haematopoietic necrosis virus (IHNV) and viral haemorrhagic septicaemia virus (VHSV), the two major causes of mass mortality (Bootland and Leong, 1999; Smail, 1999). Although a lot of research is being done for the control of these diseases, only one live attenuated vaccine for VHSV is available in Germany (Enzmann, Tübingen), and Novartis Animal Health (Switzerland) commercialised a DNA vaccine against IHNV (Apex®-IHNV) for use in Canada.

Although alternative methods to the traditional formulations such as live virus vaccines (Sommerset et al., 2005), recombinant DNA vaccines (Lorenzen et al., 2002) or live recombinant virus (Biacchesi et al., 2002; Romero et al., 2008) have been tried to generate vaccines, their development is limited by safety concerns for the consumer and for the environment. Using reverse genetics methodology, a new line of live recombinant IHNV strains were developed by Biacchesi et al. (Biacchesi et al., 2000a; Biacchesi et al., 2000b; Biacchesi et al., 2002) and tested in vaccination trials (Romero et al., 2005; Novoa et al., 2006; Romero et al., 2008). Two of the six IHNV genes were modified in the recombinant virus used in the present study (rIHNV-Gvhsv GFP). One of them was the G glycoprotein gene which encodes the G protein, involved in viral pathogenicity and capable of eliciting protective antibody production against various IHNV strains (Engelking and Leong 1989). It was replaced with the gene from VHSV. The amino acid

69 homology between the G protein of IHNV and VHSV indicates a high degree of
70 structural and functional similarity between the two fish rhabdovirus
71 glycoproteins (Lorenzen et al., 1993). Although the G-VHSV protein is able to
72 induce nonspecific protection against IHNV in experimental challenges in fish
73 (Lorenzen et al., 1998; Kim et al., 2000) the production of specific antibodies is
74 restricted by the epitope structure of VHSV (Engelking and Leong, 1989;
75 Lorenzen et al., 1990). The other gene modified was the small non-virion protein
76 Nv, which has been proven to be nonessential for recombinant IHNV, although its
77 deletion affects replication in cell culture (Thoulouze et al., 2004). Therefore, the
78 Nv gene can be used as a site of insertion for foreign genes and can serve as
79 vector for expressing additional antigens. This non-structural Nv gene was
80 replaced with the green fluorescent protein (GFP). We have previously
81 demonstrated that this recombinant virus was apathogenic for zebrafish and
82 rainbow trout. Moreover, vaccination trials showed that it was able to induce
83 protective responses against experimental infections with IHNV or VHSV in both
84 species (Novoa et al., 2006; Romero et al., 2008). However, we observed that the
85 non-specific protective response analyzed by measuring the gene expression level
86 of some antiviral genes, and the specific immune response evaluated through the
87 antibodies production, did not appear to be involved in this protection, and it was
88 suggested that other immune mechanisms could be responsible for the protection
89 conferred by the rIHNV-Gvhsv GFP virus (Novoa et al., 2006; Romero et al.,
90 2008).

91 Leukocytes are target cells for the replication of IHNV (Chilmonczyk and
92 Winton, 1994). The viral replication occurs fast in cell culture being detected the
93 first ultrastructural changes of the cytoplasm as early as 24-36 h post-infection by
94 electron microscopy (Björklund et al., 1997; Kazachka et al., 2007). Moreover,
95 the viral titre of IHNV peaked at 2 days post-infection in rainbow trout leucocytes
96 (Chilmonczyk and Winton, 1994). The cycle of infection occurs by series of well
97 described events in the following order: adsorption, penetration and uncoating,
98 transcription, translation, replication, assembly and budding (Bootland and Leong,
99 1999). The morphogenesis and replication cycle of the recombinant virus rIHNV-
100 Gvhsv GFP was analysed using transmission electron microscopy (TEM) and
101 compared with wild type IHNV to assess if the changes introduced in the viral
102 genome modified the efficacy of viral replication.

103 Leukocytes constitute an important part of the cellular defence against
104 bacterial and viral infections in fish (Secombes, 1994) by secreting reactive
105 oxygen and nitrogen intermediates and by their phagocytic capacity (Marletta et
106 al., 1988; Nathan and Hibbs, 1991; Secombes and Fletcher, 1992). Taking this
107 into account, we analysed in primary cell cultures enriched with kidney
108 macrophages different immune mechanisms triggered by the viral infection
109 (IHNV or rIHNV Gvhsv GFP), such as NO production, respiratory burst activity
110 and also the induction of apoptosis, to better understand the mechanisms
111 underlying the protection induced by the rIHNV-Gvhsv GFP virus.

112

113 **2. Materials and Methods**

114

115 **2.1. Virus titration**

116 IHNV (French isolates 32/87) (Laurencin, 1987) and the recombinant viruses
117 rIHNV-Gvhsv GFP, rIHNV GFP (Novoa et al., 2006) and rGvhsv (Romero et al.,
118 2005) were propagated in the fish epithelial cell line EPC, which is derived from
119 common carp (*Cyprinus carpio*) (Tomasec and Fijan, 1971). EPC cells were
120 cultured in Eagle's minimum essential medium (MEM Invitrogen, GIBCO)
121 supplemented with 10% foetal bovine serum (FBS Invitrogen, GIBCO), penicillin
122 (100 IU/mL) (Invitrogen, GIBCO), and streptomycin (100 µg/mL) (Invitrogen,
123 GIBCO), and buffered with 7.5% sodium bicarbonate (Invitrogen, GIBCO), and
124 were incubated at 20 °C. The viruses were inoculated on EPC cells grown in
125 MEM with antibiotics and 2% FBS at 15 °C. When the cytopathic effect (CPE)
126 was complete, the supernatants were harvested and centrifuged to eliminate cell
127 debris. Viruses were then titrated according to Reed and Muench (1938).

128

129 **2.2. Primary cell cultures enriched with kidney macrophages**

130 Primary cell cultures enriched with macrophages from rainbow trout (mean
131 weight 22 g) were obtained (Secombes, 1990). Briefly, the anterior kidney was
132 removed aseptically and passed through a 100-µm nylon mesh using Leibovitz
133 medium L-15 (Invitrogen, GIBCO) supplemented with penicillin (100 IU/mL),
134 streptomycin (100 µg/mL), heparin (10 U/mL) (Invitrogen, GIBCO) and 2% FBS.
135 The resulting cell suspension was placed on a 34%-51% Percoll density gradient
136 (GE Healthcare) and centrifuged at 500 g for 30 min at 4 °C. The interface cells

137 were collected and washed twice in L-15 containing 0.1% FBS, spinning at 500 g
138 for 5 min. The viable cell concentration was determined by trypan blue exclusion.
139 Cells were resuspended in L-15 with 0.1% FBS and dispensed into 24-well plates
140 at a concentration of 10^6 cells/mL. Adherent cells were attached to the bottom of
141 the wells by incubating 3h at 18 °C. After this period supernatants and non-
142 adherent cells were removed. All the animal experiments were reviewed and
143 approved by the CSIC National Committee on Bioethics.

144

145 ***2.3. Cell infections***

146 Primary cell cultures enriched with kidney macrophages were infected with
147 the rIHNv-Gvhsv GFP virus or wild type IHNv at a multiplicity of infection of 1
148 (MOI 1). After 30 min of adsorption, cells were washed and incubated at 15 °C in
149 L-15 medium supplemented with 2% FBS. Confluent EPC cell cultures were also
150 infected with IHNv or rIHNv-Gvhsv GFP virus as positive control groups.
151 Infected cell cultures were sampled at 24, 48 and 72 h post-infection (p.i.).
152 Samples were frozen and thawed twice to release the viral particles inside the
153 cells. After centrifugation at 12000 g for 5 min, the supernatants were stored at -
154 80 °C until use. Titration of supernatants was measured in triplicate according to
155 the protocol described by Reed and Muench (1938). Results were expressed as the
156 mean $\pm$ standard deviation (SD) and were compared using a t test ($p < 0.05$).

157 In addition, infected cultures were counterstained with DAPI (Sigma Aldrich)
158 according to the manufacturer's instructions to observe the CPE and the

159 morphological changes with high-resolution spectral confocal microscopy (Leica
160 TCS SPE).

161

162 ***2.4. Viral morphogenesis***

163 The study of the viral morphogenesis by electron microscopy was conducted
164 in EPC cells 1, 2, 3, 5 and 6 days post-infection. Cells were fixed for 1 h in 2%
165 glutaraldehyde in 0.1M cacodylate buffer. After two washes, cells were fixed in
166 2% osmium tetroxide (Invitrogen) and included in 0.25% technical agar
167 (Cultimed). Small squares of agar were processed for Poly/Bed 812 and Araldite
168 grade 502 (Polysciences, Inc.) resin blocking. Thin sections (1 μ m) stained with
169 0.5% toluidine blue were analysed by light microscopy. Ultrathin sections were
170 then stained with uranyl-acetate (FLUKA) and lead citrate (FLUKA) and
171 observed with a Philips CM 100 transmission electron microscope (TEM).

172

173 ***2.5. Analysis of innate immune cell parameters***

174 The different immune responses induced by macrophages against
175 experimental infections with recombinant rIHNV-Gvhsv GFP virus were analysed
176 and compared with the responses induced by wild type IHNV.

177 Primary cultures enriched with head kidney macrophages were obtained from
178 eight adult fish (mean weight 22 g), placed in 96-well plates (10^6 cells/mL) and
179 infected with rIHNV-Gvhsv GFP virus, rIHNV GFP virus or rGvhsv at different
180 MOI (1 and 0.01) in L-15 with 0.1% FBS at 15 °C. Primary cultures were also
181 infected with wild type IHNV or treated with culture medium as positive and

negative control groups, respectively. At 24, 48 and 72 h p.i., 50 µL of the supernatants were removed from the wells, and a nitrite assay was conducted. The remaining cells were grouped in three pools, and the respiratory burst activity was measured by the chemiluminescence (CL) method.

2.5.1. Nitrite assay

Nitric oxide (NO) production of head kidney macrophages was assayed using the method described by Neumann et al. (1995). Briefly, after incubation of macrophages at 15 °C for 24, 48 and 72 h, 50 µL of the supernatants were removed and placed in a separate 96-well plate. One-hundred microlitres of 1% sulphanilamide (Sigma) was added to each well followed by 100 µL of 0.1% N-naphthyl-ethylene-diamine (Sigma). Optical density (O.D.) was measured at 540 nm, and the molar concentration of nitrite was determined from standard curves generated using known concentrations of sodium nitrite. All the treatments were assayed in triplicate for each fish. Data were compared using a t test ($p < 0.05$). Results are expressed as the mean $\pm$ SD.

2.5.2. Chemiluminescence assay

The reactive oxygen species (ROS) production was measured 24 h, 48 h and 72 h p.i. in infected macrophages (MOI 1 and 0.01) as the emission of Relative Luminescence Units (RLUs) after cell membrane stimulation with phorbol myristate acetate (PMA, Sigma) and amplification by 5-amino-2,3-dihydro-1,4-phthalazinedione (Luminol, Sigma) according to the protocol

described by Rodriguez et al. (2008). Triplicate wells were used in all experiments. To determine if the viral infection could trigger macrophage respiratory burst activity, cells were infected as described above, and the O.D. at 550 nm was measured after 30 min without PMA stimulation. Data were compared using a t test ($p < 0.05$). ROS production index were calculated by dividing the values obtained in infected cells by the values obtained in controls. Results are expressed as the mean $\pm$ SD.

2.5.3. Apoptosis assay

Primary cultures enriched with head kidney macrophages from four rainbow trout (22 g mean weight) were obtained by the method described above. The cell concentration was adjusted to 10^6 cells/mL, and cells were placed on 24-well tissue culture plates (Falcon). Cells were infected with IHNV or rIHNV-Gvhsv GFP at a MOI 1. A negative control group was treated with culture medium, and a positive control group was treated with UV light for 30 min. Samples were taken at 24 and 72 h p.i., and Annexin V-PE (BD Biosciences) and 7AAD (BD Biosciences) staining were measured by flow cytometry as described by Romero et al. (2008). Data obtained from the four replicates were compared using a t test ($p < 0.05$).

3. Results

3.1. Viral replication

228 The recombinant rIHNV-Gvhsv GFP virus replicated as effectively as wild
229 type IHNV in both the macrophage cell culture and the EPC cell line (Fig. 1). The
230 viral titre increased along with time, changing from 3×10^4 TCID₅₀/mL to 2×10^5
231 TCID₅₀/mL in cell cultures enriched with kidney macrophages. Similar increases
232 were recorded for IHNV, the titres of which reached 5×10^5 TCID₅₀/mL in
233 macrophages 72 h p.i. The viral titre also increased in EPC cells infected with
234 IHNV and rIHNV-Gvhsv GFP, reaching 5×10^5 TCID₅₀/mL at 72 h p.i. No
235 significant differences were observed between wild type and recombinant virus-
236 infected samples in all cell types and time-points examined.

237 Infected EPC cells and infected macrophages showed a typical “bunch of
238 grapes” appearance upon IHNV infection (Fig. 2A and 2B). Although association
239 between damaged cells and GFP fluorescence was observed, no morphological
240 changes were observed in fluorescent virus-infected cells either in the cytoplasm
241 or in the nucleus until 48 h p.i. Confocal microscopy revealed that only a small
242 percentage of the cell population was actively infected by the rIHNV-Gvhsv GFP
243 virus (Fig. 2C and 2D). The green fluorescence emitted by the viral GFP protein
244 was focused in the cytoplasm. Green cytoplasmic extensions were also observed
245 (Fig. 2C and 2D). Virus-induced CPE was first observed in infected cells 24 h p.i.
246 using light microscopy on semi-thin sections (Fig. 3). The earliest ultrastructural
247 changes in virus-infected cultures involved a reduction of cell volume. This stage
248 was rapidly followed by a condensation of chromatin around the nuclear
249 membrane. Vacuolization of the cytoplasm and condensation of the chromatin
250 into one or several dense bodies was evident. Total CPE was observed at day 6

251 p.i., when the cytoplasm was completely vacuolated and dead cells and cellular
252 debris were observed.

253

254 **3.2. Viral morphogenesis**

255 The replication cycle of the rIHNV-Gvhsv GFP virus was described by
256 electron microscopy observations. The virions of the recombinant virus were
257 typically bullet-shaped, with approximate measurements of 110 nm X 70 nm (Fig.
258 4A). At 24h p.i., viral particles were attached to the cell surface (Fig. 4B).
259 Subsequent internalization by endocytosis of the viral particles into the cytoplasm
260 was observed (Fig. 4C). Endocytic compartments with an electron-dense granular
261 material inside were frequently observed at 24 h (Fig. 4D). At the same time (24 h
262 p. i.) cells with destructive changes in the cytoplasm were observed. Cytoplasm
263 inclusion bodies were observed as the first sign of virus replication (Fig. 4E and
264 4F). Chains of ribosome-bound mRNAs were detected in the cytoplasm (Fig. 4G).
265 Different stages of maturation were detectable at cellular membrane (Fig. 4H).
266 The next visible form of the virus replication (also at 24h p.i.) was the presence of
267 high amount of viruses into cytoplasmatic vacuoles (Fig. 4I). They induced the
268 disintegration of the cell structure and caused the release of the viral particles.
269 Also individual viruses were budded from the cell surface (Fig. 4J). At 48-72 h
270 after viral infection we observed many cells collapsed into apoptotic bodies with
271 different sizes. After 72 h p. i. almost all cells of the monolayer were completely
272 destroyed by the infection.

273

274 **3.3. Analysis of innate immune cell parameters**

275 **3.3.1. NO production**

276 The levels of nitrite measured in kidney macrophages infected with the
277 rIHNV-Gvhsv GFP virus or rIHNV GFP were significantly different from the
278 levels measured for cells infected with IHNV or rGvhsv, regardless of the MOI
279 used (Fig. 5). In both experimental conditions (MOI 1 and 0.01), the levels of
280 nitrite induced by the recombinant viruses rIHNV-Gvhsv GFP and rIHNV GFP
281 were similar to the levels recorded in the control group. However, infection with
282 wild type IHNV and rGvhsv always induced values of nitrite lower than the GFP
283 recombinant viruses. Statistically significant differences were obtained under all
284 of the experimental conditions except for the sample taken at 24 h p.i. at MOI 1

286 **3.3.2. ROS production**

287 Infection of macrophages with the recombinant viruses rIHNV-Gvhsv GFP,
288 rIHNV GFP, rGvhsv and the IHNV wild type virus at MOI 1 induced significant
289 increases in respiratory burst activity with respect to the controls at 24 h, 48 h and
290 72 h p.i. (Figure 6). However, at MOI 0.01 only infected cells with the
291 recombinant viruses rIHNV-Gvhsv GFP and rIHNV GFP induced a significant
292 increase with respect to controls during all the experiment. The ROS levels
293 induced by the recombinant virus rIHNV-Gvhsv GFP was always significantly
294 higher than the levels induced by the IHNV wild type at all samples and time-
295 points (Fig. 6). IHNV and rIHNV-Gvhsv GFP could not trigger respiratory burst

296 activity in infected macrophages 30 min after infection in the absence of PMA
297 (data not shown).

298

299 **3.3.3. Apoptosis assay**

300 The most important morphological changes in the population structure were
301 observed in cells treated with UV light at 72 h (Fig. 7A). The recombinant
302 rIHNV-Gvhsv GFP virus induced similar apoptotic and necrotic levels in infected
303 kidney macrophages as the wild type IHNV at 24 and 72 h p.i. (Fig. 7B). At 72 h,
304 the percentage of apoptotic macrophages (stained with Annexin V-PE) induced by
305 IHNV was the same as that induced by rIHNV-Gvhsv GFP virus (23%). The
306 percentage of necrotic cells (stained with 7AAD) was always lower than 10%. UV
307 light treatment induced the highest amount of cellular damage (apoptosis and
308 necrosis) in both primary cultures. An increase in the number of apoptotic
309 macrophages was observed from 24 to 72 h, changing from less than 10% at 24 h
310 p.i. up to 80% at 72 h p.i. (Fig. 7B). Although the percentage of macrophages
311 infected by the recombinant virus (as measured by GFP using the FL-1 channel,
312 530 nm) increased from 24 to 72 h p.i., less than 13% of the cell population
313 supported the viral replication (data not shown).

314

315 **4. Discussion**

316 IHNV is able to replicate in a variety of established cell lines (Ristow and
317 DeAvila, 1994; Bootland and Leong, 1999; Lorenzen et al., 1999) as well as in
318 many fish tissues (Wolf, 1988; Yamamoto et al., 1990), resulting in the presence

319 of virus in body fluids and mucus (Mulcahy et al., 1982; LaPatra et al., 1989).
320 Haematopoietic tissues are particularly susceptible to IHNV infection (Yasutake
321 and Amend, 1972; Estepa and Coll, 1994), and peripheral blood and kidney
322 leukocytes are involved in the pathogenesis of IHNV (Chilmonczyk and Winton,
323 1994). Our results showed that the recombinant rIHNV-Gvhsv GFP virus
324 replicated in leukocyte populations derived from the kidney of rainbow trout as
325 effectively as the wild type IHNV. Under all conditions studied, the viral titre
326 increased along with time and followed the same kinetics as IHNV, reaching
327 similar values at 72 h p.i. These data revealed that macrophage populations could
328 serve as target cells in the initial phase of rIHNV-Gvhsv GFP infection. Although
329 Novoa et al. (2006) and Romero et al. (2008) suggested that the changes
330 introduced in the genome decreased virulence in experimental infections *in vivo*,
331 no changes in replication efficiency were observed in infected cell cultures.
332 However, only a small percentage of the macrophage population (less than 13%)
333 supported viral replication *in vitro* as observed by confocal microscopy and flow
334 cytometry. Similar results were reported by Tafalla et al. (1998) using
335 immunofluorescence assays, in which only 8% of trout monolayer macrophages
336 expressed VHSV-specific fluorescence three days p.i. Estepa et al. (1992)
337 obtained a variable percentage of positive cells, ranging between 20-50%, five
338 days p.i. by flow cytometry analysis.

339 The recombinant rIHNV-Gvhsv GFP virus did not produce morphological
340 changes in the cytoplasm or in the nucleus until 48 h p.i., which coincides with
341 observations in VHSV-infected macrophages (Estepa and Coll, 1991). This virus-

342 induced CPE appeared as focal areas resembling those induced by wild type
343 IHNV, and cells had the morphology of apoptotic cells. Similar changes were also
344 described by Björklund et al. (1997) in an EPC cell line infected with the spring
345 viraemia of carp virus (SVCV), suggesting that apoptosis could be a generalised
346 cell killing mechanism after viral infections. It is interesting to note that although
347 the recombinant rIHNV-Gvhsv GFP virus was able to induce CPE in macrophage
348 populations, no histological lesions were observed in kidney, liver, spleen or brain
349 (Novoa et al., 2006; Romero et al., 2008).

350 The rIHNV-Gvhsv GFP virus initiated the replication cycle by attaching to the
351 cell surface, most probably using different cellular receptors, such as fibronectin-
352 like protein complex, phosphatidylserine, sialic acid, and other cell adhesion
353 molecules (Schlegel et al., 1983; Haywood, 1994; Broughan and Wunner, 1995;
354 Bearzotti et al., 1999), and its viral glycoprotein (G) spikes (Wagner, 1987).
355 Although the rIHNV-Gvhsv GFP virus had the G protein replaced by the G
356 protein from VHSV, no changes were observed in its ability to infect cultured
357 cells, and viral particles attached to the surface were frequently observed in TEM.
358 Also, many cytoplasmic vesicles with viral particles inside were observed,
359 suggesting that the virus entered the cell by endocytosis as has been described
360 with other enveloped viruses (Lenard and Miller, 1982; Matlin et al., 1982;
361 Marsh, 1993). Finally, the final assembly of rIHNV-Gvhsv GFP virus occurred
362 predominantly at the cell membrane and sometimes at the membranes of the Golgi
363 cisternae. After assembly, new virions were found adsorbed to neighbouring cell
364 membranes. Taking the data from the experimental titrations together with the

365 microscopy observations, we suggest that neither viral morphogenesis nor the
366 replication ability of the recombinant virus were affected by the genome
367 modification.

368 With regard to the immune cell parameters, it is well known that macrophages
369 are key cells in the first stages of a viral infection. Positive feedback mechanisms
370 and synergistic interactions intensify the immune response and give rise to potent
371 bactericidal and antiviral mechanisms, such as NO and ROS production (Reiss
372 and Komatsu, 1998; Ellermann-Eriksen, 2005). NO is usually produced by
373 macrophages in response to proinflammatory cytokines, bacterial
374 lipopolysaccharide (LPS), parasites or viruses (Marletta et al., 1988; Nathan and
375 Hibbs, 1991; Tafalla et al., 1999; Tafalla et al., 2000; Tafalla et al., 2001).
376 However, NO production has been shown to be ineffective for protecting against
377 viral infection in human cell lines (López-Guerrero and Carrasco, 1998). We
378 observed that IHNV and the pathogenic recombinant virus rGvhsv were able to
379 significantly inhibit NO production, probably as a viral mechanism to overcome
380 the host immune defences. This has also been described for VHSV (Tafalla et al.,
381 2001), as the virus was able to suppress NO production in infected macrophages
382 from turbot. In contrast, the recombinant virus rIHNV-Gvhsv GFP and rIHNV
383 GFP were not able to inhibit this production, and the nitrite values recorded were
384 similar to the levels registered in the control group. This NO concentration might
385 inhibit the earliest stages of viral replication and thus prevent viral spread,
386 promoting viral clearance and recovery of the fish. These results may indicate that
387 the Nv protein could be important in the inhibition of NO production.

388 Viral infections may also trigger the production of ROS (Akaike et al., 1998).
389 The role of ROS in apoptosis induction during viral infections has been reported
390 in mammals (Staal et al., 1990; Roederer et al., 1992; Kohno et al., 1996;
391 Skulachev, 1997; Skulachev, 1998; Kulms et al., 2000). Moreover, it has been
392 described that high levels of NO and ROS can modulate and activate apoptotic
393 cell death (Brüne et al., 1998; Tripathi and Hildeman, 2004). However, the results
394 obtained in fish so far have been contradictory. We previously observed (Tafalla
395 et al., 1998) no changes in ROS production in macrophages from rainbow trout
396 and turbot infected *in vitro* with VHSV. This lack of respiratory burst activity
397 could be correlated with the small number of macrophages that were actively
398 infected. Similar results were later observed by Chilmonczyk and Monge (1999)
399 and Tafalla and Novoa (2001). However, Stohlman et al. (1982) described that in
400 some cases, ROS production mediates the antiviral activity of macrophages, and
401 Siwicki et al. (2003) reported a significant reduction in the response of
402 macrophages isolated from rainbow trout and infected with VHSV. The decrease
403 in ROS production could be a mechanism induced by the virus to evade the host
404 immune response. Moreover, it could prevent the activation of other pathways to
405 eliminate infected cells, such as apoptosis (Skulachev, 1998). In contrast to these
406 previous results, we reported here a significant increase in the respiratory burst
407 activity of macrophages infected with rIHNV-Gvhsv GFP. Moreover, the
408 respiratory burst levels induced by this recombinant virus were significantly
409 higher than the levels induced by the wild type virus. Although the level of cell
410 infection described in this work was the same as the percentage observed by

411 Tafalla et al. (1998), macrophages infected with IHNV and rIHNV-Gvhsv GFP
412 were able to induce a high respiratory burst activity. However, the recombinant
413 virus was not able to trigger the response by itself, in agreement with the results
414 obtained by Tafalla et al. (1998) with VHSV. In any case, the high levels of
415 respiratory burst activity observed here reflected the activation of the macrophage
416 population against the viral infection.

417 The effects of rIHNV-Gvhsv GFP on the induction of apoptosis are of
418 particular interest because it is well known that this virus is able to replicate in
419 macrophages. Apoptosis has an important role in many viral infections
420 (Thoulouze et al., 1997; Gadaleta et al., 2002; Blaho, 2003; Blaho, 2004) and has
421 been shown to be involved in the cell death caused by rhabdovirus in cell lines
422 and tissues (Björklund et al., 1997; Chiou et al., 2000; Eléouët et al., 2001; Du et
423 al., 2004). Romero et al. (2008) described that rIHNV-Gvhsv GFP induced higher
424 apoptosis levels than wild type IHNV in the EPC cell line. However, in the
425 present work, a low level of apoptosis induction was recorded in primary cell
426 cultures infected with the recombinant or the wild type virus. To a certain extent,
427 this result was expected, as less than the 13% of the population supported the viral
428 replication. Our results are in accordance with previous results obtained by
429 Thoulouze et al. (1997), who analysed apoptosis induction by an attenuated rabies
430 virus strain in mouse and human lymphocytes and obtained low levels of
431 apoptosis.

432 The rIHNV-Gvhsv GFP virus did not appear to have a clear effect on NO
433 production or apoptosis in macrophages. However, the macrophage population

seemed to be stimulated by the recombinant virus, given the observed respiratory burst activation. In a previous study, MacKenzie et al. (2008) analysed the response in trout head kidney after intraperitoneal challenge with the recombinant rIHNV-Gvhsv GFP virus, IHNV and LPS. The results showed that infection with the recombinant virus induced a similar gene expression pattern to infection with the native IHNV at 24 h p.i. However, a divergence in the viral response was observed after 72 h, with the recombinant virus showing a recovery response more similar to that observed for the LPS treatment, suggesting that different mechanisms of activation were induced.

The results obtained in the present work may indicate that the Nv protein could be important in the modulation of NO and ROS production on infected cells. Moreover, the good levels of protection against experimental infection with IHNV and VHSV conferred by the recombinant virus rIHNV-Gvhsv GFP could be the result of stimulation or activation of the cellular innate immune system, although more efforts should be made to further clarify the interaction between the virus and the fish immune system.

Acknowledgements

The authors would like to thank B. Villaverde and P. Balseiro for their assistance and cooperation. This work was supported by the projects FAIRCT 98-4398 from the European Union and BIO 2000-0906 from the Spanish Ministerio de Ciencia y Tecnología. A. Romero acknowledges the CSIC for the I3P fellowship.

458 **References**

459

460 Akaike, T., Moritaka, S., Maeda, H., 1998. Free radicals in viral pathogenesis:
461 Molecular mechanisms involving superoxide and NO. Soc. Exp. Biol. Med.
462 217, 64-73.

463 Bearzotti, M., Delmas, B., Lamoureux, A., Loustau, A.M., Chilmonczyk, S.,
464 Bremont, M., 1999. Fish rhabdovirus cell entry is mediated by fibronectin.
465 J. Virol. 73, 7703-7709.

466 Biacchesi, S., Béarzotti, M., Bouguyon, E., Brémont, M., 2002. Heterologous
467 exchanges of the glycoprotein and the matrix protein in a Novirhabdovirus.
468 J. Virol. 76, 2881-2889.

469 Biacchesi, S., Thoulouze, M.I., Béarzotti, M., Yu, Y.X., Brémont, M., 2000a
470 Recovery of NV Knockout Infectious Hematopoietic Necrosis Virus
471 expressing foreign genes. J. Virol. 74, 11247-11253.

472 Biacchesi, S., Yu, Y.X., Béarzotti, M., Tafalla, C., Fernandez-Alonso, M.,
473 Bremont, M., 2000b. Rescue of synthetic salmonid rhabdovirus
474 minigenomes. J. Gen. Virol. 81, 1941-1945.

475 Björklund, H.V., Johansson, T.R., Rinne, A., 1997. Rhabdovirus-induced
476 apoptosis in a fish cell line is inhibited by a human endogenous acid
477 cysteine proteinase inhibitor. J. Virol. 71, 5658-5662.

478 Blaho, J.A., 2003. Virus infection and apoptosis (Issue I): An introduction. Int.
479 Rev. Immunol. 22, 321-236.

480 Blaho, J.A., 2004. Virus infection and apoptosis (Issue II) An introduction:
 481 Cheating death or death as a fact of life? *Int. Rev. Immunol.* 23, 1-6.

482 Bootland, L.M., Leong, J.C., 1999. Infectious Haematopoietic Necrosis Virus. In:
 483 Woo, P.T.K., Bruno, D.W. (Eds), *Fish Diseases and Disorders, Viral,*
 484 *Bacterial and Fungal Infections*, Academic Press, London, vol. 3.

485 Broughan, J.H., Wunner, W.H., 1995. Characterization of protein involvement in
 486 rabies virus binding to BHK-21 cells. *Arch. Virol.* 140, 75-93.

487 Brüne, B., Sandau, K., VonKnethen, A., 1998. Apoptotic cell death and nitric
 488 oxide: Activating and antagonistic transducing pathways. *Biochem.* 63, 817-
 489 966.

490 Chilmonczyk, S., Winton, J.R., 1994. Involvement of rainbow trout leucocytes in
 491 the pathogenesis of infectious hematopoietic necrosis. *Dis. Aquat. Org.* 19,
 492 89-94.

493 Chilmonczyk, S., Monge, D., 1999. Flow cytometry analysis as a tool for
 494 assessment of the fish cellular immune response to pathogens. *Fish Shellfish*
 495 *Immunol.* 9, 319-333.

496 Chiou, P., Kim, C., Ormonde, P., Leong, J.A., 2000. Infectious hematopoyetic
 497 necrosis virus matrix proteins inhibits host-directed gene expression and
 498 induces morphological changes of apoptosis in cell cultures. *J. Virol.* 74,
 499 7619-7627.

500 Du, C., Zhang, Q., Li, C., Miao, D., Gui, J., 2004. Induction of apoptosis in a carp
 501 leucocyte cell line infected with turbot rhabdovirus. *Virus Res.* 101, 119-
 502 126.

503 Eléouët, J.F., Druesne, N., Chilmonczyk, S., Monge, D., Dorson, M., Delmas, B.,
 504 2001. Comparative study of in situ cell death induced by the viruses of viral
 505 haemorrhagic septicemia (VHS) and infectious pancreatic necrosis (IPN) in
 506 rainbow trout. *J. Comp. Pathol.* 124, 300-307.

507 Ellermann-Eriksen, S., 2005. Macrophages and cytokines in the early defence
 508 against herpes simplex virus. *J. Virol.* 2, 59.

509 Engelking, H.M., Leong, J.C., 1989. The glycoprotein of infectious hematopoietic
 510 necrosis virus elicits neutralizing antibody and protective response. *Virus*
 511 *Res.* 13, 213–230.

512 Estepa, A., Coll, J.M., 1991. Infection of mitogen-stimulated trout leucocytes with
 513 salmonid viruses. *J. Fish. Dis.* 14, 555-562.

514 Estepa, A., Coll, J.M., 1994. Replication of rhabdovirus in trout hematopoietic
 515 cells. *Invest. Agr. Prod. Sanid. Anim.* 9, 37-44.

516 Estepa, A., Frias, D., Coll, J.M., 1992. Susceptibility of trout kidney macrophages
 517 to viral hemorrhagic septicaemia virus. *Viral Immunol.* 5, 283-292.

518 Gadaleta, P., Vacotto, M., Coulombié, F., 2002. Vesicular stomatitis virus induces
 519 apoptosis at early stages in the viral cycle and does not depend on virus
 520 replication. *Virus Res.* 86, 87-92.

521 Haywood, A.M., 1994. Virus receptors: binding, adhesion strengthening and
 522 changes in viral structure. *J Virol.* 68, 1-5.

523 Kazachka, D., Chikova, V., Ilieva, D., Christova, V., Jeleva, S., 2007.
 524 Morphogenesis of spring viraemia of carp virus in cell culture. *Biotechnol.*
 525 *& Biotechnol. Eq.* 21, 186-189.

526 Kim, C.H., Johnson, M.C., Drennan, J.D., Simon, B.E., Thomann, E., Leong, J.A.,
 527 2000. DNA vaccines encoding viral glycoproteins induce nonspecific
 528 immunity and Mx protein synthesis in fish. *J Virol.* 74, 7048–7054.

529 Kohno, T., Yamada, Y., Hata, T., Mori, H., Yamamura, M., Tomonaga, M. et al.,
 530 1996. Relation of oxidative stress and glutathione synthesis to
 531 CD95(Fas/APO-1)-mediated apoptosis of adult T cell leukemia cells. *J.*
 532 *Immunol.* 156, 4722-4728.

533 Kulms, D., Zeise, E., Poppelmann, B., Schwarz, T., 2000. DNA damage, death
 534 receptor activation and reactive oxygen species contribute to ultraviolet
 535 radiation-induced apoptosis in an essential and independent way. *Oncogene*
 536 21, 5844-5851.

537 LaPatra, S., Rohovec, J., Fryer, J., 1989. Detection of infectious hematopoietic
 538 necrosis virus in fish mucus. *Fish Pathol.* 24, 197-202.

539 Laurencin, F.B., 1987. IHNV in France. *Bull. Eur. Ass. Fish Pathol.* 7, 104.

540 Lenard, J., Miller, D.K., 1982. Uncoating of enveloped viruses. *Cell* 28, 5-6.

541 López-Guerrero, J.A., Carrasco, L., 1998. Effect of nitric oxide on poliovirus
 542 infection of two human cell lines. *J. Virol.* 72, 2538-2540.

543 Lorenzen, E., Carstensen, B., Olesen, N.J., 1999. Inter-laboratory comparison of
 544 cell lines for susceptibility to three viruses: VHSV, IHNV and IPNV. *Dis.*
 545 *Aquat. Organ.* 37, 81-88.

546 Lorenzen, N., Lorenzen, E., Einer-Jensen, K., Heppell, J., Wu, T., Davis, H.,
 547 1998. Protective immunity to VHS in rainbow trout (*Oncorhynchus mykiss*,

548 Walbaum) following DNA vaccination. *Fish Shellfish Immunol.* 8, 261–
 549 270.

550 Lorenzen, N., Lorenzen, E., Einer-Jensen, K., LaPatra, S.E., 2002. DNA vaccines
 551 as a tool for analysing the protective immune response against
 552 rhabdoviruses in rainbow trout. *Fish Shellfish Immunol.* 12, 439-453.

553 Lorenzen, N., Olesen, N.J., Jorgensen, P.E., 1990. Neutralization of Egtved virus
 554 pathogenicity to cell cultures and fish by monoclonal antibodies to the viral
 555 G protein. *J. Gen. Virol.* 71, 561–567.

556 Lorenzen, N., Olesen, N.J., Jorgensen, P.E.V., Etzerodt, M., Holtet, T.L.,
 557 Thogersen, H.C., 1993. Molecular cloning and expression in *Escherichia*
 558 *coli* of the glycoprotein gene of VHS virus, and immunization of rainbow
 559 trout with the recombinant protein. *J. Gen. Virol.* 74, 623-630.

560 MacKenzie, S., Balasch, J., Novoa, B., Ribas, L., Roher, N., Krasnov, A.,
 561 Figueras, A., 2008. Comparative analysis of the acute response of the trout,
 562 *O. mykiss*, head kidney to in vivo challenge with virulent and attenuated
 563 infectious hematopoietic necrosis virus and LPS-induced inflammation.
 564 *BMC Genomics* 9, 141.

565 Marletta, M.A., Yoon, P.Y., Iyengar, R., Leaf, C.D., Wishnok, J.S., 1988.
 566 Macrophage oxidation of L-arginine to nitrite and nitrate: nitric oxide is the
 567 intermediate. *Biochem.* 27, 8706-8711.

568 Marsh, M., 1993. Biochemical and morphological assays of virus entry. *Methods*
 569 *Enzymol.* 220, 249-261.

- 570 Matlin, K.S., Reggio, H., Helenius, A., Simon, K., 1982. Pathway of vesicular
571 stomatitis virus entry leading to infection. J. Mol. Biol. 156, 609-631.
- 572 Mulcahy, D., Burke, J., Pascho, R., Jenes, C., 1982. Pathogenesis of infectious
573 hematopoietic necrosis virus in adult sockeye salmon (*Oncorhynchus*
574 *nerka*). Can. J. Fish Aquat. Sci. 39, 1144-1149.
- 575 Nathan, C.F., Hibbs, J.B. Jr., 1991. Role of nitric oxide synthesis in macrophage
576 antimicrobial activity. Curr. Opin. Immunol. 3, 65-70.
- 577 Neumann, N.F., Fagan, D., Belosevic, M., 1995. Macrophage activating factor(s)
578 secreted by mitogen stimulated goldfish kidney leucocytes synergize with
579 bacterial lipopolysaccharide to induce nitric oxide production in teleost
580 macrophages. Dev. Comp. Immunol. 19, 473-482.
- 581 Novoa, B., Romero, A., Mulero, V., Rodriguez, I., Fernandez, I., Figueras, A.,
582 2006. Zebrafish (*Danio rerio*) as a model for the study of vaccination
583 against viral haemorrhagic septicemia virus (VHSV). Vaccine 24, 5806-
584 5816.
- 585 Reed, L.J., Muench, H., 1938. A simple method of estimating fifty per cent end-
586 points. Am. J. Hyg. 27, 493-497.
- 587 Reiss, C., Komatsu, T., 1998. Does Nitric Oxide Play a Critical Role in Viral
588 Infections? J. Virol. 72, 4547-4551.
- 589 Ristow, S., DeAvila, J., 1994. Susceptibility of four new salmonid cell lines to
590 Infectious Hematopoietic Necrosis Virus. J. Aquat. Anim. Health 6, 260-
591 265.

592 Rodríguez, I., Novoa, B., Figueras, A., 2008. Immune response of zebrafish
 593 (*Danio rerio*) against a newly isolated bacterial pathogen *Aeromonas*
 594 *hydrophila*. Fish Shellfish Immunol. 25, 239-249.

595 Roederer, M., Ela, S.W., Staal, F.J., Herzenberg, L.A., Herzenberg, L.A., 1992.
 596 N-acetylcysteine: a new approach to anti-HIV therapy. AIDS Res. Hum.
 597 Retroviruses 8, 209-217.

598 Romero, A., Figueras, A., Tafalla, C., Thoulouze, M.I., Bremont, M., Novoa, B.
 599 2005. Virulence, histological and serological studies on experimentally
 600 infected rainbow trout with different recombinant infectious hematopoietic
 601 necrosis viruses. Dis. Aquat. Org. 68, 17-28.

602 Romero, A., Figueras, A., Thoulouze, M.I., Bremont, M., Novoa, B., 2008.
 603 Infectious Hematopoietic Necrosis recombinant viruses induced protection
 604 for rainbow trout (*Oncorhynchus mykiss*). Dis. Aquat. Org. 80, 123-135.

605 Schlegel, R., Tralka, T.S., Willingham, M.C., Pastan, I., 1983. Inhibition of VSV
 606 binding and infectivity by phosphatidylserine: is phosphatidylserine a
 607 VSV-binding site? Cell 32, 639-646.

608 Secombes, C.J., Fletcher, T.C., 1992. The role of phagocytes in the protective
 609 mechanism of fish. Ann. Rev. Fish Dis. 2, 53-71.

610 Secombes, C.J., 1994. Cellular defences of fish: an update. In: Pike, A.W., Lewis,
 611 J.W., (Eds). Parasitic diseases of fish, Samara, Dyfed, pp. 209-224.

612 Secombes, C.J., 1990. Isolation of salmonid macrophages and analysis of their
 613 killing activity. In: Stolen, J.S., Fletcher, T.C., Anderson, D.P., Roberson,

614 B.S., van Muiswinkel, W.B., (Eds). Techniques in fish immunology, SOS
615 Publications, Fair Haven, pp. 137-155.

616 Siwicki, A.K., Morand, M., Kazun, B., Trapkowska, S., Pozet, F., 2003. In vitro
617 immunomodulating influence of methisoprinol on the head kidney
618 phagocyte and lymphocyte activity after suppression induced by VHSV in
619 rainbow trout (*Oncorhynchus mykiss*). Pol. J. Vet. Sci. 6, 51-53.

620 Skulachev, V.P., 1997. Membrane-linked systems preventing superoxide
621 formation. Biosci. Report 17, 347-366.

622 Skulachev, V.P., 1998. Possible role of reactive oxygen species in antiviral
623 defense. Biochem. 63, 1438-1440.

624 Smail, D.A., 1990. Viral Haemorrhagic Septicemia. In: Woo, P.T.K., Bruno,
625 D.W., (Eds). Fish Diseases and Disorders, Viral, Bacterial and Fungal
626 Infections, Academic Press, London, vol. 3.

627 Sommerset, I., Krossoy, B., Biering, E., Frost, P., 2005. Vaccines for fish in
628 aquaculture. Expert Rev. Vaccines 4, 89-101.

629 Staal, F.J.T., Roederer, M., Herzenberg, L.A., Herzenberg, L.A., 1990.
630 Intracellular thiols regulate activation of nuclear factor κ B and transcription
631 of human immunodeficiency virus. Proc. Nat. Acad. Sci. 87, 9943-9947.

632 Stohlman, S.A., Woodward, J.G., Frelinger, J.A., 1982. Macrophages antiviral
633 activity: extrinsic versus intrinsic activity. Infect. Immunol. 36, 672-677.

634 Tafalla, C., Figueras, A., Novoa, B., 1998. In vitro interaction of viral
635 haemorrhagic septicaemia virus and leucocytes from trout (*Oncorhynchus*

636 *mykiss*) and turbot (*Scophthalmus maximus*). Vet. Immunol. Immunopathol.
637 62, 359-366.

638 Tafalla, C., Figueras, A., Novoa, B., 1999. Role of nitric oxide on the replication
639 of viral haemorrhagic septicaemia virus (VHSV), a fish rhabdovirus. Vet.
640 Immunol. Immunopathol. 72, 249-256.

641 Tafalla, C., Figueras, A., Novoa, B., 2001. Viral haemorrhagic septicaemia virus
642 alters turbot (*Scophthalmus maximus*) macrophages nitric oxide production.
643 Dis. Aquat. Org. 47, 101-107.

644 Tafalla, C., Novoa, B., 2000. Requirements for nitric oxide production by turbot
645 (*Scophthalmus maximus*) head kidney macrophages. Dev. Com. Immunol.
646 24, 623-631.

647 Tafalla, C., Novoa, B., 2001. Respiratory burst of turbot (*Scophthalmus maximus*)
648 macrophages in response to experimental infection with viral haemorrhagic
649 septicaemia virus (VHSV). Fish Shellfish Immunol. 11, 727-734.

650 Thoulouze, M., Lafage, M., Montano-Hirose, J., Lafon, M., 1997. Rabies virus
651 infects mouse and human lymphocytes and induces apoptosis. J. Virol. 71,
652 7372-7380.

653 Thoulouze, M.I., Bouguyon, E., Carpentier, C., Brémont, M., 2004. Essential role
654 of the NV protein of Novirhabdovirus for pathogenicity in rainbow trout. J.
655 Virol. 78, 4098-4107.

656 Tomasec, J., Fijan, N., 1971. Virusne balesti riba (viral disease of fish). Final
657 report on research under a part of project, 6n/ 1966, Zagreb, 1971.

658 Tripathi, P., Hildeman, D., 2004. Sensitization of T cells to apoptosis. A role for
659 ROS? Apoptosis 9, 515-523.

660 Wagner, R.R., 1987. Rhabdovirus biology and infection. In: Wagner, R.R., (Eds).
661 The Rhabdoviruses, Plenum Press, New York, pp. 9-61.

662 Wolf, K., 1988. Infectious hematopoietic necrosis. In: Wolf, K., (Eds). Fish
663 Viruses and Fish Viral Diseases, Cornell University Press, Ithaca, New
664 York, pp. 83-114.

665 Yamamoto, T., Arakawa, C., Batts, N., Winton, J., 1990. Multiplication of
666 infectious hematopoietic necrosis virus in rainbow trout following
667 immersion infection: wholebody assay and immunochemistry. J. Aquat.
668 Anim. Health 27, 1-28.

669 Yasutake, W.T., Amend, D.F., 1972. Some aspects of pathogenesis of infectious
670 hematopoietic necrosis (IHN). J. Fish Biol. 4, 261-264.

671

672 **Figure legends**

673

674 **Fig. 1.** Time course of the viral titre in head kidney macrophages infected with
675 rIHNV-Gvhsv GFP or IHNV. Infected EPC cell lines were the positive control.
676 Values represent the mean and SD from three different experiments. Numbers on
677 the x axis indicate hours post-infection.

678

679 **Fig. 2.** Morphological changes induced by the rIHNV-Gvhsv GFP virus in the
680 EPC cell line and the macrophage cell culture (A and B, respectively) at 48 h p.i.
681 under confocal microscopy (MOI 1). A correlation between damaged cells and
682 GFP fluorescence was observed in the EPC cell line and the macrophage cell
683 culture (C and D, respectively). Green: recombinant virus on infected cells. Blue:
684 nuclear DAPI staining. Scale bar 250 μ m.

685

686 **Fig. 3.** Morphological changes observed under light microscopy in semi-thin
687 sections of EPC cell line and macrophages infected with rIHNV-Gvhsv GFP
688 virus. The virus-induced CPE was first apparent at 24 h p.i. Vacuolization of the
689 cytoplasm and condensation of the chromatin into one or several dense bodies was
690 observed at 48 h p.i. After six days, the cytoplasm was completely vacuolated,
691 and dead cells and cellular debris were observed.

692

693 **Fig. 4.** Viral morphogenesis and replication in the EPC cell line under electron
694 microscopy. (A and B) Infection by rIHNV-Gvhsv GFP and IHNV was initiated

695 by the attachment of the viruses to the cell surface. CS: cell surface (scale bar 100
696 nm). (C) Internalization of the recombinant virus (arrow head) by endocytosis
697 (scale bar 200 nm). (D) The appearance of endocytic compartments with the virus
698 inside was frequent (scale bar 200 nm). (E and F) Viral cores released into the
699 cytoplasm (arrows) (scale bar 200 nm). (G) Chains of ribosome-bound mRNAs of
700 the recombinant virus in the cytoplasm (arrows) (scale bar 200 nm). (H) Newly
701 synthesized viral proteins associated in the cytoplasm with *de novo* replicated
702 genomic RNA to form ribonucleoprotein cores (arrows) (scale bar 200 nm). (I)
703 Release of the new viral progeny (rIHNV-Gvhsv GFP) by cytoplasmic vesicles
704 (arrows) and disintegration of the cell structure (scale bar 200 nm). (J) The
705 budding of individual viruses from the cell surface was also observed (arrows)
706 (scale bar 100 nm).

707

708 **Fig. 5.** Production of nitrite radicals in kidney macrophages infected with rIHNV-
709 Gvhsv GFP, rIHNV GFP, rGvhsv or IHNV at different MOI (1 and 0.01) at 24,
710 48 and 72 h p.i. Results represent the mean $\pm$ SD of three replicates. (*)
711 Significant differences regarding to controls ($p < 0.05$).

712

713 **Fig. 6.** Respiratory burst activity induced by the recombinant rIHNV-Gvhsv GFP
714 virus, rIHNV GFP, rGvhsv and IHNV (MOI 1 and 0.01) in primary cultures
715 enriched with macrophages at 24 h, 48 h and 72 h p.i. Results represent the mean
716 $\pm$ SD of three replicates. (a and b) Significant differences regarding to values
717 obtained in control group and in infected cells with IHNV, respectively ($p < 0.05$).

718

719 **Fig. 7.** (A) FSC/SSC density plots of the cell population at 72 h post-treatment.

720 (B) Percentage of apoptotic and necrotic kidney macrophages infected with

721 rIHNV-Gvhsv GFP or IHNV or treated with UV light as a positive control 24 and

722 72 h p.i. The apoptotic and necrotic cells were stained with Annexin V-PE and 7-

723 AAD, respectively. Results represent the mean $\pm$ SD of four replicates. (*)

724 Significant differences ($p < 0.05$).

725

726

727

Figure 1

728

729

Figure 2

730

731

Figure 3

732

733

734

Figure 4

735

736

Figure 5

737

738

739

Figure 6

740

741

Figure 7

742