**Effects of continuous light on the reproductive system of European sea bass as
gauged by alterations of circadian variations during their first reproductive
cycle**

María José Bayarri, Silvia Zanuy, Ozlem Yilmaz and Manuel Carrillo*

Department of Fish Physiology and Biotechnology, Instituto de Acuicultura de Torre la Sal,
Consejo Superior de Investigaciones Científicas (CSIC), Castellón, Spain

* Author to whom correspondence should be addressed:

Prof. Manuel Carrillo Estévez

Instituto de Acuicultura de Torre la Sal (CSIC)

12595 Ribera de Cabanes (Castellón), Spain

Tel: + 34 964 319500

Fax: + 34 964 319509

E-mail: carrillo@iats.csic.es

Results have been partially presented at the 6th Conference of the Iberian Association of
Comparative Endocrinology (Asociación Ibérica de Endocrinología Comparada), held in Cádiz
(Spain) in September 2007.

1 **ABSTRACT**

2 The European sea bass is a short-day breeder, a characteristic that is highly valued in aquaculture. A
3 high percentage of males of this species mature precociously before reaching commercial size,
4 resulting in economic losses for fish farmers. We investigated the effects of continuous light (LL)
5 on the circadian variations of several reproductive hormones in males of this species in order to
6 understand how the presumed absence of a melatonin rhythm caused by LL affects their daily
7 profile. The study was conducted during four critical stages of the sea bass reproductive cycle (pre-
8 spermatogenesis (PSpg), spermatogenesis (Spg), spermiation (Spm), and post-spermiation (PSpm)).
9 Every 3 h during a complete 24 h cycle, six fish kept under a natural photoperiod (NP) and six
10 under LL were anaesthetized, measured, weighed, and bled. The pituitary was removed and frozen
11 at -80°C. The pituitary content of sea bream gonadotrophin-releasing (sbGnRH) and luteinizing
12 hormone (LH), as well as plasma content of LH, testosterone, and 11-ketotestosterone (11-KT)
13 were analyzed by ELISA. The percentage of spermiating males (precocity) per group was
14 determined by periodic abdominal massages of the animals. Our results confirm LL treatment,
15 maintained from the early stages of development onward, effectively reduces the percentage of
16 precocious male sea bass. As has already been described for caged sea bass, plasma LH showed a
17 clearly marked nocturnal rise near midnight during Spg and Spm during NP, but which was absent
18 under LL. Pituitary sbGnRH and LH content and plasma LH concentration, under both NP and LL
19 conditions, increased during the second half of the reproductive cycle, while sexual steroids were
20 higher at the beginning of the cycle. LL inhibited steroid secretion, especially testosterone secretion,
21 during Spg. In summary, without photoperiod cue, as accomplished by continuous exposure to LL,
22 circadian variations of reproductive hormones appeared altered, causing irregularities in the
23 reproductive process of male sea bass. These findings may have a practical application in
24 aquaculture, namely by applying LL treatment in an effort to reduce the presence of precocious
25 males in a stock. (E-mail correspondence: carrillo@iats.csic.es)

26

1 *Key words:* photoperiod manipulation, *Dicentrarchus labrax*, reproductive hormones, daily and
2 seasonal rhythms, *in vivo*.

3

1 INTRODUCTION

2 The success of current marine aquaculture is based on a sound knowledge of animal biology,
3 the application of new technologies, the development of specific diets, and control over diseases,
4 genetics, and reproduction in fish farming. Controlling reproduction is an important aspect in order
5 to obtain spawns from brood stocks to prevent over-exploitation of wild stocks and to produce fish
6 with faster growth rates by manipulating their natural reproductive rhythms, thus allowing the
7 farmer to obtain higher profits.

8 The reproductive process, like many others, is rhythmic. The physiological functions of
9 living organisms have adapted to occur at those moments of the day, and also year, when the
10 probability of success is the highest (Foster & Kreitzman, 2004). Cyclic changes in environmental
11 factors are responsible for the generation of biological rhythms in organisms, with photoperiod
12 being the most important cue in entraining these rhythms as exemplified by a recent study involving
13 light synchronization of the circadian rhythm of spawning in gilthead sea bream (Mesequer et al.,
14 2008). Therefore, the study of reproductive hormone circadian variations may provide interesting
15 information that could contribute to a better understanding of reproductive function of fish and
16 other species.

17 The European sea bass (*Dicentrarchus labrax*) is a teleost fish species that is highly valued
18 in Mediterranean aquaculture. A high percentage of the males of this species mature precociously
19 during their first year of life, before reaching commercial size, which in this species occurs at
20 around 18 months of age (Carrillo et al., 1995). This circumstance results in important economic
21 losses for fish farmers, since the nutrients and energy obtained by feeding during this period are
22 invested in gonadal development, at the expense of somatic growth. Moreover, flesh quality
23 decreases while sexual maturation is taking place (Bromage et al., 2001).

24 Reproductive control, therefore, is crucial to optimizing or extending the somatic growth
25 period of male sea bass, and photoperiod manipulation seems to be an effective method to achieve
26 such. . A number of studies have been conducted on several cultured fish species with the objective

1 of delaying or advancing gonadal maturation and spawning, using photoperiod manipulation,
2 among other techniques (Bromage et al. 1995). Artificial photoperiods have been reported to be an
3 effective method for delaying sea bass puberty (Zanuy et al., 2001) and/or altering spawning time
4 (Carrillo et al., 1993). In later studies (Rodríguez et al., 2001a; Bayarri et al, 2004), the application
5 of long photoperiods starting in the early stages of development proved effective for controlling
6 reproduction. Bayarri et al. (2004) studied melatonin and reproductive hormone circadian variations
7 during the first year of life in sea bass raised in sea cages and subjected to an artificially long
8 photoperiod (18L:6D). This study provided the first evidence of the alteration of circadian
9 variations caused by long photoperiod, affecting several different hormones in this species, such as
10 pituitary sbGnRH or plasma LH. Moreover, the final consequence of this light treatment was
11 delayed gonadal development and spawning at the time of puberty (Carrillo, Begtashi, Rodríguez,
12 & Zanuy, unpublished results). Long-term exposure of sea bass to LL proved highly effective in
13 inhibiting early precocity (Beghtashi et al., 2004), and for dramatically reducing the levels of
14 several reproductive hormones (Rodríguez et al., 2005). However, and despite their importance, no
15 data are yet available concerning the daily rhythms of reproductive hormones in fish kept under LL.
16 This is especially relevant considering the great differences that exist between the reproductive
17 effects of long photoperiods, such as that applied in the study by Bayarri et al. (2004), and LL,
18 which delays or inhibits puberty and reproductive hormones, respectively.

19 The characterization and temporal appearance of male sea bass gonadal stages, as well as
20 their endocrine regulation, have been widely studied (Prat et al., 1990, 1999; Zanuy et al., 1999;
21 Rodríguez et al., 2000a, 2000b, 2001a, 2001b, 2004, 2005; Beghtashi et al., 2004; Molés et al.,
22 2007). These studies have provided valuable information for selecting the relevant times for
23 sampling and studying hormonal daily rhythms, i.e., pre-spermatogenesis (PSpg) in September,
24 with low hormonal levels; spermatogenesis (Spg) in November, with the first significant surge of
25 hormones and morphological indexes; spermiation (Spm) in February, with maximum hormone
26 levels; and post-spermiation (PSpm) in May, with basal hormonal levels.

1 In a previous study (Bayarri et al., 2004), we tested the effects of a long photoperiod
2 (18L:6D) applied beginning in the early stages of development on male European sea bass. The aim
3 of the present study was to test the effects of LL, a treatment that presumably disrupts the melatonin
4 rhythm, on the circadian variation of a number of reproductive hormones of the brain-pituitary-
5 gonad axis (gonadotrophin releasing hormone, luteinizing hormone, and sexual steroids) during
6 four critical stages of the reproductive cycle of this species: PSpg, Spg, Spm, and PSpm.

7

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

MATERIALS AND METHODS

Animals and housing

This study took place at the Instituto de Acuicultura de Torre la Sal (Castellón, Spain), 40°N, 0°E. Six-month-old sea bass fingerlings (approx. 3.5g) obtained from L’Ecloserie Marine (Gravelines, France) were distributed into four identical 2000-liter, light-proof fiberglass tanks, supplied with well-aerated running sea water (salinity 37‰), and exposed to either a simulated natural photoperiod (NP) or constant light (LL) from the moment of their arrival in May. (At this latitude, the photoperiod reaches a maximum of 15L:9D in June, and a minimum of 9L:15D in December). The fish were kept under a natural temperature regime throughout the experiment (13-25°C), with the daily oscillation within a range of 0.5°C. Light in each tank was supplied by tungsten bulbs (PAR38Pro, Philips, Madrid, Spain) providing 650-700 lux at the water’s surface, and the simulated natural photoperiod was controlled by an electronic clock (ORBIS, Madrid, Spain), programmed weekly according to local geographical coordinates.

Fish from both light treatments were fed a commercial diet (Proaqua, Dueñas, Palencia, Spain) *ad libitum* by hand twice a day (i.e, in the morning, around 08:30 and in the afternoon around 13:30 h).

The handling of fish and conduct of the experimental procedures were always performed according to the international ethical standards as outlined in Portaluppi et al. (2008).

Experimental procedure

The objective of this experiment was to analyze the effects of LL on the daily variation of reproductive hormones and precocity of male European sea bass during their first year of life.

Every 3 h during a 24 h period, six male fish reared under the conditions prescribed for each treatment (NP and LL) were simultaneously anaesthetized with 2-phenoxyethanol (0.3 ppm) and then weighed and measured (n=48/treatment), at four critical stages of their reproductive cycle:

1 PSp_g, Sp_g, Sp_m, and PSp_m. Blood was collected by caudal puncture, using heparinized syringes,
2 and plasma was separated by centrifugation and preserved in aliquots at -80°C until the time of
3 analysis. The pituitary was removed and frozen separately in liquid nitrogen.

4 *Pituitary sbGnRH Immunoassays*

5 The pituitary was individually homogenized with a syringe in 250 µl of phosphate buffered
6 saline tween-20 (PBST), and extracted for 10 min at 80°C by adding 2N acetic acid to each
7 microcentrifuge tube containing pituitary homogenate. After centrifugation (13000 g for 30 min at
8 4°C), the supernatants were dried in a speed vacuum and reconstituted in EIA buffer for analysis.
9 The pituitary sbGnRH content was measured in reconstituted samples using a competitive enzyme-
10 linked immunosorbent assay (ELISA) similar to that described by Holland et al. (1998), which was
11 modified for sea bass. The sensitivity of the assay was 6 pg/well, and cross-reactivity with cII and
12 sGnRH was calculated to be 0.6 and 0.4%, respectively.

13 *LH Immunoassay*

14 A homologous ELISA (Mateos et al., 2006) was used to analyze LH pituitary content and
15 plasma levels. The sensitivity of the assay was approximately 0.6 ng/ml, and the intra and inter-
16 assay coefficients of variation were approximately 11% and 13%, respectively.

17 *Steroid assays*

18 A specific enzyme immunoassay (EIA) developed by Rodríguez et al. (2000a) for sea bass
19 was used to determine plasma testosterone levels. Plasma 11-ketotestosterone levels were analyzed
20 using an EIA originally developed for Siberian sturgeon (Cuisset et al., 1994), which was modified
21 for sea bass, and using a final dilution of 1:320000 for primary antibodies and 1:10 for the tracer
22 (Cayman Chemicals, MI, USA) (Rodríguez et al., 2001a).

23 *Determination of precocity*

24 Every 15 days, from February to May, the fish from both groups were subjected to an
25 abdominal massage to determine how many of them released sperm.

26 *Data analysis*

1 Data are expressed as mean $\pm$ SEM values. The statistical differences between the groups
2 were determined by a one-way analysis of variance (ANOVA), followed by a Tukey's test, with $p <$
3 0.05 considered the threshold for statistical significance.

4

5

1 **RESULTS**

2 Continuous light had an inhibiting effect on precocity (data not shown), as indicated by the
3 lower percentage of precocious animals found in the group exposed to LL (2.08% in LL vs. 12.50%
4 in NP) at the end of the experiment.

5 Under NP conditions, the pituitary sbGnRH content only exhibited statistically significant
6 daily variations during PSpm, with smooth elevations at 17:30 and 11:30 h (Figure 1). The
7 application of LL induced the appearance of peaks on the Spg and PSpm daily profiles at 23:30 and
8 05:30 h, respectively. When both light treatments were compared, significant sbGnRH pituitary
9 content differences were found for PSpg and Spg at 08:30 h, and for PSpm at 14:30, 02:30, 05:30,
10 and 11:30 h. Regardless of the light regime, lowest pituitary sbGnRH levels for the entire
11 reproductive cycle corresponded to PSpg; they increased during the following stages, reaching
12 especially high values in the case of Spm under LL.

13 The pituitary LH daily profile did not exhibit circadian oscillations under NP in any of the
14 four stages analyzed (Figure 2). On the other hand, significant daily differences appeared under LL
15 during PSpg and PSpm, with maximum LH concentrations occurring at 23:30 and 14:30 h for PSpg
16 and PSpm, respectively. Significant differences were found between both light regimes in the late
17 afternoon and during the first half of the night (from 17:30 to 23:30 h) during PSpg, at 14:30 h
18 during Spm, and at 11:30 h during PSpm. The only significant difference in pituitary LH content
19 among the different stages of the reproductive cycle in the group exposed to NP was observed at
20 08:30 h, with the highest value being recorded during Spm. In contrast, fish exposed to LL
21 displayed significant differences at most time points, although highest LH levels were also recorded
22 during Spm.

23 The daily plasma LH concentration profiles in fish maintained under NP conditions
24 displayed statistically significant peaks at 23:30 h during Spg and Spm, and 3 h earlier (20:30 h)
25 during PSpm (Figure 3). Exposure to LL, on the other hand, induced the appearance of daily
26 variations during PSpg, with the peak at 23:30 h, a pattern that was not seen under NP. Besides this,

1 the daily profiles in fish subjected to LL displayed a significant morning elevation in plasma LH
2 during Spm, and a peak at 17:30 h, which was statistically significant with respect to the 23:30 h
3 time-point value during PSpm. Comparison between the two light treatments yielded significant
4 plasma concentration differences for PSpg around midday, Spg at midnight, Spm at the first time
5 points in the morning, and PSpm at 05:30 h. Finally, the comparison between the different stages
6 within a given light regime also yielded significant differences at certain time points. In this case,
7 highest LH values in the NP group were recorded at night (23:30 and 02:30 h) during Spg, and at
8 11:30 h during PSpm, whereas under LL the maximum concentration was seen at 08:30 h during
9 Spm.

10 Significant daily variation in 11-KT was found in the plasma of fish kept under NP only
11 during PSpg and Spg, with peak values being recorded at 11:30 and 14:30 h, respectively (Figure
12 4). Under LL, the 11-KT concentration profile displayed significant circadian variation during all
13 four stages, with smooth elevations at 14:30 (PSpg), 20:30 (Spg), 02:30 (Spm), and 14:30 and 20:30
14 h (PSpm). Significant differences were seen at several time points during the daily cycle in the
15 concentration of this hormone in fish exposed to LL vs. NP conditions. Thus, 11-KT levels were
16 significantly higher in the NP group at 02:30, 08:30, and 11:30 h during PSpg, 11:30 h during Spg,
17 and during the last half of the scotophase and first half of the photophase in PSpm (i.e., 02:30,
18 05:30, 08:30, 11:30 h). The highest levels of the plasma 11-KT annual variation for both light
19 treatments occurred during the first half of the reproductive cycle, i.e., PSpg and Spg.

20 Plasma testosterone levels of fish subjected to NP exhibited significant daily variation
21 during PSpg, with a peak at 11:30 h, and also during PSpm, with a maximum concentration at 02:30
22 h, followed by a smooth decrease (Figure 5). Under LL, however, it was only during Spg that a
23 significant daily rhythm was found, with the lowest values being recorded at 23:30, 02:30, and
24 05:30 h. The testosterone levels of fish exposed to LL were significantly lower than those of the NP
25 group at 11:30 h during PSpg, on the entire Spg daily cycle, throughout the entire Spm scotoperiod,
26 and at 20:30, 02:30, 08:30, and 11:30 h during PSpm. Finally, when testosterone levels within each

1 group and throughout the entire reproductive cycle were considered, significantly higher values
2 were found at all time points during PSpg and Spg, regardless of the light regime.

3

1 **DISCUSSION**

2 The effect of LL on the circadian pattern of several reproductive hormones has been
3 analyzed for the first time during four critical stages of the reproductive cycle of European male sea
4 bass, corroborating the effectiveness of this treatment for inhibiting precocity, as previously
5 described for this species (Begtashi et al., 2004; Rodríguez et al., 2005). In carp, however, LL
6 resulted in precocious maturation of the testis during the pre-spawning phase, probably due to the
7 age and sexual status of the fish (Bhattacharya et al., 2007). In the present study, some reproductive
8 hormones, particularly plasma LH, appeared to be related to the photoperiod, as evidenced by the
9 daily profile differences detected between the LL and NP treatments. The presence of daily
10 variations under LL conditions leads us to conclude that the light treatment did not inhibit the
11 functioning of circadian oscillators, perhaps due to the persistence of other periodical cues, such as
12 temperature, which were still present. However, the 24 h rhythmicity did appear to be altered.

13 In the present experiment, pituitary sbGnRH levels under NP conditions did not vary over
14 the 24 h cycle, except in May, i.e., during the PSpm stage. Therefore, our results do not agree with
15 those of Bayarri et al. (2004), who found significant daily variations near spermatogenesis
16 (December) in male sea bass kept in sea cages. This discrepancy might suggest that the daily
17 pituitary sbGnRH profile is not relevant for completing the reproductive process of male sea bass,
18 and that other hormones are ultimately responsible. In the fish that we exposed to LL, pituitary
19 sbGnRH significantly peaked at 23:30 and 05:30 h during Spg and PSpm, respectively, in both
20 cases during the subjective night, demonstrating absence of a photoperiod cue alters this
21 neurohormone circadian variation. Regarding the annual cycle, the pituitary GnRH content has been
22 shown to be related to gonadal maturation (Okuzawa et al., 1990; Amano et al., 1992, 1993;
23 Holland et al., 1998; Andersson et al., 2001; Collins et al., 2001). In the present study, highest
24 pituitary sbGnRH values were found during Spm, especially in fish exposed to LL. Under NP
25 conditions, sbGnRH increased from November onward, reaching maximum level in February, and
26 remaining elevated even in May. This is in contrast with the findings of other studies (Rodríguez et

al., 2000b; 2004), where maximum values were recorded in November, during sexual differentiation and the first spawning season, with a smaller peak occurring in February. In these studies, however, samples were taken at one single time point during the day, where the relative position on the daily profile is not known and, therefore, does not necessarily reflect the daily sbGnRH maximum.

Pituitary LH, in those fish subjected to NP, failed to exhibit statistically significant daily rhythms at any stage of the reproductive cycle. However, the shape of November (Spg) daily profile was somewhat comparable (although slightly shifted) to that reported by Bayarri et al. (2004) for the same reproductive stage (December), with significant decreases at 16:00 and 01:00 h. Neither LL in the present study nor the long photoperiod used in a previous study by Bayarri et al. (2004) induced the appearance of significant pituitary LH rhythms during Spg. In our experiment, the circadian variation in LH under LL was only significant during PSpg (September) and PSpm (May). The highest values throughout the reproductive cycle were recorded during Spm (10-15 $\mu\text{g}/\text{mg prot}$) under both light treatments. The lack of studies regarding daily and annual pituitary LH rhythms in other species makes it difficult to undertake any further comparisons with our results.

Plasma LH peaked under NP at 23:30 h during Spg and Spm, the most critical stages of the reproductive cycle. Similar results were reported by Bayarri et al. (2004), who found a relationship between the onset of the scotophase and the appearance of the nocturnal LH rise in December, when a long photoperiod (18L:6D) was tested on sea bass. In fish exposed to this long photoperiod, the LH peak appeared after a 5 h delay, coinciding with the delay in the beginning of the dark period, with the ultimate consequence being a delay in puberty (Carrillo et al., unpublished results). In our case, no nocturnal peaks were detected during the Spg or Spm stages in fish exposed to LL. The lack of plasma LH daily rhythmicity under LL at such important reproductive times may partially explain the inhibition of precocity, since LL prevented the hormonal cascade from being completed. The nocturnal LH surges found in pre-pubertal sea bass (Bayarri et al., 2004) may resemble in certain aspects the LH pattern described in pre-pubertal humans, according to a study

1 conducted by Wu (1995). This pattern of nocturnal plasma LH peaks appears again in the results we
2 have now obtained for male sea bass during their first year of life. According to Zohar (1988), the
3 profile of the fish LH surge depends on the physiology of the gonad. For instance, in female sea
4 bream (*Sparus aurata*), a species with non-synchronous ovarian development, a daily pre-ovulatory
5 plasma LH peak was reported to appear 6 h before spawning (Zohar, 1988). In male carp, on the
6 other hand, some authors have described a time interval of 12 h between gonadotropin (GtH) rise
7 and spermiation (Courtois et al., 1986; Saad & Billard, 1985). The lowest concentration of LH over
8 the reproductive cycle was found during PSpg in September. This is similar to the findings of
9 Rodríguez et al. (2000b, 2001a, 2004), who observed lowest plasma LH concentrations at their first
10 sampling point (October-November). In female sea bass, Navas et al. (2004) once again found
11 lowest plasma LH values in October, and highest levels during ovulation, corroborating the general
12 agreement that LH could be the maturational hormone, whose secretion induces the production of
13 maturation-inducing steroids (Swanson et al., 1989; Swanson, 1991; Prat et al., 1996). In mammals,
14 such as the mink (*Mustela vison*), the lowest values for amplitude, frequency, and mean LH
15 secretion occur during the months of quiescence (Jallageas et al., 1994). These findings reinforce
16 the role of LH during the last stage of the reproductive cycle in short-day breeders.

17 Under NP, daily plasma 11-KT variations were found only during PSpg and Spg, at the
18 beginning of the reproductive cycle. The maximum concentrations in both stages were detected near
19 midday. In fish exposed to LL, significant circadian variations were evident during all four stages,
20 with a progressive delay in the appearance of the peaks throughout the reproductive cycle, similar to
21 what occurs in a free-running rhythm. The concentration of this hormone tended to increase at the
22 beginning of the day under NP, with values being significantly higher than those obtained under
23 long photoperiods (Bayarri et al., 2004), and under LL (present results). This steroid is of
24 importance in reproduction; 11-KT appears to be higher in dominant male rainbow trout than in
25 subordinate fish (Cardwell et al., 1996), and it induces typical male-type spawning behavior in male
26 goldfish (Kobayashi & Nakanishi, 1999). Therefore, absence of high levels of 11-KT during part of

1 the daily cycle under LL might probably be sufficient to modify sea bass reproduction, inhibiting
2 precocity, for example. Maximum 11-KT concentration in our study was detected at the beginning
3 of the cycle, during PSpg and Spg, which agrees with previous published results, and indicating the
4 maximum effect of 11-KT on this species occurs during spermiogenesis (Rodríguez et al., 2000b).
5 Under LL, 11-KT levels were also high during PSpg, although significantly lower than under NP.

6 Under NP conditions, testosterone showed circadian variation during PSpg and PSpm. In
7 fish exposed to LL, however, a daily rhythm was found only during Spg, with lowest values
8 attained during the subjective night. Bayarri et al. (2004) also described a significant Spg rhythm
9 under both natural and long photoperiods. In that study, none of the rhythms was affected by light,
10 since both showed a peak around 07:00 h, coinciding with the minimum water temperature. In our
11 case, testosterone peaks under NP conditions were detected near midday during PSpg, and near
12 midnight during PSpm, showing an undefined pattern of rhythmicity. Daily testosterone variations
13 have been also described for other teleost species, such as catfish (Lamba et al., 1983), carp (Santos
14 et al., 1986), and Japanese eel char (Yamada et al., 2002). Moreover, , the highest testosterone
15 concentration over the course of the reproductive cycle occurred during PSpg and Spg, that is, at
16 the beginning of the reproductive cycle, which is similar to that found for 11-KT. Rodríguez et al.
17 (2004) also found highest concnetration of this hormone in November, during the period of
18 testicular differentiation and growth(the first reproductive season for sea bass). Differences between
19 treatments were especially appreciable during Spg, when all time points throughout the day proved
20 to be statistically different.

21 In conclusion, our results derived under a natural photoperiod showed certain similarities
22 with those found in previous studies on European sea bass, especially regarding the daily variation
23 in plasma LH (Bayarri et al., 2004) as well as the concentrations of plasma LH throughout the
24 reproductive cycle (Rodríguez et al. 2000b, 2001a, 2001b, 2004, 2005). However, in contrast with
25 the study by Bayarri et al. (2004), which was performed during spermatogenesis, differences were
26 detected in the daily profile of certain reproductive hormones under NP conditions. Therefore, it

1 might be concluded that the circadian variation in those hormones, whose daily profile differs in the
2 two studies, are not very important for sexual development. Nevertheless, plasma LH levels under
3 NP peaked during the first half of the night in both studies, which demonstrates the relevance of this
4 peak for sexual maturation. In fact, when this peak shifted to the second half of the night in fish
5 housed under the long photoperiod (Bayarri et al. 2004), the consequence was delayed puberty
6 (Carrillo et al., unpublished results), and when it was not present under LL (as in the present study),
7 precocity was inhibited. Exposure to LL, on the other hand, altered the circadian and annual profile
8 of several reproductive hormones, with the ultimate consequence being reduction in the incidence
9 of precocity. As a result, it would appear that LL treatment would have a practical application in
10 Aquaculture.

11

1 **ACKNOWLEDGEMENTS**

2 This research was funded by the Spanish MEC project No. AGL 2006-04672 and by a grant
3 awarded by the Generalitat Valenciana (AE05/066 and AE07/098) to M.C.E. M.J.B.C. is the
4 recipient of a Juan de la Cierva contract from the Spanish Ministerio de Educación y Ciencia. The
5 authors would like to thank I. Martínez, M. Kiewek and V. Gracia for their help during sampling,
6 and X. Beltrán for helping in the laboratory.

7

REFERENCES

- Amano M, Aida K, Okumoto N, Hasegawa Y. (1992). Changes in salmon GnRH and chicken GnRH-II contents in the brain and pituitary, and GTH contents in the pituitary in female masu salmon, *Oncorhynchus masou*, from hatching through ovulation. *Zool. Sci.* 9:375–386.
- Amano M, Aida K, Okumoto N, Hasegawa Y. (1993). Changes in levels of GnRH in the brain and pituitary and GTH in the pituitary in male masu salmon, *Oncorhynchus masou*, from hatching to maturation. *Fish Physiol. Biochem.* 11:233–240.
- Andersson E, Fjellidal PG, Klenkeb U, Vikingstad E, Taranger GL, Zohar Y, Stefansson SO. (2001). Three forms of GnRH in the brain and pituitary of the turbot, *Scophthalmus maximus*: immunological characterization and seasonal variation. *Comp. Biochem. Physiol. Part B* 129:551-558.
- Bayarri MJ, Rodríguez L, Zanuy S, Madrid JA, Sánchez-Vázquez FJ, Kagawa H, Okuzawa K, Carrillo M. (2004). Effect of photoperiod manipulation on the daily rhythms of melatonin and reproductive hormones in caged European sea bass (*Dicentrarchus labrax*). *Gen. Comp. Endocr.* 136:72-81.
- Beghtashi I, Rodríguez L, Zanuy S, Carrillo M. (2004). Long-term exposure to continuous light inhibits precocity in juvenile male European sea bass (*Dicentrarchus labrax*, L.). I. Morphological aspects. *Aquaculture* 241:539-559.
- Bhattacharya S, Chattoraj A, Maitra SK. (2007). Melatonin in the regulation of annual testicular events in carp *Catla catla*: evidence from the studies on the effects of exogenous melatonin, continuous light, and continuous darkness. *Chronobiol. Int.* 24: 629-650.
- Bromage NR, Randall CF, Porter MJR, Davies B. (1995). How do photoperiod, the pineal gland, and circannual rhythms interact to coordinate seasonal reproduction in salmonid fish. In Goetz FW, Thomas P (eds). *Reproductive Physiology of Fish*, Austin, TX: Proceedings of the 5th International Symposium, pp. 164–166.

- 1 Bromage N, Porter M, Randall C. (2001). The environmental regulation of maturation in farmed
2 finfish with special reference to the role of photoperiod and melatonin. *Aquaculture* 197:63-
3 98.
- 4 Cardwell JR, Sorensen PW, Van Der Kraak GJ, Liley NR. (1996). Effect of dominance status on
5 sex hormone levels in laboratory and wild-spawning male trout. *Gen. Comp. Endocr.*
6 101:333-341.
- 7 Carrillo M, Zanuy S, Prat F, Serrano R, Bromage NR. (1993). Environmental and hormonal
8 control of reproduction in sea bass. In Muir JF, Roberts RJ (eds). *Recent advances in*
9 *aquaculture*. Oxford: Blackwell Science, pp. 43-54.
- 10 Carrillo M, Zanuy S, Prat F, Cerda J, Ramos J, Mañanós E, Bromage N. (1995). Sea bass
11 (*Dicentrarchus labrax*). In Bromage NR, Roberts RJ (eds). *Broodstock Management and Egg*
12 *and Larval Quality*. Oxford: Blackwell Science, pp. 138–168.
- 13 Collins PM, O’Neil DF, Barron BR, Moore RK, Sherwood NM. (2001). Gonadotropin-releasing
14 hormone content in the brain and pituitary of male and female grass rock-fish (*Sebastes*
15 *rastrelliger*) in relation to seasonal changes in reproductive status. *Biol. Reprod.* 65:173-179.
- 16 Courtois F, Takashima F, and Billard R. (1986). Stimulation of spermiation following repeated
17 injection of carp pituitary homogenates in the carp. *Bull. Jap. Soc. Sci. Fish* 52:995-997.
- 18 Cuisset B, Pradelles P, Kime DE, Kiihn ER, Babin P, Davail S, Le Menn F. (1994). Enzyme
19 immunoassay for 11-ketotestosterone using acetylcholinesterase as label: application to the
20 measurement of 11-ketotestosterone in plasma of Siberian sturgeon. *Comp. Biochem. Physiol.*
21 *Part. C* 108:229-241.
- 22 Foster RG, Kreitzman L. (2004). *Rhythms of life: The biological clocks that control the daily lives*
23 *of every living thing*. Profile Books, London.
- 24 Holland MCH, Gothilf Y, Meiri I, King JA, Okuzawa K, Elizur A, Zohar Y. (1998). Levels of the
25 native forms of GnRH in the pituitary of the gilthead seabream, *Sparus aurata*, at several
26 characteristic stages of the gonadal cycle. *Gen. Comp. Endocr.* 112:394-405.

- Jallageas M, Mas N, Boissin J, Maurel D, Ixart G. (1994). Seasonal variations of pulsatile luteinizing hormone release in the mink (*Mustela vison*). *Comp. Biochem. Physiol. Part C* 109:9-20.
- Kobayashi M, Nakanishi T. (1999). 11-Ketotestosterone induces male-type sexual behaviour and gonadotropin secretion in gynogenetic crucian carp, *Carassius auratus langsdorfii*. *Gen. Comp. Endocr.* 115:178-187.
- Lamba VJ, Goswami SV, Sundararaj BI. (1983). Circannual and circadian variations in plasma levels of steroids (cortisol, estradiol-17 β , estrone, and testosterone) correlated with the annual gonadal cycle in the catfish, *Heteropneustes fossilis* (Bloch). *Gen. Comp. Endocr.* 50:205-225.
- Mateos J, Mañanós E, Swanson P, Carrillo M, Zanuy S. (2006). Purification of luteinizing hormone (LH) in the sea bass (*Dicentrarchus labrax*) and development of a specific immunoassay. *Ciencias Marinas* 32:271-283.
- Meseguer C, Ramos J, Bayarri MJ, Oliveira C, Sánchez-Vázquez FJ. (2008). Light synchronization of the daily spawning rhythms of gilthead sea bream (*Sparus aurata* L.) kept under different photoperiod and after shifting the LD cycle. *Chronobiol. Int.* 25: 666-679.
- Molés G, Carrillo M, Mañanós E, Mylonas CC, Zanuy S. (2007). Temporal profile of brain and pituitary GnRHs, GnRH-R and gonadotropin mRNA expression and content during early development in European sea bass (*Dicentrarchus labrax* L.). *Gen. Comp. Endocr.* 150:75-86.
- Navas JM, Zauly S, Segner H, Carrillo M. (2004). β -Naphthoflavone alters normal plasma levels of vitellogenin, 17 β -estradiol and luteinizing hormone in sea bass broodstock. *Aquat. Toxicol.* 67:337-345.
- Okuzawa K, Amano M, Kobayashi M, Aida K, Hanyu I, Hasegawa Y, Miyamoto K. (1990). Differences in salmon GnRH and chicken GnRH-II contents in discrete brain areas of male and female rainbow trout according to age and state of maturity. *Gen. Comp. Endocr.* 80:116-126.

- Portaluppi F, Touitou Y, Smolensky M.H. (2008). Ethical and methodological standards for laboratory and medical biological rhythm research. *Chronobiol. Int.* 25: 999-1016.
- Prat F, Zanuy S, Carrillo M, De Mones A, Foster A. (1990). Seasonal changes in plasma levels of gonadal steroids of sea bass, *Dicentrarchus labrax*. *Gen. Comp. Endocr.* 78:361-373.
- Prat F, Sumpter JP, Tyler CR. (1996). Validation of radioimmunoassays for two salmon gonadotropins (GtH I and GtH II) and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biol. Reprod.* 54:1375-1382.
- Prat F, Carrillo M, Zanuy S, Bromage NR. (1999). Effects of constant short and long photoperiod regimes on the spawning performance and sex steroid levels of female and male sea bass (*Dicentrarchus labrax* L.). *J. Fish Biol.* 54:125–137.
- Rodríguez L, Begtashi I, Zanuy S, Carrillo M. (2000a). Development and validation of an enzyme immunoassay for testosterone: effects of photoperiod on plasma testosterone levels and gonadal development in male sea bass (*Dicentrarchus labrax*, L.) at puberty. *Fish Physiol. Biochem.* 23:141-150.
- Rodríguez L, Carrillo M, Sorbera LA, Soubrier MA, Mañanós E, Holland MCH, Zohar Y, Zanuy S. (2000b). Pituitary levels of three forms of GnRH in the male European sea bass (*Dicentrarchus labrax*, L.) during sex differentiation and first spawning season. *Gen. Comp. Endocr.* 120:67-74.
- Rodríguez L, Begtashi I, Zanuy S, Shaw M, Carrillo M. (2001a). Changes in plasma levels of reproductive hormones during first sexual maturation in European male sea bass (*Dicentrarchus labrax* L.) under artificial day lengths. *Aquaculture* 202:235-248.
- Rodríguez L, Zanuy S, Carrillo M. (2001b). Influence of daylength on the age at first maturity and somatic growth in male sea bass (*Dicentrarchus labrax*, L.). *Aquaculture* 196 (1-2):159-175.
- Rodríguez L, Carrillo M, Sorbera LA, Zohar Y, Zanuy S. (2004). Effects of photoperiod on pituitary levels of three forms of GnRH and reproductive hormones in the male European sea

- 1 bass (*Dicentrarchus labrax*, L.) during testicular differentiation and first testicular
2 recrudescence. *Gen. Comp. Endocr.* 136:37-48.
- 3 Rodríguez L, Begtashi I, Zanuy S, Carrillo M. (2005). Long-term exposure to continuous light
4 inhibits precocity in European male sea bass (*Dicentrarchus labrax*, L.): hormonal aspects.
5 *Gen. Comp. Endocr.* 140:116–125.
- 6 Saad A, Billard R. (1985). Amount of collected sperm related to spermatogenetic production and
7 the possibility of short-term storage in carp. Symposium on the aquaculture of carp and
8 related species. Evry, France.
- 9 Santos AJG, Furukawa K, Kobayashi M, Bando K, Aida K, Hanyu I. (1986). Plasma
10 gonadotropin and steroid hormone profiles during ovulation in the carp *Cyprinus carpio*. *Bull.*
11 *Jap. Soc. Sci. Fish* 52:1159-1166.
- 12 Swanson P. (1991). Salmon gonadotropins: reconciling old and new ideas. In Scott AP, Sumpter
13 JP, Kime DE, Rolfe MS (eds). *Proceedings of the Fourth International Symposium on the*
14 *Reproductive Physiology of Fish*. Sheffield: Fish Symp., pp. 2–7.
- 15 Swanson P, Bernard M, Nozaki M, Suzuki H, Kawauchi H, Dickhoff WW. (1989). Gonadotropins
16 I and II in juvenile coho salmon. *Fish Physiol. Biochem.* 7:169–176.
- 17 Wu FCW. (1995). GnRH pulse generator activity during human puberty. In Plant TM, Lee PA
18 (eds). *The Neurobiology of Puberty*. Bristol: J. Endocrinol. Ltd., pp. 185–197.
- 19 Yamada H, Satoh RI, Ogoh M, Takaji K, Fujimoto Y, Hakuba T, Chiba H, Kambegawa A, Iwata
20 M. (2002). Circadian changes in serum concentrations of steroids in Japanese eel char
21 *Salvelinus leucomaenis* at the stage of final maturation. *Zool. Sci.* 19:891-898.
- 22 Zanuy S, Carrillo M, Mateos J, Trudeau V, Kah O. (1999). Effects of sustained administration of
23 testosterone in pre-pubertal sea bass (*Dicentrarchus labrax* L). *Aquaculture* 177:21-35.
- 24 Zanuy S, Carrillo M, Felip A, Rodríguez L, Blázquez M, Ramos J, Piferrer F. (2001). Genetic,
25 hormonal and environmental approaches for the control of reproduction in the European sea
26 bass (*Dicentrarchus labrax* L). *Aquaculture* 202:187-203.

1 Zohar Y. (1988). Gonadotropin releasing hormone in spawning induction in teleosts: basic and
2 applied considerations. In INRA (ed). *Reproduction in Fish – Basic and applied aspects in*
3 *endocrinology and genetics*. Paris: INRA, pp. 47-62.

1 **FIGURE LEGENDS**

2 Figure 1. Daily pituitary sbGnRH content variations at various stages of the reproductive cycle in
3 sea bass maintained under NP (black symbols) and LL (white symbols) conditions. Horizontal
4 white and black bars represent day and night time, respectively. Different lower case and capital
5 letters indicate significant differences (ANOVA, Tukey's test, $p < 0,05$) among different time
6 points throughout the 24 h period, and among the different stages of the reproductive cycle,
7 respectively. Significant differences between treatments at each time point are indicated by an
8 asterisk (*). PSp_g = pre-spermatogenesis, Sp_g = spermatogenesis, Sp_m = spermiation, PSp_m =
9 post-spawning.

10

11 Figure 2. Pituitary LH content variations over the 24 h cycle at various stages of the reproductive
12 cycle in sea bass maintained under NP (black) and LL (white) conditions. See the legend in Figure 1
13 for details.

14

15 Figure 3. Plasma LH concentration changes over the 24 h cycle at various stages of the reproductive
16 cycle in sea bass maintained under NP (black) and LL (white) conditions. See the legend in Figure 1
17 for details.

18

19 Figure 4. Concentration of 11-ketotestosterone over the 24 h cycle at various stages of the
20 reproductive cycle in sea bass maintained under NP (black) and LL (white) conditions. See the
21 legend in Figure 1 for details.

22

23 Figure 5. Daily plasma testosterone variation in fish maintained under NP (black) and LL (white)
24 conditions at various stages of the reproductive cycle. See the legend in Figure 1 for details.

25

26

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5