Condition and growth of anchovy late larvae during different seasons in the Northwestern Mediterranean

D. Costalago^{a,*}, S. Tecchio^a, I. Palomera^a, I. Álvarez-Calleja^a, A. Ospina-Álvarez^a, S. Raicevich^b

^a Instituto de Ciencias del Mar-CSIC, 08003 Barcelona, Spain.

^b Istituto Superiore per la Ricerca e Protezione Ambientale, 30015 Chioggia (Ve), Italy.

ABSTRACT.

The fishery of European anchovy *Engraulis encrasicolus* in the Mediterranean needs several ecological approaches to be properly managed. Several surveys were carried out within this frame to study the ecology of larvae and juveniles of this species, which is known to reproduce during the warmest period of the year (May through September) in the Gulf of Lions. Special effort was made in the study of late larvae (15 mm total length until metamorphosis), since other authors have focused on larvae below that size. Unexpectedly, we collected anchovy late larvae during the December 2007 survey as well, whose range in size corresponded to a later spawning period than previously reported. Differences in the nutritional condition of these larvae were assessed by comparing indices of lipid composition and estimating growth rates from otolith measurements to infer information on the probability of survival between the two groups. The analysis of fatty acids, used as tracers of trophic relationships, indicates that these larvae fed mainly on zooplankton. Nutritional conditions of summer and late autumn larvae were very similar. In contrast, growth rates were higher for August larvae, probably due to the different temperatures in the two seasons. Our results are especially relevant in an ecological context where the rising of mean water temperatures in the Western Mediterranean could favour the extension of the anchovy spawning period up to late-Autumn months.

Keywords: *Engraulis encrasicolus*; late larvae; Northwestern Mediterranean; condition; growth.

1. Introduction

The European anchovy *Engraulis encrasicolus* is a highly valued fishery resource in the Western Mediterranean Sea (García et al., 1996; Perterra et al., 1996; Barange et al., 2009). Several studies have been carried out regarding diverse aspects of the biology and the ecology of the species in this area to improve the available tools for its fishery management (Palomera et al., 2007).

The Northwestern Mediterranean is one of the most productive areas in this sea due to the cyclonic current that flows southwards over the slope of the Gulf of Lions, carrying a significant nutrient load from the Rhône River (Salat, 1996). The Gulf of Lions also displays remarkable environmental differences between seasons, which directly influence low trophic level species (Calbet et al., 2001). The anchovy population could then be easily compromised by any sort of alteration that additionally impinges on these organisms, especially during the early development stages (i.e. eggs, larvae and juveniles) when they are particularly sensitive to any change (Palomera et al., 2007).

The anchovy spawning period in the Gulf of Lions extends from April to late September, with a peak in late June (Palomera, 1992). Therefore late larvae are expected to be found mainly during the summer, while juveniles should emerge in the autumn and winter.

During the SARDONE project, devoted to improve the management strategies of the European anchovy, two cruises were undertaken in August and December of 2007

in the Gulf of Lions in order to collect anchovy late larvae and juveniles, respectively. Unexpectedly, and in spite of the allegedly unfavorable conditions, an important number of anchovy late larvae were caught in December (a mean of 26.75 larvae/tow, against 34.20 larvae/tow in summer), well after the reported end of the spawning period.

The growth and feeding ecology of the species are among the most important issues to work on because the knowledge obtained through them is essential to understand how the population temporally and spatially develops in the environment. The relationship between this population and the plankton community is not a straightforward subject but some approaches have been attempted (Isari et al., 2008; Morais et al., 2010), especially those concerning how the zooplankton affects the population strength. Indeed, the growth rates and the nutritional condition of a population of several species, including European anchovy, have been key subjects for the study of the recruitment strength in many works (Butler, 1991; Ward et al., 2006; Palomera et al., 2007; Hidalgo et al., 2008; Islam and Tanaka, 2009; La Mesa et al., 2009).

In the present case we analyze for the first time in the Mediterranean the nutritional condition of anchovy late larvae (19-35 mm) in the field during both summer and late-Autumn seasons via lipid composition studies, and we also study their diet through the fatty acids found both in larvae and in zooplankton, which is the basic prey of anchovy at all the development stages (Plounevez and Champalbert, 1999; Bacha and Amara, 2009; Pasquaud et al., 2008; Borme et al., 2009; Catalán et al., 2010; Morote et al., 2010). Growth rates were also studied for both larvae populations to determine whether the different environmental conditions affect the early stage development of the anchovy. This work will thus help to explain to what extent the general conditions of an

unexpected December late larvae population differs from the August late larvae and whether these two populations have similar viability.

2. Materials and methods

2.1 Sample collection

Samples were collected during two cruises carried out in the Gulf of Lions (Northwestern Mediterranean sea; figure 1) in 2007 on board the *R/V L'Europe* (IFREMER, France). The first cruise (PELMED07) was conducted in the summer from the 28th of July to the 9th of August 2007 and the second cruise (JUVALION07) was carried out in late autumn, from the 8th to the 21st of December 2007.

Temperature and salinity of the water column from sea surface to 50 m depth were measured via a Seabird 19 CTD at each station. Also data of sea surface temperature in September to December since 1982 to early 2011 of Gulf of Lions area from NOAA were acquired to study possible trends.

Zooplankton samples for an analysis of biomass and plankton composition were collected using a standard WP2 net with a mesh size of 200 μm and sieved through a 3000 μm plankton mesh to obtain the 200-3000 μm mesozooplankton fraction, and by means of a scaled-down version WP2 net, with a mesh size of 53 μm and a mouth diameter of 25 cm, and sieved through a 200 μm plankton mesh to obtain the 53-200 μm fraction of microplankton. All zooplankton samples were split up with a Motoda plankton splitter (Motoda, 1959) and one-half was preserved in formalin to carry out subsequent qualitative analysis, while the other half was frozen on board for biomass

measurements and lipids analysis. As the purpose of comparatively assess the nutritional condition of the two different temporal larvae populations came up during the December cruise, plankton samples for lipids analysis were not collected during the first cruise.

Late larvae of anchovy were caught with a pelagic trawling net, towed at an average speed of 3.6 knots over a 30-40 min time span. This trawling time might seem too long to obtain larvae in proper conditions for biochemical analysis; however, the alteration of lipids composition in muscular tissue of fish larvae is small within a time up to 3 h after death (Lochmann et al., 1996).

Samples were immediately frozen in liquid nitrogen after sorting on board and transferred into a -80 °C freezer just after the arrival in the laboratory..

2.2 Lipid and fatty acid analysis

Wet weight and standard length of each larva were measured in the laboratory to the nearest 10 µg and 0.1 mm, respectively, before removing the head for otoliths analysis and the gut, as recommended by Lochmann et al. (1996). The empiric relationship between larval wet and dry weights was calculated via linear regression from other larvae of the same cruises and size ranges.

Microplankton and mesozooplankton samples from each station were pooled together before proceeding with the fatty acids extraction.

Lipid extraction was performed according to the method of Folch et al. (1957). The quantification of lipids content was carried out following the protocol of Olsen and Henderson (1989) via high-performance thin-layer chromatography (HPTLC), which

was followed by quantitative densitometry in visible light with a Bio-Rad Gel Doc XR densitometer, using Quantity One 4.6.2 software.

The evaluation of the nutritional condition of the *E. encrasicolus* late larvae was accomplished by comparing the triacylglycerol/cholesterol (TAG/CHOL) index (Håkanson, 1993) and the ratio between the percentage of total lipids and the dry mass (Norton et al., 2001). Fulton's condition index (FCI) was calculated with wet weight (W, in g) and standard length (SL, in mm) data, following the equation:

$$FCI = W \times 100 / SL^3$$

Fatty acids extraction and trans-methylation was accomplished following the protocol of Christie (1989) as modified by Li & Watkins (2001).

Four out of seven samples of larvae from August 2007, together with five larvae samples and three zooplankton samples from December 2007, were feasible to read by the gas-chromatographer.

Gas chromatographic (GC) analysis of fatty acid methyl esters (FAMES) were then performed using a Thermofisher Scientific GC8060 gas-chromatograph coupled with a MD800 mass-spectrometer. The apparatus was fitted with a BPX-70 capillary column (30 m x 0.25 mm i.d. x 0.25 µm.). Helium was used as carrier gas, with a speed of 1 ml/min. The programmed oven temperature was 60 °C (1 min.) to 260 °C (10 min) with an increment of 8 °C min⁻¹. The injector temperature was set at 270 °C and the injector split was set at 35 ml min⁻¹. Mass-spectrometry was conducted with an ion source temperature of 200 °C and an interphase temperature at 260 °C. Ionization was performed by electron impact at 70 eV, and the weight range analyzed was 50 – 550 Da.

FAMES were identified by comparing their retention times with those of the standard mixture, Supelco 37 Component FAME mix. The quantification of the identified FAMES was calculated through GC peak areas integration.

The diet of late larvae was evaluated according to the indices based on fatty acids relations 16:1(ω -7)/16:0, 18:1(ω -9)/18:1(ω -7) and EPA/DHA, and on PUFA/SFA relation (St. John and Lund., 1996; Auel et al., 2002; Rossi et al., 2006).

2.3 Growth analysis

A total of 61 larvae from August and 44 larvae from December, ranging in size from 19-31 mm (SL) and from 20-27 mm (SL), respectively, were used for a growth analysis. Both sagitta otoliths were extracted from the head of the anchovy larvae under a Leica dissection microscope (Wild M12) equipped with polarizing filters and mounted in Crystalbond 509 Amber on labeled glass slides. The otolith growth analysis was undertaken at 100X magnification under transmitted light with a microscope (Zeiss Axiospoth) coupled to a digital video recorder, while otolith nucleus was analyzed at 1000X magnification. Otolith radius (OR) and increment width (IW) (μ m) were measured to the nearest 0.1 μ m using Image-Pro Plus 5.0. The increments were measured along the longest radius, from the middle of one D-zone to the middle of the next D-zone. Following the results of Aldanondo et al. (2008), for the same species in the Bay of Biscay and under experimental conditions, increments were assumed to be daily (DI) being the first increment laid down at hatching. Until the paper of Aldanondo et al. (2008), previous works on European anchovy growth had assumed that the first increment deposition took place at the beginning of exogenous feeding, i.e. two days after hatching, as proposed by Palomera et al. (1988) in the first paper on anchovy larval

otoliths. All otoliths were read twice by two different persons, and only if the DI differed by 1 daily increment were they accepted.

Taking into account the narrow range of lengths that we are analyzing, we have assumed linear growth in agreement with the results of La Mesa et al. (2009), that is the first paper on anchovy that analyzes the growth at the metamorphic period, as is the case of our samples. Accordingly, the individual growth rate (IGR, mm d⁻¹) from the time of hatching until the time of capture was then calculated by using the equation proposed by Takahashi and Watanabe (2005).

$$IGR = (SL - SL_0) / Age$$

where SL is the measured larvae standard length corrected by using Theilacker's method (Theilacker, 1980), SL₀ is the larvae standard length at hatching, estimated to be 2.5 mm according to laboratory studies on the studied species (Regner, 1985), and Age = DI.

2.4 Data analysis

Seasonal differences between water oceanographic parameters as well as Fulton's condition index, lipids total content, lipid classes and proportion of fatty acids in larvae and plankton were assessed by means of Mann-Whitney non-parametric tests for independent samples (Dytham, 2003), and for oceanographic data ANOVA tests were performed.

Fatty acids percentage compositions were pairwise compared between larvae of both cruises, and between larvae and zooplankton collected during the December 2007 cruise by using the former test. Similarities in the fatty acid composition between

samples were measured by Euclidean distances (Legendre and Legendre, 1998). A non-metric multidimensional scaling (nMDS) was carried out on the samples similarity matrix to visually describe overall patterns.

Statistical analyses were carried over with software STATISTICA 6.0 by Statsoft, Inc., and PRIMER-E 6 software. Significance level for all tests was adopted at $p < 0.05$.

3. Results

3.1 Oceanographic data and zooplankton composition

The mean temperature within the water column (0-50 m) and the mean surface temperature (0-5 m) were significantly higher ($p < 0.0001$) in August (mean $\pm$ standard deviation values of $19.14\text{ }^{\circ}\text{C} \pm 1.32$ for surface temperature and $16.75\text{ }^{\circ}\text{C} \pm 1.02$ for average temperature) compared to December (mean $\pm$ standard deviation values of $12.64\text{ }^{\circ}\text{C} \pm 0.91$ for surface temperature and $13.04\text{ }^{\circ}\text{C} \pm 0.66$ for average temperature), while no significant differences were observed for salinity between the two cruises (mean $\pm$ standard deviation values of 37.74 ± 0.20 and 37.35 ± 1.48 for surface salinity and 37.94 ± 0.06 and 37.93 ± 0.40 for average salinity in August and December, respectively).

Data of temperature acquired from NOAA (Reynolds et al., 2002) were monthly averaged in the Gulf of Lions (area comprised between 2.5° W to 6.5° W and 41.3° N to 45.5° N), from September 1981 to December 2010, showing a positive trend in sea surface temperature during the last 4 months of every year (16.5° in September 1981 to

17.2°C in December 2010; $SST = 14.97 + 0.54 \times 10^{-4} \times \text{Serial_date}$; where Serial_date is the number of days since the January 1st of 1900).

The non-parametric Mann-Whitney U test revealed that there were significant differences for microplankton biomass between August and December cruises ($p < 0.05$), with higher values recorded during the summer (mean $\pm$ standard deviation values of $201.59 \pm 283.25 \text{ mg m}^{-3}$ and $22.76 \pm 24.18 \text{ mg m}^{-3}$, respectively).

The same pattern was observed for the mesozooplankton, where pairwise multiple comparisons (t-test) revealed that August 2007 samples had significantly higher biomass ($p < 0.05$) compared to those collected in December 2007 (mean $\pm$ standard deviation values of $33.65 \pm 10.34 \text{ mg m}^{-3}$ and $14.07 \pm 10.65 \text{ mg m}^{-3}$, respectively).

The analysis of both microplankton and mesozooplankton composition revealed a dominance of copepods, especially calanoids and cyclopoids, within the community during the two seasons (Fig. 2). Nevertheless, in August cladocerans were also important, while they were not recorded in December cruise (Fig. 3).

3.2 Growth rate

Otolith growth was significantly different between the two periods with the otoliths of the December larvae being smaller compared to those of the August larvae of the same age (t-test, $p < 0.001$) (Fig. 4). For both groups, increment width increased continuously, although for larvae caught in December, that ranged on age from 33 to 54 days, the largest increment width reached was half that of the maximum increment width in the August samples, that ranged from 22 to 44 days ($4 \text{ vs. } 8 \text{ } \mu\text{m day}^{-1}$, respectively).

This pattern was matched by mean individual growth rates (IGR) from hatching until the time of capture, that ranged between 0.50-0.93 mm d⁻¹ (mean $\pm$ standard deviation values of 0.74 ± 0.09 mm d⁻¹) for August larvae and 0.43-0.74 mm d⁻¹ (mean $\pm$ standard deviation values of 0.59 ± 0.07 mm d⁻¹) for December larvae, indicating that IGR was significantly higher in the warmer period (t-test, $p < 0.001$).

3.3 Nutritional condition

A single linear regression between larvae dry mass and wet weight was estimated for both cruises (Fig. 5) because the one-way ANCOVA did not show significant differences between August and December ($p > 0.05$), considering wet weight as covariant.

Subsequently, the empirical equation obtained from that relationship was applied to estimate the larvae dry mass (mean $\pm$ standard deviation values of 23.0 ± 7.0 and 18.0 ± 7.0 mg/ larva for August and December late larvae, respectively) as well as in further analysis regarding the dry mass of lipid contents in the samples.

Mean standard lengths did not significantly differ between the two cruises (mean $\pm$ standard deviation values of 27.3 ± 3.0 mm and 27.0 ± 3.4 mm for August 2007 and December 2007, respectively).

On the contrary, Fulton's condition index of larvae was significantly higher ($p < 0.05$) in the August individuals (mean $\pm$ standard deviation values of 0.598 ± 0.131 and 0.489 ± 0.153 in August and December, respectively).

Pools of 4-6 larvae each from August and December cruises were processed for lipid extraction.

Lipids content in larvae did not show any statistical differences between the two seasons ($p = 0.123$), and no significance was found regarding triacylglycerol ($p = 1.000$), cholesterol ($p = 0.705$), free fatty acid ($p = 0.570$) or polar lipid content ($p = 1.000$) within the anchovy samples (Table 1).

The TAG/CHOL ratio was determined to range between 0.53 – 0.72 for August larvae and 0.60 – 0.82 for December larvae, and it did not exhibit significant differences between cruises as well ($p = 0.186$).

3.4 Fatty acids analysis

Of the 23 fatty acids identified, 16:0, eicosapentaenoic acid or EPA (20:5(ω -3)) and docosahexanoic acid or DHA (22:6(ω -3)) made up 58-75 % of total fatty acids in the zooplankton and larvae samples (Table 2), with DHA and 16:0 being the most common fatty acids found in both August and December larvae and within the zooplankton. The other abundant fatty acids found were 14:0, 18:0, 16:1(ω -7), 18:1(ω -9) and 18:1(ω -7).

The proportion of PUFA was higher than any other type of FFAA among larvae of both cruises, while SFA were the most abundant in the zooplankton.

A multi-dimensional scaling (MDS), with a stress coefficient <0.01 , shows the similarity in fatty acid composition among the larvae of August and December and the zooplankton of December. Three groups can be differentiated on the plot, specifically a group comprised of 4 out of 5 samples of the December anchovy larvae, the 4 samples of August larvae, and the 3 samples of zooplankton (Fig. 6). ANOSIM test confirmed the presence of significant differences in the multivariate fatty acid composition between all three groups ($R^2 = 0.83$ at $p < 0.0001$).

Table 3 shows the mean values of the fatty acid indices estimated from the data obtained in this work, apart from those related to the summer zooplankton, which were acquired from (Rossi et al., 2006). It is important to notice that the composition of the zooplankton during the summer of 2007 was equal to that described by Rossi et al. (2006).

The indices 18:1(ω -9)/18:1(ω -7) and PUFA/SFA indicate the degree of carnivory in late anchovy larvae and zooplankton. No statistically significant difference was observed among the larvae of the two cruises or among the December larvae and the December zooplankton, although the overall values of these indices were relatively high (Auel et al., 2002).

The indices 16:1(ω -7)/16:0 and EPA/DHA, which estimate the importance of diatoms in the diet of larvae, show significant differences among August and December larvae, with both indices being higher in August. There is also a statistically significant difference between December larvae and zooplankton, with the larvae having a lower ratio than the zooplankton.

4. Discussion

Engraulis encrasicolus in the Northwestern Mediterranean has been intensively exploited (Palomera et al., 2007); therefore, alterations of any factor (e. g. temperature, salinity, currents, predation, food availability and overexploitation) affecting early stages (i.e. eggs, larvae and juveniles) of engraulids could have vital importance for the strength of recruitment and thus for the future of the population, due to their high larval

growth and mortality rates (Houde, 1989; Takahashi and Watanabe, 2005; Ruiz et al., 2006).

The evidences proving that European anchovy larvae feed on plankton (Conway et al., 1999; Tudela et al., 2002; Bacha and Amara, 2009; Morote et al., 2010) lead us to assume that changes in the plankton community affect the feeding habits of larvae, thus influencing their nutritional condition and, possibly, their survival (Fuiman and Cowan Jr., 2003).

Zooplankton biomass and taxonomic composition showed clear differences between August and December 2007, with the summer being the period when a higher presence of these organisms was recorded. Nevertheless, neither the TAG/CHOL index (Fraser et al., 1987) nor the lipids percentage nor the polar lipids content (Norton et al., 2001) indicates differences between seasons regarding the nutritional condition of anchovy late larvae. Håkanson (1989, 1993) pointed out that TAG/CHOL values below 0.2 - 0.3 indicate a poor nutritional condition, thus according to our results that showed TAG/CHOL average values around 0.64-0.70 in both seasons, it can be stated that both larvae in August and December presented a satisfactory nutritional condition. For this reason the observed differences in food availability cannot be assumed to affect the biochemical condition of both larvae populations.

According to Fulton's index, statistically significant differences can be observed between August and December larvae concerning physical condition, showing that weight at size is higher in summer larvae compared to those collected in late autumn. The apparent discrepancy between biochemical nutritional condition and physical condition data can be rationalized by considering that morphometric condition indices take longer to show the effects of food intake (Catalán et al., 2007). In addition, taking into account that information from both FCI and growth rates show effects of a previous

period compared to biochemical indices, it is not surprising that results of FCI agree with those of growth analysis. As the condition of our larvae needs to be evaluated for a short period of time, we consider that Fulton's index should not be taken into account to assess the nutritional condition in this study.

In this sense, the absence of differences between biochemical conditions of anchovy larvae observed in our study could be explained to a certain extent by the observed seasonal differences in sea water temperature. Moreover temperature is recognized to substantially influence the metabolic rates in marine organisms (McLaren, 1963; Ikeda, 1985) and in the specific zooplankton composition.

Indeed, during the summertime, larvae must compensate the higher metabolic rate imposed by the higher temperature, with an energetically richer diet. On the other hand, larvae could simply eat more. However, this assumption could not be enough to compensate the energy expense since the more preys are chased and caught the higher the energetic cost is.

In summer, despite the higher zooplankton biomass, a significant presence of cladocerans was recorded (Fig. 3). Cladocerans are considered of less energetic value than copepods (Boldt and Haldorson, 2002), in agreement with previous findings of Champalbert (1996) and Calbet et al. (2001). Moreover, stomach contents analysis carried out on juvenile stages of anchovy of the investigated area show a high preference for cladocerans in summer (Costalago, unpublished data). Although we cannot be sure that the studied late larvae feed on the exact same prey as juveniles do, feeding preferences of larvae are based on zooplankton rather than on any other group (Tudela et al., 2002; Borme et al., 2009; Catalán et al., 2010; Morote et al., 2010) due to their inability to catch smaller preys without a proper development of gill rakers (van der Lingen et al., 2006). Therefore it is very likely that anchovy late larvae are also

frequently feeding on cladocerans during the summer, as it happens in the Adriatic Sea (Borme et al., 2009).

On the other hand, the December larvae population is supposed to feed predominantly on copepods, which are higher energetic zooplankters. The differences in metabolic rates as well as in the energy content of diet could explain the similar nutritional condition observed in the seasonal pools of late larvae.

Masuda (2003) demonstrated that a lack of docosahexanoic acid could produce an ineffective feeding behavior, and even a higher mortality rate within a population. The finding that DHA was present and showed similar concentrations in both populations further confirm the occurrence of similar nutritional conditions in anchovy late larvae in both cruises. This result could also be explained by the fact that, in summer, lower DHA concentration was expected because a high UVB radiation can negatively affect DHA production by algae and August is the month with the highest solar radiation in the Western Mediterranean. In addition, Kainz et al. (2004) stated that the cladoceran community shows lower DHA concentrations. The higher plankton biomass available during August would help to compensate for the DHA deficit of larvae and result in equal conditions compared to winter.

Several studies regarding the effect that essential fatty acids (EFAs) have on fish development have been devised for reared larvae (Morais et al., 2007), but only a few examine natural populations at sea (Reuss and Poulsen, 2002). The present study intended to ascertain the role that FFAA play as trophic markers between anchovy late larvae and plankton in the Gulf of Lions, and to determine the main planktonic groups that are part of the diet of the larvae.

Anchovy fatty acid composition suffers seasonal variation (Zlatanov and Laskaridis, 2007). In particular, saturated fatty acids (SFAs) 16:0, 20:0, 22:0 and 24:0

can be more easily synthesized by all aquatic organisms (Dalsgaard et al., 2003), while PUFAs, which are the first preference for fish lipases (Lie and Lambertsen, 1985) and must be obtained from the diet, are present in low concentration in oligotrophic seas (Fahl and Kattner, 1993). In addition, PUFAs are critical for survival of marine organisms (Brett and Müller-Navarra, 1997). For these reasons, trophic indices used in this study were based basically on unsaturated fatty acids.

The higher proportion of PUFAs compared to other types obtained in this study can be easily explained when taking into account the accumulation of these FFAA with age in anchovy larvae (Rossi et al., 2006).

The absolute values of the indices 16:1(ω -7)/16:0 and EPA/DHA are comparatively low (Auel et al., 2002), and confirm a clear tendency towards a non-diatom diet (Table 3). The low value of the C16/C18 index corroborates the mentioned low preference for diatoms.

Although limited, diatom consumption appears significantly higher during summer. This should be expectable because there is a spring bloom of diatoms in this area (Leblanc et al., 2003). As shown by the low overall values of the above mentioned indices and by some other works (e.g. Conway et al., 1999; Tudela et al. 2002; Islam and Tanaka. 2009; Morote et al. 2010) anchovy larvae normally do not feed on phytoplankton. Thus, it is likely that any trace of phytoplankton in the larvae comes either indirectly from the consumed zooplankton or from accidentally ingested phytoplankton.

On the contrary, our find of moderately high levels of 18:1(ω -9) and long-chain monounsaturated fatty acids points out that calanoid copepods are important prey for anchovy late larvae (Werner and Auel, 2005), as showed in other studies with the

carnivory indices 18:1(ω -9)/18:1(ω -7) and PUFA/SFA (Cripps and Atkinson, 2000; Garrido et al., 2008).

To date, knowledge of European anchovy reproductive behavior in the western Mediterranean indicates that the spawning period begins in the spring, when the water starts to heat up and reaches 14 to 15 °C, and finishes in late-September, when water temperature starts to drop (Palomera, 1992).

However, the growth rates estimated in this work, together with otoliths analysis and age determination, indicate that anchovy larvae gathered in December 2007 were hatched approximately between the end of October and mid- November, which is well after the end of the spawning period previously known for this species (Palomera, 1992; Palomera et al., 2007). Thus, a prolongation of the spawning period took place that year, likely brought about by a process of sea surface heating. During the last decades, mean temperatures in the Western Mediterranean have been rising (Salat and Pascual, 2002; Reynolds et al., 2002). If the autumn-winter anchovy larvae population is becoming a norm in the biology of the species, we could likely assume that the spawning period of *E. encrasicolus* is being extended, favoring a wider spawning period and perhaps the survival of these late larvae of European anchovy in colder months.

Our analysis regarding the growth of both August and December cohorts shows that growth rates were significantly different among cohorts, being higher in August. This difference could be due to water temperature or to food availability (Takahashi and Watanabe, 2005; Aldanondo et al., 2008). Mean temperature during August 2007 was 19°C, while during December cruise it was 12 °C, a difference that could cause statistically different growth rates, yet still in agreement with the similar nutritional condition recorded in this study, as explained above in terms of different energetic expenses.

Feeding activities have a positive correlation to water temperature (Houde, 1989), so lower growth rates in December may be due to a reduction in food intake and metabolic rates, rather than to food shortage (Takahashi and Watanabe, 2005), as the good nutritional condition observed in this period seems to confirm. Moreover, the Gulf of Lions is a rich environment in terms of food availability, favouring a proper nutritional condition in anchovy larvae, as exposed by García et al. (1998).

The noted lack of significant differences both in the TAG/CHOL index and in the polar lipids content between seasons in the anchovy larvae would imply a major relevance of temperature in the differences found in growth rates. These differences are also confirmed by Takasuka and Aoki (2006), who found a direct relationship between temperature and growth rate in Japanese anchovy larvae.

Considering the results of this work, we cannot assure that either of the populations of larvae has a greater probability of survival than the other. Further research concerning the recruitment success of these late larvae populations is needed.

Some studies suggest that mortality of anchovy larvae of the Mediterranean is inversely related to growth rates (Allain et al., 2003; Palomera et al., 2007; La Mesa et al., 2009), supported by the idea that larvae with slower growth rate remain as larvae longer (“stage-duration” mechanism, see Chambers and Leggett (1987)) and extends the exposure to predation. In this sense, we think that the “bigger is better” hypothesis, as described by Leggett and Deblois (1994), is a paradigm that fits to our results.

Takasuka et al. (2003) also found that larvae of Japanese anchovy with lower growth rates were proportionally more abundant within predators’ stomachs than in the sea, inferring that were more vulnerable to predation. If this holds true, anchovy larvae found in December 2007 would be more vulnerable to predation and have less probability to reach the adult stage. However, further studies regarding the biology of

predators that feed on anchovy larvae should be carried out, in order to estimate how their feeding activity could affect the probability of mortality of anchovy larvae.

Acknowledgments

This research was conducted within the European project SARDONE (FP6 – 44294). The authors gratefully acknowledge the collaboration of the IFREMER scientific staff and of the captain and crew of the RV L'Europe for their help during the cruises. We also thank the collaboration of Itziar Álvarez and Juan Pablo Beltrán in the analysis of the otoliths and Guiomar Rotllant and the people of IRTA for their guidelines regarding lipids extraction.

References

- Aldanondo, N., Cotano, U., Etxebeste, E., Irigoien, X., Alvarez, P., Murguía, A.M. de, Herrero, D., 2008. Validation of daily increments deposition in the otoliths of European anchovy larvae (*Engraulis encrasicolus* L.) reared under different temperature conditions. *Fisheries Research* 93, 257–264.
- Allain, G., Petitgas, P., Grellier, P., Lazure, P., 2003. The selection process from larval to juvenile stages of anchovy (*Engraulis encrasicolus*) in the Bay of Biscay investigated by Lagrangian simulations and comparative otolith growth. *Fisheries Oceanography* 12, 407–418.
- Auel, H., Harjes, M., Rocha, R. da, Stübing, D., Hagen, W., 2002. Lipid biomarkers indicate different ecological niches and trophic relationships of the Arctic hyperiid amphipods *Themisto abyssorum* and *T. libellula*. *Polar Biology* 25, 374–383.
- Bacha, M., Amara, R., 2009. Spatial, temporal and ontogenetic variation in diet of anchovy (*Engraulis encrasicolus*) on the Algerian coast (SW Mediterranean). *Estuarine, Coastal and Shelf Science* 85 (2), 257–264.
- Barange, M., Coetzee, J., Takasuka, A., Hill, K., Gutierrez, M., Oozeki, Y., van der Lingen, C., Agostini, V., 2009. Habitat expansion and contraction in anchovy and sardine populations. *Progress In Oceanography* 83, 251–260.
- Boldt, J.L., Halderson, L.J., 2002. A bioenergetics approach to estimating consumption of zooplankton by juvenile pink salmon in Prince William Sound, Alaska. *Alaska Fish. Res. Bull* 9, 111–127.
- Borme, D., Tirelli, V., Brandt, S., Fonda Umani, S., Arneri, E., 2009. Diet of *Engraulis encrasicolus* in the northern Adriatic Sea (Mediterranean): ontogenetic changes and feeding selectivity. *Marine Ecology Progress Series* 392, 193–209.
- Brett, M., Müller-Navarra, D.C., 1997. The role of essential fatty acids in aquatic food web processes. *Freshwater Biology* 38, 483–499.
- Butler, J., 1991. Mortality and Recruitment of Pacific Sardine, *Sardinops sagax caerulea*, Larvae in the California Current. *Canadian Journal of Fisheries and Aquatic Sciences* 48, 1713–1723.
- Calbet, A., Garrido, S., Saiz, E., Alcaraz, M., Duarte, C.M., 2001. Annual zooplankton succession in coastal NW Mediterranean waters: the importance of the smaller size fractions. *Journal of Plankton Research* 23, 319–331.
- Catalán, I., Berdalet, E., Olivar, M. P., Roldán, C., 2007. Response of muscle-based biochemical condition indices to short-term variations in food availability in post-flexion reared sea bass *Dicentrarchus labrax* (L.) larvae. *Journal of Fish Biology* 70, 391–405.

- Catalán, I., Folkvord, A., Palomera, I., Quílez-Badía, G., Kallianoti, F., Tselepidis, A., Kallianotis, A., 2010. Growth and feeding patterns of European anchovy (*Engraulis encrasicolus*) early life stages in the Aegean Sea (NE Mediterranean). *Estuarine, Coastal and Shelf Science* 86, 299-312.
- Chambers, R.C., Leggett, W.C., 1987. Size and age at metamorphosis in marine fishes: an analysis of laboratory-reared winter flounder (*Pseudopleuronectes americanus*) with a review of variation in other species. *Canadian Journal of Fisheries and Aquatic Sciences* 44, 1936-1947.
- Champalbert, G., 1996. Characteristics of zooplankton standing stock and communities in the Western Mediterranean Sea: Relations to hydrology. *Scientia Marina* 60, 97-113.
- Christie, W.W., 1989. *Gas Chromatography and Lipids: A practical guide*. The Oily Press, Dundee, UK, 307 pp.
- Conway, D., Coombs, S., Smith, C., 1999. Feeding of anchovy *Engraulis encrasicolus* larvae in the northwestern Adriatic Sea in response to changing hydrobiological conditions. *Marine Ecology Progress Series* 175, 35-49.
- Cripps, G., Atkinson, A., 2000. Fatty acid composition as an indicator of carnivory in Antarctic krill, *Euphausia superba*. *Canadian Journal of Fisheries and Aquatic Sciences* 57, 31-37.
- Dalsgaard, J., St John, M., Kattner, G., Müller-Navarra, Dörthe, Hagen, W., 2003. Fatty acid trophic markers in the pelagic marine environment. *Advances in marine biology* 46, 225-340.
- Dytham, C., 2003. *Choosing and using statistics: a biologist's guide*. Wiley-Blackwell, U.K., 264 pp.
- Fahl, K., Kattner, G., 1993. Lipid Content and fatty acid composition of algal communities in sea-ice and water from the Weddell Sea (Antarctica). *Polar Biology* 13, 405-409.
- Folch, J., Lees, M., Sloane-Stanley, G., 1957. A simple method for the isolation and purification of total lipids from animal tissues. *The Journal of Biological Chemistry* 226, 497-509.
- Fraser, A.J., Sargent, J., Gamble, J., MacLachlan, P., 1987. Lipid class and fatty acid composition as indicators of the nutritional condition of larval Atlantic herring. *American Fisheries Society, Proceedings 10th Annual Larval Fish Conference*. Miami, Florida, USA, pp. 129-143.
- Fuiman, L.A., Cowan Jr, J.H., 2003. Behavior and recruitment success in fish larvae: repeatability and covariation of survival skills. *Ecology* 84, 53-67.
- García, A., Palomera, I., 1996. Anchovy early life history and its relation to its surrounding environment in the Western Mediterranean basin. *Scientia Marina* 60, 155-166.
- García, A., Cortés, D., Ramírez, T., 1998. Daily larval growth and RNA and DNA content of the NW Mediterranean anchovy *Engraulis encrasicolus* and their relations to the environment. *Marine Ecology Progress Series* 166, 237-245.
- Garrido, S., Rosa, R., Ben-Hamadou, R., Cunha, M. E., Chicharo, A., van der Lingen, C.D., 2008. Spatio-temporal variability in fatty acid trophic biomarkers in stomach contents and muscle of Iberian sardine (*Sardina pilchardus*) and its relationship with spawning. *Marine Biology* 154, 1053-1065.
- Håkanson, J.L., 1993. Nutritional condition and growth rate of anchovy larvae (*Engraulis mordax*) in the California Current: two contrasting years. *Marine Biology* 115, 309-316.
- Håkanson, J., 1989. Condition of larval anchovy (*Engraulis mordax*) in the Southern California Bight, as measured through lipid analysis. *Marine Biology* 102, 153-159.
- Hidalgo, M., Tomas, J., Hoeie, H., Morales-Nin, B., Ninnemann, U.S., 2008. Environmental influences on the recruitment process inferred from otolith stable isotopes in *Merluccius merluccius* off the Balearic Islands. *Aquatic Biology* 3, 195-207.
- Houde, E.D., 1989. Comparative growth, mortality, and energetics of marine fish larvae: Temperature and implied latitudinal effects. *Fishery Bulletin* 87, 471-495.
- Ikeda, T., 1985. Metabolic rates of epipelagic marine zooplankton as a function of body mass and temperature. *Marine Biology* 85, 1-11.
- Isari, M., Fragopoulou, N., Somarakis, S., 2008. Interannual variability in horizontal patterns of larval fish assemblages in the northeastern Aegean Sea (eastern Mediterranean) during early summer. *Estuarine, Coastal and Shelf Science* 79, 607-619.
- Islam, M.S., Tanaka, M., 2009. Diet and prey selection in larval and juvenile Japanese anchovy *Engraulis japonicus* in Ariake Bay, Japan. *Aquatic Ecology* 43, 549-558.
- Kainz, M., Arts, M.T., Mazumder, A., 2004. Essential fatty acids in the planktonic food web and their ecological role for higher trophic levels. *Limnology and Oceanography* 49, 1784-1793.
- Kruskal, J.B., Wish, M., 1978. *Multidimensional scaling*. Issues 7-11. SAGE University Paper, London, 93 pp.
- La Mesa, M., Donato, F., Giannetti, G., Arneri, E., 2009. Growth and mortality rates of European anchovy (*Engraulis encrasicolus*) in the Adriatic Sea during the transition from larval to juvenile stages. *Fisheries Research* 96, 275-280.
- Leblanc, K., Quéguiner, B., Garcia, N., Rimmelin, P., Raimbault, P., 2003. Silicon cycle in the NW Mediterranean Sea: seasonal study of a coastal oligotrophic site. *Oceanologica Acta* 26, 339-355.
- Legendre, P., Legendre, L., 1998. *Numerical Ecology*. Elsevier, Amsterdam. 853 pp.
- Leggett, W.C., DeBlois, E., 1994. Recruitment in marine fishes: Is it regulated by starvation and predation in the egg and larval stages? *Netherlands Journal of Sea Research* 32, 119-134.
- Li, Y., Watkins, B.A., 2001. Analysis of fatty acids in food lipids. In Wrolstad, R.E., (ed.), *Currents Protocols in Food Analytical Chemistry*. John Wiley, New York, p. D1.2.1.
- Lie, Ø., Lambertsen, G., 1985. Digestive lipolytic enzymes in cod (*Gadus morhua*): fatty acid specificity. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 80, 447-450.

- Lochmann, S., Maillet, G., Taggart, C., Frank, K., 1996. Effect of gut contents and lipid degradation on condition measures in larval fish. *Marine Ecology Progress Series* 134, 27-35.
- Masuda, R., 2003. The critical role of docosahexaenoic acid in marine and terrestrial ecosystems: from bacteria to human behavior. In *The Big Fish Bang. Proceedings 26th Annual Larval Fish Conference*. Institute of Marine Research, Bergen, Norway. pp. 249-256.
- McLaren, I.A., 1963. Effects of temperature on growth of zooplankton and the adaptive value of vertical migration. *Journal of the Fisheries Research Board of Canada* 20, 685-727.
- Morais, S., Conceicao, L., Ronnestad, I., Koven, W., Cahu, C., Zamboninoinfante, J., Dinis, M., 2007. Dietary neutral lipid level and source in marine fish larvae: Effects on digestive physiology and food intake. *Aquaculture* 268, 106-122.
- Morais, P., Babaluk, J., Correia, A.T., Chicharo, M.A., Campbell, J.L., Chicharo, L., 2010. Diversity of anchovy migration patterns in an European temperate estuary and in its adjacent coastal area: Implications for fishery management. *Journal of Sea Research* 64 (3), 295-303.
- Morote, E., Olivar, M.P., Villate, F., Uriarte, I., 2010. A comparison of anchovy (*Engraulis encrasicolus*) and sardine (*Sardina pilchardus*) larvae feeding in the Northwest Mediterranean: influence of prey availability and ontogeny. *ICES Journal of Marine Science* 67, 897-908.
- Motoda, S., 1959. Devices of simple plankton apparatus. *Memoirs of the Faculty of Fisheries, Hokkaido University* 7, 73-94.
- Norton, E., MacFarlane, R., Mohr, M., 2001. Lipid class dynamics during development in early life stages of shortbelly rockfish and their application to condition assessment. *Journal of Fish Biology* 58, 1010-1024.
- Olsen, R., Henderson, R., 1989. The rapid analysis of neutral and polar marine lipids using double-development HPTLC and scanning densitometry. *Journal of Experimental Marine Biology and Ecology* 129, 189-197.
- Palomera, I., Morales-Nin, B., Lleonart, J., 1988. Larval growth of anchovy, *Engraulis encrasicolus*, in the Western Mediterranean Sea. *Marine biology* 99, 283-291.
- Palomera, I., 1992. Spawning of anchovy *Engraulis encrasicolus* in the Northwestern Mediterranean relative to hydrographic features in the region. *Marine Ecology Progress Series* 79, 215-223.
- Palomera, I., Olivar, M., Salat, J., Sabates, A., Coll, M., Garcia, A., Morales-Nin, B., 2007. Small pelagic fish in the NW Mediterranean Sea: An ecological review. *Progress In Oceanography* 74, 377-396.
- Pasquaud, S., Elie, P., Jeantet, C., Billy, I., Martinez, P., Girardin, M., 2008. A preliminary investigation of the fish food web in the Gironde estuary, France, using dietary and stable isotope analyses. *Estuarine, Coastal and Shelf Science* 78 (2), 267-279.
- Pertierra, J.P., Lleonart, J., 1996. NW Mediterranean anchovy fisheries*. *Scientia Marina* 60 (S2), 257-267.
- Plounevez, S., Champalbert, G., 1999. Feeding behaviour and trophic environment of *Engraulis encrasicolus* (L.) in the Bay of Biscay. *Estuarine, Coastal and Shelf Science* 49 (2), 177-191.
- Regner, S., 1985. Ecology of planktonic stages of the anchovy, *Engraulis encrasicolus* (Linnaeus, 1758), in the central Adriatic. *Acta Adriatica* 26, 1-113.
- Reuss, N., Poulsen, L., 2002. Evaluation of fatty acids as biomarkers for a natural plankton community. A field study of a spring bloom and a post-bloom period off West Greenland. *Marine Biology* 141, 423-434.
- Reynolds, R.W., Rayner, N.A., Smith, T.M., Stokes, D.C., and Wang, W., 2002. An improved in situ and satellite SST analysis for climate. *Journal of Climate* 15, 1609-1625.
- Rossi, S., Sabatés, A., Latasa, M., Reyes, E., 2006. Lipid biomarkers and trophic linkages between phytoplankton, zooplankton and anchovy (*Engraulis encrasicolus*) larvae in the NW Mediterranean. *Journal of Plankton Research* 28, 551.
- Ruiz, J., García-Isarch, E., Huertas, E., Prieto, L., Juárez, A., Muñoz, J.L., Sánchez-Lamadrid, A., Rodríguez-Gálvez, S., Naranjo, J.M., Baldó, F., 2006. Meteorological and oceanographic factors influencing *Engraulis encrasicolus* early life stages and catches in the Gulf of Cadiz. *Deep Sea Research Part II: Topical Studies in Oceanography* 53, 1363-1376.
- Salat, J., 1996. Review of hydrographic environmental factors that may influence anchovy habitats in northwestern Mediterranean. *Scientia Marina* 60, 21-32.
- Salat, J., Pascual, J., 2002. Tracking long term hydrological change in the Mediterranean Sea. In *CIESM Workshop Series* 16, pp. 39-32.
- St. John, M., Lund, T., 1996. Lipid biomarkers: linking the utilization of frontal plankton biomass to enhanced condition of juvenile North Sea cod. *Marine Ecology Progress Series* 131, 75-85.
- Takahashi, M., Watanabe, Y., 2005. Effects of temperature and food availability on growth rate during late larval stage of Japanese anchovy (*Engraulis japonicus*) in the Kuroshio-Oyashio transition region. *Fisheries Oceanography* 14, 223-235.
- Takasuka, A., Aoki, I., Mitani, I., 2003. Evidence of growth-selective predation on larval Japanese anchovy *Engraulis japonicus* in Sagami Bay. *Marine Ecology Progress Series* 252, 223-238.
- Takasuka, A., Aoki, I., 2006. Environmental determinants of growth rates for larval Japanese anchovy *Engraulis japonicus* in different waters. *Fisheries Oceanography* 15, 139-149.
- Theilacker, G., 1980. Changes in body measurements of larval northern anchovy, *Engraulis mordax*, and other fishes due to handling and preservation. *Fishery Bulletin* 78, 685-692.
- Tudela, S., Palomera, I., Quilez, G., 2002. Feeding of anchovy *Engraulis encrasicolus* larvae in the north-west Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* 82, 349-350.
- van der Lingen, C. D., Hutchings, L., Field, J.G., 2006. Comparative trophodynamics of anchovy *Engraulis encrasicolus* and sardine *Sardinops sagax* in the southern Benguela: are species alternations between small pelagic fish trophodynamically mediated? *African Journal of Marine Science* 28, 465-477.

Ward, T.M., Mcleay, L.J., Dimmlich, W.F., Rogers, P.J., McClatchie, S., Matthews, R., Kampf, J., Van Ruth, P.D., 2006. Pelagic ecology of a northern boundary current system: effects of upwelling on the production and distribution of sardine (*Sardinops sagax*), anchovy (*Engraulis australis*) and southern bluefin tuna (*Thunnus maccoyii*) in the Great Australian Bight. Fisheries Oceanography 15, 191-207.

Werner, I., Auel, H., 2005. Seasonal variability in abundance, respiration and lipid composition of Arctic under-ice amphipods. Marine Ecology Progress Series 292, 251-262.

Zlatanov, S., Laskaridis, K., 2007. Seasonal variation in the fatty acid composition of three Mediterranean fish: sardine (*Sardina pilchardus*), anchovy (*Engraulis encrasicolus*) and picarel (*Spicara smaris*). Food Chemistry 103, 725-728.

Fig. 1. Map of the study area, with the positions of plankton stations and trawls in August and December 2007.

Fig. 2. Relative abundances of the main copepods species within micro and mesozooplankton samples of both August and December 2007 campaigns.

Fig. 3. Pie charts illustrating the mean percentage composition of the main plankton groups in August and December 2007 cruises.

Fig. 4. Anchovy larvae mean increment width by estimated age (days) for larvae caught in August and December 2007. (Error bars: standard deviation).

Fig. 5. Relationship between dry mass (DM) and wet weight (WW) of anchovy late larvae. Linear regression fitted by $DM = 0.1692 \cdot WW + 1.8262$ ($r^2 = 0.9578$).

Fig. 6. Nonmetric Multi-dimensional scaling (nMDS) plot of similarities in the fatty acid composition of anchovy late larvae and zooplankton. *Data of zooplankton summer from Rossi et al. (2006).

Figure 1.

Figure 2.

July/August 2007

December 2007

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Table 1

Total lipid content, lipid class values and TAG/CHOL index ($\mu\text{g}/\text{larva}^{-1}$), presented as Mean $\pm$ SD.

	August 2007 (N = 7 pools of 4–6 larvae each)	December 2007 (N = 5 pools of 4–5 larvae each)
Total lipid content	770.4 $\pm$ 275.1	664.5 $\pm$ 92.7
% lipid/dry weight	4.2 $\pm$ 0.8	4.4 $\pm$ 0.2
<i>Neutral lipids</i>		
Triacylglycerol (TAG)	922.2 $\pm$ 379.4	873.8 $\pm$ 242.2
Cholesterol	1414.4 $\pm$ 476.6	1227.8 $\pm$ 249.9
Free fatty acid	1406.0 $\pm$ 425.6	1228.2 $\pm$ 221.7
Steryl ester	–	–
Polar lipids	230.6 $\pm$ 65.9	223.1 $\pm$ 37.8
TAG/CHOL index	0.64 $\pm$ 0.06	0.70 $\pm$ 0.09

Table 2

Fatty acids composition of anchovy late larvae and zooplankton, presented as mean $\pm$ SD.

	Engraulis encrasicolus		Zooplankton
	August 2007 (N = 4)	December 2007 (N = 5)	December 2007 (N = 3)
<i>Fatty acids</i>			
14:0 ^{a,b}	5.06 $\pm$ 0.39	1.71 $\pm$ 0.32	8.94 $\pm$ 2.10
15:0 ^{a,b}	0.61 $\pm$ 0.02	0.49 $\pm$ 0.06	0.88 $\pm$ 0.37
16:0	23.39 $\pm$ 1.47	25.63 $\pm$ 7.17	24.31 $\pm$ 4.83
17:0 ^a	0.51 $\pm$ 0.02	0.69 $\pm$ 0.03	0.67 $\pm$ 0.19
18:0 ^b	4.22 $\pm$ 0.52	4.32 $\pm$ 0.12	5.73 $\pm$ 0.48
20:0 ^a	0.06 $\pm$ 0.01	0.04 $\pm$ 0.003	0.11 $\pm$ 0.02
22:0	0.05 $\pm$ 0.01	0.04 $\pm$ 0.01	–
24:0 ^a	0.14 $\pm$ 0.03	0.08 $\pm$ 0.02	0.07 $\pm$ 0.00
Total saturated	34.0	33.0	40.7
15:1	0.04 $\pm$ 0.01	0.04 $\pm$ 0.01	0.11 $\pm$ 0.00
16:1 (ω -7) ^{a,b}	3.66 $\pm$ 0.38	1.42 $\pm$ 0.09	6.47 $\pm$ 1.74
18:1 (ω -9) ^{a,b}	5.16 $\pm$ 0.33	5.57 $\pm$ 0.13	6.82 $\pm$ 1.48
18:1 (ω -7)	1.91 $\pm$ 0.28	1.92 $\pm$ 0.16	1.84 $\pm$ 0.36
20:1 (ω -9)	0.20 $\pm$ 0.15	0.13 $\pm$ 0.01	0.42 $\pm$ 0.10
22:1 (ω -9)	–	–	0.08 $\pm$ 0.00
24:1 (ω -9) ^a	0.12 $\pm$ 0.03	0.27 $\pm$ 0.05	0.08 $\pm$ 0.00
Total monounsaturated	11.1	9.3	15.8
18:2 (ω -6) ^a	1.19 $\pm$ 0.11	0.62 $\pm$ 0.12	0.83 $\pm$ 0.29
18:3 (ω -6)	0.55 $\pm$ 0.09	0.40 $\pm$ 0.05	0.64 $\pm$ 0.08
18:3 (ω -3) ^{a,b}	0.85 $\pm$ 0.15	1.18 $\pm$ 0.15	1.89 $\pm$ 0.05
20:3 (ω -6)	0.04 $\pm$ 0.00	0.03 $\pm$ 0.01	–
20:3 (ω -3)	0.05 $\pm$ 0.01	0.07 $\pm$ 0.01	–
20:4 (ω -6)	0.48 $\pm$ 0.06	0.55 $\pm$ 0.09	0.31 $\pm$ 0.05
20:5 (ω -3)	11.29 $\pm$ 0.64	10.79 $\pm$ 0.85	16.57 $\pm$ 6.34
22:6 (ω -3) ^b	34.25 $\pm$ 2.96	39.28 $\pm$ 4.47	17.35 $\pm$ 6.01
Total polyunsaturated	48.7	52.9	37.6
Others	6.2	4.8	5.9
Total	100	100	100

^a Statistical difference between August and December 2007 larvae, $p < 0.05$.

^b Statistical difference between December 2007 larvae and zooplankton, $p < 0.05$.

Table 3

Fatty acids trophic indices in August and December 2007 anchovies larvae and December 2007 zooplankton, presented as mean $\pm$ SD.

	Engraulis encrasicolus		Zooplankton	
	August 2007	December 2007	June 2000 ^a	December 2007
16:1(ω -7)/16:0	0.16 $\pm$ 0.01	0.06 $\pm$ 0.01	0.96	0.27 $\pm$ 0.08
20:5 (ω -3)/22:6 (ω -3)	0.33 $\pm$ 0.01	0.27 $\pm$ 0.02	0.53	0.96 $\pm$ 0.42
C16/C18	1.79 $\pm$ 0.38	1.94 $\pm$ 0.55	1.70	1.76 $\pm$ 0.30
18:1(ω -9)/18:1 (ω -7)	2.70 $\pm$ 0.30	2.89 $\pm$ 0.23	3.15	3.71 $\pm$ 0.62
PUFA/SFA	1.43 $\pm$ 0.21	1.60 $\pm$ 0.42	0.65	0.92 $\pm$ 0.42

^a June 2000 zooplankton taken as reference level from Rossi et al. (2006).