

1 Production of bioavailable and refractory dissolved organic matter by coastal heterotrophic
2 microbial populations

3 Christian Lønborg^{a,b*}, Xosé A. Álvarez-Salgado^b, Keith Davidson^a & Axel E.J. Miller^a

4 ^aScottish Association for Marine Science, Dunstaffnage Marine Laboratory Oban, Argyll,
5 PA37 1QA, United Kingdom

6 ^bCSIC, Instituto de Investigacións Mariñas, Eduardo Cabello 6, 36208 Vigo, Spain

7

8 *Corresponding author:

9 Tel. +34 986 231 930

10 Fax. +34 986 292 762

11 Email: clonborg@gmail.com

12

Abstract

Production of dissolved organic matter (DOM) by heterotrophic microbial communities isolated from Loch Creran (Scotland) was studied in time course incubations in which cells were re-suspended in artificial seawater amended with variable proportions of glucose, ammonium and phosphate. The incubation experiments demonstrated that microheterotrophs released part of the substrate as new DOM, with a production efficiency of 11 ± 1 % for DOC, 18 ± 2 % for DON and 17 ± 2 % for DOP. Estimating the impact of this production in Loch Creran, showed that from 3 ± 1 % (DOC) to 72 ± 16 % (DOP) of DOM could originate from the heterotrophic microbial community. The produced DOM (PDOM) was both bioavailable (BDOM) and refractory (RDOM). Bioavailability as assessed by the difference between the maximum and the end DOM concentration, was generally higher than found in natural systems, with DOP (73 ± 15 %, average \pm SD) more bioavailable than DON (70 ± 15 %), and DON than DOC (34 ± 13 %).

The stoichiometry of PDOM was linked to both nutrient uptake and BDOM ratios. Absorption and fluorescence of DOM increased significantly during the incubation time, indicating that microheterotrophs were also a source of coloured DOM (CDOM) and that they produce both bioavailable protein-like and refractory humic-like fluorophores.

Key words: DOM, coloured DOM, bioavailable DOM, refractory DOM, stoichiometry, microbial heterotrophy

34 **1. Introduction**

35 River inputs of inorganic nutrients support high primary production rates in coastal
36 marine ecosystems (Lalli & Parsons, 1997), which lead to the release of significant
37 amounts of autochthonous dissolved organic matter (DOM) (Nagata, 2000). Although
38 heterotrophic microbial communities are the major consumers of this DOM (Seitzinger &
39 Sanders, 1997), considerably less attention has been paid to their ability to produce DOM.
40 Some studies have suggested that microheterotrophs could be a significant but hardly
41 quantified source of bioavailable and refractory DOM in marine systems (Ogawa et al.,
42 2001; Kawasaki & Benner, 2007).

43 Release of DOM by phytoplankton has been repeatedly observed in both laboratory
44 and field studies, with exudation rates, stoichiometry and availability being related to
45 nutrient levels (e.g. Obernosterer & Herndl, 1995). Some studies have suggested that
46 bacteria are able to alter their C:N:P biomass ratios depending on the organic substrate
47 ratios (e.g. Tezuka, 1990), while others find constant biomass ratios (Golman et al. 1987).
48 However, the effect of changing C:N:P substrate ratios on the heterotrophic microbial
49 production of DOM is largely unknown.

50 Heterotrophic microbes have been shown to produce chromophoric DOM (CDOM)
51 during mineralization processes (e.g. Rochelle-Newall & Fisher, 2002). CDOM absorbs
52 light in the ultraviolet (UV) region declining to near-zero levels in the red region of the
53 spectrum (Stedmon & Markager 2001). A part of the absorbed light is reemitted at longer
54 wavelengths (FDOM), with two main DOM fluorophores being identified: the protein- and
55 humic-like (Coble et al. 1990; 1996). Protein-like fluorescence (FDOM_t) has been

suggested as an indicator of freshly produced DOM, while humic-like fluorophores (FDOMm) characterize older, more refractory DOM (Coble et al. 1996).

We hypothesised that marine heterotrophic microbial communities could be a quantitative important source of DOM in coastal marine waters, with the stoichiometry of the produced DOM depending on the substrate ratios. This was investigated by isolating coastal heterotrophic microbial populations from Loch Creran, Scotland and growing the community in 0.2 µm filtered artificial seawater, amended with carbon (glucose), inorganic nitrogen (ammonium) and phosphorus (phosphate) as growth media. Over the time course of the experiment dissolved organic carbon (DOC), nitrogen (DON), and phosphorus (DOP) concentrations, as well as DOM fluorescence were monitored as described below.

2. Material and methods

2.1. Incubation experiments

Samples for the microbial inoculum were obtained during winter and spring (16 January and 20 March 2007) from a depth of 5 m in the Scottish fjord Loch Creran. These sampling dates were chosen to investigate the production of DOM by microbial heterotrophic populations collected under different biological and hydrological conditions (See Lønborg et al. 2009).

The artificial seawater was prepared with water treated in a Milli-Q ultraviolet (UV) plus purification unit, which resulted in low carbon and nutrient content. NaCl, KCl, NaHCO₃, Na₂SO₄, MgCl₂·6H₂O, CaCl₂·2H₂O, KBr, H₃BO₃, NaF, SrCl₂·6H₂O, Na₂SiO₃·6H₂O, FeCl₃·6H₂O, MnSO₄·6H₂O, ZnSO₄·9H₂O and CoSO₄·9H₂O modified from Harrison et al. 1980, were added to the Milli-Q water to reach salinity levels equal to the

79 microbial inoculum, 28.2 (16 January) and 28.3 (20 March) repectively. NaCl, Na₂SO₄, and
80 KCl were combusted (450°C for 4 hours) before use to remove organic carbon. Glucose
81 (C₆H₁₂O₆), ammonium (NH₄Cl) and phosphate (KH₂PO₄) were added as substrate and pH
82 was adjusted to 8.0 with either HCl or NaOH. The microbial inoculum was prepared by
83 gravity filtering the seawater from Loch Creran two times through precombusted GF/F
84 filters. The GF/F filtrate was thereafter inoculated into the media in a 1: 50 dilution. A total
85 of 16 incubations were conducted in duplicate in two litre amber glass bottles and incubated
86 at a constant temperature of 14°C in the dark. The DOC concentration of the artificial
87 seawater, 20 µmol L⁻¹, originated from the added salts and the inoculum it was considered
88 refractory and was therefore subtracted from all samples. If we consider that the 20 µmol
89 L⁻¹ is 100% bioavailable and that the microbes would convert 11% (as found in this study)
90 of this DOC into new DOC, this would lead to an underestimation of the microbial
91 produced pool of ~ 2µmol L⁻¹ and 1.5µmol L⁻¹ for the refractory pool (considering a
92 bioavailability of 33%).

93 Dissolved inorganic nitrogen (DIN) and phosphorus (DIP) concentrations of the
94 artificial seawater were below the detection limit. Glucose, ammonium and phosphate
95 concentrations added to the medium ranged from 163 to 867 µmol L⁻¹ of C, 21.4 to 157.8
96 µmol L⁻¹ of N, and 1.3 to 13.6 µmol L⁻¹ of P (Table 1). C:N ratios of the added substrate
97 varied between 2 and 17, C:P ratios from 32 to 311 and 5 to 57 for N:P. These ratios were
98 chosen to vary widely around the average C:N:P ratio of bacterial biomass (50:10:1)
99 (Fagerbakke et al. 1996).

100 Since dissolved organic nitrogen (DON) and phosphorus (DOP) concentrations were
101 below the detection limit at the beginning of the experiments, it has been assumed that the
102 DON and DOP produced during the experiment was of microbial origin, and was named as

103 produced DOM (PDOM). Sub-samples were collected at days 0, 3, 11, 30, 50 and 100 after
 104 the beginning of the experiments to follow: dissolved organic matter absorbance (CDOM)
 105 and fluorescence (FDOM), glucose, DOC, DIN, DIP, total dissolved nitrogen (TDN) and
 106 phosphorus (TDP), total organic carbon (TOC), total nitrogen (TN) and phosphorus (TP).
 107 Samples for the dissolved phase were filtered through prewashed (> 1 litre of sterile filtered
 108 Milli-Q water) 0.2 µm polycarbonate membranes. Samples for CDOM analysis were stored
 109 in amber glass bottles at 4°C in the dark. Glucose and FDOM samples (20 mL) were
 110 collected in pre-combusted (450°C for 6 hours) glass ampoules and stored frozen (-20°C)
 111 in the dark until analyzed. The effect of freezing the FDOM samples was measured by
 112 taking samples over time with no major impact found. Samples (10 ml) for TOC, DOC, TN
 113 and TDN analysis were collected in pre-combusted (450°C for 6 hours) glass ampoules and
 114 preserved by adding 10 µL 85 % H₂PO₄. Samples for DIN and DIP were stored in acid
 115 washed polyethylene bottles and kept frozen (-20°C). TDP and TP samples were collected
 116 in glass bottles and frozen (-20°C) until analysis. All glassware used was acid-washed (2
 117 µmol L⁻¹ HCl) for 24 hours and washed three times with Milli-Q water before use.

118 2.3. Sample analysis

119 The absorption of CDOM was measured in four replicates in a Thermo Nicolet
 120 Evolution 300 Turn spectrophotometer using Milli-Q water as a blank. Before analysis
 121 samples and Milli-Q water were warmed to room temperature. The absorption was
 122 measured in a 1 cm quartz cuvette at 375 nm. The absorption coefficients (α_λ) were
 123 calculated as follows:

$$\alpha_\lambda = \frac{2.303 \cdot A_\lambda}{L}$$

124

125 Where A_λ is the optical density and the denominator L is the cell path-length in metres
126 (Stedmon & Markager, 2001).

127 FDOM was measured in duplicate in a Perkin Elmer LS 55 luminescence
128 spectrophotometer. The instrument was equipped with a xenon discharge lamp, using a 1
129 cm quartz fluorescence cell. Milli-Q water was used as a reference, and subtracted the
130 FDOM measurements. Discrete excitation/emission (Ex/Em) pair measurements were
131 performed at peaks T (aromatic amino acids, average Ex/Em, 280/350 nm; FDOMt) and M
132 (marine humic substances, average Ex/Em 320/410 nm; FDOMm), according to Nieto-Cid
133 et al. (2006). Following Nieto Cid et al. (2006), fluorescence units were expressed in ppb
134 equivalents of tryptophan (ppb Trp) for FDOMt and ppb equivalents of quinine sulphate
135 (ppb QS) for FDOMm.

136 Glucose was measured in triplicate at days 0, 3, 11 and 30, using the enzyme assay
137 described by Hicks & Carey (1968). Concentrations were determined using a four-point
138 standard curve following the subtraction of a Milli-Q blank. DOC, TOC, TDN and TN
139 were measured in at least triplicate, using a nitrogen-specific Antek 7020 nitric oxide
140 chemiluminescence detector, coupled in series with the carbon-specific infrared gas
141 analyser of a Shimadzu TOC-5000 organic carbon analyser. Three to five replicate
142 injections of 150 μ l were performed per sample. Concentrations were determined by
143 subtracting a Milli-Q blank and dividing by the slope of a daily standard curve. DON
144 concentrations were obtained by subtracting DIN from TDN ($\text{DON} = \text{TDN} - \text{DIN}$), with
145 the standard error calculated as $\text{SE}_{\text{DON}}^2 = \text{SE}_{\text{TDN}}^2 + \text{SE}_{\text{DIN}}^2$. DIN (ammonium and
146 nitrate+nitrite) was measured in four replicates with a Lachat (quickchem 500) auto
147 analyzer using standard protocols (Hansen & Grasshoff 1983). Triplicate samples were
148 analyzed for DIP by the standard molybdenum blue technique using a 5 cm cuvette

(Koroleff, 1983). TDP and TP were measured in triplicate by oxidation to soluble reactive phosphorous with the addition of sulphuric acid and persulphate (Koroleff, 1983), following autoclaving at 100°C for 90 min. The standard error for DOP was estimated as: $SE^2_{DOP} = SE^2_{TDP} + SE^2_{SRP}$. POM concentrations were calculated as the difference between TOC and DOC for POC, TN and TDN for PON and TP and TDP for POP. The corresponding standard errors were calculated as $SE^2_{POC} = SE^2_{TOC} + SE^2_{DOC}$, $SE^2_{PON} = SE^2_{TN} + SE^2_{TDN}$ and $SE^2_{POP} = SE^2_{TP} + SE^2_{TDP}$, respectively. Regression model II was used to examine the linear relationship between pairs of variables (Sokal & Rohlf 1995). In the cases where the intercept was not significantly different from zero, the intercept was set to zero and a new slope was calculated. The confidence level was set at 95% with the statistical analysis conducted in Statistica 6.0.

3. Results

3.1. *Production of DON and DOP by heterotrophic microbial populations*

TN (Fig. 1b) and TP (Fig. 1c) remained approximately constant throughout the incubations, TOC decreased rapidly during the first phase of all the cultures (initial 30 days) and thereafter more slowly (Fig. 1a). Particulate organic matter (POM) reached maximum values at day 10 with values between 12 and 138 $\mu\text{mol L}^{-1}$ for POC (Fig. 1a), 4.7 to 49.3 $\mu\text{mol L}^{-1}$ for PON (Fig. 1b) and 0.6 to 4.4 $\mu\text{mol L}^{-1}$ for POP (Fig. 1c). DIN and DIP concentrations were not significantly different from TN and TP at the beginning of the incubations. The time profile of nutrients concentrations was V-shaped, with a pronounced decrease to a minimum during the first and an increase during the second phase of the cultures (Figs 1b, c). Nitrate+nitrite concentrations were undetectable at the beginning of the incubations, but increased from day 50 onwards in all experiments, concomitantly with

the corresponding decrease of ammonium (Fig. 1d). The total uptake of DIN (UDIN) and DIP (UDIP) (Table 1) was calculated as $UDIN = DIN_0 - LDIN$ and $UDIP = DIP_0 - LDIP$, where DIN_0 and DIP_0 are the initial nutrient concentrations and LDIN, LDIP are the minimum DIN and DIP concentrations obtained at the end of the first phase of the cultures. DOM maximum was found at day 30 of the incubations (Fig. 2), associated with complete depletion of the added glucose (data not shown). The produced DOM (PDOM) reached concentrations of 38 to 81 $\mu\text{mol L}^{-1}$ for PDOC, 2.2 to 9.1 $\mu\text{mol L}^{-1}$ for PDON, and 0.24 to 0.78 $\mu\text{mol L}^{-1}$ for PDOP (Table 1). Calculating the bioavailability (BDOM) as the difference between PDOM and the refractory DOM (RDOM) at the end of the incubations, it resulted that $33 \pm 12 \%$ (average \pm SD) of PDOC, $70 \pm 16 \%$ of PDON and $79 \pm 14 \%$ of PDOP were bioavailable. The PDOM C:N ratios varied between 6 and 16, C:P ratios from 20 to 224 and 5 to 21 for N:P. Resulting C:N:P ratios varied between 20: 5: 1 and 225: 14: 1 (average 103: 11: 1). The stoichiometry of PDOM was linearly related with the C: N: P ratios of UDOC, UDIN and UDIP (Fig 3a, c, e).

3.3. Production of chromophoric DOM

Table 3a shows how a_{375} increased from being undetectable at the beginning of the incubations reaching an average value of $0.47 \pm 0.26 \text{ m}^{-1}$ at the end. In parallel to the production of CDOM, FDOM also increased during the incubations above background levels in all cases, with a mean (\pm SE) of $12.9 \pm 5.6 \text{ ppb Trp}$ for FDOMt and $6.4 \pm 3.8 \text{ ppb QS}$, for FDOMm (Table 3b, c). The amounts of FDOMt and FDOMm produced increased with substrate uptake and produced DOM (Table 4).

4. Discussion

194 Single-substrate approaches as used in this study are an obvious simplification of
195 reality, and will not fully represent the natural microbial growth substrate. These
196 experiments should therefore be seen as a first trial to quantify heterotrophic microbial
197 production of DOM.

198 Two contrasting phases have been identified during all the incubations (Fig. 1) : i) an
199 initial net-anabolic phase, when the added glucose was still detectable, as characterised by
200 the net consumption of the glucose, DIN and DIP and a net production of DOM; and ii) a
201 final net-catabolic phase, where glucose was undetectable, characterised by the net
202 production of DIN and DIP and the net consumption of the DOM produced during the
203 anabolic phase of the cultures. Goldman and Dennet (1991) also observed in their cultures
204 that ammonium was taken up as long as a readily assimilable carbon source such as glucose
205 was present, whilst nutrient regeneration was evident only after glucose was completely
206 utilized.

207 3.1 Production of DON and DOP by heterotrophic microbial populations

208 POM and inorganic nutrients concentrations showed opposite patterns, POM
209 increased during the net-anabolic phase of the cultures, reaching maximum values at day 10
210 (Fig. 1a-c) and subsequently decreased. The increase in nitrate+nitrite following the
211 decrease in ammonium concentrations (Fig. 1d), suggested, that ammonium was consumed
212 i) as a substrate during the net-anabolic phase of the cultures (Kirchman, 1994); and ii) as
213 an energy source during the net-catabolic phase of the cultures on basis of the complete
214 oxidation of ammonium to nitrate, i.e. nitrification (Wada & Hattori, 1991).
215 In the net-anabolic phase, the use of glucose and inorganic nutrients by the cultured
216 community of microheterotrophs was accompanied by increases in DON and DOP

217 concentrations (Fig 1a, c). We hypothesise that bacteria take up the dissolved substrate,
218 converting it into organic matter which is subsequently released as DOM by direct extra
219 cellular release and/or by protist grazing and viral lysis (Caron et al., 1985; Riemann &
220 Middelboe, 2002; Kawasaki & Benner, 2007). A DOM maximum was found at day 30 of
221 the incubations (Fig. 2), associated with complete depletion of the added glucose (data not
222 shown). The efficiency of DOM production was found using the slope of the linear
223 regressions between UDOC (glucose) and PDOC ($11 \pm 1 \%$), UDIN and PDON ($18 \pm 3 \%$),
224 and UDIP and PDOP ($17 \pm 3 \%$) (Table 2), which was comparable findings in similar
225 experiments (Kawasaki & Benner, 2007).

226 In the net-catabolic phase of the incubations, DOM concentrations decreased until
227 the end of the experiments (Fig. 1; Fig. 2). The calculated bioavailability (BDOM) suggest
228 that PDOC and PDON were more bioavailable than found in natural marine waters (9–30
229 %, DOC; 30–40 %, DON; Hopkinson et al., 2002; Lønborg & Søndergaard, 2008), while
230 PDOP bioavailability was comparable with natural systems (70–80%; Nausch & Nausch,
231 2006). Thereby PDOP was more bioavailable than PDON, and PDON more than PDOC
232 consistently with findings in natural marine systems (Hopkinson et al., 2002). The slopes of
233 the linear regressions between BDOM and PDOM indicate that most of the variations in
234 PDOM were due to the bioavailable fraction (Table 2). The significant origin intercepts of
235 those regressions together with the significant DOM end levels (Table 1, Fig. 2) showed
236 that part of the PDOM persisted at the end of the incubations, demonstrating that the
237 microbial communities also produce refractory DOM (RDOM) in agreement with previous
238 studies (Ogawa et al., 2001; Kawasaki & Benner, 2007). Estimating the environmental
239 impact of the microbial DOM production, we can use published DOC and nutrient data
240 from Loch Creran (Lønborg et al. 2009), together with average efficiency of DOC (11 ± 1

241 %), DON (18 ± 3 %), and DOP (17 ± 3 %) production found in this study. These
242 calculation showed that the heterotrophic microbial community on average could produce 5
243 ± 2 μM DOC, 0.7 ± 0.5 μM DON and 0.08 ± 0.04 μM DOP, corresponding to 3 ± 1 %, $8 \pm$
244 7 % and 72 ± 16 % of the measured DOC, DON and DOP in Loch Creran.

245 The average PDOM C:N:P stoichiometry (103: 11: 1), was generally richer in
246 carbon than typically found for bacterial biomass (50: 10: 1; Goldman et al., 1987;
247 Fagerbakke et al., 1996), and more N and P rich than for both algae produced (170: 6.5: 1;
248 Conan et al., 2007) and marine bulk DOM (300: 22: 1; Benner, 2002). The stoichiometry of
249 PDOM was linearly related with the C: N: P ratios of UDOC, UDIN and UDIP (Fig 3a, c,
250 e). The slopes of the linear regressions show that PDOM had C:N and C:P ratios which
251 were 63 ± 13 % and 68 ± 14 % lower than the substrate, while the N:P ratios (105 ± 15 %)
252 were similar, indicating that the stoichiometry of PDOM is linked to the substrate ratios.
253 The fact that the C:N and C:P ratios of PDOM were 30– 40 % lower than the corresponding
254 substrate ratios can be attributed to the bacterial respiration of glucose, which previously
255 has been found to represent 30–60 % of the total consumption (Bianchi et al., 1998). The
256 stoichiometry of BDOM was similar to ranges reported for bacterial C: N: P ratios
257 (Goldman et al., 1987; Fagerbakke et al., 1996). The slopes of the correlation between
258 PDOM and BDOM stoichiometries (Fig. 3b, d, f) indicate that BDOM had C:N and C:P
259 ratios that were 53 ± 19 % and 44 ± 8 % lower respectively, while N:P ratios (92 ± 25 %)
260 were not significantly different from PDOM.

261 4.2. *Production of chromophoric DOM*

262 Recent studies suggest that marine bacteria produce CDOM and FDOM in coastal
263 ecosystems depending on the chemical composition of the substrate (e.g. Rochelle-Newall

264 & Fisher, 2002; Yamashita & Tanoue, 2008). Table 3a shows how a_{375} increased both
265 during the net-anabolic and net-catabolic phase from being undetectable at the beginning of
266 the incubations reaching higher values (average - $0.47 \pm 0.26 \text{ m}^{-1}$) at the end. This increase
267 indicates microbial production of coloured high molecular weight aromatic DOM (Pages &
268 Gadel, 1990). In parallel to the production of CDOM, FDOM also increased during both
269 the net-anabolic and net-catabolic phase above background levels in all cases (Table 3b, c),
270 in agreement with previous studies (Kramer & Herndl, 2004; Yamashita & Tanoue, 2004).
271 The amounts of FDOM_t and FDOM_m produced increased with substrate uptake and
272 produced DOM (Table 4). Although protein-like fluorophores have been suggested to
273 represent freshly produced DOM (Yamashita & Tanoue, 2004), our study found links
274 between FDOM_t and DOM bioavailability, but also the production of refractory FDOM_t. It
275 firstly indicates that protein-like fluorophores not only provide a measure of labile
276 material; and secondly suggests that biological processes could be a source of refractory
277 FDOM_t. FDOM_m was related linearly with both BDOC and RDOC, confirming it as a
278 product of microbial degradation processes (Yamashita & Tanoue, 2004; 2008; Nieto-Cid
279 et al., 2006).

280 **Acknowledgement**

281 This study was funded by a fellowship to C.L from the Early Training site ECOSUMMER
282 (Marie Curie action 20501). The crew of Seol Mara is acknowledged for help with the
283 collection of field samples used in this study.

284

285 **References**

- 286 Benner, R., 2002. Chemical composition and reactivity. In: Hansell D.A., Carlson, C.
287 (Eds.), Biogeochemistry of Marine Dissolved Organic Matter. Academic Press,
288 USA pp. 59–90
- 289 Bianchi, A., Van Wambeke, F., Garcin, J., 1998. Bacterial utilization of glucose in the
290 water column eutrophic to oligotrophic pelagic areas in the eastern Atlantic Ocean.
291 Journal of Marine Systems 14, 45–55.
- 292 Blough, N.V., Del Vecchio, R., 2002. Chromophoric DOM in the coastal environment. I In:
293 Hansell D.A., Carlson, C. (Eds.), Biogeochemistry of Marine Dissolved Organic
294 Matter. Academic Press, USA, pp. 509– 546.
- 295 Coble, P.G., Green S.A., Blough N.V., Gasgolian R.B., 1990. Characterization of dissolved
296 organic matter in the Black Sea by fluorescence spectroscopy. Nature 348, 432–435.
- 297 Coble, P.G., 1996. Characterization of marine and terrestrial DOM in seawater using
298 excitation–emission matrix spectroscopy. Marine Chemistry 51, 325–346.
- 299 Caron, D.A., Goldman, J.C., Andersen, O.K, Dennett M.R., 1985. Nutrient cycling in a
300 microflagellate food chain: II. Population dynamics and carbon cycling. Marine
301 ecology progress Series 24, 243–254.
- 302 Conan, P., Søndergaard, M., Kragh, T., Thingstad, F., Pujo–Pay, M., Williams, P.J. Le B.,
303 Markager, S., Cauwet, G., Borch, N.H., Evans, D., Riemann, B., 2007. Partitioning
304 of organic production in marine plankton communities: The effects of inorganic
305 nutrient ratios and community composition on new dissolved organic matter.
306 Limnology and Oceanography 52, 753–765.

- 307 Fagerbakke, K.M., Heldal, M., Norland, S., 1996. Content of carbon, nitrogen, oxygen,
308 sulphur and phosphorus in native aquatic and cultured bacteria. *Aquatic Microbial*
309 *Ecology* 10, 15–27.
- 310 Goldman, J.C., Caron, D.A., Dennett, M.R. 1987. Regulation of gross growth efficiency
311 and ammonium regeneration in bacteria by substrate C : N ratio. *Limnology and*
312 *Oceanography* 32, 1239-1252.
- 313 Goldman, J.C., Dennett, M.R., 1991. Ammonium regeneration and carbon utilization by
314 marine grown on mixed substrates. *Marine Biology* 109, 369–378.
- 315 Hansen, H. P., Grasshoff, K., 1983. Automated chemical analysis. In: Grasshoff, K.,
316 Ehrhardt, M., Kermling, K. (Eds.), *Methods of seawater analysis* Verlag Chemie,
317 Germany, pp. 347–395.
- 318 Harrison, P.J., Waters, R.E., Taylor, F.J.R., 1980. A broad spectrum artificial seawater
319 medium for coastal and open ocean phytoplankton. *J. Phycol.* 16, 28–35.
- 320 Hicks, S.E., Carey, F.G., 1968. Glucose determination in natural waters. *Limnology and*
321 *Oceanography* 13, 361–363.
- 322 Hopkinson, C.S., Vallino, J.J., Nolin, A., 2002. Decomposition of dissolved organic matter
323 from the continental margin. *Deep–Sea Research part II* 49, 4461–4478.
- 324 Kawasaki, N., Benner, R., 2006. Bacterial release of dissolved organic matter during cell
325 growth and decline: Molecular origin and composition. *Limnology and*
326 *Oceanography* 51, 2170– 2180.
- 327 Kirchman, D.L. 1994. The uptake of inorganic nutrients by heterotrophic bacteria.
328 *Microbial Ecology* 28, 255–271.

329 Koroleff, F., 1983. Simultaneous oxidation of nitrogen and phosphorus compounds by
 330 persulfate. In: Grasshoff, K., Ehrhardt, M., Kermling, K. (Eds.), *Methods of*
 331 *seawater analysis* Verlag Chemie, Germany, pp. 168–169.

332 Kramer, G.D., Herndl, G.J., 2004. Photo– and bioreactivity of chromophoric dissolved
 333 organic matter produced by marine bacterioplankton. *Aquatic Microbial Ecology*
 334 36, 239–246.

335 Lalli, C.M., Parsons, T.R., 1997. *Biological Oceanography: An Introduction*, 2nd .
 336 Butterworth–Heinemann, Oxford, 314 pp.

337 Lønborg, C., Søndergaard, M., 2009. Microbial availability and degradation of dissolved
 338 organic carbon and nitrogen in two coastal areas. *Estuarine, Coastal and Shelf*
 339 *Science*. 81, 513–520.

340 Lønborg, C., Davidson K., Álvarez-Salgado X.A., Miller A.E.J., 2009. Bioavailability and
 341 bacterial degradation rates of dissolved organic matter in a temperate coastal area
 342 during an annual cycle. *Marine Chemistry* 10.1016/j.marchem.2009.02.003

343 Nagata, T., 2000. Production mechanisms of dissolved organic matter. In: Kirchman, D.L.
 344 (Ed.), *Microbial ecology of the oceans*. Wiley–Liss, USA, pp. 121–152.

345 Nausch, M., Nausch, G., 2006. Bioavailability of dissolved organic phosphorus in the
 346 Baltic Sea. *Marine Ecology Progress Series* 321, 9–17.

347 Nieto–Cid, M., Álvarez–Salgado, X.A., Perez, F.F., 2006. Microbial and photochemical
 348 reactivity of fluorescent dissolved organic matter in a coastal upwelling system.
 349 *Limnology and Oceanography* 51, 1381–1390.

350 Obernosterer, I., Herndl, G.J., 1995. Phytoplankton extra–cellular release and bacterial
 351 growth: dependence on the inorganic N:P ratio. *Marine Ecology Progress Series*
 352 116, 247–257.

353 Ogawa, H., Amagai, Y., Koike, I., Kaiser, K., Benner, R., 2001. Production of refractory
 354 dissolved organic matter by bacteria. *Science* 292, 917–920.

355 Pages, J., Gadel, F., 1990. Dissolved organic matter and UV absorption in a tropical
 356 hyperhaline estuary. *The Science of the Total Environment* 99, 173–204.

357 Riemann, L., Middelboe, M., 2002. Viral lysis of marine bacterioplankton: Implications for
 358 organic matter cycling and bacterial clonal composition. *Ophelia* 56, 57–68.

359 Rochelle-Newall, E.J., Fisher, T.R., 2002. Production of chromophoric dissolved organic
 360 matter fluorescence in marine and estuarine environment: an investigation into the
 361 role of phytoplankton. *Marine Chemistry* 77, 7–21.

362 Sokal, F.F., Rohlf, F.J., 1995. *Biometry*. Freeman, New York.

363 Seitzinger, S.P., Sanders, R.W., 1997. Contribution of dissolved organic nitrogen from
 364 rivers to estuarine eutrophication. *Marine Ecology Progress Series* 159, 1–12.

365 Stedmon, C.A., Markager, S.S., 2001. The optics of chromophoric dissolved organic matter
 366 (CDOM) in the Greenland sea: An algorithm for differentiation between marine and
 367 terrestrially derived organic matter. *Limnology and Oceanography* 46, 2087–2093.

368 Tezuka, Y., 1990. Bacterial regeneration of ammonium and phosphate as affected by the
 369 carbon : nitrogen : phosphorus ratio of organic substrates. *Microbial Ecology* 19,
 370 227–238.

371 Wada, E., Hattori, A. 1991. *Nitrogen in the Sea: Forms, Abundances, and Rate Processes*.
 372 CRC Press, 208 pp.

373 Yamashita, Y., Tanoue, E., 2004. In situ production of chromophoric dissolved organic
 374 matter in coastal environments. *Geophysical Research Letters* 31,
 375 doi:10.1029/2004GL01973.

376 Yamashita, Y., Tanoue, E., 2008. Production of bio-refractory fluorescent dissolved
377 organic matter in the ocean interior. *Nature Geoscience* 1, 579–582.
378

379 **Figure legends**

380 Figure. 1. Examples of time evolution (Exp. 1 shown in table 1), of a) total (TOC)
381 particulate (POC) and dissolved organic carbon (DOC), b) total nitrogen (TN), particulate
382 organic nitrogen (PON), dissolved inorganic (DIN) and organic (DON) nitrogen; c) total
383 phosphorus (TP), particulate organic phosphorus (POP), dissolved inorganic (DIP) and
384 organic (DOP) phosphorus; d) nitrate/nitrite ($\text{NO}_3^-/\text{NO}_2^-$) and ammonium (NH_4^+). Error
385 bars represent standard errors.


386

387 Figure. 2. Production of dissolved organic nitrogen (DON) and phosphorus (DOP) in the
388 16 experiments, a),b) started in January 2007 and c),d) in March 2007. The experimental
389 numbers 1 to 16 represents the same experiment numbers as shown in table 1.

390

391 Figure. 3. Significant regressions between substrate uptake (UDOC, UDIN, UDIP) and
392 produced DOM (PDOM) stoichiometry and the X–Y plots of linear relation between
393 PDOM and bioavailable DOM (BDOM) stoichiometry. With a) UDOC:UDIN with
394 PDOC:PDON, b) PDOC:PDON versus BDOC:BDON, c) UDOC:UDIP versus
395 PDOC:PDOP, d) PDOC:PDOP with BDOC:BDOP, e) UDIN:UDIP versus PDON:PDOP
396 and f) PDON:PDOP with BDON:BDOP. Slope, intercept, and standard error are values found
397 by Model II regression. R^2 = coefficient of determination, p = significant levels.

398



399

400 Fig.1.





Fig.2.



403

404 Fig.3.

Table 1. Concentrations of DOC (UDOC), DIN (UDIN) and DIP (UDIP) utilized during the experiments, DOC (PDOC), DON (PDON), and DOP (PDOP) produced during the experiments, and residual DOC (RDOC), DON (RDON) and DOP (RDOP) at the end of the experiments. Experiments 1 to 8 were commenced in January and experiments 9 to 16 in March 2007. Values are means of three replicates \pm standard error given in $\mu\text{mol L}^{-1}$.

Exp.	DOC	DIN	DIP	UDOC	UDIN	UDIP	PDOC	PDON	PDOP	RDOC	RDON	RDOP
1	396 \pm 1	80.6 \pm 0.6	4.06 \pm 0.02	396 \pm 1	40.7 \pm 0.9	2.48 \pm 0.10	52 \pm 1	5.3 \pm 1.4	0.59 \pm 0.13	33 \pm 1	1.6 \pm 0.8	0.14 \pm 0.04
2	440 \pm 6	80.2 \pm 0.4	1.42 \pm 0.02	440 \pm 6	20.8 \pm 0.4	1.08 \pm 0.08	51 \pm 1	5.7 \pm 0.8	0.11 \pm 0.04	31 \pm 1	0.4 \pm 0.1	0.02 \pm 0.05
3	425 \pm 1	25.1 \pm 0.1	4.18 \pm 0.03	425 \pm 1	19.5 \pm 0.3	2.32 \pm 0.01	47 \pm 2	5.1 \pm 1.0	0.43 \pm 0.04	35 \pm 1	1.2 \pm 0.2	0.17 \pm 0.02
4	163 \pm 1	77.6 \pm 0.3	4.45 \pm 0.02	163 \pm 1	8.6 \pm 0.4	1.19 \pm 0.02	26 \pm 2	2.2 \pm 0.9	0.25 \pm 0.04	21 \pm 1	1.6 \pm 0.4	0.07 \pm 0.05
5	789 \pm 3	93.4 \pm 0.2	4.73 \pm 0.03	789 \pm 3	62.8 \pm 0.3	3.78 \pm 0.08	84 \pm 1	11.7 \pm 0.8	0.67 \pm 0.05	42 \pm 1	1.1 \pm 0.5	0.07 \pm 0.02
6	411 \pm 6	157.5 \pm 0.4	4.47 \pm 0.04	411 \pm 6	32.7 \pm 0.6	3.00 \pm 0.04	45 \pm 4	5.3 \pm 0.9	0.61 \pm 0.06	24 \pm 1	2.4 \pm 0.5	0.10 \pm 0.01
7	396 \pm 2	79.2 \pm 0.1	8.11 \pm 0.06	396 \pm 2	34.2 \pm 0.3	3.66 \pm 0.08	48 \pm 1	4.3 \pm 0.8	0.67 \pm 0.08	28 \pm 1	1.5 \pm 0.6	0.20 \pm 0.02
8	433 \pm 1	88.1 \pm 0.4	4.59 \pm 0.04	433 \pm 1	18.1 \pm 0.8	2.35 \pm 0.10	42 \pm 1	4.8 \pm 0.8	0.52 \pm 0.09	28 \pm 1	1.3 \pm 0.8	0.06 \pm 0.04
9	543 \pm 2	84.0 \pm 0.6	7.06 \pm 0.07	543 \pm 2	29.4 \pm 0.7	3.95 \pm 0.15	74 \pm 1	5.7 \pm 0.8	0.55 \pm 0.10	38 \pm 1	1.9 \pm 0.9	0.12 \pm 0.07
10	440 \pm 7	82.1 \pm 0.3	4.09 \pm 0.02	440 \pm 7	35.0 \pm 0.8	3.90 \pm 0.06	49 \pm 3	6.3 \pm 0.9	0.52 \pm 0.07	39 \pm 1	1.4 \pm 0.5	0.09 \pm 0.03
11	411 \pm 9	51.0 \pm 0.4	7.69 \pm 0.06	411 \pm 9	41.1 \pm 0.8	5.10 \pm 0.07	48 \pm 2	6.3 \pm 0.8	0.78 \pm 0.08	38 \pm 1	1.6 \pm 0.3	0.18 \pm 0.04
12	409 \pm 3	88.5 \pm 0.2	7.57 \pm 0.07	409 \pm 3	24.6 \pm 0.6	2.91 \pm 0.15	45 \pm 1	4.8 \pm 1.0	0.48 \pm 0.09	32 \pm 1	1.8 \pm 0.4	0.06 \pm 0.02
13	867 \pm 2	84.1 \pm 0.3	7.43 \pm 0.10	867 \pm 2	45.1 \pm 0.6	5.37 \pm 0.10	93 \pm 2	6.2 \pm 0.8	0.72 \pm 0.06	54 \pm 3	2.5 \pm 0.4	0.05 \pm 0.03
14	449 \pm 2	164.7 \pm 0.4	7.28 \pm 0.05	449 \pm 2	34.1 \pm 1.0	2.91 \pm 0.09	41 \pm 2	6.2 \pm 0.9	0.55 \pm 0.13	34 \pm 1	2.4 \pm 1.1	0.11 \pm 0.04
15	442 \pm 1	74.8 \pm 0.4	13.64 \pm 0.05	442 \pm 1	30.9 \pm 0.5	1.33 \pm 0.09	50 \pm 3	6.0 \pm 1.0	0.28 \pm 0.12	42 \pm 1	2.2 \pm 0.5	0.17 \pm 0.03
16	448 \pm 3	96.5 \pm 0.9	7.57 \pm 0.03	448 \pm 3	27.0 \pm 1.6	2.06 \pm 0.08	43 \pm 1	5.8 \pm 1.1	0.36 \pm 0.08	27 \pm 2	2.3 \pm 0.9	0.04 \pm 0.02

410 Table 2. Significant linear regressions between used DOC (UDOC) and produced DOC (PDOC);
 411 bioavailable (BDOC) and PDOC; used dissolved inorganic nitrogen (UDIN) and produced DON (PDON);
 412 bioavailable DON (BDON) and PDON); used phosphate (UDIP) and produced DOP (PDOP);, and
 413 bioavailable DOP (BDOP) and PDOP. Slope, intercept, and standard error (SE) are values found by Model II
 414 regression. R^2 = coefficient of determination, p = level of significance, n.s. – not significant.
 415

X	Y	Slope (\pm SE)	Intercept (\pm SE)	R^2	p
UDOC	PDOC	0.11 ± 0.01	n.s	0.90	<0.00001
BDOC	PDOC	1.4 ± 0.2	27 ± 3	0.80	<0.00001
UDIN	PDON	0.18 ± 0.02	n.s	0.85	<0.0001
BDON	PDON	1.0 ± 0.2	1.7 ± 0.3	0.92	<0.001
UDIP	PDOP	0.17 ± 0.02	n.s	0.79	<0.00001
BDOP	PDON	1.1 ± 0.1	0.08 ± 0.03	0.82	<0.00001

416

417 Table 3. a) Absorption coefficients (a_{375}) at day 0 (a_{375_0}), 30 ($a_{375_{30}}$) and 100 ($a_{375_{100}}$). b) Initial
 418 (FDOM_{t0}), produced (PFDOM_{t0}) and refractory (RFDOM_{t0}) protein-like fluorescence; c) Initial (FDOM_{m0}),
 419 produced (PFDOM_m) and refractory (RFDOM_m) marine humic-like fluorescence; and Experiments 1 to 8
 420 were commenced in January and experiments 9 to 16 in March 2007, corresponding to the experiments
 421 described in table 1. Values are means of two replicates \pm standard error.

Experiment	$a_{375_0} (m^{-1})$	$a_{375_{30}} (m^{-1})$	$a_{375_{100}} (m^{-1})$	a)
1	0.05 ± 0.05	0.44 ± 0.08	0.67 ± 0.02	
2	0.01 ± 0.02	0.27 ± 0.03	0.61 ± 0.09	
3	0.06 ± 0.06	0.25 ± 0.10	0.92 ± 0.12	
4	0.07 ± 0.08	0.32 ± 0.08	0.60 ± 0.09	
5	0.01 ± 0.01	0.36 ± 0.03	0.52 ± 0.10	
6	0.07 ± 0.08	0.50 ± 0.05	0.84 ± 0.17	
7	0.01 ± 0.04	0.38 ± 0.10	0.45 ± 0.18	
8	0.02 ± 0.02	0.13 ± 0.07	0.64 ± 0.09	
9	0.07 ± 0.08	0.31 ± 0.08	0.41 ± 0.04	
10	0.08 ± 0.09	0.42 ± 0.10	0.69 ± 0.07	
11	0.08 ± 0.09	0.24 ± 0.05	0.34 ± 0.07	
12	0.03 ± 0.04	0.24 ± 0.04	0.38 ± 0.05	
13	0.02 ± 0.02	0.32 ± 0.08	0.41 ± 0.16	
14	0.04 ± 0.04	0.39 ± 0.04	0.53 ± 0.03	
15	0.01 ± 0.01	0.34 ± 0.05	0.34 ± 0.09	
16	0.01 ± 0.01	0.52 ± 0.03	1.33 ± 0.09	

422

Experiment	FDOM _{t0} (ppb Trp)	PFDOM _{t0} (ppb Trp)	RFDOM _{t0} (ppb Trp)	b)
1	3.5 ± 0.6	17.7 ± 0.7	21.1 ± 0.1	
2	3.2 ± 0.6	10.0 ± 0.9	13.2 ± 0.3	
3	2.6 ± 0.4	13.0 ± 1.0	15.6 ± 0.7	
4	2.3 ± 0.2	6.1 ± 0.7	8.4 ± 0.5	
5	2.5 ± 0.2	19.9 ± 0.7	22.4 ± 0.5	
6	3.9 ± 0.1	17.7 ± 0.6	21.5 ± 0.5	
7	3.0 ± 0.3	21.2 ± 0.4	24.2 ± 0.1	
8	1.3 ± 0.5	10.9 ± 0.7	12.2 ± 0.2	
9	2.3 ± 0.6	16.3 ± 0.8	18.6 ± 0.2	
10	3.2 ± 0.3	7.9 ± 0.9	11.1 ± 0.5	
11	2.6 ± 0.3	10.6 ± 0.4	13.2 ± 0.2	
12	2.3 ± 0.6	7.9 ± 1.1	10.2 ± 0.5	
13	4.5 ± 0.9	15.2 ± 1.0	19.7 ± 0.1	

423

14	2.9 ± 0.1	11.4 ± 0.3	14.3 ± 0.2
15	1.0 ± 0.2	14.5 ± 0.6	15.5 ± 0.4
16	2.3 ± 0.2	9.4 ± 0.6	11.7 ± 0.4

Experiment	FDOMm ₀ (ppb Trp)	PFDOMm (ppb Trp)	RFDOMm (ppb Trp)	c)
1	2.58 ± 0.3	9.0 ± 0.5	11.7 ± 0.1	
2	2.5 ± 0.5	10.6 ± 0.8	13.1 ± 0.3	
3	2.0 ± 0.6	8.9 ± 0.7	10.8 ± 0.1	
4	2.5 ± 0.3	4.7 ± 0.7	7.2 ± 0.5	
5	2.9 ± 0.1	14.0 ± 0.2	16.8 ± 0.2	
6	2.2 ± 0.3	5.0 ± 0.9	7.2 ± 0.6	
7	1.9 ± 0.4	5.4 ± 0.5	7.3 ± 0.2	
8	1.8 ± 0.2	4.8 ± 0.2	6.6 ± 0.2	
9	2.8 ± 0.3	7.8 ± 1.0	10.5 ± 0.7	
10	2.3 ± 0.5	3.7 ± 0.6	6.0 ± 0.1	
11	1.8 ± 0.6	4.0 ± 0.7	5.8 ± 0.2	
12	2.4 ± 0.3	3.2 ± 1.1	5.6 ± 0.8	
13	2.6 ± 0.1	13.1 ± 0.2	15.7 ± 0.1	
14	2.2 ± 0.3	2.8 ± 0.5	5.0 ± 0.3	
15	2.0 ± 0.4	5.2 ± 0.5	7.2 ± 0.1	
16	1.8 ± 0.2	5.1 ± 0.3	6.9 ± 0.1	

424

425

426 Table 4. Matrix of the correlation coefficient (R^2) of the significant ($p < 0.05$) linear regressions
 427 between DOM and the produced protein- (PFDOM_T) and marine humic-like (PFDOM_M)
 428 flourophores. n.s. – not significant.

X-Y	FDOM _T	FDOM _M
UDIN	0.60	0.33
UDOC	0.53	0.64
RDOC	n.s.	0.45
BDOC	0.77	0.53
BDON	0.30	0.38
BDOP	0.44	n.s.

430