Seed shape in model legumes: approximation by a cardioid reveals differences in ethylene insensitive mutants of *Lotus japonicus* and *Medicago truncatula*

Emilio Cervantes^{1*}, José Javier Martín¹, Pick Kuen Chan², Peter M. Gresshoff², Ángel Tocino³

1: IRNASA – CSIC. Salamanca. Apartado 257. 37080. Salamanca. Spain

2: ARC Centre of Excellence for Integrative Legume Research. The University of Queensland. St. Lucia, Brisbane QLD 4072. Australia

3: Dpto. de Matemáticas, Universidad de Salamanca. Spain.

*: Corresponding author.

INTRODUCTION

The description of plant form and shape is useful to understand evolutionary relationships and quantify differences between species, as well as to define the processes of plant development over space and time. The scientific discipline of *Plant Morphology* (Kaplan, 2001), however, needs to be mathematically and not solely descriptively based. Seed spacing in a sunflower head, or rings swirls on a pineapple fruit, whose distribution follows Fibonacci series, are good examples, while others remain awaiting discovery. Computational modeling of plant development has been applied to legume lateral root development (Han et al, 2011) as well as nodule pattern (Han et al, 2010b). Such mathematical and computational basis permits the *in silico* testing of thousands of experimental conditions to achieve outcomes which then need to be verified in ‘real world’ experiments.

Recently we described a mathematical-geometrical model to quantify the shape of *Arabidopsis* seeds. It consisted in the comparison of the outline of the longitudinal equatorial section of the seed with a transformed cardioid (Cervantes *et al.*, 2010). The cardioid is the trajectory described by a point of a circle that rolls around another fixed circle with the same radius, and transforming it by multiplying the x-axis by the scaling factor φ (≈ 1.61803399), the so-called ‘Golden Ratio’, the figure obtained resembled the image of an *Arabidopsis* dry seed. The similarity was evaluated by a factor, the G index, allowing the precise morphological description of genotypes and opening new ways for the association between genetic changes and alterations in the morphology, *i.e.*, a classical genome-phenome analysis.

Modeling the *Arabidopsis* seed shape by a transformed cardioid allowed comparison between genotypes showing that ethylene insensitive *etr1-1* mutants presented alterations from the wild type and suggesting that ETR1 (one of the histidine protein kinase ethylene receptors) function goes beyond its role in ethylene signal transduction (Cervantes *et al.*, 2010). The functional connection between ethylene perception and seed shape is unclear but may involve diverse aspects of seed development as well as drying rates.

Legume seeds offer an excellent material to expand this type of analysis in search of accurate morphological descriptions. Legumes are a large family containing members of great agronomic interest (*e.g.*, soybean, pea, common bean, medics, clovers as well as trees such as *Acacias*, *Robinia* and *Pongamia*; Biswas *et al.*, 2011). In the context of seed morphology, legume species are characterized by remarkable variation, ranging from small seeds such as *Lotus japonicus* to large-sized seeds of *Phaseolus* or *Vicia* and gigantic ones (7-8 cm long) in Black Bean Tree (*Castanospermum australe*) in Australia.

Interestingly, *Lotus japonicus* and *Medicago truncatula*, two model species intensively used in genomic analysis (Jiang and Gresshoff, 1997; Sato *et al.*, 2008; Cannon *et al.*, 2006; Young *et al.*, 2005), are amenable to morphological analysis based on a simple model resembling the one described for *Arabidopsis*, *i.e.*, the comparison with a cardioid. Whereas the seed dimensions of *Lotus japonicus* match well to a cardioid, in *Medicago truncatula* accurate adjustment is obtained by the simple transformation of scaling the vertical axis by a factor equal to the Golden Ratio. As in the case of *Arabidopsis*, the application of these models allows the description of alterations in the ethylene response mutants.

MATERIALS AND METHODS

Plant material

Seeds of *Lotus japonicus* of the ecotypes Gifu and Miyakojima (MG-20) were used. MG-20 is a Japanese ecotype from a more tropical habitat than Gifu. The former comes from the Miyakojima Islands south of the Japanese island chain, and the latter from the Gifu prefecture close to Tokyo. They differ in their ABA sensitivity to germination and other photoperiod related traits (Biswas *et al.*, 2009). *LjETR1-1* is a transgenic line obtained from the stable transformation of Gifu with the dominant *Arabidopsis* ethylene receptor gene *ETR1-1* (Lohar *et al.*, 2009). Mutant *enigma-1* is one of three independent mutants of the *Lotus EIN2a* gene in the MG-20 background produced by EMS mutagenesis and M2 selection for ethylene insensitivity in seedling root growth (Chan *et al.*, 2012); *Ljhar1-1* is a mutant of the Gifu variety in *HAR1*, a gene involved in the systemic regulation of nodulation and structurally related to the *Arabidopsis thaliana* *CLAVATA1* (*CLV1*) gene (Krusell *et al.*, 2002; Nishimura *et al.*, 2002; Wopereis *et al.*, 2000). Seeds of *Medicago truncatula* wild type A17 as well as the ethylene insensitive and increased nodulation mutant *sickle* (Penmetsa *et al.*, 2008) were used. Of interest is the fact that both *sickle* and *enigma-1* are mutated in *EIN2*, an integral component of the *Arabidopsis* ethylene sensing pathway.

Photography and image analysis

The seeds were observed with a Nikon ‘SMZ-2T’ stereo-microscope. Photographs of longitudinal views were taken with a digital camera Nikon ‘Coolpix 950’ and analyzed

with the software image Photoshop CS4 (Adobe), specialized for image processing. In this process, graph paper was used to convert pixel into μm .

Size and shape analysis

The area of the seed images was obtained with Photoshop CS4 and used as the measure of seed size. Circularity index (Schwartz, 1980), given by

$$I = 4\pi \frac{\text{area}}{(\text{perimeter})^2},$$

is a measure of the similarity of a plane figure to a circle. It ranges from 0 to 1 giving the value of 1 for circles and it is a useful magnitude as a first approximation to seed shape. Eight independent experiments with 10 seeds for each variety or genotype were used in *Lotus japonicus*. For *Medicago truncatula* three experiments with 10 seeds for each genotype were done.

Quantification of the image adjustments to a cardioid

The cardioid is the trajectory described by a point of a circle that rolls around another fixed circle with the same radius, see figure 1A (left). This common radius is the only parameter needed to define a particular cardioid. In Cervantes *et al.* (2010), the image of *Arabidopsis* seeds was shown to adjust well to a cardioid modified by elongating its horizontal axis by a factor equal to the Golden Ratio ϕ . The pictures of seeds were combined with cardioid (figure 1A; *Lotus*) or modified cardioid (figure 1B; *Medicago*) images.

Quantification of the adjustment was done in each seed by the comparison of the areas in two regions: The common region in the cardioid and the seed image, and the regions not shared between both areas (see figure 1). The index of adjustment (J) is defined by:

$$J = \frac{\text{area (C)}}{\text{area (C)} + \text{area(D)}} \times 100,$$

where C represents the common region and D the regions not shared. Note that J is a measure of seed shape, not of its area. It ranges between 0 and 100 decreasing when the size of the non-shared region grows, and equals 100 when cardioid and seed image areas coincide, *i.e.*, when area (D) is zero.

The adjustment was done with an unmodified cardioid for *Lotus japonicus*. In *Medicago* seeds, the adjustment was improved by elongating the y-axis by a factor of ϕ (figure 1B). Shape comparison among genotypes was done by the statistical contrasting of their J index values in dry seeds as well as after 24 hours of imbibition.

Statistical analysis

ANOVA was used for the comparison between different groups. Post-hoc analysis was carried out using Scheffé test for samples of heterogeneous size. Tuckey test was used for samples of similar sizes. Statistical analysis was done with SPSS.

RESULTS

Tables 1 to 4 contain the results from experiments with *Lotus japonicus*. The mean values for seed image area (table 1A), circularity index (table 1B), and J index (table 1C) in the varieties and genotypes used are shown. Tables 2 and 3 present the contrast tests for seed image area and J index, while table 4 contains a schematic representation of results. Table 5 contains values for area, circularity index and index J of dry and imbibed seeds in *Medicago truncatula* genotypes. The results are represented in figure 3.

Adjustment of wild-type Lotus japonicus seeds to the cardioid shape and comparison of size and shape between varieties

The images of Lotus seeds adjust to an unmodified cardioid both in dry seeds (figure 1A) as well as for seeds imbibed for 24 hours. There was no difference between J index in dry seeds of Gifu (92.6 %) and MG-20 (92.7%). In contrast, after imbibition, J index was smaller in Gifu (90.9%) than in MG-20 (93.5%; tables 1C and 3; figure 3), perhaps reflecting differential ABA sensitivity in germination described by Biswas *et al.* (2009). Seeds of Gifu were smaller than MG-20, both in dry status and after imbibition (tables 1A and 2; figure 3).

Variation in size and form in Lotus japonicus mutant genotypes

Size and form of seeds in genotypes Gifu, *LjETR1-1*, MG-20, *enigma-1* and *Ljhar1-1* were compared. MG-20 is the parent for *enigma-1*; Gifu, for *LjEtr1-1* and *Ljhar1-1*.

Dry seeds

Table 2A shows the results of the ANOVA test concerning the areas of dry seeds. Gifu seeds were larger than those of *Ljhar1-1*, reflecting differences in plant vigour (see Wopereis *et al.*, 2000). Mutant seeds in the ethylene pathway in both varieties were smaller than their corresponding wild-type sources, that is, *LjETR1-1* seeds were smaller than Gifu, and *enigma-1* smaller than MG-20. No differences were detected in the values of circularity index between wild-type varieties or between each of the varieties and their respective mutants (ANOVA not shown). The J-index was smaller in *enigma-1* than MG-20 (table 3A, figure 3). No differences in J-index were found between *Ljhar1-1* and *LjETR1-1* with their parental variety (Gifu).

Imbibed seeds

Analysis of imbibed seeds of Lotus genotypes revealed differences in size between genotypes (table 2B): Gifu seeds were larger than their mutant genotypes *Ljhar1-1* and *LjETR1-1*. Seeds of MG-20 were larger than *enigma-1*, which is consistent with difference in pod and plant size (Chan et al, 2012); no differences were observed between genotypes in their circularity index. After imbibition, the J-index values were smaller for mutant genotypes in the ethylene sensing pathway: *enigma-1* had smaller values than MG-20 and *LjETR1-1* was smaller than Gifu (table 3B). There was no difference between Gifu and *Ljhar1-1*.

Comparison between dry and imbibed seeds

Area and circularity indexes increased in all varieties and genotypes upon seed imbibition (table 1A and B), as indicated by T-test ($P < 0.05$; not shown). The J-Index decreased with imbibition in genotypes *LjETR1-1*, Gifu and *enigma-1* ($P < 0.05$; not shown). No differences were detected in MG-20 and *Ljhar1-1* (not shown).

Adjustment of wild-type Medicago seeds to a cardioid

For the adjustment of Medicago seeds, departing from a cardioid the vertical axis was scaled by a factor equal to the Golden Ratio $\phi \approx 1.618$ (figure 2B). The mean values for J-index in *Medicago truncatula* A17 were 87.1% and 86.8% for dry and imbibed seeds respectively (table 5). No differences were found between treatments.

Variation in size and form in Medicago genotypes

Dry seeds

Differences between wild type A17 and the *sickle* mutant were found in size (area) and circularity index, but not in the J-index (table 5).

Imbibed seeds

After imbibition no differences were observed between both *Medicago* genotypes either in area, circularity or J-index (not shown).

Comparison between dry and imbibed seeds

Imbibition resulted in increased areas and reduced circularity index values in both genotypes ($P < 0.05$; not shown). There was no change in J-index with imbibition (not shown).

DISCUSSION

The analysis of plant development requires an accurate and quantitative description of morphology. Space, volume, shape and surface interact as biophysical constraints with gene activity. Changes in size and shape may occur independently as the results of developmental programs in a “regular” environment or in response to changes (stress) in environmental conditions (Cervantes and Tocino, 2009). The morphological analysis of seeds in model species may reveal quantitative measures and give new parameters useful for the identification of genetic elements responsible for seed shape, a field of research that is independent from size, and that has remained virtually unexplored.

In a previous study, the image of a transformed cardioid was used as a model for the shape of *Arabidopsis* seeds. The transformation resulted of elongating the cardioid along the x-axis (*i.e.*, horizontally) by a factor of ϕ , the Golden Ratio (Cervantes *et al.*, 2010). The model based in a cardioid curve fits better the shape of *Arabidopsis* seeds than a former model based on a prolate spheroid (Robert *et al.*, 2008) and it is here applied to seeds of the model legumes *Lotus japonicus* and *Medicago truncatula*.

Lotus seeds are rounded and their shape is best described by the model of an unmodified cardioid, whereas for *Medicago* the cardioid needed to be elongated by a factor of ϕ in the vertical direction to provide a fit. The seeds of *Arabidopsis* and *Medicago* present an elongated form according to their growth inside the pods. The final seed shape resembles a modified cardioid. In *Arabidopsis*, the seed presents the micropyle in one pole and the cardioid is elongated in the horizontal direction. In *Medicago truncatula*, called Australian Barrel Medic, the pod is compacted into a fused spiral, and the seed develops inside leaving the micropyle in a central position (figure 2B), so the resulting cardioid is elongated in a vertical direction. In both species, *Medicago truncatula* and *Arabidopsis thaliana*, similar physical restrictions during seed development result in an optimized adjustment modifying the cardioid by a factor of ϕ . Lotus seeds, as many other legumes (soybean, pea, chickpea and lentil) develop independently inside an elongated pod (figure 2A), but are more rounded than those of *Medicago* or *Arabidopsis*.

In three model species (*Arabidopsis*, *Medicago*, *Lotus*) seed shape can be well-adjusted to simple geometric models: unmodified or slightly modified cardioid curves. Departing from these curves it may be possible to design new geometric models adapted to other species in their families. Model species share important characteristics (short life cycles, small and abundantly produced seeds, lack of specialized forms). They represent examples of organisms with “type r” strategies and, thus, simple, easily described seed shape may be an important feature of model plants (Begon *et al.*, 2006; Cervantes, 2008).

The comparisons between genotypes demonstrated differences in seed shape (based on the cardioid), as well as in size. Differences in ABA sensitivity between varieties Gifu and MG-20 have been previously reported (Biswas *et al.*, 2009). Here, we have described that MG-20, an ecotype more tropical and less sensitive to ABA during seed germination than Gifu, has larger seeds, thus suggesting that in *Lotus japonicus*, an inverse relationship may exist between seed size, latitude of origin and ABA sensitivity. An inverse relationship between seed size and latitude has been reported in general for plants (Moles *et al.*, 2007), as well as in the legume family for seeds of *Glycine* species (Murray *et al.*, 2003).

Antagonistic roles have been found for ABA and ethylene in seeds. In *Arabidopsis*, ethylene-insensitive *ein2* mutants have enhanced sensitivity to ABA (Beaudoin *et al.*, 2000; Ghassemian *et al.*, 2000). Seeds of *enigma* mutants have enhanced sensitivity to ABA (Chan *et al.* 2012) that is associated with reduced seed size. ABA is known to promote quiescence and suppress differentiation in the *Arabidopsis* primary root meristem (Zhang *et al.*, 2010); thus, in *Lotus japonicus*, reduced seed size may result from limited cell division as a consequence of increased ABA sensitivity. Also, reduced size results in an increased ‘surface/volume’ ratio that may be responsible for higher ABA penetration in the cells. Finally, ABA insensitivity may involve a delay in pod abscission and maturation resulting in larger seeds.

Arabidopsis ethylene insensitive mutants (*etr1-1*) presented differences in seed shape when compared to the wild type: mutant seeds resemble less a cardioid having lower values of G (equivalent to J-index) index than wild type (Cervantes *et al.*, 2010). The

genotypes mutated in the ethylene sensing pathway in *Lotus* and *Medicago* presented a more complex range of alterations in seed size and shape.

In *L. japonicus*, differences in seed size were found both in dry seeds and after imbibition. In addition to the differences between varieties (MG-20 has larger seeds than Gifu), in general seeds of the wild-type varieties were larger than their corresponding mutants, resulting in a broad range of diversity for seed size in *Lotus japonicus*. Reduced seed size in the mutants may be associated with altered meristematic activity, this being a direct effect of the mutation in *Ljhar1-1* (Nishimura *et al.*, 2002; Krusell *et al.*, 2002), or a consequence of altered hormonal sensitivity (ABA and ethylene in the ethylene-insensitive mutants).

Differences in shape were also found. Dry seeds of MG-20 fitted better to a modified cardioid than seeds of mutant *enigma-1* (J-index was smaller in *enigma-1* than the parent). The effect was maintained in imbibed seeds. Similarly, in imbibed seeds, the J-index was also smaller in *LjETRI-1* than in its wild type parent Gifu. Thus, in *enigma-1*, reduced seed size was associated with altered shape both in both dry and imbibed seeds (table 4 and figure 3). In *LjETRI-1* both effects were associated in imbibed seeds (dry seeds of *LjETRI-1* were smaller than Gifu, but no differences in shape were observed). Finally, seeds of *LjHar1-1*, although of reduced size did not present differences in shape with the parent (Gifu). Thus, the results allow discrimination between changes in size and shape during seed germination, and show different effects of the mutants in both characters. The effect of *enigma-1* mutants goes beyond seed size and involves shape alterations.

Mutants in *EIN2* gene presented alterations in size and shape both in dry and imbibed seeds in *Lotus*, but only in size and circularity index (not in J-index), in dry seeds of *Medicago*. Mutants in this gene have also different effects in nodulation. *Medicago* mutant (*sickle*) developed increased nodule numbers per plant while *Lotus* mutant (*enigma*) had reduced or equivalent values (dependent on the allele). These nodulation differences may be accounted for by the presence of a second copy of *Lotus EIN2* (*LjEIN2b*; Desbrosses and Stougaard, 2011) on *Lotus* chromosome 5, suggesting pathway diversification after gene duplication. Our results add new data to the contrasting nodulation phenotypes indicating divergences between both model species.

Our present results allow us to distinguish two processes during seed germination, *i.e.*, seed growth and the development of shape (table 4). Mutations in *Lotus EIN2a* gene (*enigma-1*) have effects on seed size and shape. The fact that other mutations (*LjHar1-1*), have effects only on seed size, but not on shape indicates that both processes may involve different regulatory mechanisms. For example, seed size may be the result of cell size, seed filling, and cell division activity, processes being under the control of distinct metabolic pathways, whereas seed shape could be submitted to systemic regulation, cell extension, physical constraint, and regulation of polarity. The combination of these processes represents an interesting system for the analysis of hormonal action in plant morphology.

ACKNOWLEDGEMENTS

PMG thanks the ARC for a Centre of Excellence grant, which supported PKC. Dr Bandana Biswas is thanked for the *enigma-1* mutant isolation and characterization. Dongxue Li and Ning Chen are thanked for technical support.

REFERENCES

- Beaudoin N, Serizet C, Gosti F, Giraudat J. Interactions between abscisic acid and ethylene signaling cascades. *Plant Cell* 2000; 12: 1103-1115.
- Begon M, Townsend CA, Harper JL. *Ecology: From Individuals to Ecosystems*, 4th ed. Blackwell, Oxford, UK. 2006.
- Biswas B, Chan PK, Gresshoff PM. A novel ABA insensitive mutant of *Lotus japonicus* with a wilted phenotype displays unaltered nodulation regulation. *Molecular Plant* 2009; 2: 487-499.
- Biswas B, Scott TE, Gresshoff PM. Tree legumes as feedstock for sustainable biofuel production: Opportunities and challenges. *J Plant Physiol.* 2011; 168: 1877-1884.
- Cannon SB, Sterck L, Rombauts S, Sato S, Cheung F, Gouzy J, Wang X, Mudge J, Vasdewani J, Schiex T, Spannagl M, Monaghan E, Nicholson C, Humphray SJ, Schoof H, Mayer KF, Rogers J, Quétier F, Oldroyd GE, Debellé F, Cook DR, Retzel EF, Roe BA, Town CD, Tabata S, Van de Peer Y, Young ND. Legume genome evolution viewed through the *Medicago truncatula* and *Lotus japonicus* genomes, *Proc Natl Acad Sci USA* 2006; 103 (40): 14959-14964.
- Cervantes E. Challenging *Arabidopsis thaliana* as the ultimate model species: Can seed germination be the Achilles' heel? *The European Journal of Plant Science and Biotechnology* 2008; 2 (1): 106-109.
- Cervantes E, Tocino A. Ethylene, free radicals and the transition between stable states in plant morphology. *Plant Signalling and Behaviour* 2009; 4(5): 367-371.
- Cervantes E, Martín JJ, Ardanuy R, de Diego JG, Tocino A. Modeling the *Arabidopsis* seed shape by a cardioid: efficacy of the adjustment with a scale change with factor equal to the Golden Ratio and analysis of seed shape in ethylene mutants. *J Plant Physiol.* 2010; 167: 408-410.

- Chan PK, Biswas B, Gresshoff PM. Classical ethylene insensitive mutants of the *Arabidopsis EIN2* orthologue lack the expected 'hypernodulation' response in *Lotus japonicus*. J. Integrated Plant Biology 2012; (in review).
- D'Arcy Thompson, W. On Growth and Form. Cambridge University Press, Cambridge, UK. 1917.
- Desbrosses GJ, Stougaard J. Root nodulation: a paradigm for how plant-microbe symbiosis influences host developmental pathways. Cell Host Microbe. 2011; 10:348-58.
- Ghassemian M, Nambara E, Cutler S., Kawaide H, Kamiya Y, and McCourt P. Regulation of abscisic acid signaling by the ethylene response pathway in *Arabidopsis*. Plant Cell 2000; 12: 1117-1126.
- Gielis J, Gerats T. A botanical perspective on plant shape modeling. Proceedings of the International Conference on Computing, Communications and Control Technologies Vol. VI, 2004, 265-272.
- Han L, Gresshoff PM, Hanan J. A functional-structural modelling approach to autoregulation of nodulation. Annals of Botany 2011; 107: 855-863.
- Han L, Hanan J, Gresshoff PM. Computational Complementation: A modelling approach to study signalling mechanisms during legume autoregulation of nodulation. PLOS Computational Biology 2010; 6 :e1000685.
- Jean, RV. Phyllotaxis: a Systemic Study in Plant Morphogenesis. Cambridge University Press, Cambridge, UK. 1994.
- Jiang Q, Gresshoff PM. (1997). Classical and molecular genetics of the model legume *Lotus japonicus*. Mol. Plant Microbe Interactions 1997; 10: 59-68.
- Kaplan, D R. The science of plant morphology: definition, history, and role in modern biology. American Journal of Botany 2001; 88:1711-1741.

Krusell L, Madsen LH, Sato S, Aubert G, Genua A, Szczygłowski K, Duc G, Kaneko T, Tabata S, de Bruijn F, Pajuelo E, Sandal N, Stougaard J. Shoot control of root development and nodulation is mediated by a receptor-like kinase. *Nature* 2002; 420: 422-426.

Lohar D, Stiller J, Kam J, Stacey G, Gresshoff PM. Ethylene insensitivity conferred by a mutated *Arabidopsis* ethylene receptor gene alters nodulation in transgenic *Lotus japonicus*. *Annals of Botany* 2009; 104: 277–285.

Moles A, Ackerly DD, Tweddle JC, Dickie JB, Smith R, Leishman MR, Mayfield MM, Pitman A, Wood JT, Westoby M, Murray BR, Brown AHD, Grace JP. Geographic gradients in seed size among and within perennial Australian *Glycine* species. *Australian Journal of Botany* 2003; 51 (1): 47 - 56

Nishimura R, Hayashi M, Wu GJ, Kouchi H, Imaizumi-Anraku H, Murakami Y, Kawasaki S, Akao S, Ohmori M, Nagasawa M, Harada K, Kawaguchi M. HAR1 mediates systemic regulation of symbiotic organ development. *Nature* 2002; 420: 426–429.

Penmetsa RV, Uribe P, Anderson J, Lichtenzveig J, Gish JC, Nam YW, Engstrom E, Xu K, Sckisel G, Pereira M, Baek JM, Lopez-Meyer M, Long SR, Harrison MJ, Singh KB, Kiss GB, Cook DR. The *Medicago truncatula* ortholog of the *Arabidopsis* EIN2 gene, sickle, is a negative regulator of symbiotic and pathogenic microbial interactions. *Plant Journal* 2008; 55: 580–595.

Robert C, Noriega A, Tocino A, Cervantes E. Morphological analysis of seed shape in *Arabidopsis thaliana* reveals altered polarity in mutants of the ethylene signaling pathway. *J Plant Physiol* 2008; 165: 911-919.

Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, Sasamoto S, Watanabe A, Ono A, Kawashima K, Fujishiro T, Katoh M, Kohara M, Kishida Y, Minami C, Nakayama S, Nakazaki N, Shimizu Y, Shinpo S, Takahashi C, Wada T, Yamada M,

Ohmido N, Hayashi M, Fukui K, Baba T, Nakamichi T, Mori H, Tabata S. Genome structure of the legume, *Lotus japonicus*. *DNA Research* 2008; 15: 227-239.

Schwarz, H. Two-dimensional feature-shape indices. *Mikroskopie (Wien)* 1980; 37 (Suppl.): 64-67.

Vogel H. A better way to construct the sunflower head. *Mathematical Biosciences* 1979; 44: 179-189.

Wopereis J, Pajuelo E, Dazzo FB, Jiang Q, Gresshoff PM, De Bruijn FJ, Stougaard J, and Szczyglowski K. Short root mutant of *Lotus japonicus* with a dramatically altered symbiotic phenotype. *Plant Journal* 2000; 23: 97-114.

Young ND, Cannon SB, Sato S, Kim D, Cook DR, Town CD, Roe BA, Tabata S. Sequencing the genespaces of *Medicago truncatula* and *Lotus japonicus*. *Plant Physiol* 2005; 137:1174–1181.

Zhang H, Han W, De Smet I, Talboys P, Loya R, Hassan A, Rong H, Jürgens G, Paul Knox J, Wang MH. ABA promotes quiescence of the quiescent centre and suppresses stem cell differentiation in the Arabidopsis primary root meristem. *Plant Journal* 2010; 64(5):764-74.

Figure legends

Figure 1

A. Adjustment of the bi-dimensional image of dry seeds of *Lotus japonicus* var. Gifu to a cardioid. The cardioid curve is shown on the left. In the centre, photographic images of *Lotus* seeds were superimposed to the cardioid. For the quantification of the adjustment areas in two regions were compared: The common region to the cardioid and the seed image and the regions not shared between both figures (figure on the right, striped).

B. Adjustment of the bi-dimensional image of dry seeds of *Medicago truncatula* var. Jemalong (A-17) to a cardioid scaled in the vertical axis by a factor of ϕ . The cardioid curve and modifications are shown on the left. In the centre, photographic images of *Medicago truncatula* seeds were superimposed to the cardioid. Quantification of the adjustment was as for Lotus (see legend to figure 1A).

Figure 2

Close up photo of pods and seeds: A. *Lotus japonicus* (scale bar equals 1 cm). B. *Medicago truncatula* (scale bar equals 5 mm).

Figure 3

Graphical representation of Area, Circularity index (I), and J-index in genotypes of *Lotus japonicus* (A) and *Medicago truncatula* (B).

Tables

A)

	Wild Type		Mutants		
	MG-20	Gifu	<i>enigma-1</i>	<i>Ljetr-1-1</i>	<i>Ljhar1-1</i>
Dry seeds	1.6	1.5	1.5	1.3	1.3
Imbibed seeds	3.8	3.2	3.1	2.6	2.8

B)

	Wild Type		Mutants		
	MG-20	Gifu	<i>enigma-1</i>	<i>Ljetr-1-1</i>	<i>Ljhar1-1</i>
Dry seeds	0.82	0.82	0.82	0.82	0.82
Imbibed seeds	0.86	0.85	0.85	0.85	0.86

C)

	Wild Type		Mutants		
	MG-20	Gifu	<i>enigma-1</i>	<i>Ljetr-1-1</i>	<i>Ljhar1-1</i>
Dry seeds	92.7	92.6	90.9	92.6	91.5
Imbibed seeds	93.5	90.9	89.8	88.5	92.4

Table 1

A) Mean values for area (mm²) of dry and imbibed seeds in *Lotus japonicus* genotypes are given.

B) Mean values for circularity index I of dry and imbibed seeds in *L. japonicus* genotypes.

C) Mean values for J-index of dry and imbibed seeds in *L. japonicus* genotypes.

A)

Genotype	N	Subset $\alpha = 0.05$		
		1	2	3
<i>Ljhar1-1</i>	94	1.28		
<i>LjETR1-1</i>	92	1.31		
<i>Gifu</i>	94		1.46	
<i>enigma-1</i>	95		1.47	
<i>MG20</i>	89			1.56
Significance		0.524	0.994	1.00

B)

Genotype	N	Subset $\alpha = 0.05$		
		1	2	3
<i>Ljhar1-1</i>	33	2.63		
<i>LjETR1-1</i>	31	2.84	2.84	
<i>Gifu</i>	26		3.15	
<i>enigma-1</i>	43		3.16	
<i>MG20</i>	32			3.81
Significance		0.574	0.135	1.00

Table 2

Contrast tests after ANOVA in *Lotus japonicus* genotypes. A) Tuckey test showing differences between groups for the area of dry seeds and B) Scheffé test showing differences between groups for the area of imbibed seeds.

A)

Genotype	N	Subset $\alpha = 0.05$	
		1	2
<i>enigma-1</i>	31	90.90	
<i>Ljhar1-1</i>	30	91.49	91.49
<i>LjEtr1-1</i>	30		92.61
Gifu	30		92.64
MG-20	30		92.74

B)

Genotype	N	Subset $\alpha = 0.05$			
		1	2	3	4
<i>LjEtr1-1</i>	33	88.48			
<i>enigma-1</i>	26	89.78	89.78		
Gifu	43		90.94	90.94	
<i>Ljhar1-1</i>	31			92.37	92.37
MG20	32				93.53
Significance		0.394	0.510	0.299	0.517

Table 3

A) Tuckey test showing differences between groups after ANOVA for J-index of dry seeds in *L. japonicus* genotypes.

B) Scheffé test showing differences between groups after ANOVA for J-index of imbibed seeds in *L. japonicus* genotypes.

	Size		Shape	
	dry	imbibed	dry	imbibed
Gifu vs MG-20	+	+	-	+
<i>LjEtr1-1</i> vs Gifu	+	-	-	+
<i>Ljhar1-1</i> vs Gifu	+	+	-	-
<i>enigma-1</i> vs MG-20	+	+	+	+

Table 4

Differences in seed size and shape between varieties of Lotus (Gifu vs MG-20), or between mutants and their respective parental genotypes. MG-20 is the parental for *enigma-1*; Gifu, for *LjEtr1-1* and *Ljhar1-1*. (+) represents difference; (-), no difference.

	Area		Circularity index		J-index	
	<i>A 17</i>	<i>sickle</i>	<i>A 17</i>	<i>sickle</i>	<i>A 17</i>	<i>sickle</i>
Dry seeds	5.3	4.7	0.72	0.70	87.1	86.0
Imbibed seeds	10.1	9.1	0.70	0.69	86.8	86.4

Table 5

Mean values for area, circularity index and index J of dry and imbibed seeds in *Medicago truncatula* genotypes.

Figure 1
[Click here to download high resolution image](#)

A

B

Figure 2
[Click here to download high resolution image](#)

A

B

Figure 3

Fig 3