Reactivity of unsaturated 5(4*H*)-oxazolones with Hg(II) acetate.

Synthesis of methyl *N*-benzoylamino-3-arylacrylates

Gheorghe-Doru Roiban,^{a,d,*} Tatiana Soler,^b Maria Contel,^c Ion Grosu,^a Carlos Cativiela^d
and Esteban P. Urriolabeitia^e

^a*Organic Chemistry Department, Faculty of Chemistry and Chemical Engineering, Babes Bolyai University, Arany Janos 11, 400028, Cluj Napoca, (Romania)*

^b*Servicios Centrales de Investigación, Facultad Ciencias, 03690 San Vicente de Raspeig, Alicante, (Spain)*

^c*Chemistry Department, Brooklyn College and The Graduate Center, The City University of New York, New York 11210, (USA)*

^d*Departamento de Química Orgánica, Instituto de Ciencia de Materiales de Aragón, CSIC-Universidad de Zaragoza, 50009 Zaragoza, (Spain)*

^e*Departamento de Compuestos Organometálicos, Instituto de Ciencia de Materiales de Aragón, CSIC-Universidad de Zaragoza, 50009 Zaragoza, (Spain)*

Correspondence e-mail: gdroiban@chem.ubbcluj.ro

ABSTRACT

An efficient and high-yield procedure to prepare methyl *N*-benzoylamino-3-arylacrylates from unsaturated (*Z*)-2-aryl-4-arylidene-5-(4*H*)-oxazolones and Hg(OAc)₂ in methanol is described herein. The observed reactivity of mercury(II) acetate here is different to its usual metallating behaviour, since it cleaves the unsaturated oxazolone ring without change of stereochemistry.

Keywords: 5-(4*H*)-Oxazolones; azlactone ring opening, alcoholysis; mercury(II) acetate.

INTRODUCTION

The most important reaction in the chemistry of oxazolones is the nucleophilic opening of the heterocyclic ring,^[1] hydrolysis and alcoholysis of 5(4*H*)-oxazolones giving the respective *N*-acylamino acids and esters, which are used to prepare a variety of new synthetic amino

acids.^[2] The use of *N*-acylamino alkyl esters as the starting α -dehydroamino acid derivatives also enable the selective formation of the corresponding isoquinolines.^[3] Some esters can be used as plant growth regulators,^[4] while other acrylic acid derivatives have herbicidal activity^[5] or are useful intermediates for the synthesis of the antiinflammatory agents.^[6] The methods used to synthesize methyl *N*-benzoylamino-3-arylacrylates include alcoholysis of 5(4*H*)-oxazolones in presence of MeONa,^[7] KOH,^[8] TEA,^[3,9] NEt₃,^[10] P(OEt)₃,^[11] or cyclooligosaccharides.^[12] In some cases, sonication has been also used to promote the nucleophilic ring opening.^[13] Recently we have shown that unsaturated 5(4*H*)-oxazolones can be *ortho*-palladated to give dimers which evolve after photocycloaddition to unprecedented oxazolones.^[14] Palladated oxazolones can be also regioselectively functionalized, the process leading to a series of clean products that are easy to isolate.^[15] Still, certain unsaturated oxazolone substrates proved to resist to the action of different palladation agents, while for other harsher conditions^[14] were needed. To extend oxazolones C-H bond activation and functionalization we focused on finding alternative milder synthetic routes involving a metal activator such as mercuriation. As it is known many aromatic substrates which refuse to react with Pd(OAc)₂ were derivatized using organomercury route followed by transmetallation.^[16] Mercury's unique abilities to mediate organic transformations and to transfer organic ligands to different transition metal centers provide a convenient entry to different substrates, difficult to prepare using classic synthetic methods.^[17] In particular Hg(OAc)₂ is known to attack easily aromatic compounds,^[17] and gives many interesting complexes by direct mercuriation.^[18] However, the increased Lewis acid character of Hg compared to Pd may induce sometime a different reactivity behavior function of metal as it was already noticed in metallation of 1,5-bis(dimethylamino)naphthalene, when Hg(OAc)₂ and Pd(OAc)₂ gave different metallated products.¹⁹ Another example is 3,4,5-trimethoxybenzaldehyde which does not react with Pd(OAc)₂ but it is easily mercuriated in the presence of mercury(II) acetate.^[20] To our knowledge, the treatment of unsaturated oxazolones with mercuric derivatives has not been investigated until now, therefore we

report here the reactivity of 5(4*H*)-oxazolones towards mercury(II) acetate.

RESULTS AND DISCUSSION

With this aim we prepared (*Z*)-2-phenyl-4-arylidene-5(4*H*)-oxazolones **1a-e** according to known literature procedures (**Equation 1**).^[9] Two geometric isomers are possible for **1a-e**, but the Erlenmeyer synthesis proceeds stereoselectively, favoring the thermodynamically more stable (*Z*) isomer.^[21] Arguments are based on NMR studies,^[22] chromatographic behavior^[23] and previous X-ray determinations of molecular structures of related oxazolones.^[24] With azlactones **1a-e** in hand, the next step was the study of their reactivity towards Hg(OAc)₂ in methanol, being known that different C,*N*-complexes have been prepared in these conditions.^[18] Therefore, substrate **1a**^[25] was refluxed in methanol in the presence of equimolar quantities Hg(OAc)₂. However, after workup only 2-benzoylamino-3-phenylmethylacrylate **2a**^[26] was isolated in a very good 98% yield (**Equation 1**). The presence of the expected metallated product could not be detected.

Equation 1

The same reaction occurred when oxazolones **1b**^[27] and **1c**^[28] were treated with Hg(OAc)₂, since ester derivatives **2b** and **2c**^[29] (**Table I**) were obtained in high yields. (*Z*)-2-phenyl-4(4-methoxybenzylidene)-5(4*H*)-oxazolone **1d**^[30] afforded, after treatment with Hg(OAc)₂, 2-benzoylamino-3-(4-methoxyphenyl)methyl acrylate **2d**.^[31] It is evident that in these processes the Hg(OAc)₂ did not behave as a metallation agent, but hydrolysed the oxazolone ring affording methyl *N*-benzoylamino-3-arylacrylates **2a-d**. Curiously the electron activated substrate **1e**^[32] could not be hydrolyzed in the above conditions even if the reaction time was increased to 24 hours. In addition, even it is widely accepted that the mercuriation occurs mainly through an electrophilic substitution S_EAr pathway^[33] no metallating product was detected after workup when activated substrate **1e** was employed.

It should be noted that compound **1a** remained practically unchanged when refluxed in methanol alone for 20 hrs, as noted also by Shabana et al.^[11] Under the same reaction conditions Pd(OAc)₂ leave the oxazolone ring intact as we observed in our initial attempts to get cyclopalladated derivatives.^[15] In order to see if the reaction can be catalytically driven **1d** was refluxed in MeOH using 3% molar of Hg(OAc)₂. However, the observed conversion of **1d** into **2d** was only (***)% after more than 36 hours.

Table I

In the ¹H NMR spectrum of **2d**, the shielding of the aromatic protons shows that the delocalization of the electron density occurs in a lower extent in comparison with starting oxazolone. The presence of a broad peak around 7.70 ppm, in **2d**, assigned to the acidic proton from the NH group is in keeping with the oxazolone cleavage. In the aliphatic area of the spectrum the existence of the methyl ester group is confirmed by a sharp peak at δ 3.79 ppm. Methyl *N*-benzoylamino-3-arylacrylates **2a-d** have been also characterized using IR spectroscopy. Absorptions assigned to the stretch of the C=O group were detected at about 1700 cm⁻¹, shifted to lower energy in all studied cases with respect to the two-component band appearing in the 1740-1780 cm⁻¹ region in starting oxazolones.^[14] Bands due to NH vibrations appear around 3300 cm⁻¹ in products **2a-d** while the C=N stretching vibration appear at 1652 cm⁻¹. Mass spectra performed on compounds **2a-d** are strongly in the favor of the drawn structures. When compound **2b** is analyzed under MS-ES conditions the protonated molecule is detected at *m/z* 359.9, 361.8, while the cationic [MNa]⁺ species appear at *m/z* 381.8, 383.8.

The structure of 2-benzylamino-3-(4-chlorophenyl)methyl acrylate (**2c**) was also confirmed by the determination of its molecular structure by X-ray diffraction methods. From the structure of **2c** it is obvious the cleavage of the heterocyclic ring and the incorporation of a methanol molecule (**Figure 1**). In comparison with starting oxazolones **1a-d**, which are planar,^[24] the molecule planarity is destroyed by the ring opening. The crystal structure of **2c**

displays a *Z* configuration about the C8C9 double bond. This is the same configuration than that found in the oxazolone **1c**, this meaning that the reaction occurs with stereochemical retention of the configuration.

Figure 1

The molecular conformation is stabilized by an intramolecular and several intermolecular hydrogen bonds. The intramolecular H-bonds are formed between the proton on the aromatic bond C13H13 and the N1 atom [H13A...N1 2.5266 (14) Å] and has similar parameters to those found in isobutyl *N*-benzoylamino-3-(4-chlorophenyl) acrylate^[34] and *N*-benzoylamino-3-(4-chlorophenyl) acrylic acid.^[35] In addition to that, intermolecular H-bonds involving the aromatic CH groups, carbonyl O atoms, Cl atoms of the chlorophenyl groups and NH protons also stabilize the molecule.

In conclusion we observed that reaction of unsaturated 5(4*H*)-oxazolones with Hg(OAc)₂ in methanol did not lead to any metallated product. Instead mercury(II) acetate showed low Lewis acid behavior when reacted with unsaturated oxazolones. In the given reaction conditions oxazolone ring was cleaved, methyl *N*-benzoylamino-3-arylacrylates being obtained in high yields. The reaction did not occur in catalytic amounts of mercury(II) acetate while electron activated substrates could not be hydrolyzed.

EXPERIMENTAL

Chemicals of commercial grade were used without further purification. ¹H and ¹³C{¹H} NMR spectra were recorded in CDCl₃, at room temperature, on a Bruker Avance 400 spectrometer (δ in ppm, *J* in Hz) at ¹H operating frequency of 400.13 MHz. ¹H and ¹³C NMR spectra were referenced using the solvent signal as internal standard. MS-ES mass spectra were recorded using an Esquire 3000 ion-trap mass spectrometer (Bruker Daltonic GmbH) equipped with a standard ESI/APCI source. Infrared spectra (4000-380 cm⁻¹) were recorded

on a Perkin-Elmer Spectrum One IR spectrophotometer, using nujol mulls between polyethylene sheets. Data collection was performed at room temperature on an Oxford Diffraction Xcalibur2 diffractometer using graphite-monocromated Mo-K α radiation (λ = 0.71073 Å). An hemisphere of data was collected based on three ω -scan or ϕ -scan runs. The diffraction frames were integrated using the program CrysAlis RED^[36] and the integrated intensities were corrected for absorption with SADABS.^[37] *Structure Solution and Refinement.* The structure was solved and developed by Patterson and Fourier methods.^[38] All non-hydrogen atoms were refined with anisotropic displacement parameters. The hydrogen atoms were placed at idealized positions and treated as riding atoms. Each hydrogen atom was assigned an isotropic displacement parameter equal to 1.2 times the equivalent isotropic displacement parameter of its parent atom. The structure was refined to F_o^2 , and all reflections were used in the least-squares calculations.^[39] Elemental analyses (CHNS) were carried out on a Perkin-Elmer 2400-B microanalyser.

Synthesis of the starting compounds

Unsaturated 5-(4*H*)-oxazolones **1a**,^[25] **1b**,^[27] **1c**,^[28] **1d**^[30] and **1e**^[32] were previously prepared and characterised before. Also alcoholysis products **2a**,^[26] **2c**^[29] and **2d**^[31] have been also spectroscopically characterised elsewhere.

General Procedure for preparation of methyl *N*-benzoylamino-3-arylacrylates

(*Z*)-2-Arylidene-4-phenyl-5(4*H*)-oxazolone (1.0 equiv) and mercury(II) acetate (1.0 equiv) were added in MeOH (10-30 mL). The mixture was heated for 3 to 4 hours at 70 °C until the monitoring of the reaction using TLC showed completion and then allowed to cool at room temperature. The solvent was evaporated and the residue was purified on a chromatographic column (Ethyl Acetate) to give **2a-d**.

Synthesis of methyl *N*-benzoylamino-3-(2-bromophenyl)acrylate

(*Z*)-2-(2-Bromobenzylidene)-4-phenyl-5(4*H*)-oxazolone **1b** (1.27 g, 3.89 mmol) and mercury(II) acetate (1.24 g, 3.89 mmol) were added in MeOH (30 mL). The mixture was heated for 3 hours at 70 °C and after reaction completion monitored on TLC, allowed to cool at room temperature. The solvent was evaporated to dryness and then the solid washed

three times with cold diethyl ether. The united ether phases were evaporated to give **2b** as a white solid (Yield 1.34 g, 96%). ^1H NMR (400 MHz, CDCl_3) δ ppm: 3.85 (s, 3H, COOCH_3), 7.12 (td, $^3J = 7.7$ Hz, $^4J = 1.7$ Hz, 1H, $\text{H}_{4''}$), 7.18 (t, $^3J = 6.8$ Hz, 1H, $\text{H}_{5''}$), 7.39 (t, $^3J = 7.6$ Hz, 2H, $\text{H}_{3'}$, $\text{H}_{5'}$), 7.47 (m, 2H, $\text{H}_{4'}$, $\text{H}_{6''}$), 7.52 (s, 1H, $\text{H}_{7''}$), 7.57-7.62 (dd, $^3J = 8.0$ Hz, $^4J = 1.2$ Hz, 1H, $\text{H}_{3''}$), 7.77 (d, $^3J = 7.4$ Hz, 2H, $\text{H}_{2'}$, $\text{H}_{6'}$), 7.97 (broad s, 1H, NH). ^{13}C NMR (101 MHz, CDCl_3) δ ppm: 52.84 (1C, COOCH_3), 124.34, 126.06, 133.21, 134.54 (4C, $\text{C}_{1''}$, $\text{C}_{2''}$, $\text{C}_{1'}$, C_2), 127.14 (1C, $\text{C}_{5''}$), 127.33 (2C, $\text{C}_{2'}$, $\text{C}_{6'}$), 128.56 (2C, $\text{C}_{3'}$, $\text{C}_{5'}$), 129.34, 129.37 (2C, $\text{C}_{4'}$, $\text{C}_{7''}$), 130.09 (1C, $\text{C}_{4''}$), 132.07 (1C, $\text{C}_{6''}$), 132.83 (1C, $\text{C}_{3''}$), 165.38, 165.43 (2C, CO, COOCH_3). MS-ES m/z : 359.9, 361.8 $[\text{MH}]^+$, 381.8, 383.8 $[\text{MNa}]^+$. Anal. Calc. for $\text{C}_{17}\text{H}_{14}\text{BrNO}_3$: C, 56.69; H, 3.92; N, 3.89. Found: C, 56.84; H, 3.76; N, 3.56.

ACKNOWLEDGMENTS

Funding from CNCSIS TD87/2007 (Romania), the Ministerio de Ciencia e Innovacion (Projects CTQ2008–01784 and CTQ2007–62245, Spain), and Gobierno de Aragon (PI071–09) is gratefully acknowledged. DR thanks also to WFS for financial support.

Equation 1

Table I. Alcoholysis of (*Z*)-2-aryl-4-arylidene-5(4*H*)-oxazolones **1a-d**.

Oxazolone	X	Product	Yield %
1a	H	2a	98
1b	2-Br	2b	96
1c	4-Cl	2c	97
1d	4-OMe	2d	94
1e	3,4-OMe	-	-

Figure 1. ORTEP plot of **2c** with thermal ellipsoids drawn at the 50% probability level

REFERENCES

- 1 Cativiela, C.; Díaz-de-Villegas, M. D. *The Chemistry of Heterocyclic Compounds*; Wiley: New York; Palmer, D. C. Ed.; 2004, Vol. 60, pp. 129-330.
- 2 Liang, J.; Ruble, J. C.; Fu, G. C. Dynamic kinetic resolutions catalyzed by a planar-chiral derivative of DMAP: Enantioselective synthesis of protected α -amino acids from racemic azlactones. *J. Org. Chem.* **1998**, 63, 3154-3155.
- 3 Hoshina, H.; Tsuru, H.; Kubo, K.; Igarashi, T.; Sakurai, T. Formation of isoquinoline derivatives by the irradiation of *N*-acetyl- α -dehydrophenylalanine ethyl ester and its derivatives. *Heterocycles* **2000**, 53, 2261-2274.
- 4 Keith, D. D.; Weigele, M. Butenoic acid derivatives. U.S. Patent 4216008, 1980; *Chem. Abstr.* **1981**, 94, 16083.
- 5 Bakos, J.; Gebhardt, I.; Lukacs, L.; Kiraly, J.; Bihari, F.; Heil, B.; Durko, M.; Radvany, B.; Boros, P.; Eifert, G. Preparation of acrylic acid derivatives as herbicides. Hungarian Patent HU 47203, 1989; *Chem. Abstr.* **1991**, 114, 116927.
- 6 Slouka, J. Preparation of 2-acetylamino-3-[1-phenyl-3-(4-chlorophenyl)-4-pyrazolyl]acrylic acid as an intermediate for Ionazolac. Czechoslovakian Patent CS 267956, 1990; *Chem. Abstr.* **1991**, 114, 185493.
- 7 Jursic, B. S.; Sagiraju, S.; Ancalade, D. K.; Clark, T.; Stevens, E. D. Practical preparation of *Z*- α -(*N*-acetylamino)- and *Z*- α -(*N*-benzoylamino)- α,β -unsaturated acids. *Synthetic Commun.* **2007**, 37, 1709-1714.
- 8 Gagey, N.; Emond, M.; Neveu, P.; Benbrahim, C.; Goetz, B.; Aujard, I.; Baudin, J.-B.; Jullien, L. Alcohol uncaging with fluorescence reporting: evaluation of *o*-acetoxypheyl methyloxazolone precursors. *Org. Lett.* **2008**, 10, 2341-2344.
- 9 (a) Rao, Y. S.; Filler, R. Geometric isomers of 2-aryl(aralkyl)-4-arylidene(alkylidene)-5(4*H*)-oxazolones. *Synthesis* **1975**, 749-764; (b) Rzeszotarska, B.; Karolak-Wojciechowska, J.; Broda, M. A.; Galdecki, Z.; Trzezwinska, B.; Koziol, A. E. Conformational investigation of

α,β -dehydropeptides. Part VI. Molecular and crystal structure of benzyloxycarbonylglycyl-(*Z*)-dehydrophenylalanine. *Int. J. Pept. Protein Res.* **1994**, *44*, 313-319.

10 Maekawa, K.; Hishikawa, N.; Kubo, K.; Igarashi, T.; Sakurai, T. Preferential formation of cis-4,5-dihydrooxazole derivatives via photoinduced electron transfer-initiated cyclization of *N*-acyl- α -dehydroarylalanine alkyl esters. *Tetrahedron* **2007**, *63*, 11267-11281.

11 Shabana, R.; Sidky, M. M.; El-Kateb, A. A.; Soliman, F. M. Studies on organophosphorus compounds. Part IX. The action of di-, and trialkyl phosphites and thio acids on some enamino lactones. *Phosphorus Sulfur* **1987**, *34*, 57-61.

12 Lee, S.; Jung, S. Cyclosophoraose as a catalytic carbohydrate for methanolysis. *Carbohydr. Res.* **2004**, *339*, 461-468.

13 Rao, C. C.; Lalitha, N. A new methodology for the synthesis of *N*-acylaminocinnamates and amides from azlactones. *Indian J. Chem., Sect. B* **1994**, *33*, 3.

14 Roiban, D.; Serrano, E.; Soler, T.; Grosu, I.; Cativiela, C.; Urriolabeitia, E. P. Unexpected [2 + 2] C-C bond coupling due to photocycloaddition on orthopalladated (*Z*)-2-aryl-4-arylidene-5(4*H*)-oxazolones. *Chem. Commun.* **2009**, 4681-4683.

15 Roiban, G. D.; Serrano, E.; Soler, T.; Contel, M.; Grosu, I.; Cativiela, C.; Urriolabeitia, E. P. ortho-Palladation of (*Z*)-2-aryl-4-arylidene-5(4*H*)-oxazolones. *Organometallics* **2010**, *29*, 1428-1435.

16 (a) Constable, E. C.; Leese, T. A.; Tocher, D. A. Polynuclear intermediates in the synthesis of cyclometallated complexes; the crystal and molecular structure of (2-pyridylphenyl)mercury(II) chloride tetramer. *J. Chem. Soc., Chem. Commun.* **1989**, 570-571;

(b) Clark, A. M.; Rickard, C. E. F.; Roper, W. R.; Wright, L. J. Electrophilic substitution reactions at the phenyl ring of the chelated 2-(2'-pyridyl)phenyl ligand bound to ruthenium(II) or osmium(II). *Organometallics* **1999**, *18*, 2813-2820; (c) Parish, R. V.; Wright, J. P.; Pritchard, R. G. Mercury(II) and gold(III) derivatives of 2-phenyl pyridines and 2-phenyl-4-(methylcarboxylato)quinoline. *J. Organomet. Chem.* **2000**, *596*, 165-176; (d) Clark, A. M.; Rickard, C. E. F.; Roper, W. R.; Wright, L. J. Bromination and nitration reactions of

metallated (Ru and Os) multiaromatic ligands and crystal structures of selected products. *J. Organomet. Chem.* **2000**, 598, 262-275.

17 Gabbaï Melaimi, F. P.; Burress, C. N.; Melaimi, M. -A.; Taylor, T. J. *Comprehensive Organometallic Chemistry III*; Crabtree, R. H.; Mingos, D. M. P., Eds.; Elsevier: Amsterdam, U.K.; 2007; Vol. 2, pp. 419-474.

18 (a) Urriolabeitia, E. P. *Palladacycles: Synthesis, Characterization and Applications*; Wiley: Weinheim, Dupont, J.; Pfeffer, M., Eds.; 2008, pp. 35-67; (b) Soro, B.; Stoccoro, S.; Minghetti, G.; Zucca, A.; Cinellu, M. A.; Gladiali, S.; Manassero, M.; Sansoni, M. Synthesis of the first C-2 cyclopalladated derivatives of 1,3-bis(2-pyridyl)benzene. Crystal structures of [Hg(N-C-N)Cl], [Pd(N-C-N)Cl], and [Pd₂(N-C-N)₂(μ-OAc)]₂[Hg₂Cl₆]. Catalytic activity in the Heck reaction. *Organometallics* **2005**, 24, 53-61.

19 Phillips, I. G.; Steel, P. J. Cyclometallated compounds : VIII. Double cyclopalladation of diacetylbenzene dioximes and 1,5-bis(dimethylamino)naphthalene. *J. Organomet. Chem.* **1991**, 410, 247-255.

20 Vicente, J.; Abad J. -A.; Stiakaki, M. -A.; Jones, P. G. Palladium-assisted rearrangement of HC(O)C₆H₂(OMe)₃-3,4,5 to HC(O)C₆H₂(OMe)₃-2,3,4. X-Ray structures of [PdR(PPh₃)(bipy)](CF₃SO₃)[R = C₆H(CHO)-6,(OMe)₃-2,3,4 and C₆H(CHO)-2,(OMe)₃-3,4,5; bipy = 2,2'-bipyridyl]. *J. Chem. Soc., Chem Commun*, **1991**, 137-139.

21 Kumar, K.; Phelps, D. J.; Carey, P. R. Resonance Raman labels: spectroscopic studies on cis- and trans-4-benzylidene-2-phenyl-Δ²-oxazolin-5-one and an isotopically substituted analog. *Can. J. Chem.* **1978**, 56, 232-239.

22 (a) Venkataratnam, R. V.; Suryakumari, A. Configurational assignment to geometric isomers of 2-phenyl-4-(3-phenyl-2-propenylidene)-5(4*H*)-oxazolone. *Indian J. Chem., Sect. B* **1988**, 27, 832-833; (b) Prokof'ev, E. P.; Karpeiskaya, E. I. The proton coupled carbon-13 NMR direct determination of *Z*-, *E*-configuration of 4-benzylidene-2-phenyl(methyl)-Δ²-oxazolin-5-ones and products of their solvolysis. *Tetrahedron Lett.* **1979**, 737-740; (c) Cativiela, C.; Sanchez-Ferrando, F. Measurement of long-range heteronuclear selective

carbon-13 {proton} NOE enhancements: 4-(α -arylethylidene)-2-phenyl-5(4*H*)-oxazolones. *Magn. Reson. Chem.* **1985**, 23, 1072-1075.

23 (a) Kavrakova, I.; Koleva, V.; Kurtev, B. On the relative chromatographic mobility of *Z*- and *E*-4-arylidene-2-phenyl-5-oxazolones. *Dokl. Bolg. Akad. Nauk* **1984**, 37, 1495-1496; *Chem. Abstr.* **1985**, 103, 37396; (b) Palamareva, M.; Kavrukova, I. Microcomputer aided choice of mobile phases for thin-layer chromatographic separation of some *E*- and *Z*-oxazolones and related cinnamates on silica. *Izv. Khim.* **1988**, 21, 218-225; *Chem. Abstr.* **1989**, 111, 247239; (c) Palamareva, M.; Kurtev, B.; Kavrakova, I. Chromatographic behavior of diastereoisomers. X. Thin-layer chromatographic retentions on silica of some (*E*)- and (*Z*)-oxazolones and related cinnamates as a function of mobile phase effects or Hammett constants. *J. Chromatogr.* **1991**, 545, 161-175.

24 (a) Haasbroek, P. P.; Oliver, D. W.; Carpy, A. J. M. Evidence of the structure of 4-(4'-acetoxybenzylidene)-2-methyl-5-oxazolone and its phenylpropenoic acid derivatives. *J. Mol. Struct.* **2003**, 648, 61-67; (b) Buseti, V.; Mayoral, J. A.; Cativiela, C.; Diaz de Villegas, M. D.; Ajo, D. Crystal structure of cyclic dehydroaminoacid derivatives. I. Tetrasubstituted (*Z*)-5(4*H*)-oxazolones. *Z. Kristallogr.* **1989**, 189, 65-76.

25 (a) King, S. W.; Riordan, J. M.; Holt, E. M.; Stammer, C. H. Synthesis of racemic (*E*)- and (*Z*)-1-amino-2-phenylcyclopropanecarboxylic acid: (*E*)- and (*Z*)-cyclopropylphenylalanine. *J. Org. Chem.* **1982**, 47, 3270-3273; (b) Hamidian, H.; Tikdari, A. M. Novel synthesis of unsaturated 5(4*H*)-oxazolone derivatives with using palladium(II) acetate as a catalyst and microwave irradiation in solvent-free condition. *Heterocycl. Commun.* **2006**, 12, 29-34.

26 Pines, S. H.; Kozlowski, M. A.; Karady, S. Mechanism and stereochemical considerations in the reaction of some arylserine derivatives with thionyl chloride. *J. Org. Chem.* **1969**, 34, 1621-1627.

27 Mesaik, M. A.; Rahat, S.; Khan, K. M.; Zia-Ullah; Choudhary, M. I.; Murad, S.; Ismail, Z.; Atta-ur-Rahman; Ahmad, A. Synthesis and immunomodulatory properties of selected oxazolone derivatives. *Bioorg. & Med. Chem.* **2004**, 12, 2049-2057.

-
- 28 (a) Cativiela, C.; Diaz de Villegas, M. D.; Melendez, E. On the synthesis of geometric isomers of 2-methyl (or phenyl)-4-[α -arylethylidene]-5(4*H*)-oxazolones. *J. Heterocycl. Chem.* **1985**, *22*, 1655-1657; (b) Kidwai, M.; Sapra, P.; Bhushan, K. R.; Misra, P. Microwave-assisted solid-support synthesis of pyrazolino/iminopyrimidino/thioxopyrimidino imidazolines. *Synthesis* **2001**, 1509-1512.
- 29 Cativiela, C.; Diaz de Villegas, M. D.; Melendez, E. Synthesis of methyl (*Z/E*)-2-acetamido(or benzamido)-3-aryl-2-butenates. *Tetrahedron* **1986**, *42*, 583-590.
- 30 (a) Yadav, L. D. S.; Singh, A. Microwave activated solvent-free cascade reactions yielding highly functionalized 1,3-thiazines. *Tetrahedron Lett.* **2003**, *44*, 5637-5640; (b) Ghosh, S.; Datta, D. B.; Sen, N. Synthesis of *N*-(*E*)-(4-methoxystyryl)benzamide (alataamide). *Synthetic Commun.* **1987**, *17*, 299-307; (c) Bautista, F. M.; Campelo, J. M.; Garcia, A.; Luna, D.; Marinas, J. M.; Romero, A. A. Study on dry-media microwave azalactone synthesis on different supported KF catalysts: influence of textural and acid-base properties of supports. *J. Chem. Soc., Perkin Trans. 2* **2002**, 227-234.
- 31 Ghosh, S.; Datta, D. B.; Sen, N. Synthesis of *N*-(*E*)-(4-methoxystyryl)benzamide (alataamide). *Synthetic Commun.* **1987**, *17*, 299-307.
- 32 O'Reilly, N. J.; Derwin, W. S.; Lin, H. C. Optically pure (*S*)-6,7-dimethoxy-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid and asymmetric hydrogenation studies related to its preparation. *Synthesis*, **1990**, 550-556.
- 33 Berger, A.; de Cian, A.; Djukic, J. -P.; Fischer, J.; Pfeffer, M. Syntheses of ortho-mercured and -palladated (η^6 -arene)tricarbonylchromium complexes. *Organometallics* **2001**, *20*, 3230-3240.
- 34 Su, G. F.; Wang, Z. C.; Huang, W. Y.; Wang, Z. X.; Chen, Z. L. (*Z*)-Isobutyl 2-benzamido-3-(4-chlorophenyl)acrylate. *Acta Cryst. E* **2009**, *65*, o2890.
- 35 Palmer, R. A.; Richards, A. J.; Lisgarten, J. N. 2-Benzamido-3-(*p*-chlorophenyl)propenoic acid (antrex). *Acta Cryst. C* **1995**, *51*, 2338-2340.

-
- 36 Oxford Diffraction. *CrysAlis RED*. version 1.171.27p8. 2005, release CrysAlis171 .NET, Abingdon, Oxford, England.
- 37 Sheldrick, G. M. *SADABS*, 2008, University of Göttingen, Germany.
- 38 Sheldrick, G. M. SHELXS-86, *Acta Crystallogr.* **1990**, *A46*, 467.
- 39 Sheldrick, G. M. SHELXL-97: FORTRAN program for the refinement of crystal structures from diffraction data. Göttingen University, 1997. Molecular graphics were done using the commercial package SHELXTL-PLUS, Release 5.05/V: © 1996, Siemens Analytical X-ray Instruments, Inc., Madison, Wisconsin.