Title

Effects of phosphorus deficiency and genetic variation on leaf terpene contents and emission rates in *Pinus pinaster* seedlings susceptible and resistant to the attack of the pine weevil *Hylobius abietis* L.

Authors and affiliations

Josep-Salvador Blanch¹, Luis Sampedro², Joan Llusà¹, Xoaquin Moreira², Rafael Zas³, Josep Peñuelas¹

¹ Global Ecology Unit CREAM-CEAB-CSIC, Edifici C, Universitat Autònoma de Barcelona, 08193, Bellaterra, Catalunya, Spain

² Centro de Investigación Forestal de Lourizán – Unidad Asociada MBG-CSIC. Apartado 127. 36080 Pontevedra, Spain

³ Misión Biológica de Galicia. Consejo Superior de Investigaciones Científicas. Apdo 28. 36080 Pontevedra, Spain

Short title

Phosphorus deficiency and genetic variation on *Pinus pinaster* terpene emissions and contents

Corresponding author

Josep-Salvador Blanch
Global Ecology Unit CREAM-CEAB-CSIC
Edifici C, Universitat Autònoma de Barcelona
08193
Bellaterra, Catalunya, Spain
e-mail:js.blanch@creaf.uab.cat

Keywords:

Maritime pine, nutrient stress, *Hylobius abietis*, Galicia.

Abstract

We studied the effects of phosphorus fertilization on foliar terpene concentrations and on foliar volatile terpene emission rates in different half-sib families of *Pinus pinaster* Ait. seedlings. Half of them appeared to be resistant to the attack of the pine weevil *Hylobius abietis* L., a generalist phloem feeder, and the other half appeared to be susceptible to this insect. We hypothesized that P stress could modify the terpene concentration in the needles and thus derive to altered terpene emission patterns relevant to plant-insect signalling. The total concentrations and emission rates ranged between 5732 and 13995 $\mu\text{g g}^{-1}$ d.m. and between 2 and 22 $\mu\text{g g}^{-1}$ d.m. h^{-1} respectively. The storage and emission were dominated by the isomers α and β -pinene (77.2 % and 84.2 % of the total terpene amount respectively). P stress caused in both resistant and susceptible families an increase of 31% of the foliar terpene concentrations with an associated 5-fold decrease of the terpene emission rates. Those higher contents would indicate an allocation of the “excess of carbon” generated due to growth being limited because of P scarcity, to terpene emissions. Sensible families showed a higher increase of terpene emission rates, which could be related to plant-animal communication.

Introduction

Phosphorus has many roles in plant growth and metabolism. One of the principal functions of phosphorus is energy transfer: through the action of adenosine triphosphate (ATP). ATP and its derivatives, ADP and AMP, are involved in all aspects of energy transfer in every part of plant growth. Apart from this global function, phosphorus is also necessary for assembling nucleic acids (DNA and RNA), proteins, enzymes and carbohydrates. It plays an essential role in photosynthesis and is involved in the formation of sugars and starch. The various roles of phosphorus denote the fact that it is important in the formation of seeds and the development of roots. It also speeds plant maturity and helps the plant resist stresses (Urbano, 1999).

However, fertilization of young pine seedlings and the subsequent boosting of primary growth rates could lead to increased susceptibility to pests and diseases due to altered allocation patterns of energy to growth and defence and/or improved tissue quality for the insects. In this sense, in a field study Zas *et al.* (2006a; 2008) found that traditional silvicultural practices such as phosphorus fertilization could lead to

greater susceptibility to the attack by the pine weevil *Hylobius abietis* L. in seedlings of *P. pinaster* and *P. radiata*, which may be at least partially explained by a reduction in resistance (Moreira *et al.* 2008). The pine weevil *Hylobius abietis* L. is a generalist phloem-feeder that constitute a major pest in conifer plantations in all Europe, where causes important regeneration problems due to the fact that adults feed the bark of young seedlings (Conord *et al.* 2006; Leather *et al.* 1999). The susceptibility of *P. pinaster* to *H. abietis* attack has been found to present an intense genetic variation, where some families were consistently more damaged than others (Zas *et al.* 2005).

Greater nutrient availability could directly increase the nutritional value of the plant tissues and thus increase the preference by the insects (Ayres & Lombardero 2000; Moreira *et al.* 2009). Phosphorus fertilization on P stressed pine seedlings may diminish the allocation of energy to constitutive and induced defences by favouring the growth rates. Several models of plant defence suggest altered patterns of allocation to chemical defences in environments with increased nutrient availability. The Carbon nutrient balance (Bryant *et al.* 1983) stated that when growth is limited by nutrients, plants allocate the “excess carbon” to the production of secondary metabolites. The Growth differentiation balance (Lorio 1986) recognizes that all secondary metabolites have an ontogenetically determined phenology and that their synthesis is emphasized during periods of plant differentiation. Growth dominates during favourable conditions, and differentiation is at a maximum only when conditions are suboptimal for growth. This could be more evident in tree species with predetermined growth such as pine trees. The Optimal allocation model (Tuomi *et al.* 1991) predicts decreasing investment in defence with increasing resource availability, because reduced costs of tissue production could compensate higher risks of herbivore predation. Higher phosphorus availability could also lead to a higher appearance of the fertilized plants to the insect. Fertilization could change the amount of emitted and leaf-contained volatile organic compounds as it may affect the secondary metabolism as stated by “excess carbon” hypotheses (Peñuelas & Estiarte 1998).

Maritime pine (*Pinus pinaster* Ait.) has been widely chosen as forestation species in Galicia (NW Spain) since the XVIIIth century. Despite being partly replaced in the last decades by species with higher productions like *Pinus radiata* and *Eucalyptus globulus*, *P. pinaster* is still the most important forest tree species in Galicia. According to the last forest survey (DGCN 2000), Galicia has 389,489 ha of

monospecific stands and 243,735 ha of mixed stands with eucalyptus or broad-leaved species. Thus, *P. pinaster* is present in 44% of the total Galician wooded area. The intensive silviculture applied to *P. pinaster* entails short rotations (15 to 45 years), in which there is an important extraction of nutrients of the system (Merino *et al.* 2003).

Conifer plantations in Galicia commonly suffer important nutrient deficiencies (Martins *et al.* 2009). These plantations are usually located on acid soils with few nutrients, especially phosphorus. Moreover, the loss of nutrients due to harvesting can lead to decreased reserves of soil available limiting nutrients (Dambrine *et al.* 2000; Merino *et al.* 2005). Under those conditions, phosphorus stress is commonly found in conifer and especially in *P. pinaster* stands in NW Spain (Martins *et al.* 2009).

We hypothesized that P stress could modify the terpene concentration in the needles and the photosynthetic activity of *P. pinaster* thus leading to altered terpene emission patterns relevant to plant-insect signalling. The objective of the present study was to test this hypothesis. With this aim and the additional aims of studying the effect of genetic variation and the relationships with the resistance to pests, we analyzed the effect of P fertilization on terpene concentrations and on terpene emission rates in half-sib families of *P. pinaster* seedlings cultivated under controlled conditions, previously found to be resistant or susceptible to the large pine weevil in field conditions in Galicia forests.

Material and Methods

Experimental design and plant material

We performed a two factorial experiment with different pine genetic entries and phosphorus fertilization treatments under controlled conditions. The experimental layout was a randomized split-plot design replicated in four blocks, with four phosphorus fertilization treatments acting as the whole factor and six genetic entries as the split factor. In total, we sampled 72 pine seedlings, corresponding to 3 blocks (randomly selected within the 4 blocks) $\times$ 4 phosphorus fertilization treatments $\times$ 6 genetic entries nested into two susceptibility groups, ‘susceptible’ and ‘resistant’ families.

P. pinaster families belonged to six different open-pollinated families with common mother tree (half-sibs). Three families were previously recognized to be susceptible to the attack by the pine weevil (*Hylobius abietis*) in an extensive field study, while the other three families appeared to be more resistant to this plague (Zas *et al.* 2005).

Damage (debarked area by the pine weevil) to the susceptible families in that field study was more than two-fold greater than that suffered by the resistant families (Zas *et al.* 2005). All progenies are native from the coastal region of Galicia (NW Spain).

Plant material, greenhouse conditions and experimental fertilization

On 7 February 2006, *P. pinaster* seeds were individually sown in 2 L. pots containing perlite in a greenhouse with controlled temperature (10-22 °C at night and day, respectively), low photosynthetic photon flux density and daily water irrigation.

On 15 March 2006, we started to apply the fertilization treatments by sub-irrigation (every two days) with four different fertilization treatments. The complete balanced fertilization (“P20”) was prepared according optimum requirements for maritime pine tree growth, containing 100 ppm of N, 20 ppm of P, 40 ppm of K, 10 ppm of Ca, 20 ppm of Mg, and the necessary amounts of micronutrients and trace elements. The other phosphorus fertilization (“P10”, “P5” and “P2”) differed only in the concentration of phosphorus, which were unbalanced to limit the growth promoting a P limitation, with a concentration of 10, 5 and 2 ppm of phosphorus in the fertilizer solution, respectively. The pH values were adjusted to 6.5 in all the solutions. Fertilizer solutions were replaced every two weeks. The experiment was carried out in a glass greenhouse (36.5 m long and 15 m wide) belonging to the facilities of CIF Lourizán (Pontevedra, NW Spain, UTM coordinates 29T 42°24’33’’ N 8°39’47’’W).

Photosynthetic activity and terpene emission collection

On 24-27 July 2006, measurements of net photosynthetic rates, stomatal conductance and terpene emissions were conducted. These measurements were done at controlled standard conditions (30°C and 1000 $\mu\text{mol m}^{-2} \text{h}^{-1}$ PAR). CO₂ exchange was measured using a non-dispersive infra-red gas analyzer (IRGA), model ADC-LCPro+ (ADC Inc. Hoddesdon, Hertfordshire, England) connected to a conifer leaf chamber (ADC Inc. Hoddesdon, Hertfordshire, England). CO₂ uptake (*A*) and stomatal conductance (*g_s*) were measured in lateral shoots on *P. pinaster*. *A* and *g_s* values were expressed on a projected leaf area basis measured with Li-Cor 3100 Area Meter (Li-Cor Inc., Nebraska, USA).

In order to sample terpene emissions, a T-system was installed outside the cuvette of the IRGA-porometer. Part of the air passed through cartridges (8 cm long

and 0.3 cm internal diameter) manually filled with terpene adsorbents Carboxen 1003, and Carboxen 1003, and Carboxen Y (Supelco, Bellefonte, Pennsylvania) separated by plugs of quartz wool by using a Q_{max} air sampling pump (Supelco, Bellefonte, Pennsylvania) at constant flow. The hydrophobic properties of the tubes were supposed to minimize sample displacement by water. In these tubes, terpenes did not suffer chemical transformations as checked with standards (α -pinene, β -pinene, camphene, myrcene, p-cymene, limonene, sabinene, camphor, and dodecane). Prior to use, these tubes were conditioned for 10 min at 350 °C with a stream of purified helium. The sampling time was 5 min, and the flow varied between 470 and 500 ml min⁻¹ depending on the tubes' adsorbent and quartz wool packing. A calibrated air sampling pump was used to trap isoprenoids. The trapping and desorption efficiency of liquid and volatilized standards such as α -pinene, β -pinene or limonene was practically 100%. In order to eliminate the problem of memory effect of previous samples, blanks of 5-min air sampling without plants were carried out immediately before and after each measurement. The glass tubes were stored in a portable fridge at 4 °C and taken to the laboratory where they were stored at -28 °C until analysis (within 24-48 h). There were no observable changes in terpene concentrations after storage of the tubes as checked by analyzing replicate samples immediately and after 48-h storage. Emission rate calculations were made on mass balance basis and by subtracting the control values (without plants) from the values of samples with plants.

Seedling harvesting and nutrient analyses

On 1 August 2006, we measured height and basal diameter (mean of two measures) and then we destructively sampled all the pine seedlings. For the analysis of foliar terpene content, a composite sample of primary needles from different parts of each tree was collected, deep frozen and preserved at -80 °C into close-tight glass vials. To estimate final seedling biomass we separated into roots, stems, braquiblasts and needles and immediately put in a drying oven for 72 h at 65 °C. After being removed from the oven, we weighed samples to the nearest 0.001 g. All the samples were finely grounded, labelled and preserved for nutrient analysis.

For the analysis of nitrogen and phosphorus content, 0.3 g of phloem, roots, braquiblasts and needles were digested in a mixture of selenous sulphuric acid and hydrogen peroxide (Walinga *et al.* 1995). Nitrogen was colorimetrically analysed in diluted aliquots of this digestion using a BioRad 680 microplate reader (California,

USA) at $\lambda = 650$ nm according the method proposed by Sims *et al.* (1995). Phosphorus was analysed in the same diluted aliquots by inductively coupled plasma optical emission spectroscopy (ICP-OES) using a Perkin-Elmer Optima 4300DV (Massachusetts, USA) in the central laboratory facilities at Universidade de Vigo – CACTI (www.uvigo.es/webs/cactiweb/). Nitrogen and phosphorus concentration were expressed in mg g^{-1} dried weight of tissue.

Terpene analysis

Tubes with trapped emitted monoterpenes were inserted in an OPTIC3 injector (ATAS GL International BV 5500 AA Veldhoven, The Netherlands) where they were desorbed at 250 °C during 3 min. Terpenes were separated using a 30m x 0.25mm x 0.25 μm film thickness capillary column (SPB TM-5 Fused Silica Capillary column, Supelco Inc., Bellefonte, PE, USA). After sample injection, the initial temperature (40°C) was increased at 30 °C min^{-1} up to 60 °C, and thereafter at 10 °C min^{-1} up to 150 °C maintained for 3 min, and thereafter at 70 °C min^{-1} up to 250 °C, which was maintained for another 5 min. Helium flow was 1 mL min^{-1} . The identification of terpenes was conducted by GC-MS and comparison with standards from Fluka (Buchs, Switzerland), literature spectra and GCD Chemstation G1074A HP with the Wiley275 library. Terpene calibration curves (for 4 different terpene concentrations) were always significant ($r^2 > 0.99$). The most abundant terpenes had very similar sensitivity (differences were less than 5%). Terpene concentration was referred to needle dried weight (d.w.).

For extraction of resin terpenoids in the needles, three-four needles were grounded under liquid nitrogen in Teflon tubes with a Teflon embolus. Then, we added 1 mL of pentane as extractant and 0.1 μL of dodecane, a non-terpenoid internal standard. Teflon tubes with pentane samples were centrifuged in an ultrasonic bath for 5 minutes at 5000 rpm and 5-10 °C to separate the liquid and solid phases. Pentane extracts were immediately recovered and transferred to chromatography glass vials. After recovering the pentane extract, the mass of the needle pellet was determined by oven-drying at 65 °C for 4 days. Terpenes in the extract were analyzed using a Hewlett Packard HP59822B GC-MS (Palo Alto, CA, USA) with a robotic sample processor (FOCUS) (ATAS GL International BV 5500 AA Veldhoven, The

Netherlands). Separation, quantification and identification were performed as described above.

Statistical analyses

All traits were analyzed by the following model: $H_{ijk} = \mu + B + P + R + G(R) + P*G(R) + P*R + B*R + B*P + \epsilon_{ijk}$, where H_{ijk} is the variable of the trait, μ is the overall mean, B, P, R and G are the main effects of block, phosphorus fertilization, resistance and genotype, and ϵ_{ijk} is the experimental error. Genotype was nested in resistance G(R). B*P was considered a random factor for properly analyze the split plot design (Littell *et al.* 2006). A MIXED procedure of SAS was used. When main effects were significant, differences among treatment means were tested for significance using the LSMEAN statement.

A PROC GLM procedure of SAS (Littell *et al.* 2006) was used for the MANOVA analyses; Wilk's Lambda statistics were used.

Results

Plant growth and needle nutrient concentrations

Fertilization treatments strongly affected plant growth ($F = 20.82$, $P < 0.001$) and phosphorus concentration in plant tissues ($F = 141.39$, $P < 0.0001$) (Table 1). Plants with complete fertilization (20 ppm) produced 2.5-fold greater biomass than plants with lower fertilization (Fig. 1). P concentration in needles was strongly influenced by fertilization, showing increasing values accordingly to the P fertilization, where plants under balanced fertilization exhibited P concentrations 3-fold greater than P stressed plants (Fig. 1). The only treatment that drove P concentration in needles under critical levels was 2 ppm; therefore, this treatment was the one that generated the clearest P deficiency.

Nitrogen concentration in needles was only slightly greater, but significant ($F = 5.97$, $P < 0.05$) in complete fertilization than in P stressed plants (Table 1, Fig.1).

Those families with a resistant behaviour at field showed slightly higher concentrations of P ($F = 7.79$, $P < 0.01$) in leaf tissues than susceptible families, but no differences in terms of N concentrations ($F = 3.16$, $P < 0.1$) and total biomass ($F = 2.73$, $P = 0.1$) were detected (Table 1, Fig.1).

Photosynthesis (A), stomatal conductance (g_s) and transpiration rates (E)

Fertilization treatments decreased photosynthesis ($F = 4.48$, $P < 0.05$) and transpiration ($F = 6.12$, $P < 0.05$) (Table 1, Fig. 2): complete fertilization (20 ppm) produced lower A and E than the lowest fertilization treatment (2 ppm) (Fig. 2). However, these effects were different in resistant families than in sensible families: there was a strong interaction $P \times R$ (Table 1) for A , g_s and E : sensible families showed the lowest values of A and E at 10 ppm of fertilization, and resistant families showed the lowest values of A and E at 20 ppm of fertilization.

Different families had significant differences in photosynthesis ($F = 2.72$, $P < 0.05$) and stomatal conductance ($F = 10.23$, $P < 0.0001$) (Table 1).

Volatile terpenes

Several mono- and sesquiterpenes were found in both leaf concentrations and in terpene emissions. The relative percentages in the total amount is shown in Table 2. The isomers α and β -pinene dominate the production (77.2 %) and emission (84.2 %) of the total terpene amount. Δ^3 -carene is also present with high percentage: 14.3% of the concentrations, and 5% of the emission rates. The rest of the compounds appeared in smaller percentages.

MANOVA analysis for the individual compounds showed significant differences for phosphorus deficiency ($\lambda = 0.15$, $P < 0.01$), resistance ($\lambda = 0.44$, $P < 0.01$) and genotype ($\lambda = 0.03$, $P < 0.0001$), but there was not significant effect for $P \times res$, which indicates that the different treatments influenced not the individual but the whole terpene profile of our samples (Table 3).

Terpene concentrations significantly increased with phosphorus deficiency ($F = 4.25$, $P < 0.05$) (Table 1, Fig. 3). On the contrary, total terpene emission rates significantly decreased with phosphorus fertilization ($F = 9.76$, $P < 0.01$) (Table 1, Fig. 3). This increase was much higher in sensible species than in resistant families (Fig. 3).

There was a high effect of Genotype ($F = 26.78$, $P < 0.0001$) in terpene emission rates: different families showed different behaviours. Thus, total terpene emission rates significantly increased ($F = 19.48$, $P < 0.0001$) in sensible families at higher P doses (10 and 20 ppm).

Discussion

Terpene compounds

The mean concentration values ranged from an average of 7.9 mg g^{-1} (with 20 ppm of P addition) to an average of 12.6 mg g^{-1} (with 2 ppm of P addition) (Fig. 3), which are lower than other literature values for the same species (Arrabal *et al.* 2005) or in other pine species (e.g. Blanch *et al.* 2009).

P. pinaster is not considered a big isoprenoid emitter (Kesselmeier & Staudt 1999). However, the mean emission rates values ranged from an average of $2.5 \text{ } \mu\text{g g}^{-1} \text{ h}^{-1}$ (with 2 ppm of P addition) to an average of $16 \text{ } \mu\text{g g}^{-1} \text{ h}^{-1}$ (with 20 ppm of P addition) (Fig. 3) which is much higher than literature values: Simon *et al.* (1994) found emission rates of $0.2 \text{ } \mu\text{g g}^{-1} \text{ h}^{-1}$ in *P. pinaster*.

The difference of values could be due to the differences in climate during the measures, which were warmer in our location. Moreover, emission rates depend on several other factors such as ontogeny, the qualitative and quantitative composition of the terpenes, the terpene pool position, the pathway of diffusion of the terpene within the plant, the morphology of the vegetal surface, the vapour pressure of terpenes, the temperature, the relative humidity, the water stress, and mechanical as well as chemical injuries of the plant. These lower concentration values and the higher emission rate suggest that in the site of measurement (Galicia) and the season of the year considered (summer), *P. pinaster* tends to emit the monoterpenes instead of keeping them in the terpene pools.

Our results where α - and β -pinene were the 77.2% and 84.2% of the total emission rates and concentrations respectively (Table 2) agree with previous studies that have shown that α - and β -pinene are the principal terpenes emitted (Simon *et al.* 1994) and contained (Arrabal *et al.* 2005; Ormeño *et al.* 2009) by *P. pinaster*. The vapour pressure of these two compounds is two to three times higher than those of other terpenes, whereas the rate constants of their reactions with O_3 , $\text{OH}^\cdot$ and $\text{NO}_3^\cdot$ are lower (Atkinson 1990).

Phosphorus and genetic effects

P concentration in leaves was above the P deficiency levels proposed for field studies (Bonneau 1995) in all cases. That is, our fertilization ranged from high levels to low levels, but always within the regular physiological margin. According to Fig. 1, the

fertilization stress treatment was significantly effective: the higher the fertilization dose, the higher the concentration of P in the plant. The increase of nitrogen and phosphorus leaf concentrations with P fertilization has been previously reported (Keay *et al.* 1968). Moreover in agreement with previous authors that have reported the effect that P fertilization increases the growth of *P. pinaster* (Zas *et al.* 2006b), P fertilization increased the amount of biomass in fertilized plants (Fig. 1).

There was a general effect of P deficiency on photosynthesis rates, stomatal conductance and transpiration which showed certain tendency to increase with lower P doses (Fig. 3). The most fertilized seedlings showed a decrease of A , g_s and E in comparison with the least fertilized seedlings. Previous authors have also reported negative correlations between P fertilization and A (Cheaib *et al.* 2005; Loustau *et al.* 1999). Warren & Adams (2002) suggested that the lack of photosynthetic response to P supply was the result of a deficiency of N induced by high P supply.

Resistant and non-resistant species showed contrary responses to initial P deficiency. Genetic differences in nutrient use efficiency in response to fertilization in many tree species have been previously reported (i.e. Zas *et al.* 2006b; 2008).

Regarding the production and emission rates of terpenes, the most P-stressed conditions (doses 1 and 2 ppm) showed higher leaf terpene concentrations and lower terpene emission rates (Fig. 3). P deficient plants seem to store the produced terpenes in storing pools instead of emitting them. Higher amounts of terpenes were emitted in the less stressed conditions (doses 10 and 20 ppm) in comparison with the most stressed conditions, especially in sensible species (Fig. 3). These higher emission rates can be explained by many of the theories based on the “excess carbon” hypothesis (Peñuelas & Estiarte 1998) such as the Carbon-Nutrient Balance theory (Bryant *et al.* 1983) and the Growth Differentiation theory (Lorio 1986), which state that when the resources are higher than the needs for growth, plants use those resources to produce carbon based secondary metabolites (Peñuelas & Estiarte 1998).

An increase of P would also increase the attack of *H. abietis* due to the fact that fertilization increases the amount of α -pinene emitted and that *H. abietis* is attracted by the monoterpene α -pinene (Moreira *et al.* 2008). As a result, the amount of debarked area in young seedlings would increase with higher P availability (Zas *et al.* 2006b). The fact that sensible species emit more terpenes with higher P concentrations than resistant species could be explained as a defence of these plants to

H. Abietis: sensible plants emit more terpenes under fertilized conditions (Fig.3) in order to attract natural predators of the weevil that could attack them under those high fertilized conditions. Degenhardt (2008) showed that terpene emissions can attract nematodes to the plant that emits those terpenes, and nematodes are natural predators of *H. abietis* (Dillon *et al.* 2006).

In conclusion, higher phosphorus availability altered the plant physiology (higher biomass, higher nutrient concentrations), increased the production of monoterpenes, and decreased the of *P. pinaster* (storing species). There was a genetic effect: different families showed different responses in physiology and in terpene production and emission. The higher terpene emission rates of sensible families could explain a greater attraction of predators of the pine weevil.

Acknowledgements

We thank Patricia Martins for her superb technical assistance in the experimental setup, and Chema Mendaña and other collaborators for their assistance with the greenhouse. This research was supported by INIA grants, and by funding from Spanish Government grants (RTA 2005-173, RTA07-100 and CGL2006-04025/BOS) and Consolider-Ingenio Montes (CSD2008-0040) and from the Catalan Government grant SGR2009/458. LS and XM received financial support from DOC-INIA and PREDOC-INIA grant programs respectively.

References

- Arrabal, C., Cortijo, M., de Simon, B. F., Vallejo, M. C. G. and Cadahia, E. (2005) Differentiation among five Spanish *Pinus pinaster* provenances based on its oleoresin terpenic composition. *Biochemical Systematics and Ecology*. **33**, 1007-1016
- Ayres, M.P. and Lombardero, M.J. (2000) Assessing the consequences of global change for forest disturbance from herbivores and pathogens. *The Science of the Total Environment*. **262**, 263-286
- Atkinson, R. (1990) Gas-Phase Tropospheric Chemistry of Organic-Compounds - a Review, *Atmospheric Environment Part A-General Topics*. **24**, 1-41
- Bonneau, M. (1995) *Fertilisation des forêts dans les pays tempérés: théorie, bases du diagnostic, conseils pratiques, réalisations expérimentales*. ENGREF, Nancy, France.

400 Blanch, J. S., Peñuelas, J., Sardans, J. and Llusà J. (2009) Drought, warming
 401 and soil fertilization effects on leaf volatile terpene concentrations in *Pinus halepensis*
 402 and *Quercus ilex*. *Acta Physiologiae Plantarum*. **31**, 207-218

403 Bryant, J. P., Chapin, F. S. and Klein, D. R. (1983) Carbon Nutrient Balance
 404 of Boreal Plants in Relation to Vertebrate Herbivory. *Oikos*. **40**, 357-368

405 Cheaib, A., Mollier, A., Thunot, S., Lambrot, C., Pellerin, S. and Loustau, D.
 406 (2005) Interactive effects of phosphorus and light availability on early growth of
 407 maritime pine seedlings. *Annals of Forest Science*. **62**, 575-583

408 Conord, C., Lemperiere, G., Taberlet, P. and Despres, L. (2006) Genetic
 409 structure of the forest pest *Hylobius abietis* on conifer plantations at different spatial
 410 scales in Europe. *Heredity*. **97**, 46-55

411 Dambrine, E., Vega, J. A., Taboada, T., Rodriguez, L., Fernandez, C., Macias,
 412 F. and Gras, J. M. (2000) Budgets of mineral elements in small forested catchments in
 413 Galicia (NW Spain), *Annals of Forest Science*. **57**, 23-38

414 Delzon, S., Bosc, A., Cantet, L. and Loustau, D. (2005) Variation of the
 415 photosynthetic capacity across a chronosequence of maritime pine correlates with
 416 needle phosphorus concentration. *Annals of Forest Science*. **62**, 537-543

417 Degenhardt, J. (2008) Ecological roles of vegetative terpene volatiles. In: A.
 418 Schaller (Ed). *Induced Plant Resistance to Herbivory*. Springer, Dordrecht,
 419 Netherlands: 433-442.

420 DGCN (2000) *Tercer Inventario Forestal Nacional, 1997-2006: Galicia.*,
 421 Ministerio de Medio Ambiente, Dirección General de Conservación de la Naturaleza,
 422 Madrid, Spain.

423 Dillon, A.B., Ward, D., Downes, M.J. and Griffin, C.T. (2006) Suppression of
 424 the large pine weevil *Hylobius abietis* (L.) (Coleoptera : Curculionidae) in pine
 425 stumps by entomopathogenic nematodes with different foraging strategies. *Biological*
 426 *Control*. **38**:217-226

427 Keay, J., Turton, A. G. and Campbell, N. A. (1968) Some Effects of Nitrogen
 428 and Phosphorus Fertilization of *Pinus Pinaster* in Western Australia. *Forest Science*.
 429 **14**: 408-417

430 Kesselmeier, J., and Staudt, M. (1999) Biogenic volatile organic compounds
 431 (VOC): An overview on emission, physiology and ecology. *Journal of Atmospheric*
 432 *Chemistry*. **33**: 23-88

433 Leather, S. R., Day, K. R. and Salisbury, A. N. (1999) The biology and
 434 ecology of the large pine weevil, *Hylobius abietis* (Coleoptera : Curculionidae): a
 435 problem of dispersal? *Bulletin of Entomological Research*. **89**: 3-16
 436 Littell, R.C., Milliken, G.A., Stroup, W.W., Wolfinger, R.D. and
 437 Schabenberger, O. (2006) *SAS System for mixed models, second edition*. SAS
 438 Institute, Cary, NC.
 439 Lorio, P. L. (1986) Growth-Differentiation Balance - a Basis for
 440 Understanding Southern Pine-Beetle Tree Interactions. *Forest Ecology and*
 441 *Management*. **14**: 259-273
 442 Loustau, D., Ben Brahim, M., Gaudillere, J. P. and Dreyer, E. (1999)
 443 Photosynthetic responses to phosphorus nutrition in two-year-old maritime pine
 444 seedlings. *Tree Physiology*. **19**: 707-715
 445 Martins, P., Sampedro, L., Moreira, X. and Zas, R. (2009) Nutritional status
 446 and genetic control of phenotypic plasticity to nutrient availability in *Pinus pinaster*.
 447 A multisite field study in NW Spain. *Forest Ecology and Management*. **258**: 1429-
 448 1436
 449 Mattson, W. J. and Haack, R. A. (1987) The Role of Drought in Outbreaks of
 450 Plant-Eating Insects. *Bioscience*. **37**: 110-118
 451 Merino, A., Balboa, M. A., Soalleiro, R. R. and Gonzalez, J. G. A. (2005)
 452 Nutrient exports under different harvesting regimes in fast-growing forest plantations
 453 in southern Europe. *Forest Ecology and Management*. **207**: 325-339
 454 Merino, A., Rey, C., Brañas, J. and Rodríguez-Soalleiro, R. (2003) Biomasa
 455 arbórea y acumulación de nutrientes en plantaciones de *Pinus radiata* D. Don en
 456 Galicia, *Investigación agraria: Sistemas y recursos forestales*. **2**: 85-98
 457 Moreira, X., Costas, R., Sampedro, L. and Zas, R. (2008) Short
 458 communication. A simple method for trapping *Hylobius abietis* (L.) alive in northern
 459 Spain. *Investigación agraria: Sistemas y recursos forestales*. **17**: 188-192
 460 Moreira, X., Sampedro, L. and Zas, R., (2009) Defensive responses of *Pinus*
 461 *pinaster* seedlings to exogenous application of methyl-jasmonate: Concentration
 462 effect and systemic response. *Environmental and Experimental Botany*. **67**: 94-100
 463 Ormeño, E., Céspedes, B., Sanchez, I. A., Velasco-Garcia, A., Moreno, J. M.,
 464 Fernandez, C. and Baldy, V. (2009) The relationship between terpenes and
 465 flammability of leaf litter. *Forest Ecology and Management*. **257**: 471-482

- Peñuelas, J., and Estiarte, M. (1998) Can elevated CO₂ affect secondary metabolism and ecosystem function? *Trends in Ecology & Evolution*. **13**: 20-24
- Simon, V., Clement, B., Riba, M. L. and Torres, L. (1994) The Landes Experiment - Monoterpenes Emitted from the Maritime Pine. *Journal of Geophysical Research-Atmospheres*. **99**: 16501-16510
- Sims, J.T., Vasilas, B.T., Gartley, K.L., Milliken, B. and Green, V. (1995), Evaluation of soil and plant nitrogen tests for maize on manured soils of the Coastal Plain. *Agronomy Journal*. **87**: 213-222
- Tuomi, J., Fagerstrom, T. and Niemela, P. (1991) Carbon allocation, phenotypic plasticity, and induced defences. In: D. W. Tallamy and M. J. Raupp (Eds). *Phytochemical Induction by Herbivores*. John Wiley & Sons, New York: 85-104.
- Urbano, P. (1999) *Tratado de fitotecnica general*, Mundi-Prensa ed., Barcelona.
- Walinga, I.; Van der Lee, J. J.; Houba, V. J. and Van Vark, W. (1995) *Plant analysis manual*, Kluwer Academic Publ., Dordrecht, the Netherlands.
- Warren, C. R. and Adams M. A. (2002) Phosphorus affects growth and partitioning of nitrogen to Rubisco in *Pinus pinaster*. *Tree Physiology*. **22**: 11-19
- Zas, R., Sampedro, L., Moreira, X. and Martins, P. (2008) Effect of fertilization and genetic variation on susceptibility of *Pinus radiata* seedlings to *Hylobius abietis* damage. *Canadian Journal of Forest Research*. **38**: 63-72
- Zas, R., Sampedro, L., Prada, E., Lombardero, M. J. and Fernández-López, J. (2006a) Fertilization increases *Hylobius abietis* L. damage in *Pinus pinaster* Ait. Seedlings. *Forest Ecology and Management*. **222**: 137-144
- Zas, R., Pichel, F., Martins, P. and Fernández-López, J. (2006b) Fertilization x genotype interaction in *Pinus radiata* open pollinated families planted in three locations in Galicia (NW Spain). *New Forests*. **32**: 253-263
- Zas, R., Sampedro, L., Prada, E. and Fernández-López, J. (2005) Genetic variation of *Pinus pinaster* Ait. seedlings in susceptibility to the pine weevil *Hylobius abietis* L. *Annals of Forest Science*. **62**: 681-688

Tables

Table 1. Summary of the split-plot model for P and N concentration in needles, total biomass, net photosynthetic rates ($\mu\text{g m}^{-2} \text{ s}^{-1}$), stomatal conductance ($\text{mol m}^{-2} \text{ s}^{-1}$), transpiration rates ($\mu\text{mol m}^{-2} \text{ s}^{-1}$), Total Terpene Contents ($\mu\text{g g}^{-1}[\text{d.m.}]$) and Total Terpene Emission Rates ($\mu\text{g g}^{-1}[\text{d.m.}] \text{ h}^{-1}$) of *P. pinaster*. B P R and G are the main effects of block, fertilization, resistance and genotype. Genotype was nested in resistance G(R).

	DF num	DF den	P needles		N needles		Total biomass	
			F	p	F	p	F	p
B	3	9	1,19	0,3661	0,89	0,4841	1,81	0,2151
P	3	9	141,39	<.0001	5,97	0,0160	20,82	0,0002
R	1	33	7,79	0,0087	3,16	0,0846	2,73	0,1081
G(R)	4	33	1,15	0,3526	0,78	0,5483	2,95	0,0345
P*G(R)	12	33	1,85	0,0793	0,89	0,5632	1,94	0,0659
P*R	3	33	3,75	0,0201	1,45	0,2462	4,53	0,0091
B*R	3	33	0,21	0,8917	0,68	0,5697	1,52	0,2271

	DF num	DF den	Net photosynthetic rate		Stomatal conductance		Transpiration Rate	
			F	p	F	p	F	P
B	3	9	0,01	0,9981	0,07	0,9730	0,17	0,9151
P	3	9	4,48	0,0347	2,58	0,1179	6,12	0,0148
R	1	33	1,30	0,2623	0,28	0,5977	0,19	0,6685
G(R)	4	33	2,72	0,0460	10,23	<.0001	1,76	0,1603
P*G(R)	12	33	2,03	0,0539	2,70	0,0118	1,49	0,1788
P*R	3	33	7,35	0,0007	6,17	0,0019	5,08	0,0053
B*R	3	33	0,79	0,5106	0,88	0,4595	0,38	0,7658

	DF num	DF den	Total terpene concentration		Terpene emission rate	
			F	p	F	p
B	3	9	0,48	0,7038	1,61	0,2544
P	3	9	4,25	0,0396	9,76	0,0034
R	1	33	0,22	0,6456	19,48	0,0001
G(R)	4	33	4,16	0,0078	16,78	<.0001
P*G(R)	12	33	1,99	0,0584	3,56	0,0028
P*R	3	33	0,94	0,4329	5,32	0,0046
B*R	3	33	1,05	0,3823	2,47	0,0813

Table 2. Individual and total terpene concentrations (n=68) and emission rates (n=70) for all families and all treatments.

	Terpene concentration		Terpene emission	
	$\mu\text{g g}^{-1}$	%	$\mu\text{g g}^{-1}\text{h}^{-1}$	%
cis-ocimene	14,66	0,16		
α -pinene	4203,48	46,65	4,33	46,80
camphene	63,36	0,70	0,34	3,70
β -pinene	2757,90	30,60	3,46	37,45
myrcene	133,80	1,48	0,05	0,57
Δ^3 -carene	1288,85	14,30	0,47	5,05
sabinene	296,44	3,29	0,06	0,67
β -phellandrene			0,25	2,69
terpinolene	30,39	0,34	0,04	0,39
α -fenchene	27,68	0,31		
trans-caryophyllene	65,16	0,72		
α -humulene	29,49	0,33		
germacrene	50,92	0,57		
limonene+b-phellandrene			0,17	1,80
other compounds	47,59	0,53	0,08	0,89

Table 3. Summary of the Multivariate Analysis for Total Terpene Contents ($\mu\text{g g}^{-1}\text{d.m.}$) and Total Terpene Emission Rates ($\mu\text{g g}^{-1}\text{d.m. h}^{-1}$) for *P. pinaster*. B P R and G are the main effects of block, fertilization, resistance and genotype. Genotype was nested in resistance G(R).

Manova hipótesis	Wilk's Lambda	p-value
Non-general P effects	0.15194675	0.0036
Non-general R effects	0.44274740	0.0060
Non-general G(R) effects	0.03232137	<.0001
Multivariate analysis	0.28185726	0.3472

Figure legends

Fig.1 Nitrogen and Phosphours concentrations (mg g^{-1} d.m.) in needles and total biomass (mg), for fertilization and resistance treatments. Vertical bars indicate standard error of the mean ($n=12$ in the two upper panels, $n=24$ in the lower panel). Different letters indicate significant statistical differences among fertilization levels.

Fig.2 Net photosynthetic rates ($\mu\text{g m}^{-2} \text{s}^{-1}$), stomatal conductance ($\text{mol m}^{-2} \text{s}^{-1}$) and transpiration rates ($\text{mmol m}^{-2} \text{s}^{-1}$) for fertilization and resistance treatments. Vertical bars indicate standard error of the mean ($n=12$). Different letters indicate significant statistical differences among fertilization levels.

Fig 3. Total Terpene Contents ($\mu\text{g g}^{-1}$ [d.m.]) and Total Terpene Emission Rates ($\mu\text{g g}^{-1}$ d.m. h^{-1}) for fertilization and resistance treatments. Vertical bars indicate standard error of the mean ($n=12$). Different letters indicate significant statistical differences among fertilization levels.

537
538

Fig. 1

Fig. 2.

Fig. 3