1 **Effects of light availability on mixotrophy and microzooplankton grazing in an**
2 **oligotrophic plankton food web: evidences from a mesocosm study in Eastern**
3 **Mediterranean waters**

4 Albert Calbet^{1*}, Rodrigo A. Martínez¹, Stamatina Isari¹, Soultana Zervoudaki², Jens
5 C. Nejstgaard³, Paraskevi Pitta⁴, Andrey F. Sazhin⁵, Despoina Sousoni^{4, 6}, Ana
6 Gomes¹, Stella A. Berger^{3,7}, Tatiana M. Tsagaraki⁴, Robert Ptacnik⁸

7 ¹ Institut de Ciències del Mar (CSIC). Ps. Marítim de la Barceloneta 37-49. 08003.
8 Barcelona, Spain.

9 ² Institute of Oceanography, Hellenic Centre for Marine Research, P.O. Box 712,
10 19013 Anavyssos, Athens, Greece

11 ³ Skidaway Institute of Oceanography, 10 Ocean Science Circle, Savannah, GA,
12 31411 USA.

13 ⁴ Institute of Oceanography, Hellenic Centre for Marine Research, P.O. Box 2214,
14 71003 Heraklion, Crete, Greece

15 ⁵ P.P.Shirshov Institute of Oceanology RAS, 36 Nahimovski prospect, 117997
16 Moscow, Russia

17 ⁶ Department of Biology, University of Crete, P.O. Box 2208, 71409 Heraklion,
18 Crete, Greece

19 ⁷ Department of Biology, University of Bergen, P.O. Box 7803, 5020 Bergen, Norway

20 ⁸ ICBM, University of Oldenburg. Schleusenstrasse 1, DE-26382 Wilhelmshaven.
21 Oldenburg, Germany.

22

23 * Corresponding author; email: acalbet@icm.csic.es; Phone: +34932309519; FAX:
24 +34932309500

25

26

27

1 **Abstract**

2 Plankton biomass and composition in the pelagic zone of oceans is exposed to
3 changes in availability of light and nutrients due to large-scale ocean circulation and
4 water column stratification. We hypothesized that displacement of plankton from
5 surface to deeper darker waters would not only favor heterotrophy over time, as
6 previously suggested, but also first rapidly affect the level of mixotrophy and,
7 consequently, overall microbial grazing in plankton food webs. To test this in an
8 oligotrophic marine system we incubated Eastern Mediterranean water (from 10 m
9 depth north of Crete in September 2010) in 2.8 m³ mesocosms simulating two
10 different light intensities at the sampling station, surface waters (ca. 10 m; mesocosms
11 L1) and deeper layers (ca. 50-60 m; mesocosms L4). The biomass and abundance of
12 the main planktonic groups were monitored either daily or every second day,
13 depending on the group. Microzooplankton grazing rates and the contribution of
14 mixotrophic feeding were estimated by a combination of dilution experiments and
15 incubations with live fluorescently labeled algae (LFLA). Although no nutrients were
16 added to the mesocosms the chlorophyll *a* increased during the first 2 days of the
17 experiment in both treatments. This increase resulted from phytoplankton growth in
18 the light L1-mesocosm (autotrophic biomass was ca. doubled in L1 compared to L4),
19 but was mostly due to photoadaptation of the algae in the L4-mesocosm, as indicated
20 by lower carbon to chlorophyll *a* ratios. By the end of the experiment, the total
21 biomass of protozoan and metazoan grazers in L1 was ca. twofold higher than in L4.
22 The microzooplankton responded within the first 24 h, showing different grazing
23 activity in L1 than in L4. Microzooplankton grazing rates on total Chl *a* were similar
24 in both treatments; however, phytoplankton instantaneous growth rates were higher in
25 the more illuminated mesocosm. This resulted in a closer coupling between both rates

1 in L4, where all production was grazed daily, than in L1. Nevertheless, the overall
2 flux of carbon through the microzooplankton was 33-60% higher in L1 compared to
3 L4 throughout the experiment. The fraction of mixotrophy in the ciliate community
4 varied in L1 (20-50%), but decreased over time in L4 from 50% to 30%. Our results
5 do not support studies from freshwater, postulating that reduced light and nutrient
6 limitation may increase herbivore production due to stoichiometric effects. Finally,
7 we discuss how mixotrophy may bias rate estimates in dilution experiments.

8

9

10

11 **Key words:** light, Eastern Mediterranean, microzooplankton, mixotrophy, mesocosm,
12 dilution experiments

13

1 **1. Introduction**

2 Light is a major driver of life on Earth, and therefore regulates the production and
3 distribution of phototrophic organisms. This spatial regulation is evident in all aquatic
4 ecosystems as the vertical distribution of the organisms, especially the plankton, is
5 greatly dependent on light availability. One of the most remarkable biological
6 structures related to light in aquatic systems is the establishment of phytoplankton
7 maxima at certain depths, occurring whenever both light and inorganic nutrients are
8 available in sufficient amounts. However, when thermal stratification prevents
9 continuous upwelling of nutrients, the phytoplankton maxima are usually found in
10 deeper layers, where the organisms attain a compromise between nutrient and light
11 availability (Cullen, 1982). An extreme example of this situation is the deep
12 chlorophyll maxima in oligotrophic waters, where phytoplankton can be found at
13 relatively high abundance at depths generally ranging from 50 to 100 m (Estrada et
14 al., 1993; Calbet et al., 1996; Letelier et al., 1996; Ignatiades et al., 2002; Casotti et
15 al., 2003). In these deep layers light is scarce (ca. 1% surface irradiance, Estrada et al.,
16 1993; Ignatiades et al., 2009) and nutrients, although still limiting due to the trophic
17 characteristics of these systems, occur in sufficient amounts to maintain stable
18 phytoplankton communities. On the other hand, the communities that develop in the
19 extensive mixed layers of these poorly productive ecosystems experience both severe
20 nutrient limitation and variable light levels as a result of the vertical displacement due
21 to the circulation of the water masses. In order to cope with these extreme
22 environmental challenges it has been suggested that the algae supplement their
23 nutrient acquisition with mixotrophy during periods of low dissolved nutrient
24 concentrations in the surface mixed layer (Arenovski et al., 1995). Theoretically, light
25 limitation should result in a reduced production of autotrophs, and could modify the

1 trophic characteristics of some algae (i.e., enhancement of mixotrophy); both
2 mechanisms potentially scaling up the food web. However, it is not clear how these
3 communities respond to strong changes in light intensity when vertically displaced in
4 the water column.

5 In this work we investigate the effects of an abrupt decrease in light supply on
6 natural marine plankton communities collected in light-saturated oligotrophic surface
7 waters. The light intensities were manipulated to represent the light availability at ca.
8 10 m depth in a mixed surface layer (where water was collected) and to simulate a
9 low light climate equivalent to 50-60 m water column depth, near the base of the
10 thermocline for the season (Casotti et al., 2003; Ramfos et al., 2006). To understand
11 how light modulates the response of the plankton in the oligotrophic mixed layer of
12 the Eastern Mediterranean we conducted a series of microzooplankton grazing and
13 mixotrophy incubation experiments, and analyzed the composition of the
14 microzooplankton community in a mesocosm experiment using natural water without
15 any nutrient addition. Great care was taken to keep the nutrient levels naturally low.
16 We hypothesized that i) the flow of matter through the food web would be diminished
17 by a reduction in the energy input, and ii) that the plankton community would rapidly
18 adapt to the new, darker, conditions by a fast increase of mixotrophic algae, and
19 perhaps a longer-term negative response of mixotrophic ciliates, subsequently
20 displaced by heterotrophic ciliates.

21

22 **2. Methods**

23 *2.1. Experimental set up*

1 This study was conducted as a part of a larger international experiment
2 (LightDynamix) funded by the European project MESOAQUA (<http://mesoaqua.eu/>)
3 at the Cretacosmos mesocosm facility of the Hellenic Centre for Marine Research
4 (Crete, Greece). The total LightDynamix experimental design involved 4 light
5 intensities (L1, L2, L3, and L4), duplicated in 2.8 m³ mesocosms (replicates a, b) and
6 run between 23 September and 1 October 2010. A detailed description of the overall
7 mesocosm design and plankton development in all treatments is provided in Ptacnik et
8 al. (in prep.). In short, the mesocosms were filled with water pumped from 10 m depth
9 into acid cleaned 1 m³ polyethylene (PE) containers, at a station 6 nautical miles north
10 of Heraklion, Crete (Greece). Within 2-4 h the containers were transported to land and
11 the content distributed evenly between the 8 mesocosm bags immersed in a tank with
12 water at *in situ* temperature, by gravity siphoning with acid cleaned plastic tubes.
13 Filling took altogether two days. During this time, the mesocosms were kept dark by
14 non-transparent covers. The mesocosms were made of transparent PE bags, with a
15 diameter of 1.32 m and a depth of 2.5 m, surrounded by a double layer of PE black
16 and white bags serving as an optical isolation. Optical lids for light manipulations in
17 the mesocosms were created using 4 different grades of neutral density filters (Lee
18 Filters, UK) and were installed at night after filling the mesocosms. Thus the
19 experimental manipulation started the following morning (day 0). Due to the labour
20 intensity of the incubation experiments and analysis of trophic level of the protists
21 performed in this study we were restricted to focus on two experimental (mesocosm)
22 light intensities only. We used the lightest and darkest, i.e. mesocosms L1a,b and
23 L4a,b, respectively. The grey filters covering these mesocosms attenuated
24 approximately 5 and 87% of the incoming light, resulting in average light intensities
25 of ca. 60% and 6% of incident light intensity inside each of the L1 and L4

mesocosms, respectively. Thus closely mimicking the light intensities at the sampling depth 10 m and near the base of the thermocline for the season (50-60 m, Ramfos et al., 2006). We measured vertical profiles of photosynthetically active radiation (PAR) in 0.5 m steps with a spherical quantum sensor (LI-139SA; Licor, Lincoln, NE, USA) while simultaneously measuring incident PAR (flat quantum sensor LI-190SA) above the water surface outside the mesocosm, both attached to a Licor 1500 data logger. Light inside each mesocosm was calculated as the light intensity averaged over the water column (Imix). The water in the mesocosms was gently mixed by slow bubbling through an airlift pipe as described in Ptacnik et al. (in prep.). Because ciliates and other plankton may be disturbed by bubbles or any activity creating turbulence, on the one side; but, on the other side, some type of mixing is needed to create homogenous mesocosm contents, we created a mixing system that uses a minimal amount of air, with a maximal but gentle mixing effect. It was modelled after the mixing systems that have been successfully used for over 30 years in the mesocosms at the Norwegian National Mesocosm centre, University of Bergen (see e.g. Williams and Egge, 1998; Nejstgaard et al., 2006), but down-scaled for these smaller mesocosms used here. In addition effort was made to make the bubbles big, to minimize the surface to volume ratio of the bubbles and thus the actually encounter rate between plankton and bubbles.

2.2. *Community composition and biomass*

Samples (125-250 mL) for chlorophyll *a* concentration (Chl *a*) were collected daily from each mesocosm. The water was sequentially filtered onto duplicate 10, 2, 0.65, and 0.2 µm pore-size, 47 mm diameter, polycarbonate filters (GE Water & Process

Technologies), to obtain the size-fractions: 0.2-0.65 μm , 0.65-2 μm , 2-10 μm , and > 10 μm , respectively. The filters were extracted immediately in acetone overnight at 4°C and measured on a TD Turner 700 (Turner Designs, Sunnyvale, CA), with and without acidification according to Parsons et al. (1984). We collected particulate seston daily by filtering 500 ml sample water onto GF-F filters (47 mm). The filters were dried overnight and placed in petri slides until further analyses. The amount of particulate organic carbon, phosphorus and nitrogen on the filters was analysed by x-ray fluorescence (XRF) of the specific elements using a S4 Pioner (Bruker AXS, Karlsruhe Germany) at the University of Bergen (Norway). We then calculated C:P, C:N molar ratios, and C:Chl *a* mass ratios.

Samples (50-100 ml) for analysis of the plankton community composition in the mesocosms were taken every second day, stained with primuline (Direct Yellow 59, Sigma-Aldrich Co.), fixed with 3.6% glutaraldehyde solution with 10% glycerol (final concentrations) and gently filtered onto 0.6- μm pore-size black polycarbonate membrane filters, and frozen until analysis (within 1-2 days). The method is a modification from Grebecki (1962), Hobbie et al. (1977) and Caron (1983) with the glycerol added to reduce the damage of especially small delicate protists during filtration as described in Sazhin et al. (2007). Samples were analysed by epifluorescence microscopy and cell volumes were calculated by approximation of simple geometrical 3D shapes and converted into cell carbon as described in Menden-Deuer and Lessard (2000). In addition, samples for abundance of tintinnids and larger oligotrich ciliates were collected from one replicate bag of each treatment (L1a, L4a) every other day and fixed with acid Lugol's solution (2% final concentration), subsequently settled (250 mL) for 48 h and counted on an inverted microscope at 150x magnification. In order to estimate the percentage of mixotrophic ciliates an

1 additional set of samples (250 mL) was fixed with borax-buffered formalin (final
2 concentration 2% formaldehyde), subsequently settled in the dark for 48 h and
3 counted on an inverted epifluorescence microscope at 150x magnification using blue
4 light.

5 At the end of the experiment, mesozooplankton abundance and composition in
6 the mesocosms were determined by pumping the entire volume of each mesocosm
7 through the same 48- μ m mesh submerged plankton net. Samples were preserved
8 immediately after collection in 4% borax-buffered formaldehyde solution, with the
9 exception of the L1b-sample, that was lost during the collection process. Species
10 composition analysis and the estimation of total mesozooplankton abundance were
11 carried out under a dissecting microscope in sub-samples taken with a Stempel
12 pipette. Major mesozooplankton taxa (e.g., copepods, cladocerans, doliolids,
13 appendicularians, etc.) were counted, and copepods and cladocerans were further
14 identified to the species level when possible (at least 300 copepods were counted per
15 sample). According to the latest taxonomy (Boxshall and Halsey, 2004), the order of
16 Poecilostomatoida has been transferred to the order of Cyclopoida, hence for the
17 purposes of this work have been presented together.

18 19 *2.3. Microzooplankton grazing and mixotrophy experiments*

20 Every other day we conducted standard grazing dilution experiments (Landry and
21 Hassett, 1982) in one replicate of the lightest and darkest mesocosms (L1a and L4a,
22 respectively). The water for these experiments was collected early in the morning by
23 gravity-filling of 30 L dark containers using silicon tubing. Once in the laboratory, the
24 water was gently siphoned into a 60 L bucket and carefully mixed to avoid generating

1 bubbles that could damage delicate ciliates (Gifford, 1985; Broglio et al., 2004). A
2 known volume of water was gravity-filtered through a Pall Acropak 0.8/0.2 500
3 capsule (previously flushed, including tubing, with diluted HCl and thoroughly rinsed
4 with deionised water) to obtain the dilution series at the proportions 25, 50, 75 and
5 100% of the experimental water. The dilution series was prepared in 1.3 L PC bottles,
6 which were amended with a nutrient mixture ($1\ \mu\text{M}\ \text{NH}_4\text{Cl}$, $0.07\ \mu\text{M}\ \text{Na}_2\text{HPO}_4$, and
7 $0.5\ \mu\text{M}\ \text{Na}_2\text{SiO}_3$) to guarantee that nutrients were not limiting in any treatment. To
8 assess the natural growth of algae two 100% experimental water (i.e., not diluted)
9 bottles without addition of nutrients were also prepared. Special care was taken to
10 work under dim light conditions for the darkest treatments.

11 In order to avoid adverse light effects on the mesocosms when initiating and
12 ending the incubation periods (ca. 24 h) of the grazing experiments, the bottles were
13 incubated in specially prepared mesocosm incubators equipped with an easily
14 opening-closing cover made from the same neutral density filter used for the
15 mesocosms L1 and L4, hence providing temperature ($24.3\text{-}25.7^\circ\text{C}$) and light levels
16 comparable to these mesocosms (same level attenuation; for absolute values see
17 Ptacnik et al. in prep.). The bottles were hanged from a floating wheel at 0.5 m depth,
18 which corresponded to the average light intensity of the mesocosms. The wheel
19 rotated at app. 1 r.p.m propelled by a submerged water pump. This approach assured
20 that the light conditions were homogeneous for all the bottles. However, because
21 these incubators were filled with brackish nutrient rich water from a well at the
22 HCMR, different from the nutrient poor water of the mesocosms, a denser
23 phytoplankton bloom developed at days 4-5. This had consequences for the light
24 intensity at which the experimental bottles were incubated, and is discussed below. In
25 the beginning (initial samples), as well as at the end of the incubations, we took

duplicate samples for the quantification of the different size-fractions of Chl *a*, and for the quantification of nanoflagellates (in the initial and unfertilized bottles). For the quantification of nanoflagellates in the dilution grazing experiments, 75-100 mL of glutaraldehyde preserved samples were stored at 4°C in the dark for 2h and then filtered onto 2 µm black polycarbonate filters, and stained with 4', 6'-diamidino-2-phenylindole (DAPI). We counted 200 to 500 cells under epifluorescence microscopy in each filter. The rates obtained for the cell counts of autotrophic flagellates were compared to Chl *a*-based growth rate estimates. This way we could both estimate the cellular growth rates of the phytoplankton and quantify potential artifacts in pigment based growth rates due to photoadaptation changing the cellular Chl *a* contents during the incubations.

For all experiments, we used Model I linear regressions to obtain the slope (m ; grazing mortality rate, d^{-1}) of the equation relating the fraction of undiluted water (x) and the net phytoplankton growth rates (y ; K) estimated from changes in Chl *a* concentration during incubation. Because the intercept of the equation (μ_n) would provide an overestimation of phytoplankton instantaneous growth rates (due to the addition of nutrients in the bottles), the instantaneous *in situ* growth rates (μ) in dilution grazing experiments were derived from net growth in the unfertilized bottles (K_o ; where no nutrients added) and were corrected for mortality by microzooplankton from dilution experiments when the latter was significant (Landry and Hassett, 1982).

$$\mu = K_o + m$$

When we found saturated feeding responses we used the linear relationship of the most diluted treatments to obtain the phytoplankton instantaneous growth rates with added nutrients (μ_n) according to Gallegos et al. (1989) and Dolan et al. (2000).

1 The microzooplankton grazing rate (m_n) was then calculated for the undiluted
2 nutrient-enriched bottles as

$$3 \quad m_n = \mu_n - K_n$$

4 where K_n is the phytoplankton net growth rate in the undiluted nutrient-enriched
5 bottles. This grazing rate calculated as indicated above does not completely
6 correspond to the *in situ* microzooplankton grazing rate (m). According to Moigis and
7 Gocke (2003) and Moigis (2006) the *in situ* microzooplankton grazing rate (i.e., in the
8 undiluted bottles without added nutrients) should be calculated as

$$9 \quad m = m_n * K_o * (\exp(K_n * t) - 1) / K_n * (\exp(K_o * t) - 1)$$

10 where t is the incubation time.

11 In parallel we assessed the amount of herbivory of mixotrophic algae (large
12 flagellates and dinoflagellates) by measuring the uptake of live fluorescently labeled
13 algae (LFLA) during 24 h *in situ* grazing experiments. Aiming at a better
14 representation of the nanoflagellate size spectrum of the mesocosm environment two
15 different types of algae were used to produce the LFLA stocks: *Isocrhysis galbana* (4-
16 6 μ m) and *Tetraselmis* sp. (8-10 μ m). Algal cultures grown exponentially in f/2
17 medium at 18°C in a 12:12 h light: dark cycle were labeled with the live fluorescent
18 vital stain CellTracker® Blue CMAC (7-amino 4-chloromethylcoumarin, Molecular
19 Probes Inc.) according to Teegarden (1999). The latter vital stain makes the cells
20 fluoresce bright blue when excited by UV light at 354 nm.

21 The incubations were conducted in 650 ml polycarbonate bottles, filled with
22 water from the mesocosm L1a and L4a, and lasted for ca. 24h. Two sets of triplicate
23 bottles were prepared, one set without nutrient additions and one set amended with
24 nutrients as in the dilution experiments. These bottles were used for an additional

1 estimate of the effect of nutrient limitation. Each experimental bottle was inoculated
2 with a mixture of fluorescent prey and was subsequently gently mixed. We tried to
3 keep total LFLA concentration inside the bottles below 25% of the abundance at
4 which similarly sized flagellates are typically found in oligotrophic areas of the
5 Mediterranean Sea (1.3×10^2 - 6.0×10^3 cells ml^{-1}) (Christaki et al., 1999, 2001).
6 Subsamples were always taken prior incubations to determine exact initial
7 concentration of LFLA in the bottles, which at times was more than expected (see
8 discussion). The incubations took place in the already described incubators, together
9 with the dilution experiment bottles. At the end of each incubation 75-100 ml samples
10 were fixed with glutaraldehyde (final concentration 1%) and processed as above
11 (dilution grazing experiments), with the exception that no DAPI stain was used.
12 Protozoan cells with LFLA prey inside were enumerated under an Olympus BX 40
13 inverted epifluorescence microscope at a magnification of 400 x. Mixotrophic activity
14 was calculated as the percentage of mixotrophs $>10 \mu\text{m}$ and $<10 \mu\text{m}$ found with
15 labelled prey inside.

16
17

18 **3. Results**

19 *3.1. Community composition and biomass*

20 In Fig. 1 we show the time course of the total phytoplankton biomass (measured as
21 Chl *a*) along the 8 days of experiment in mesocosms experiencing high (L1) and low
22 (L4) light treatments, as well as the contribution of the different size fractions to the
23 standing stock of Chl *a*. Both treatments and parallels showed an increase in Chl *a* the
24 first 2-3 days of experiment, from 0.08 to ca. $0.2 \mu\text{g Chl } a \text{ L}^{-1}$. While values in both
25 the L4 mesocosms were approximately $0.2 \mu\text{g Chl } a \text{ L}^{-1}$ throughout the experiment,

1 the L1 mesocosms showed a rapid decline during day 4 and 5 and ended (on day 8) at
2 concentrations near half the maximum attained. The Chl *a* fractions 0.6-2 μm and 2-
3 10 μm were of similar proportion and together made up 75-90% of the total Chl *a* at
4 all times (Fig. 1). Except for a slight increase in the contribution of Chl *a* > 10 μm
5 during the experiment in all treatments, there were no clear changes in size groups
6 over time in the mesocosms.

7 The development of the autotrophic protist biomass determined by
8 epifluorescence microscopy (Fig. 2) resembled the pattern of Chl *a* development in
9 the L1 mesocosms. However, in the L4 mesocosms the autotroph biomass showed a
10 development different from Chl *a*, with lower biomass values and less clear peaks
11 ($0.94 \pm 0.53 \text{ SE } \mu\text{gC L}^{-1}$, Fig. 2). The total biomass peak of heterotrophic eukaryotes in
12 the L1 mesocosms ($1.7 \pm 0.12 \text{ SE } \mu\text{gC L}^{-1}$) was more than twofold higher than in L4
13 ($0.75 \pm 0.07 \text{ SE } \mu\text{gC L}^{-1}$), although the temporal development was similar.

14 In all mesocosms, mixotrophic ciliates and autotrophic dinoflagellates
15 increased their biomass towards the end of the study, becoming significant
16 components to the community, otherwise dominated by autotrophic flagellates (Fig.
17 2). In mesocosms L1 heterotrophic flagellates and ciliates dominated the protozoan
18 community-biomass the second and sixth day of the experiment, respectively. In L4
19 mesocosms, however, the heterotrophic community remained quite stable with the
20 only exception of a peak of ciliates on day 6 (Fig. 2). The Lugol preserved samples
21 showed a progressive increase of ciliates in both mesocosm treatments until day 6 and
22 a slight decrease the last day (Fig. 3). The contribution of mixotrophs to the total
23 abundance of ciliates consistently decreased in L4a from 50% to 30% and was quite
24 variable in L1a (Fig. 3).

1 The mesozooplankton biomass community at the end of the experiment was
2 higher in the L1 treatment than in L4 (Fig. 4). The mesozooplankton community was
3 numerically dominated by copepods in all measured treatments with calanoida
4 (mostly *Clausocalanus* spp. and *Calocalanus* spp.) showing ca. double abundance in
5 L1a, being the most abundant group in this mesocosm. Cyclopoida (mostly *Oithona*
6 spp.) abundance was similar at both light intensities. Other important components,
7 although at much lower abundances were appendicularia and cladocera (Fig. 4).

8 The elemental ratios of C, N, and P, as well as the C:Chl *a* mass ratio are
9 presented in Fig. 5. Light did not significantly affect any of the ratios ($p > 0.05$;
10 grouped ANOVA with repeated measures), however some differences are evident in
11 the proportion of Chl *a* per unit of carbon since day 4 (Fig. 5A), and in the C:N at
12 days 1 and 2 (Fig. 5B). C:P ratios followed a similar pattern in both treatments along
13 the experiment, with peaks at days 3 and 4 for L4 and L1, respectively and an increase
14 of the quotient last day in all mesocosms (Fig. 5C).

16 3.2. Microzooplankton dilution grazing experiments

17 We conducted four dilution-grazing experiments (Landry and Hassett, 1982) in each
18 replicate "a" of the mesocosms L1 and L4. Overall, all (Chl *a*-based) phytoplankton
19 growth and mortality rates observed for total and size-fractionated Chl *a* were quite
20 high, with many saturation-feeding responses at the end of the experiment, *sensu*
21 Gallegos (1989) (Fig. 6; Table 1). During day 1 we observed consistent positive
22 slopes for total phytoplankton and all size-fractions in the low light mesocosm (L4a,
23 Fig. 6). In general, mortality rates (m) on total Chl *a* were similar in both treatments,
24 but the instantaneous growth rates (μ) were higher in L1a mesocosm. While the

1 smallest size-fractions of the phytoplankton (Chl *a*) were more heavily grazed than
2 the largest sizes in L1a, the microzooplankton community grazing severely impacted
3 the 0.65-2 μm size-fraction, but not the 0.2-0.65 μm in L4a.

4 In order to control for artifacts from potential photoadaptation processes
5 during the incubations we estimated net growth rates of the unfertilized bottles by
6 counting nanoflagellates on 2 μm pore-size filters. These data are compared to the net
7 growth rates in the same bottles for the closest Chl *a* size-fraction (2-10 μm) in Fig. 7.
8 No significant differences were observed for the first 3 experiments in L1a, and for
9 the first 2 in L4a ($p < 0.05$). However, after these dates the differences became
10 significant ($p > 0.05$). Therefore, our data on instantaneous growth rate (μ) in these
11 experiments based on Chl *a* appear to be overestimated since day 5 on.

12 Dilution experiments were also used to assess phytoplankton nutrient
13 limitation during the incubations, by comparing the net phytoplankton growth rates
14 between the fertilized and unfertilized undiluted bottles (Fig. 8). Nutrients appeared to
15 be limiting in all the L1-experiments, except the one conducted the first day (t-test; p
16 < 0.05). For L4a mesocosm the results showed no clear pattern. Differences in this
17 treatment were significant (although not always indicating higher growth in nutrient
18 amended bottles) in all experiments, except the ones at day 5 (t-test; $p < 0.05$).

19 By comparing the biomass of eukaryotic autotrophs (Fig. 2) with the grazing
20 rates from microzooplankton (Table 1) we attempted to estimate the autotrophic
21 carbon flux through microzooplankton. Due to the 1-day decoupling between grazing
22 and biomass estimates, we calculated the carbon flux by multiplying the averaged
23 biomass of autotrophs between two consecutive sampling dates (C_{average}) by the
24 corresponding grazing coefficient (m), as estimated from the dilution experiments.

1 These values can be converted into total flux by including the total volume (V) of the
2 mesocosm in the equation.

$$3 \text{ Carbon flux} = m * C_{average} * V$$

4 The results for both mesocosms are presented in Fig. 9. As a whole, the carbon flux
5 through microzooplankton was ca. 50% higher in the high light mesocosms than in
6 the low light treatments.

7

8 *3.3. Identification of major microbial herbivores and mixotrophy*

9 As previously described we identified the major microbial herbivores using live
10 fluorescently-labelled cultivated algae. The percentage of pigmented protists (ANF)
11 with labelled prey inside was low (Table 2), ranging from 3 to 6%. However, the
12 contribution of mixotrophic organisms (flagellates and dinoflagellates) to the total
13 number of grazers on phytoplankton (considering the heterotrophs with labelled prey
14 inside) was considerable (ca. 50%). We tested for significant differences (t-test) in the
15 percentages of heterotrophs vs mixotrophs for the variables light intensity and nutrient
16 addition. We only found a significant difference at the last day, when we found a
17 moderate increase of 1.6 times in the percentage of mixotrophic flagellates with
18 labelled prey in the L4 bottles compared to L1 ones (light effect), and this difference
19 was only apparent in the unfertilized treatments.

20

21 **4. Discussion**

22 *4.1. Responses of the microbial community to the light conditions*

1 Overall, one order of magnitude reduction in light in the L4 mesocosms compared to
2 the L1 ones resulted in lower peak biomass by a factor of 0.6 for autotrophs, 0.4 for
3 protozoans ($p < 0.05$; Two-way grouped ANOVA with repetition), and 0.6 for
4 mesozooplankton, and a change in the composition of the community at all trophic
5 levels. Similar reductions in the magnitudes of phytoplankton, micro- and
6 mesozooplankton bloom peaks were observed by experimentally increased
7 stratification depth (reduced light supply) in freshwater mesocosms (Berger et al.,
8 2010) indicating that light is an important driver of primary production scaling up to
9 higher trophic levels.

10 While the water used to initiate the mesocosms showed typical low Chl *a*
11 concentrations for the area (Psarra et al., 2000; Ignatiades et al., 2002), the Chl *a*
12 concentration increased faster in the darker mesocosms (L4) than in the more
13 illuminated ones (L1). Although this could suggest e.g. photoinhibition (Neale, 1987)
14 in L1 or increased growth rate of algae in L4 (for unknown reasons), the data reveal
15 that the rapid Chl *a* increase was likely due to an increase in pigments per cell, and
16 not to an increase in autotroph cell numbers in L4 (Fig. 2). This can be interpreted as
17 a photoadaptation in response to the dim light conditions in L4 (Falkowski, 1980;
18 Cullen and Lewis, 1988; Therriault et al., 1990). This response was not initially
19 evident in the community C:Chl *a* ratio, which was similar for both treatments during
20 the first days of the experiment (Fig. 6). We have to take into account that this ratio
21 includes not only autotrophs, but heterotrophic and detrital carbon as well, therefore
22 not properly describing the variations in the Chl *a* contents per cell, but rather
23 mirroring the evolution of heterotrophs and autotrophs in the mesocosms. Higher
24 proportion of heterotrophs by the end of the experiment in L1 resulted in a higher
25 C:Chl *a* ratio in this treatment. It is more difficult to interpret the variations in C:N

1 and C:P ratios; neither of them showing any clear difference between treatments.
2 Only, L1 mesocosms showed peaks of higher C:N, indicating an impoverishment of
3 the nutritional quality of the community, as theoretically expected, but the differences
4 were not significant.

5 The lower autotrophic biomass in the low light (L4) mesocosms could be
6 explained either by lower phytoplankton instantaneous growth rates or higher grazing
7 pressure than in the L1's, or both. The dilution experiments should give us both these
8 rates. Although the dilution experiments for L4a at day 1 cannot be interpreted in a
9 meaningful manner due to the positive slopes (possibly due to trophic cascades during
10 incubations as described by Calbet et al., 2011a), we can compare the rates for day 3,
11 the closest to the peak of phytoplankton biomass in all mesocosms where C:Chl *a*
12 ratios did not differ among treatments. At day 3 both mortality and growth rates of
13 total phytoplankton were slightly lower for the L1a mesocosm, indicating a slower
14 circulation of energy through the food web compared to the lower light mesocosms.
15 In L1a, both grazing and growth were similar, indicating that all the potential
16 production was consumed within the day. However, for L4a the measured mortality
17 rates exceeded the potential production. If this was the case, we should expect a
18 reduction of phytoplankton biomass in day 4 for this mesocosm. However, this was
19 not observed in the development of the Chl *a*, or in the cellular carbon. Actually, the
20 measured mortality rates exceeded the instantaneous phytoplankton growth rates in
21 several cases, especially in L4. These discrepancies have to be understood in the light
22 of other components of the community that were not considered in the dilution
23 grazing experiments, i.e. mesozooplankton. We sampled water for the dilution grazing
24 experiments from the upper part of the mesocosms, when the sun was up and the
25 mesozooplankton may have remained at the bottom of the bags. At night these

1 mesozooplankton are expected to migrate up the mesocosm and selectively graze on
2 microzooplankton (Calbet and Saiz, 2005; Saiz and Calbet, 2011), thus releasing
3 phytoplankton from their most important grazers (Calbet and Landry, 2004; Calbet et
4 al., 2008a), to a larger extent than in the bottles (Nejstgaard et al., 2001). Because the
5 mesozooplankton community in the mesocosms were rich in carnivorous species,
6 such as *Oithona* spp., they could have significantly predated on the microzooplankton,
7 and thus shaped the abundance of phytoplankton through trophic cascades.
8 Consequently, given our dilution bottles did not contain representative amounts of
9 mesozooplankton, we will refer to the estimated microzooplankton community rates
10 as “potential”, and likely on the higher side.

11 Microbial herbivores were exerting a very high potential grazing pressure on
12 most of the Chl *a* size-fractions considered, except the 0.2-0.6 μ m size-fraction in
13 L4a. We should expect to mostly find prokaryotes in this size-fraction, possibly
14 sheltered from heavy predation through trophic cascades in the food web. A similar
15 situation has been described in other oligotrophic areas, where trophic cascade effects
16 masked the grazing on the small phytoplankton (Nejstgaard et al., 1997; Calbet et al.,
17 2001; Calbet et al., 2008b). Surprisingly, we found saturated-feeding responses
18 (Gallegos, 1989) at the end of the experiment. While the concentration of prey
19 increased initially and saturated feeding rates of the microzooplankton could have
20 been expected at that point, we would not anticipate this situation for oligotrophic
21 areas. Although our data are not unique in showing saturated-feeding in dilution
22 grazing experiments of oligotrophic sites (Worden and Binder, 2003; Berninger and
23 Wickham, 2005; Calbet et al., 2008b), given the little biomass of prey attained by the
24 end of the experiment, this fact would imply microzooplankters from oligotrophic
25 sites present a faster-saturating feeding curves than those from productive ones. We

1 are not aware of studies comparing feeding kinetics of protozoans from different
2 ecosystems that can corroborate our results, and thus remains as a hypothesis to be
3 tested.

4

5 *4.2. Microbial community composition and mixotrophy*

6 We have so far discussed how the phytoplankton community as a whole progressed in
7 the different mesocosms. We will now discuss the development of the different
8 groups during the experiment. A conspicuous result was the gradual increase of
9 pigmented dinoflagellates and ciliates (Fig. 2). Both groups are larger in size than the
10 rest of components of the microbial community, and both have the potential to act as
11 mixotrophs (Stoecker, 1999; Johnson, 2011).

12 At organism level, nutrients and light intensity may trigger different feeding
13 responses in obligate mixotrophs, phagotrophic algae, and photosynthetic protozoa
14 (Bird and Kalff, 1989; Skovgaard, 1996; Jones et al., 1997; Legrand et al., 1998;
15 Stoecker, 1998; Stickney et al., 2000. However, at community level, continuous
16 darkness should favour heterotrophic metabolism because their higher efficiencies of
17 feeding and incorporation of organic matter into the cell (Tittel et al., 2003). In our
18 experiments, providing that light and nutrients were limiting, in the L4 treatment we
19 could expect a fast response of existing mixotrophic algae (organism-level response),
20 to be gradually replaced by heterotrophic grazers (community-level response).

21 However, at relatively short time intervals (ca. 24h) we did not detect any significant
22 variation on the mixotrophic contribution to overall grazing on algae (unfortunately,
23 we do not have data on mixotrophic grazing on prokaryotes). At longer time intervals
24 heterotrophic microbial biomass was below that of pigmented organisms (quotient

1 biomass heterotrophs/autotrophs < 1). This indicates dominance of autotrophic
2 processes over the microbial food web. Therefore, our premise was not met.

3 We should be aware, however, that the light intensities chosen in our study
4 were intended to simulate the water circulation in the mixed layer of an oligotrophic
5 highly illuminated sea. Even though in L4 we drastically reduced the irradiance by
6 94% compared to the surface irradiance, the light available in this treatment was still
7 above the light-saturated intensities necessary to develop deep-chlorophyll (90-100
8 $\mu\text{E m}^{-2} \text{s}^{-1}$; Harris, 1986; Ignatiades et al., 2002). Therefore, the community responses
9 of autotrophs and heterotrophs for these light intensities should not be as evident as
10 the ones expected for more limiting irradiance conditions, or even complete darkness.

11 Mixotrophs in L1 treatments should remain unaffected, given the similarity of
12 the experimental conditions to *in situ*, although we cannot disregard a gradual
13 increase in mixotrophy due to nutrient limitation during the experiment. Little is
14 known about the time frame of this process in natural systems, and we did not observe
15 any clear change in mixotrophy of flagellates in the L1 treatments, and mixotrophic
16 ciliates showed either an erratic pattern of abundance or contributed proportionally
17 less to the total abundance of ciliates.

18 Regarding the technique used to estimate mixotrophy, although we intended to
19 keep the LFLA concentration below 25% of the natural abundance of similarly-sized
20 flagellates, the LFLA were added without previous knowledge of the concentrations
21 in the experimental water. Consequently, the contribution of labelled flagellates to the
22 bulk was generally 25-50%, with one very high value of 83% (L1a, day one). These
23 data may thus not estimate ingestion rates or total carbon flux accurately.

24 Nevertheless, in accordance with Li et al. (1996, and references therein) we still argue

they show the percentage of mixotrophs within autotrophic algae, and contribution of mixotrophic grazing to total grazing, and are overall more natural than the alternative approach based on dead labelled algae, or inert microspheres.

4.3. Possible mixotrophy-related artifacts in our experimental set up

Mixotrophs, as any other grazer, may show positive slopes during dilution assays, due to higher prey encounter rates and thus higher feeding/growth rates in the least diluted treatments (Dolan et al., 2000). It can be argued this may confound Chl *a* based estimates of phytoplankton growth rates by flattening the slope of the Chl *a*-based regression and artificially reducing the estimates of microzooplankton grazing (Landry, 1995; Schlüter, 1998; Calbet et al., 2008b). Unfortunately, the impact of this process in a standard dilution experiment is quite difficult to predict because it will depend on the biomass of mixotrophs, and their growth and grazing rates. In our experiments, the contribution of mixotrophs to total number of grazers was on average 50%, which is similar to other values found for other areas (Hall et al., 1993; Safi and Hall, 1999; Unrein et al., 2007). However, this does not imply that these values are equivalent to half the microzooplankton grazing impact. It is well known that mixotrophs are less efficient in consuming prey than the heterotrophs (Stoecker, 1998; Jeong et al., 2010). For instance, for Mediterranean coastal waters, maximum specific ingestion rates of autotrophic flagellates of 5-20 μm were of the order of 3.6 bacteria h^{-1} , whereas for heterotrophic flagellates were 15.4 bacteria h^{-1} , i.e. a factor of 4 (Unrein et al., 2007). For mixotrophic dinoflagellates specific ingestion rates of prey were estimated to be 4 times lower than heterotrophic ones (Calbet et al., 2011b). If we assume a similar 4 times lower biomass specific ingestion efficiency by the

1 mixotrophs (50% of the active grazers according our data), compared to the
2 heterotroph microzooplankton in our experiments, the mixotrophs would only account
3 for ca. 12.5% of the total estimated ingestion here. Converting this relative portion of
4 microzooplankton community grazing efficiency into autotrophic biomass consumed,
5 based on a gross growth efficiency of 30%, the 12.5% of total ingestion translates into
6 3.75% of expressed mixotroph production. This is a relatively modest production that
7 would not be expected to alter the accuracy of the rates determined by the dilution
8 experiments, or to explain the positive curve in the L4a, day1 experiment (Fig. 6).

9

10 *4.4. Scaling up the results to the food web dynamics*

11 Theoretically, a limitation in light intensity should result in a reduction in the attained
12 biomass of autotrophs (Huisman, 1999; Diehl, 2002), which should have negative
13 consequences for consumers, and further on higher trophic levels. This theory was
14 practically demonstrated in comparative lake studies (e.g., Berger et al., 2006).
15 However, it has been suggested for nutrient-limited lakes that reduced light increases
16 herbivore production due to a stoichiometric effect of the balance between light and
17 nutrients (Urabe and Sterner, 1996; Urabe et al., 2002). The mechanism proposed is
18 based on an increase of the nutrient contents per alga relative to carbon within alga for
19 light-limited organisms, and the opposite for well-illuminated ones (Diehl et al.,
20 2002). Therefore, improved food quality would enhance production of grazers at
21 lower light intensities (Diehl, 2007). Even though this theory could apply also to
22 marine oligotrophic systems, the differences between lake and marine food webs may
23 make the effect less relevant. In lakes, at least in those where the hypothesis was
24 tested, primary producers were directly consumed by crustacean grazers

1 (cladocerans); therefore, very susceptible to variations in nutrient composition of the
2 prey (Urabe et al., 2002). In marine oligotrophic systems, on the other hand, the
3 complexity of the food web makes it more likely that the effect of nutrient imbalances
4 is buffered through homeostasis in intermediate trophic levels. Moreover, the
5 relevance of mixotrophy (Stoecker, 1999; Stoecker et al., 2009), feeding selection on
6 nutritious prey (Cowles et al., 1988; Saiz and Calbet, 2011), nutrient tunneling effects
7 (Thingstad and Cuevas, 2010), and the ability of many protozoans to upgrade food
8 quality (Klein Breteler et al., 1999; Broglio et al., 2003; Veloza et al., 2006) would
9 make any nutrient deficiency of algae grown under high light conditions dampened
10 within the food web.

11 In our experiments the decrease in light intensity resulted in a reduction on the
12 final biomass of both autotrophs and heterotrophs (including metazoans) by a factor
13 of ca. 0.5. This contradicts the theory proposed of a lower zooplankton production in
14 highly illuminated lakes (Urabe et al., 1996; Urabe et al., 2002). Even though we
15 could not detect a clear nutritional response of the different light treatments, the C:N
16 of the more illuminated mesocosms tended to be higher than the one of the darkest.
17 Additionally, since the fourth day of the experiment the data indicate a possible
18 decrease in nutritional quality of phytoplankton in the most illuminated mesocosms
19 (higher C:Chl *a* ratios), but this likely impoverishment of the diet was not reflected in
20 a lower final abundance of grazers (both micro- and mesozooplankton). Therefore,
21 any effect of poor nutritional value of the prey might have been dampened before
22 reaching higher order consumers. This could be due to a variety of mechanisms (e.g.,
23 intermediary trophic links, food upgrading, mixotrophy, nutrient tunnelling), many of
24 them not examined in this study, but will have to be resolved in future studies in order
25 to better understand the effect of light availability in oligotrophic aquatic food webs.

1

2 **Acknowledgements**

3 This work was funded through projects CTM2009-08783 from the Spanish Ministry
4 of Science and Innovation assigned to A.C. and projects LightDynamix, LiMic and
5 LightCopFed from MESOAQUA (EU FP7-INFRA-2008-1, 228224), assigned to R.P.
6 (LightDynamix), A.C., R.A.M., A.S., A.G. (LiMic) and S.I. (LightCopFed). We are
7 indebted to G. Piperakis for his inspired technical assistance throughout the
8 experiment, to S. Zivanovic, E. Dafnomili and Dr. M. Tsapakis for assistance with
9 chemical analyses, and to I. Santi, P.A. Gazihan-Akoglu, and P. Dimitriou for
10 conducting the Chl *a* analyses. The captain and the crew of the R/V Philia are thanked
11 for their assistance during the transportation of water to the mesocosms. We also wish
12 to thank Panagiotis Vavilis, Dimitris Apostolakis, Zacharias Grigorakis, Manolis
13 Ntoumas and Manos Pettas for their immediate and willing response to our
14 continuous requests for help with the zodiac. Finally, we want to thank the rest of
15 participants of the mesocosm experiment: K. Altoja, P. Laas, S. Moorthi, R.
16 Ptacnikova, S.J. Royer, S. Schabhueetl, M. Striebel, and T. Tanaka, for their help and
17 support.

18

1 **References**

- 2 Arenovski, A.L., Lim, E.L., Caron, D.A., 1995. Mixotrophic nanoplankton in
3 oligotrophic surface waters of the Sargasso Sea may employ phagotrophy to
4 obtain major nutrients. *J. Plankton Res.* 17(4), 801-820.
- 5 Berger, S.A., Diehl, S., Kunz, T., Albrecht, D., Oucible, M.A., Ritzer, S., 2006. Light
6 supply, plankton biomass, and seston stoichiometry in a gradient of lake
7 mixing depths. *Limnol. Oceanogr.*, 51(4), 1898–1905.
- 8 Berger, S.A., Diehl, S., Stibor, H., Trommer, G., Ruhenstroth, M., 2010. Water
9 temperature and stratification depth independently shift cardinal events during
10 plankton spring succession. *Global Change Biol.* 16, 1954-1965.
- 11 Berninger, U.-G., Wickham, S.A., 2005. Response of the microbial food web to
12 manipulation of nutrients and grazers in the oligotrophic Gulf of Aqaba and
13 northern Red Sea. *Mar. Biol.* 147(4), 1017-1032.
- 14 Bird, D., Kalff, J., 1989. Phagotrophic sustenance of a metalimnetic phytoplankton
15 peak. *Limnol. Oceanogr.* 34, 155-162.
- 16 Broglia, E., Jónasdóttir, S.H., Calbet, A., Jakobsen, H.H., Saiz, E., 2003. Effect of
17 heterotrophic versus autotrophic food on feeding and reproduction of the
18 calanoid copepod *Acartia tonsa*: relationship with prey fatty acid composition.
19 *Aquat. Microb. Ecol.* 31(3), 267-278.
- 20 Boxshall, G.A., Halsey, S.H., 2004. An introduction to copepod diversity vol. 1, pp.1-
21 421, vol. 2, pp. 423- 966, The Ray Society, London.
- 22 Broglia, E., Saiz, E., Calbet, A., Trepát, I., Alcaraz, M., 2004. Trophic impact and
23 prey selection by crustacean zooplankton on the microbial communities of an
24 oligotrophic coastal area (NW Mediterranean Sea). *Aquat. Microb. Ecol.*
25 35(1), 65-78.
- 26 Calbet, A., 2008. The trophic roles of microzooplankton in marine systems. *Ices J.*
27 *Mar. Scien.* 65(3), 325-331.
- 28 Calbet, A., Alcaraz, M., Saiz, E., Estrada, M., Trepát, I., 1996. Planktonic herbivorous
29 food webs in the Catalan Sea (NW Mediterranean): Temporal variability and

- 1 comparison of indices of phyto-zooplankton coupling based on state variables
2 and rate processes. J. Plankton Res. 18(12), 2329-2347.
- 3 Calbet, A., Bertos, M., Fuentes-Grünewald, C., Alacid, E., Figueroa, R., Renom, B.,
4 Garcés, E., 2011b. Intraspecific variability in *Karlodinium veneficum*: Growth
5 rates, mixotrophy, and lipid composition. Harmful Algae 10(6), 654-667.
- 6 Calbet, A., Landry, M.R., 2004. Phytoplankton growth, microzooplankton grazing,
7 and carbon cycling in marine systems. Limnol. Oceanogr. 49, 51-57.
- 8 Calbet, A., Saiz, E., 2005. The ciliate-copepod link in marine ecosystems. Aquat.
9 Microb. Ecol. 38(2), 157-167.
- 10 Calbet, A., Landry, M.R., Nunnery, S., 2001. Bacteria-flagellate interactions in the
11 microbial food web of the oligotrophic subtropical North Pacific. Aquat.
12 Microb. Ecol. 23(3), 283-292.
- 13 Calbet, A., Saiz, E., Almeda, R., Movilla, J.I., Alcaraz, M., 2011a. Low
14 microzooplankton grazing rates in the Arctic Ocean during a *Phaeocystis*
15 *pouchetii* bloom (Summer 2007): fact or artifact of the dilution technique? J.
16 Plank. Res. 33(5), 687-701.
- 17 Calbet, A., Trepát, I., Almeda, R., Saló, V., Saiz, E., Movilla, J.I., Alcaraz, M., Yebra,
18 L., Simó, R., 2008. Impacts of micro- and nanograzers on phytoplankton
19 assessed by standard and size-fractionated dilutions. Aquat. Microb. Ecol 50,
20 154-156.
- 21 Caron, D.A., 1983. Technique for enumeration of heterotrophic nanoplankton using
22 epifluorescence microscopy, and comparison with other procedures. Appl.
23 Environ. Microbiol. 46, 491-498.
- 24 Casotti, R., 2003. Composition and dynamics of the phytoplankton of the Ionian Sea
25 (eastern Mediterranean). J. Geophys. Res. 108(C9).
- 26 Christaki, U., Giannakourou, A., Wambeke, F.V., Grégori, G., 2001. Nanoflagellate
27 predation on auto- and heterotrophic picoplankton in the oligotrophic
28 Mediterranean Sea. J. Plank. Res. 23(2), 1297-1310.

- 1 Christaki, U., Giannakouron, A., Van Wambeke, F., Grégori, G., 2001. Nanoflagellate
2 predation on auto- and heterotrophic picoplankton in the oligotrophic
3 Mediterranean Sea. *J. Plankton Res.* 23(11), 1297-1310.
- 4 Cowles, T.J., Olson, R.J., Chisholm, S.W., 1988. Food selection by copepods:
5 discrimination on the basis of food quality. *Mar. Biol.* 100(1), 41-49.
- 6 Cullen, J.J., 1982. The deep chlorophyll maximum: comparing vertical profiles of
7 chlorophyll *a*. *Can. J. Fish. Aquat. Sci.* 39, 791-803.
- 8 Cullen, J.J., Lewis, M.R., 1988. The kinetics of algal photoadaptation in the context of
9 vertical mixing. *J. Plankton Res.* 10, 1039-1063.
- 10 Diehl, S. (2007) Paradoxes of enrichment: Effects of increased light versus nutrient
11 supply on pelagic producer-grazer systems. *Am. Nat.* 169, 173-191.
- 12 Diehl, S., Berger, S., Ptacnik, R., Wild, A., 2002. Phytoplankton, light, and nutrients
13 in a gradient of mixing depths: Field experiments. *Ecology* 83, 399-411.
- 14 Dolan, J.R., Gallegos, C.L., Moigis, A., 2000. Dilution effects on microzooplankton
15 in dilution grazing experiments. *Mar. Ecol. Prog. Ser.* 200, 127-139.
- 16 Estrada, M., Marrasé, C., Latasa, M., Berdalet, E., Delgado, M., Riera, T., 1993.
17 Variability of deep chlorophyll maximum characteristics in the Northwestern
18 Mediterranean. *Mar. Ecol. Prog. Ser.* 92, 289-300.
- 19 Falkowski, P.G., 1980. Light-shade adaptation in marine phytoplankton. In:
20 Falkowski, P. G. (ed.) *Primary productivity in the sea*. Plenum, New York, p.
21 99-111.
- 22 Gallegos, C.L., 1989. Microzooplankton grazing on phytoplankton in the Rhode River,
23 Maryland: nonlinear feeding kinetics. *Mar. Ecol. Prog. Ser.* 57, 23-33.
- 24 Gifford, D.J., 1985. Laboratory culture of marine planktonic oligotrichs (Ciliophora,
25 Oligotrichida). *Mar. Ecol. Prog. Ser.* 23, 257-267.
- 26 Grebecki, A. 1962: Adsorption des fluorochromes par le cystome des Cilies. *Bull.*
27 *Acad. Pol. Sci.* 10, 483-485.

- 1 Hall, J.A., Barrett, D.P., James, M.R., 1993. The importance of phytoflagellate,
2 heterotrophic flagellate and ciliate grazing on bacteria and picophytoplankton
3 sized prey in a coastal marine environment. *J. Plankton Res.* 15, 1075.
- 4 Harris, G.P., 1986. *Phytoplankton Ecology*. Chapman & Hall, London.
- 5 Hobbie, J. E., Daley, R. J., Jasper, S., 1977. Use of nucle- opore filters for counting
6 bacteria by fluorescence micro- scopy. *Appl. Environ. Microbiol.* 33, 1225–
7 1228.
- 8 Huisman, J., 1999. Population Dynamics of Light-Limited Phytoplankton: Microcosm
9 Experiments. *Ecology* 80, 202-210.
- 10 Ignatiades, L., Psarra, S., Zervakis, V., Pagou, K., Souvermezoglou, E.,
11 Assimakopoulou, G., Gotsis-Skretas, O., 2002. Phytoplankton size-based
12 dynamics in the Aegean Sea (Eastern Mediterranean). *J. Mar. Sys.* 36, 11-28.
- 13 Jeong, H.J., Yoo, Y.D., Kim, J.S., Seong, K.A., Kang, N.S., Kim, T.H., 2010.
14 Growth, feeding and ecological roles of the mixotrophic and heterotrophic
15 dinoflagellates in marine planktonic food webs. *Ocean Science Journal* 45(2),
16 65-91.
- 17 Johnson, M.D., 2011. The acquisition of phototrophy: adaptive strategies of hosting
18 endosymbionts and organelles. *Photosynth. Res.* 107(1), 117-132.
- 19 Jones, H., 1997. A classification of mixotrophic protists based on their behaviour.
20 *Freshwater Biol.* 37, 35-43.
- 21 Klein Breteler, W.C.M., Schogt, N., Baas, M., Schouten, S., Kraay, G.W., 1999.
22 Trophic upgrading of food quality by protozoans enhancing copepod growth:
23 role of essential lipids. *Mar. Biol.* 135(1), 191-198.
- 24 Landry, M.R., Hassett, R.P., 1982. Estimating the grazing impact of marine micro-
25 zooplankton. *Mar. Biol.* 67, 283-288.
- 26 Landry, M.R., Constantinou, J., Kirshtein, J., 1995. Microzooplankton grazing in the
27 central equatorial Pacific during February and August, 1992. *Deep-Sea Res. II*
28 42(2-3), 657-671.

- 1 Legrand, C., Granéli, E., Carlsson, P., 1998. Induced phagotrophy in the
2 photosynthetic dinoflagellate *Heterocapsa triquetra*. *Aquat. Microb. Ecol.* 15,
3 65-75.
- 4 Letelier, R.M., Dore, J.E., Winn, C.D., Karl, D.M., 1996. Seasonal and interannual
5 variations in photosynthetic carbon assimilation at Station ALOHA. *Deep-Sea*
6 *Res. II* 43(2-3), 467-490.
- 7 Li, A.S., Stoecker, D.K., Coats, D.W., Adam, E.J., 1996. Ingestion of fluorescently
8 labeled and phycoerythrin-containing prey by mixotrophic dinoflagellates.
9 *Aquat. Microb. Ecol.* 10, 139-147.
- 10 Menden-Deuer, S., Lessard, E.J., 2000. Carbon to volume relationships for
11 dinoflagellates, diatoms, and other protist plankton. *Limnol. Oceanogr.* 45(3),
12 569-579.
- 13 Moigis, A.G., 2006. The clearance rate of microzooplankton as the key element for
14 describing estimated non-linear dilution plots demonstrated by a model. *Mar.*
15 *Biol.* 149, 743-762.
- 16 Moigis, A.G., Gocke, K., 2003. Primary production of phytoplankton estimated by
17 means of the dilution method in coastal waters. *J. Plank. Res.* 25, 1291-1300.
- 18 Neale, P.J., 1987. Algal photoinhibition and photosynthesis in the aquatic
19 environment. In Kyle, D.J., Osmond, C.B., Arntzen, C.J. (eds),
20 *Photoinhibition*. Elsevier, Amsterdam, pp. 35–65.
- 21 Nejstgaard, J.C., Frischer, M.E., Verity, P.G., Anderson, J.T., Jacobsen, A., Zirbel,
22 M.J., Larsen, A., Martínez-Martínez, J., Sazhin, A.F., Walters, T., Bronk,
23 D.A., Whipple, S.J., Borett, S.R., Patten, B.C., Long, J.D. 2006. Plankton
24 development and trophic transfer in seawater enclosures with nutrients and
25 *Phaeocystis pouchetii* added. *Mar. Ecol. Prog. Ser.* 321, 99-121.
- 26 Nejstgaard, J.C., Gismervik, I., Solberg, P.T. 1997. Feeding and reproduction by
27 *Calanus finmarchicus*, and microzooplankton grazing during mesocosm
28 blooms of diatoms and the coccolithophore *Emiliana huxleyi*. *Mar. Ecol.*
29 *Prog. Ser.* 147, 197-217.

- 1 Nejstgaard, J.C., Naustvoll, L.J., Sazhin, A., 2001. Correcting for underestimation of
2 microzooplankton grazing in bottle incubation experiments with
3 mesozooplankton. Mar. Ecol. Prog. Ser. 221, 59-75.
- 4 Parsons, T.R., Maita, Y., Lalli, C.M., 1984. A manual of chemical and biological
5 methods for sea water analysis Pergamon Press, Oxford.
- 6 Psarra S., Tselepides A., Ignatiades L., 2000. Primary productivity in the oligotrophic
7 Cretan Sea (NE Mediterranean): seasonal and interannual variability. Prog.
8 Oceanogr. 46, 187-204.
- 9 Ramfos, A., Isari, S., Somarakis, S., Georgopoulos, D., Koutsikopoulos, C.,
10 Fragopoulou, N., 2006. Mesozooplankton community structure in offshore and
11 coastal waters of the Ionian Sea (eastern Mediterranean) during mixed and
12 stratified conditions. Mar. Biol. 150(1), 29-44.
- 13 Safi, K.A., Hall, J.A., 1999. Mixotrophic and heterotrophic nanoflagellate grazing in
14 the convergence zone east of New Zealand. Aquat. Microb. Ecol. 20, 83-93.
- 15 Saiz, E., Calbet, A., 2011. Copepod feeding in the ocean: scaling patterns,
16 composition of their diet and the bias of estimates due to microzooplankton
17 grazing during incubations. Hydrobiologia 666(1), 181-196.
- 18 Sazhin, A.F., Artigas, L.F., Nejstgaard, J.C., Frischer, M.E., 2007). The colonization
19 of two *Phaeocystis* species (Prymnesiophyceae) by pennate diatoms and other
20 protists: a significant contribution to colony biomass. Biogeochemistry 83,
21 137-145.
- 22 Schülter, L., 1998. The influence of nutrient addition on growth rates of
23 phytoplankton groups, and microzooplankton grazing rates in a mesocosm
24 experiment. J. Exp. Mar. Biol. and Ecol. 228, 53-73.
- 25 Skovgaard, A., 1996. Mixotrophy in *Fragilidium subglobosum* (Dinophyceae): growth
26 and grazing responses as functions of light intensity. Mar. Ecol. Prog. Ser.
27 143, 247-253.
- 28 Stickney, H., 2000. The impact of mixotrophy on planktonic marine ecosystems.
29 Ecol. Model. 125, 203-230.

- 1 Stoecker, D.K., 1998. Conceptual models of mixotrophy in planktonic protists and
2 some ecological and evolutionary implications. *Europ. J. Protistol.* 34, 281-
3 290.
- 4 Stoecker, D.K., 1999. Mixotrophy among Dinoflagellates. *Journal of Eukaryotic*
5 *Microbiology* 46(4), 397-401.
- 6 Stoecker, D.K., Johnson, M.D., deVargas, C., Not, F., 2009. Acquired phototrophy in
7 aquatic protists. *Aquat. Microb. Ecol.* 57, 279-310.
- 8 Straile, D., 1997. Gross growth efficiencies of protozoan and metazoan zooplankton
9 and their dependence on food concentration, predator-prey weight ratio, and
10 taxonomic group. *Limnol. Oceanogr.* 42(6), 1375-1385.
- 11 Teegarden, G.J., 1999. Copepod grazing selection and particle discrimination on the
12 basis of PSP toxin content. *Mar. Ecol. Prog. Ser.* 181, 163-176.
- 13 Therriault, J.C., Booth, D., Legendre, L., Demers, S., 1990. Phytoplankton
14 photoadaptation to vertical excursion as estimated by an in vivo fluorescence
15 ratio. *Mar. Ecol. Prog. Res.* 60, 97-111.
- 16 Thingstad, T.F., Cuevas, L.A., 2010. Nutrient pathways through the microbial food
17 web: principles and predictability discussed, based on five different
18 experiments. *Aquat. Microb. Ecol.* 61(3), 249251-249262.
- 19 Tittel, J., 2003. Mixotrophs combine resource use to outcompete specialists:
20 Implications for aquatic food webs. *Proceedings of the National Academy of*
21 *Sciences* 100(22), 12776-12781.
- 22 Unrein, F., Massana, R., Alonso-Sáez, L., Gasol, J.M., 2007. Significant year-round
23 effect of small mixotrophic flagellates on bacterioplankton in an oligotrophic
24 coastal system. *Limnol. Oceanogr.* 52, 456-469.
- 25 Urabe, J., Sterner, R.W., 1996. Regulation of herbivore growth by the balance of light
26 and nutrients. *Proceedings of the National Academy of Sciences of the United*
27 *States of America* 93, 8465.

- 1 Urabe, J., Kyle, M., Makino, W., Yoshida, T., Andersen, T., Elser, J.J., 2002.
2 Reduced Light Increases Herbivore Production Due To Stoichiometric Effects
3 of Light/Nutrient Balance. Ecology 83, 619-627.
- 4 Veloza, A.J., Chu, F.L., Tang, K.W., 2006. Trophic modification of essential fatty
5 acids by heterotrophic protists and its effects on the fatty acid composition of
6 the copepod *Acartia tonsa*. Mar. Biol. 148(4), 779-788.
- 7 Williams, P.J.L.B., Egge, J.K., 1998, The management and behaviour of the
8 mesocosms. Est. Coast. Shelf. Sci. 46, 3-14.
- 9 Worden, A.Z., Binder, B.J., 2003. Application of dilution experiments for measuring
10 growth and mortality rates among *Prochlorococcus* and *Synechococcus*
11 populations in oligotrophic environments. Aquat. Microb. Ecol. 30, 159-174.
- 12
- 13

Table 1. Phytoplankton instantaneous growth rates without nutrient addition (μ ; d⁻¹) and in situ mortality rates (m ; d⁻¹) obtained from dilution grazing experiments for the different size-fractions of Chl *a* in mesocosms L1a and L4a.

Mesocosm L1a

	Day 1			Day 3			Day 5			Day 7		
Chl <i>a</i> fraction	<i>m</i>	r ²	μ	<i>m</i>	r ²	μ	<i>m</i>	r ²	μ	<i>m</i>	r ²	μ
total	0.85	0.70	0.90	0.82	0.92 **	0.77	0.60	0.74	0.74	0.71	0.95 **	0.90
> 10 μm	0.96 *	0.86	1.17	ns	nd	0.13	ns	nd	0.44	0.47	0.79 **	0.52
2 to 10 μm	0.81	0.75	0.61	0.68	0.98 **	0.36	ns	nd	0.31	0.49	0.89	0.95
0.65 to 2 μm	0.85	0.55	0.99	1.07	0.85 **	1.19	0.66	0.85	0.52	0.69	0.84 **	0.89
0.2 to 0.65 μm	1.23	0.56	1.93	1.16	0.74	1.46	1.09	0.61	1.31	1.34	0.85 **	0.38

Mesocosm L4a

Chl <i>a</i> fraction	<i>m</i>	μ	<i>m</i>	r ²	μ	<i>m</i>	r ²	μ	<i>m</i>	r ²	μ
total	posit slope	nd	0.86	0.87	0.68	0.61	0.94	0.67	0.57	0.96 **	0.38
> 10 μm	posit slope	nd	0.26	0.62	0.62	0.32	0.73 **	0.32	0.49	0.65	0.39
2 to 10 μm	posit slope	nd	0.81	0.808	0.42	0.63	0.96	0.61	0.39	0.74 **	0.37
0.65 to 2 μm	posit slope	nd	1.19	0.816	1.09	0.85	0.98 **	0.94	0.40	0.85 **	0.36
0.2 to 0.65 μm	ns	-0.14	ns	nd	-0.0015	ns	nd	-0.09	0.56	0.58	-0.94

* one outlier removed; ** feeding saturation; nd = not determined; ns = not significant; posit slope = positive slope.

Table 2. Percentage of flagellates and dinoflagellates with ingested prey. The data have been obtained by multiplying the percentage of grazers with labeled prey times the proportion of labeled prey out of total prey. The experiments were conducted under the addition of excess of nutrients and without nutrients. ANF = autotrophic nanoflagellates, HNF = heterotrophic nanoflagellates, SE = standard error

Mesocosm	Nutrients	Day	% ANF	SE	% HNF	SE
L1a	No	1	2.59	0.21	7.59	0.19
L1a	No	3	5.61	1.25	11.67	0.84
L1a	No	5	4.80	0.71	8.88	2.18
L1a	No	7	2.71	0.19	8.19	0.54
L1a	Yes	1	2.44	0.41	4.59	0.73
L1a	Yes	3	4.06	0.31	8.43	1.58
L1a	Yes	5	4.02	0.62	4.85	0.78
L1a	Yes	7	2.83	0.69	12.05	1.77
L4a	No	1	4.13	0.23	8.35	0.81
L4a	No	3	5.19	0.58	9.14	1.04
L4a	No	5	4.19	1.14	7.52	0.74
L4a	No	7	4.08	0.45	9.93	1.13
L4a	Yes	1	4.56	0.37	8.59	0.76
L4a	Yes	3	5.51	1.18	10.33	0.84
L4a	Yes	5	4.10	0.92	7.85	1.30
L4a	Yes	7	3.58	0.56	11.63	1.15

Figure legends

Figure 1. Above: concentration of chlorophyll *a* (Chl *a*) along the experiment in the mesocosms L1a,b and L4a,b. Curves represent average values. Below: contribution of the different size-fractions of Chl *a*. Left panels L1, right panels L4.

Figure 2. Above: biomass of autotrophs and heterotrophs along the experiment in the mesocosms L1a,b and L4a,b (error bars are SE). Middle: contribution of the different groups of autotrophs to total autotrophic biomass. Below: contribution of the different groups of heterotrophs to total heterotrophic biomass. Left panels L1, right panels L4. Note that ciliates may not be well sampled with epifluorescence microscopy slides; however, they have been included in the plot for comparative purposes. Data are fitted to a smooth curve.

Figure 3. Abundance of ciliates (Lugol based) and contribution to mixotrophic ciliates (formaldehyde based) in the mesocosms L1a (above), and L4a (below).

Figure 4. Final abundance of the different mesozooplanktonic groups in L1a and L4a,b treatments. Note data for L1b are missing.

Figure 5. Elemental and mass ratios. A) Mass ratio of Carbon/Chl *a*, B) Molar ratio of Carbon/Nitrogen, C) Molar ratio of Carbon/Phosphorous. Data are fitted to a smooth curve. Note day 8 for C:N is missing.

Figure 6. Example of dilution grazing experiment plots corresponding to total Chl *a* in mesocosms L1a and L4a.

Figure 7. Autotroph growth rates obtained by cell counts (solid symbols) and by Chl *a* analysis of the 2 to 10 μm fraction (open symbols) in L1a and L4a mesocosms. Asterisks indicate significant differences ($p < 0.05$).

Figure 8. Total Chl *a* based phytoplankton growth rates in the nutrient amended and unamended bottles in L1a and L4a mesocosms. Error bars are SE.

Figure 9. Autotrophic carbon flux through microzooplankton in L1 and L4 mesocosms (see results); nd: not determined because positive slope in the microzooplankton grazing experiments (see results).

Figure

Fig. 1

L1

L4

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9