12-6-07

Altering the laccase functionality by *in vivo* assembly of mutant libraries with different mutational spectra

Miren Zumárraga¹, Susana Camarero², Sergey Shleev³, Arturo martinez-arias¹,
Antonio Ballesteros¹, Francisco J. Plou¹ and Miguel Alcalde^{1*}

¹ Instituto de Catálisis y Petroleoquímica (CSIC). Cantoblanco, 28049 Madrid, Spain.

² Centro de Investigaciones Biológicas, (CSIC), Ramiro de Maeztu, 9, 28040 Madrid, Spain.

³ Department of Analytical Chemistry, Lund University, 22100 Lund, Sweden.

*Correspondence: Miguel Alcalde, Instituto de Catálisis y Petroleoquímica (CSIC), Cantoblanco, 28049 Madrid, Spain. Tel: (34) 915855479, Fax: (34) 915854760. E-mail: malcalde@icp.csic.es, malcalde@cheme.caltech.edu

Keywords: *in vivo* assembly, laccase, C-terminal tail, protein folding.

Running title: altering laccase functionality by IvAM

ABSTRACT

The generation of diversity for directed protein evolution experiments shows an important bottleneck in the *in vitro* random mutagenesis protocols. Most of them are biased towards specific changes that eventually confer a predicted and conservative mutational spectrum, limiting the exploration of the vast protein space. The current work describes a simple methodology to *in vivo* recombine mutant libraries with different nucleotide bias created by *in vitro* methods. This *in vivo* assembly was based on the accurate physiology of *Saccharomyces cerevisiae*, which as host, provided its high homologous recombination frequency to shuffle the libraries in a non-mutagenic way. The fungal thermophilic laccase from *Myceliophthora thermophila* expressed in *S. cerevisiae* was submitted to this protocol under the selective pressure of high concentrations of organic solvents. Mutant 2E9 with ~3-fold better kinetics than parent type showed two consecutive amino acid changes (G614D –GGC/GAC- and E615K - GAG/AAG-) due to the *in vivo* shuffling of the mutant libraries. Both mutations are located in the C-terminal tail that is specifically processed at the Golgi during the maturation of the protein by the *Kex2* protease. Notoriously, the oxygen consumption at the T2/T3 trinuclear copper cluster was altered and the catalytic copper at the T1 site was perturbed showing differences in its redox potential and geometry. The change in the isoelectric point of C-terminal extension upon mutations seems to affect the folding of the protein at the posttranslational processing steps providing new insights in the significance of the C-terminal tail for the functionality of the ascomycete laccases.

INTRODUCTION

In the last 15 years, the idea of mimicking the darwinist concept of natural selection employing experimental protocols directed to enhance or even create novel complex functions in protein structures has opened a big window to an unexplored world.¹⁻⁵ To most of us, laboratory evolution (the generation of diversity by *in vitro* or *in vivo* methods coupled to high-throughput screening assays) represents a *shortcut* to tailor enzymes with several improved features.⁶⁻⁹ In spite of the fact that molecular biologists and protein engineers are focusing their efforts to farther push the technology of directed enzyme evolution with the main aim of limiting the protein space, *lab* evolution is still in its infancy.^{10,11} The requirement of more efficient high-throughput assays to screen bigger libraries -over 10^8 - 10^{10} variants- is one of the main hurdles.^{12,13} Moreover, the generation of diversity is unequivocally limited by the nature of the genetic code¹⁴. In this sense, computational methods and *semi-rational* approximations (i.e. combinatorial saturation mutagenesis¹⁵) or more recently the introduction of new tools such as circular permutation¹⁶, are being used to design proteins efficiently. But, yet, we have to deal with the fact that natural evolution is somehow a sophisticated engineering process from whose efficacy we are far away. In other words, the accumulation of subtle changes at genetic level not always drive to the best enzymatic adaptation, or at least not with the molecular tools that we have currently in hand. Basically there are two ways of generating diversity: the *asexual* approach (i.e. the random introduction of mutations based on error-prone PCR techniques) and the *sexual* one (DNA recombination).¹⁷ The latter has achieved a reasonable level of development enabling DNA fragments of low homology to recombine each other.¹⁸ Thus, nowadays the main concern in generation of diversity stems from unsatisfactory mutagenic methods^{19,20} which usually display limited and predicted mutational spectra because of the intrinsic bias of each DNA-polymerase.²¹

In the last few years, the use of *Saccharomyces cerevisiae* physiology as biomolecular tool is providing new alternatives to the current bottlenecks found during the construction of *in vitro* evolution libraries.²²⁻²⁵ We believe that *S. cerevisiae* accurate device may be helpful to *in vivo* recombine mutant libraries created by several means, -i.e. with different bias-, and therefore, to produce an unexpected mutational spectrum. In order to validate this hypothesis, we have chosen as scaffold the ascomycete laccase from *Myceliophthora thermophila*, MtL, that was previously evolved for functional expression in *S. cerevisiae*.²⁶ Due to its biotechnological interest (with applications in the pulp-kraft bleaching, oxidation of xenobiotics, chemical synthesis...), laccase is a promising candidate to be fuelled by laboratory evolution strategies.²⁷⁻²⁹ Laccases belong to the family of multi-copper oxidases -with one copper T1 where the reducing substrate binds and a trinuclear copper cluster T2/T3 in which the O₂ is reduced to two molecules of water-.²⁸ MtL gene constitutes a suitable and complex scaffold possessing a pre-proleader and an exclusive C-terminal tail which codes for parts of the protein that are supposed to be cleaved during post-translational stages in yeast.²⁶ The importance and processing of these polypeptides in native or heterologous hosts are not well understood yet -although they have been extensively studied in other ascomycete laccases-³⁰⁻³² and therefore have been included for mutagenesis and recombination in the current study.

This article describes the use of *S. cerevisiae* for *in vivo* recombining distinct MtL mutant libraries (from polymerases with different bias) giving rise to a variant with modifications at the C-terminal tail. Taking advantage of the eukaryotic machinery, the libraries created by several *in vitro* methodologies were *in vivo* shuffled and screened to find mutants with better fitness towards organic cosolvents -generally used to enhance the solubility of many of the laccase reducing substrates-. The role of the C-terminal tail and its action during protein folding is discussed in detail.

MATERIAL AND METHODS

All chemicals were of reagent-grade purity. Taq-DNA polymerase, ABTS (2,2'-azino-bis(3-ethybenzthizoline-6-sulfonic acid)) and *S. cerevisiae* transformation kit were purchased from Sigma-Aldrich (Madrid, Spain). Parent type laccase gene was engineered from MtLT2²⁶ through three consecutive cycles of *in vitro* evolution under the selective pressure of increasing concentrations of organic cosolvents, accumulating mutations L429V, E182K and N552H in the mature protein (unpublished material). GeneMorph PCR mutagenesis kit and *E. coli* XL2-blue competent cells were from Stratagene (La Jolla, CA). Protease deficient *S. cerevisiae* strain BJ5465 was from LGCPromochem (Barcelona, Spain). Shuttle vector pJRoC30 with auxotrophy for uracil and with ampicillin resistance gene was from Novozymes (Davis, CA). Zymoprep yeast plasmid miniprep kit, zymoclean gel DNA recovery kit, and DNA clean and concentrator TM-5 kit were from Zymo Research (Orange, CA). QIAprep spin miniprep kit was from QIAGEN (West Sussex, UK). Restriction enzymes BamHI and XhoI were from New England Biolabs (Hertfordshire, UK).

Culture media

Minimal medium contained 100 ml 6.7% sterile yeast nitrogen base, 100 ml 19.2 g/L sterile yeast synthetic drop-out medium supplement without uracil, 100 ml sterile 20% raffinose, 700 ml *dd* H₂O, 1 ml 25 g/L chloramphenicol. YP medium contained 10 g yeast extract, 20 g peptone, and *dd* H₂O to 650 ml. Expression medium contained 720 ml YP, 67 ml 1 M KH₂PO₄ pH 6.0 buffer, 10 µl 1 M CuSO₄, 111 ml 20% galactose, 1 ml 25 g/L chloramphenicol and *dd* H₂O to 1000 ml. YPAD solution contained 10 g yeast extract, 20 g peptone, 100 ml 20% sterile glucose, 100 mg adenine hemisulfate, 1 ml 25 g/L chloramphenicol and *dd* H₂O to 1000 ml. SC drop-out plates contained 6.7 g yeast nitrogen base, 100 ml 19.2 g/L sterile yeast synthetic

drop-out medium supplement without uracil, 20 g bacto agar, 100 ml 20% sterile glucose, 1 ml 25 g/L chloramphenicol, and *dd*H₂O to 1000 ml.

Production of laccase in *S. cerevisiae*

One single colony from *S. cerevisiae* clone containing parent-type or mutant laccase genes was picked from a SC drop-out plate, inoculated in 3 ml of minimal medium and incubated for 48 h at 30°C and 225 rpm (Micromagmix shaker, Ovan, Spain). An aliquot of cells was removed and inoculated into a final volume of 50 ml of minimal medium (optical density, OD₆₀₀=0.25) in a 500 ml flask. Incubation proceeded until two growth phases were completed (6 to 8 h). Thereafter, 450 ml of expression medium containing 5.6 mM CuSO₄ were inoculated with the 50 ml preculture in a 2.0 - litre baffled flask (OD₆₀₀=0.1). Incubation was stopped after 38-42 h at 30°C and 225 rpm (laccase activity was maximal reaching a plateau; OD₆₀₀=28-30). The cells were separated by centrifugation for 20 min at 3000 *g* (4°C). Supernatant was double-filtered (by both glass membrane and a nitrocellulose membrane of 0.45 µm pore size) and concentrated to 20 ml in an ultrafiltration cell (Amicon/Millipore, Barcelona, Spain) equipped with a 10 kDa cutoff membrane.

Purification of laccases

Parent and mutant laccases were purified using fast protein liquid chromatography (FPLC) equipment (LCC-500CI, Amersham Bioscience, Barcelona, Spain). Concentrated crude extract was first submitted to fractional precipitation with ammonium sulfate to 50% saturation (w/v) at 0°C and centrifuged at 17,000 *g* for 20 min. The supernatant was filtered and loaded into a hydrophobic interaction column (HIC, Hiload 16/10 Phenyl Sepharose High Performance, Amersham) equilibrated with 1.8 M (NH₄)₂SO₄ in 10 mM sodium phosphate (pH 6.1). Proteins were eluted with a linear gradient from 1.8 to 0 M (NH₄)₂SO₄. Fractions with laccase activity were pooled, concentrated and dialyzed against 10 mM sodium phosphate (pH 6.1). Semi-purified

laccase was applied to an anion exchange column (DEAE Sepharose CL-6B) preequilibrated with 10 mM sodium phosphate (pH 6.1). Proteins were eluted with a linear gradient from 0 to 0.4 M NaCl. Fractions with laccase activity were pooled, concentrated and dialyzed against 10 mM sodium phosphate (pH 6.1). Fractions throughout the purification protocol were analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) with 8 % polyacrilamide using a Miniprotean 3 Cell (BioRad, US). Proteins were stained with both Coomassie brilliant blue and silver. Protein concentration was determined using the Bio-Rad protein reagent and bovine serum albumin as a standard. Purified laccases were stored at -20°C .

***In vivo* assembly of mutant libraries with different mutational spectra (IvAM)**

1) Mutagenic PCR

Mutagenic PCR was carried out using a gradient thermocycler (Mycycler, Biorad, US). Thermal cycling parameters were: 95°C for 2 min (1 cycle), 94°C for 0.45 min, 53°C for 0.45 min, 74°C for 3 min (28 cycles), 74°C for 10 min (1 cycle). The primers used for amplification were: IG88-S sense (5'-CCTCTATACTTTAACGTCAAGG-3', binds at bp 5'-160-180-3' of pJRoC30) and IG88-R antisense (5'-GGGAGGGCGTGAATGTAAGC-3', binds at bp 5'-2139-2158-3' of pJRoC30). The size of Mtl gene is 1912 bp. The circular autonomous vector pJRoC30 with Mtl gene has a size of 12337 bp.

a) Taq/MnCl₂ libraries

Taq/MnCl₂ libraries were prepared at different concentrations of MnCl₂ to estimate an appropriate mutation rate. Additionally, the effect of balanced/unbalanced dNTPs and the concentration of template was analyzed.

a.1) Taq/MnCl₂ libraries with unbalanced dNTPs: in a 50 μl final volume, two different reactions conditions were explored (with different template concentrations). Reaction

i): 90 nM IG88-S; 90 nM IG88-R; 1.5 ng/ μ l laccase template; 0.2 mM dATP; 0.2 mM dGTP; 0.6 mM dCTP; 0.6 mM dTTP; 3 % DMSO; 0.75 mM MgCl₂; 0.05 U/ μ l Taq polymerase. Reaction ii) 90 nM IG88-S; 90 nM IG88-R; 0.1 ng/ μ l laccase template; 0.04 mM dATP; 0.04 mM dGTP; 0.1 mM dCTP; 0.1 mM dTTP; 3 % DMSO; 1.5 mM MgCl₂; 0.05 U/ μ l. The influence of MnCl₂ was studied at different final concentrations (0, 0.05mM, 0.1mM, 0.2mM).

a.2) Taq/MnCl₂ libraries with balanced dNTPs and low concentration of template: the concentrations for each reagent -in 50 μ l final volume- were as follows: 90 nM IG88-S; 90 nM IG88-R; 0.1 ng/ μ l laccase template; 0.3 mM dNTPs (0.075 mM each); 3 % DMSO; 1.5 mM MgCl₂; 0.05 U/ μ l Taq polymerase. The influence of MnCl₂ was studied at different final concentrations (0, 0.002 mM, 0.005 mM, 0.01 mM, 0.02 mM, 0.05 mM, 0.1 mM and 0.2 mM).

b) Mutazyme libraries

High and low mutation rate libraries were constructed using the Genemorph kit (Stratagene, CA). The concentration of each reagent -in 50 μ l final volume- was as follows: 370 nM IG88-S; 370 nM IG88-R; 0.8 mM dNTPs; 3 % DMSO; 0.05 U/ μ l Mutazyme DNA polymerase; 13 ng/ μ l and 4 ng/ μ l laccase templates for low mutation rate libraries; and 4 pg/ μ l laccase template for high mutation rate library.

2) Preparing libraries in *S. cerevisiae*

PCR products were cleaned and concentrated (DNA clean and concentrator TM-5 kit, Zymo Research), loaded into a low melting point preparative agarose and purified using the Zymoclean gel DNA recovery kit (Zymo Research). PCR product was cloned behind the Gal 10 promoter of the expression shuttle vector pJRoc30. Cloning was performed by replacing MtlT2 gene from pJRoc30. To remove parent gene, plasmid pJRoc30 was linearized by cutting with restriction enzymes XhoI and BamHI (New England Biolabs). Linearized plasmid was concentrated (DNA clean and

concentrator TM-5 kit) and loaded into a preparative low melting point agarose followed by an agarose gel extraction (Zymoclean gel DNA recovery kit). PCR products were mixed with linearized vector and transformed into competent yeast cells using the Yeast transformation kit (Sigma). Right before the thermal shock, 10 % DMSO was introduced in the transformation mixture to improve the transformation efficiency (above 15,000 clones per transformation reaction).

3) *In vivo* recombination of mutant libraries in *S. cerevisiae*

Taq/MnCl₂ and Mutazyme libraries were equimolarly mixed. The equimolar mixture was added to 100 ng of linearized vector (ratio vector:equimolar library tested: 1:1, 1:2, 1:4, 1:6, 1:8, 1:10) and transformed into yeast competent cells as described above.

HTPS assays: end-point colorimetric assay in the presence of organic-cosolvent.

Screening protocol was performed according to Alcalde et al ²⁹ with minor modifications. Individual clones were picked into 96-well plates (Sero-well, Staffordshire, UK) containing 50 µl of minimal medium per well. In every single plate the column number 6 was inoculated with standard (parent type), and one well (H1) was not inoculated (control). Plates were wrapped with parafilm (to prevent evaporation) and incubated at 30°C, 225 rpm and 80 % relative humidity in a humidity shaker (Minitron-INFORS, Biogen, Spain). After 48 h, 160 µl of expression medium were added to each well, and plates were incubated again. After 24 h, OD₆₀₀ was recorded to determine the cell density in each well using a microplate reader (VersaMax, Molecular Devices, Sunnyvale, CA). Plates (master plates) were centrifuged (Eppendorf 5810R centrifuge, Germany) for 5 min at 3000 *g* at 4°C. 20 µl of supernatant were transferred from master plate with the help of a robot (Liquid Handler Quadra 96-320, Tomtec, Hamden, CT, US) to a replica plate (polypropylene

solvent resistant plate). 180 μl of assay solution (final concentration in the well 3 mM ABTS, with/without 50% organic co-solvent -acetonitrile or ethanol-, 100 mM Britton and Robinson buffer pH 6.0) was added to each well of replica plate containing supernatant. Plates were stirred and the absorption at 418 nm ($\epsilon_{\text{ABTS}}^{*+}=36,000 \text{ M}^{-1} \text{ cm}^{-1}$) was recorded in the plate reader. Plates were incubated at room temperature until green color was developed, and the absorbance was measured again. Activities were calculated from the difference between absorbance after incubation and the initial absorption, divided by the OD_{600} of each well; relative activities were normalized against the parent type in the corresponding plate.

Rescreening procedures

First rescreening: aliquots of 5 μl of the best clones were used from master plates to inoculate 50 μl of minimal media in new 96-well plates. Columns 1 and 12, rows A and H, were not used to prevent the appearance of false positives. Five wells on the microplates were inoculated with the same clone. Five wells on each plate were used for the standard. The screening procedure then was the same as above.

Second rescreening: an aliquot from the wells with the most active clones of first rescreening was inoculated in 3 ml of YPAD and incubated at 30°C and 225 rpm for 24 h. Plasmids from these cultures were extracted (Zymoprep yeast plasmid miniprep kit, Zymo Research). As the product of the zymoprep was very impure and the concentration of extracted DNA was very low, the shuttle vectors were transformed into super-competent *E. coli* cells (XL2-Blue, Stratagene) and plated onto LB-amp plates. Single colonies were picked and used to inoculate 5 ml LB-amp media and were grown overnight at 37 °C and 225 rpm. Plasmids were then extracted (QIAprep spin miniprep kit, QIAGEN). *S. cerevisiae* was transformed with plasmids from the best mutants and also with parent type. Five colonies of every single mutant were picked and rescreened as described above.

DNA sequencing

Plasmid-containing variants laccase genes were sequenced at the Sequencing Core Facility of the Instituto de Investigaciones Biomedicas, CSIC, Madrid, using an Applied Biosystems 377 automated fluorescent DNA sequencer. The primers used were as follows: forward: IG88-S and mtlsq2 (5'-GAAGGGCACCAACCTGC-3', binds at bp 5'-643-659-3' of pJRoC30); reverse: mtlsq3 (5'-CGCACGTAAAAGTCGTGG-3', binds at bp 5'-1657-1673-3' of pJRoC30) and IG88-R.

Spectroelectrochemical characterization

Spectroelectrochemistry of purified laccases (MtL variants as well as *Trametes hirsuta* laccase as control) was carried out using a micro-spectroelectrochemical cell consisting of the gold capillary electrode. The design of the cell was described elsewhere³³. The potential of the gold capillary of the cell was controlled by a three-electrode potentiostat BAS LC-3E from Bioanalytical Systems, BAS (West Lafayette, IN, USA). In these measurements an Ag|AgCl|KCl reference electrode (BAS) and a platinum counter electrode were used. The absorbance spectra were monitored with PC2000-UV-VIS, a miniature fiber optic spectrometer from Ocean Optics (Dunedin, FL, USA) with an effective range between 200 and 1100 nm. The pre-treatment of the gold capillary working electrode of the spectroelectrochemical cell was carried out by washing the cell capillary with a peroxide-sulfuric acid mixture followed by rinsing with Millipore water. The redox potential of the T1 site of the enzymes was determined by mediated spectroelectrochemical redox titration, MRT, using spectroelectrochemical set-up described above. Complex mediator system containing four different mediators ($K_4[Fe(CN)_6]$, $K_4[W(CN)_8]$, $K_4[Os(CN)_6]$, and $K_4[Mo(CN)_8]$ with formal redox potentials of 430 mV, 520 mV, 640 mV, and 780 mV *vs.* NHE, respectively) was used for the MRT performed in accordance with previously published methodology³⁴.

Spectral investigations

EPR analysis of copper sites of purified laccases

Electron paramagnetic resonance –EPR– measurements were carried out with a Bruker ER200D instrument operating in the X-band ($\nu \approx 9.6$ GHz) using a DPPH standard (in a T-type double cavity) for frequency calibration. Portions of sample in potassium phosphate buffer were introduced into a spectroscopic quartz probe cell. The spectra were recorded at 77 K and were typically performed at 19.5-milliwatt microwave power (no signal saturation was apparent in independent experiments up to 40 milliwatts), 100 kHz modulation frequency, 2G modulation amplitude, 40 ms time constant, and 1×10^5 receiver gain.

Circular dichroism spectra of purified laccases

Circular dichroism (CD) spectra were recorded using a Jasco J-720 spectropolarimeter (with a scan rate of 20 nm/min and a bandwidth of 1.0 nm). The measurements were performed under N₂ atmosphere employing a 0.01-cm cell with a protein concentration of ~ 2 mg/ml. The parameters of the secondary structures were calculated using an in house computer program "Protein-CD v 1.5" (Moscow, Russia) as described in ^{35,36}.

O₂-reduction activity measurements

The O₂-reduction activity of laccases was measured by mass spectrometry with 3.0 mM ABTS as electron donor in 10 mM acetate buffer pH 4.5, at 27°C. The measurement cell had 10 ml volume and was separated from the mass spectrometer (Masstorr 200 DX quadrupole, VG Quadrupoles Ltd.) by a teflon membrane through which gasses from the solution diffuse ³⁷. The response at the spectrometer at certain mass value is proportional to the partial pressure of the gas in the solution of the cell. O₂-uptake was measured at mass 16. Calibration was done with the solution saturated with pure O₂ and background signal was estimated with the solution purged with pure Ar. O₂-consumption by the mass spectrometer (measured in absence of laccase for

each O₂ concentration) was subtracted to the activity measurements, in which 5-25 µg of laccase were added.

RESULTS AND DISCUSSION

Analysis of mutant libraries constructed by Taq/MnCl₂ and Mutazyme

Two different DNA-polymerases -Taq polymerase and Mutazyme- were used to *in vitro* create several mutant libraries of *Myceliophthora thermophila* laccase. First of all, mutation rate was tuned to introduce 1-3 amino acid changes (1-5 nucleotide substitution/laccase gene). Few exceptions aside^{38,39}, higher mutations are generally not suitable in laboratory evolution experiments, especially when one wants to distinguish beneficial mutations from those that are neutral or deleterious. Mutagenic PCR using Taq DNA polymerase takes advantage from the lack of 3'-5' proofreading exonuclease activity. Generally, the modulation of mutational rate can be accomplished by: introducing MnCl₂ in the reaction mixture; varying the ratios of nucleotides in the reaction; changing the initial template concentration or increasing concentration of MgCl₂.¹⁴ Laccase parent gene was submitted to PCR amplification with Taq/MnCl₂, unbalanced dNTPs and high concentration of template. To make a correct comparison between mutagenic PCR protocols under different conditions, small libraries (two 96 well-plates) were constructed and their landscapes provided an estimation of the mutation frequency from the fraction of inactive clones, Fig. 1. Typically an error rate resulting in a library with 30 to 45% of clones having less than 10% of the wildtype activity is suitable for achieving improvements during a laboratory evolution process.¹⁸ When unbalanced dNTPs were introduced, mutation rate was lower than expected (with only 4.6% mutants having less than 10% of the parent enzyme's activity under 0.2 mM MnCl₂). The concentrations of dNTPs and MgCl₂ were varied and the amount of template reduced to increase mutational frequency. Under these conditions -see

Material and Methods section- the mutational rate was considerably enhanced (with 14% of mutants having less than 10% of the M_tLT2 activity in absence of MnCl₂, Table I). Libraries with balanced dNTPs were further studied. The mutation frequency was smoothly tuned by just varying the MnCl₂ concentration, ranging from 34% to 81% of clones with less than 10% of the parent's activity, for Taq libraries from 0 mM to 0.05 mM of MnCl₂ (Fig. 1, Table I). Mutagenic libraries were also constructed using the Mutazyme DNA polymerase. This polymerase has a high intrinsic error rate which can be simply modulated by adjusting the initial target DNA amount for the amplification reaction -the lower the concentration of template the higher the mutational rate-. When DNA template concentration was between 13 to 4 ng/μl, libraries landscapes were appropriate (from 36.8% to 47.1% of inactive clones).

In vivo* assembly of mutant libraries (IvAM) in *S. cerevisiae

The Transition/Transversion ratio (Ts/Tv) is commonly used to calculate the bias in the mutational spectra. Both Mutazyme and Taq DNA polymerases favor transitions (purine to purine changes and pyrimidine to pyrimidine changes) over transversions (purine to pyrimidine changes and *vice versa*) (Ts/Tv ranged from 2.9 to 0.8).⁴⁰⁻⁴² Apart from this similarity, both polymerases behave in different manner. First, they do not share the same AT→GC/GC→AT ratio. Taq is biased to AT→GC changes whereas Mutazyme is biased to GC→AT changes. Moreover, Mutazyme is highly likely to incorporate A→N and T→N changes whereas Taq DNA is biased to G→N and C→N mutations.

Generally it is advisable to combine the mutation biases of these or other polymerases by alternating between them in successive rounds of evolution, or in a parallel approach just creating independent mutant libraries with different polymerases in one single generation.¹⁸ We consider that the recombination of mutant libraries with different mutational spectra can also constitute an interesting alternative during a

laboratory laccase evolution process. To this end we tackled the *in vivo* construction of an unique library in *S. cerevisiae*, from the two *in vitro* independent libraries (Taq/MnCl₂ and Mutazyme). Owing to the high homologous recombination frequency of *S. cerevisiae*, *in vivo* DNA-shuffling of mutant libraries takes place generating a mutational spectra which is derived from the own features of both polymerases. Libraries Taq-V and Mutazyme-I were selected to validate this procedure. Both libraries showed a suitable mutational rate -Table I- to reasonably build a generation where beneficial mutations could be identified and at the same time evaluated the *in vivo* recombination process.

Taq-V and Mutazyme-I libraries were prepared using a set of primers which allowed to amplify extra-homologous sequences at the 5' and 3' ends (40 bp and 66 bp respectively). These big overhangs recombined with the ends of the linearized vector forming an autonomously replicating plasmid without altering the open reading frame. Both libraries were equimolarly mixed and transformed along with the linearized vector in *S. cerevisiae* (see details in Material and Methods). To guarantee the recombination/transformation efficiency, six different linearized vector:equimolar library were investigated. Ratio 1:8 provided the best result with over 15,000 clones per transformation reaction, which is quite similar to that obtained with the circular replicating plasmid.

Screening the library constructed by IvAM in organic co-solvents

The screening was done in the presence of high concentrations of acetonitrile and ethanol to find variants with resistance towards these cosolvents of biotechnological significance and commonly employed in multiple biotransformations catalysed by laccases.^{11,29} Results were very consistent in both media showing 38.0% and 36.4% of clones with less of 10% wildtype's activity for ethanol and acetonitrile, respectively. Over 1000 clones were screened, finding several mutants with around 2-

fold better rates than parent type both in acetonitrile and ethanol, Fig. 2. Several variants were sequenced and in all cases a mutation frequency of 1-4 nucleotide/laccase gene was detected. All sequenced mutants displayed transitions over transversions. This result is not unexpected keeping in mind that both polymerases are biased towards these sort of changes. As can be appreciated in the landscapes, one variant -2E9- highlighted among remaining positive mutants. 2E9 was purified and further characterized showing ~3 fold better kinetics than parent type along with significant changes in the redox potentials at the T1 site and in the oxygen uptake at the T2/T3 copper cluster (Table II). 2E9 variant showed a good tolerance towards organic solvents keeping 51% and 30% of its initial activity under the presence of ethanol 30% (v/v) and acetonitrile 30% (v/v), respectively (Table III, Fig. 3). DNA sequencing revealed an unexpected result. 2E9 presented two mutations in consecutives codons (G614D -GGC/GAC- and E615K -GAG/AAG-). In both cases two transitions -G→A- were produced. Although these kind of changes are more biased for Mutazyme than for Taq/MnCl₂ (43.7 % vs 13.6 %) the participation of the Taq/MnCl₂ in these mutations can not be ruled out either. The likelihood of introducing two positive mutations consecutively by using conventional error prone PCR is hardly expected and to our best knowledge there are no reports of this phenomena in other protein evolution investigations. Rather than serendipity, this result suggests that the approach of *in vivo* recombining mutant libraries with different bias in *S. cerevisiae* can provide other alternatives to the standard *in vitro* protocols.

Functionality of the C-terminal tail

Mutant libraries were built throughout the whole laccase gene, including several processing sites, because of their significance in the laccase total activity.²⁶ Thus, the sequence targeted for random mutagenesis included: signal leader (22 amino acids) + pro-leader (25 amino acids) + mature protein (559 amino acids) + C-terminal tail (14

amino acids). Signal leader, pro-leader and C-terminal tail code for parts of the laccase that are cleaved during maturation. The signal leader is necessary for the secretion of the laccase, which is firstly glycosylated upon translocation to the endoplasmic reticulum. Afterwards, the yeast secretion machinery operates -through the Golgi- in processing, further glycosylating and finally secreting the enzyme. Pro-leader peptide may act as a chaperone (as happening with the α -factor mating pheromone) although its function in laccase processing is not well understood yet.⁴³ The function of C-terminal tail in the ascomycete laccases is also under the shadow of controversy.^{26,32} Kiiskinen and Saloheimo³² studied the expression of the *Melanocarpus albomyces* laccase, MaL, -with 75% of homology to MtL- in *S. cerevisiae*, suggesting that yeast apparatus was not able to process the C-terminus correctly. By contrast, we reported the correct processing of the C-terminus of MtL in *S. cerevisiae* after the introduction of a *Kex2* protease site²⁶ -which was further corroborated by C-terminal sequencing-. *Kex2* protease is placed at the Golgi, where it cleaves fusion proteins with Lys-Arg processing sites (as also happens with the α -factor prepro leader). At this point, it is worth noting that the two consecutive amino acid changes of 2E9 are located in the middle of the C-terminal tail (the eighth and ninth amino acid of the C-terminal tail). In the current study, parent type already possessed a *Kex2* protease recognition site at the C-terminal processing site of the protein, adjusting therefore the protein sequence to the different protease specificities of *S. cerevisiae*.

Since the C-terminal polypeptide is processed and therefore is not part of the mature protein, it is not straightforward to find out how those changes can modulate the function of the mature 2E9. To evaluate whether or not the mutations at the C-terminal tail might affect the geometric and electronic structure of catalytic coppers in the mature protein, spectral investigations of both parent type and 2E9 variant were performed. Comparison of the $g_{||}$ and $A_{||}$ parameters of the parallel component of the

T1 copper signal (Fig. 4a) from EPR spectroscopy reveals that the mutations induced small but significant changes in the chemical environment of this site ($g_{\parallel} = 2.204$ and $A_{\parallel} = 7.9 \times 10^{-3} \text{ cm}^{-1}$ for the parent laccase, and $g_{\parallel} = 2.210$ and $A_{\parallel} = 7.4 \times 10^{-3} \text{ cm}^{-1}$ for 2E9). Hence, the redox potential value at the T1 site (Table I) was also perturbed upon mutations. Similar alterations have been reported when several residues in the neighbourhood of the T1 site were submitted to site directed mutagenesis.⁴⁴⁻⁴⁶ The geometry of the paramagnetic copper T1 in the overall architecture of MtL comes from a complex network of interactions⁴⁷ (van der Waals forces, electrostatic interactions, hydrogen bonds and covalent linkages) throughout the laccase structure. This network is building during protein folding at the posttranslational processing steps. Circular dichroism spectra of the parent type and 2E9 mutant in the far UV region were recorded in order to compare the secondary structure of the enzymes. As shown in Fig. 4b, spectra of both laccases were very similar and had a single broad negative band with minimum at ca. 205 nm. The theoretical quantitative comparison of the content of different structure motifs (see Materials and Methods section) addressed that the secondary structure of the 2E9 mutant was not significantly altered by the mutations (data not shown).

In one of our previous work, we created a C-truncated version of *Myceliophthora thermophila* laccase by introducing a stop codon at the processing site²⁶. The catalytic efficiency of the resulting enzyme –which was processed without the C-terminal extension- was reduced 10-fold although the expression level was kept. Thus, it seems to be clear that somehow the C-terminal tail exerts a strong influence during processing steps which eventually is affecting in how mature enzyme behaves. Similar C-terminal processing has also been described for laccases from the ascomycetes *Neurospora crassa*³¹, *Podospora anserine*³⁰ and *Melanocarpus albomyces*.³² It has been suggested that carboxyl-terminal processing may play a role

in the activation of the enzyme. This hypothesis is based on the inherent basic character of the C-terminal tail (in our parent type, 4 out of 14 residues are basic ones -two Lys and two Arg- conferring to the tail an isoelectric point -pI- of 8.59). This value is considerably higher than that of the mature protein -5.29-. According to this theory³¹, the C-terminal tail forms a loop interacting with the active site to prevent binding of copper ions during processing. Interestingly, throughout the directed evolution process the pI of mature protein practically kept unalterable but not that of the C-terminal tail, Table IV. In particular, the two mutations located in 2E9 variant shifted the pI up to 1.1 units to the basic side -from 8.59 to 9.70- while keeping constant the mature protein pI at 5.46 -the pH activity profile of 2E9 did not significantly change, Fig. 5-. We assume that these mutations might contribute to a higher grade of tightness between the C-terminal extension and the main enzymatic core, which would affect to the protein folding and therefore the final mature enzyme. Hence the differences in the EPR T1 signals and in the estimated redox potentials (Fig. 4a, Table II), whereas similarity of secondary structures was kept (Fig. 4b). Our theory is further supported by the fact that also the $K_M^{O_2}$ was altered in 2E9 -oxygen uptake was reduced by ~ 1.46 fold (Table II)-. Taking into account that the oxygen molecules access to the trinuclear copper cluster through a broad channel⁴⁸, any subtle modification during the protein folding might have affected the entrance and exit of O_2 with the concomitant increase in the $K_M^{O_2}$.

Although to validate these hypothesis only crystallographic comparative analysis of parent type and 2E9 could give us new insights (currently under investigation), in view of the reported data it seems highly likely that the broad differences in pI between mature protein and C-terminal tail might help to establish stronger electrostatic interactions between both polypeptides during protein processing altering the eventual laccase functionality.

CONCLUSIONS

In this work we have taken advantage of the cellular mechanism of *S. cerevisiae*, which unlike *Escherichia coli*, displays a high homologous recombination frequency.^{23,49} This virtue, along with its ability for glycosylation/secretion make *S. cerevisiae* a suitable host for laboratory evolution of eukaryotic proteins. *S. cerevisiae* possesses a high transformation efficiency, and using an appropriate episomal vector does not integrate the foreigner plasmid into the chromosome (as *Pichia pastoris* does), enabling its further manipulation. All these features have been extensively exploited to evolve enzymes by *in vivo* shuffling without introducing additional mutations (*in vivo* shuffling is a non-mutagenic method, unlike most of *in vitro* DNA recombination protocols generally employed in bacteria).^{18,50} Here, we have used these qualities to assemble mutant libraries with different bias using a protocol that we have named IvAM. With this approach, the laccase functionality in cosolvents was explored, pointing out the significance of C-terminal extension in the function of ascomycete laccases. IvAM constitutes a robust methodology that can be straightforwardly used to simplify the process of enzyme engineering by laboratory evolution. The future use of *S. cerevisiae* machinery in the laboratory evolution picture can be useful to further push laccases and other oxidoreductases towards technological applications (bioremediation schemes, design of biofuel cells...).⁵¹

ACKNOWLEDGEMENTS

We thank Dr. A. L. de Lacey and C. Vaz (ICP, CSIC) for measuring the kinetics for O₂-reduction. We thank Prof. F.H. Arnold (Caltech) for her support during this investigation. The authors also thank Dr. J. M. Andreu from the Centro de Investigaciones Biologicas, (CSIC, Madrid, Spain) and M. Gorbacheva (Institute of Biochemistry, Moscow, Russia) for CD determinations and calculations of the secondary

structures. This material is based upon work founded by Spanish Ministry of Education and Science Projects VEM2004-08559 and CTQ2005-08925-C02-02/PPQ; Comunidad de Madrid Project GR/AMB/0690/2004; CSIC Project 200580M121; the Swedish Research Council, EU Project NMP2-CT-2006-026456 and Ramón y Cajal Program. MZ thanks Gobierno Vasco for a fellowship.

REFERENCES

1. Arnold FH. Fancy footwork in the sequence space shuffle. *Nat Biotechnol* 2006;24:328-330.
2. Bloom JD, Meyer MM, Meinhold P, Otey CR, MacMillan D, Arnold FH. Evolving strategies for enzyme engineering. *Curr Opin Struc Biol* 2005;15:447-452.
3. Farinas ET, Bulter T, Arnold FH. Directed enzyme evolution. *Curr Opin Biotech* 2001;12:545-551.
4. Arnold FH. Combinatorial and computational challenges for biocatalyst design. 2001; *Nature* 409:253-257.
5. Tao H, Cornish VW. Milestones in directed enzyme evolution. *Curr Opin Chem Biol* 2002;6:858-864.
6. Cherry JR, Fidantsef AL. Directed evolution of industrial enzymes: an update. *Curr Opin Biotech* 2003;14:438-443.
7. Jaeger KE, Eggert T. Enantioselective biocatalysis optimized by directed evolution. *Curr Opin Biotech* 2004;15:305-313.
8. Umeno D, Arnold FH. Evolution of a pathway to novel long-chain carotenoids. *J Bacteriol* 2004;186:1531-1536.
9. Glieder A, Farinas ET, Arnold FH. Laboratory evolution of a soluble, self-sufficient, highly active alkane hydroxylase. *Nat Biotechnol* 2002;20:1135-1139.

10. Lutz S, Patrick WM. Novel methods for directed evolution of enzymes: quality, not quantity. *Curr Opin Biotech* 2004;15:291-297.
11. Alcalde M, Ferrer M, Plou FJ, Ballesteros A. Environmental biocatalysis: from remediation with enzymes to novel green processes. *Trends Biotechnol* 2006;24:281-287.
12. Wahler D, Reymond JL. High-throughput screening for biocatalysts. *Curr Opin Biotech* 2001;12:535-544.
13. 2003. Directed Enzyme Evolution: screening and selection methods. F.H. Arnold and G. Georgiou (eds). Humana Press, Totowa, New Jersey.
14. Wong TS, Zhurina D, Schwaneberg U. The diversity challenge in directed protein evolution. *Comb Chem High T Scr* 2006;9:271-288.
15. Chica RA, Doucet N, Pelletier JN. Semi-rational approaches to engineering enzyme activity: combining the benefits of directed evolution and rational design. *Curr Opin Biotech* 2005;16:378-384.
16. Qian Z, Lutz S. Improving the catalytic activity of *Candida antarctica* lipase B by circular permutation. *J Am Chem Soc* 2005;127:13466-13467.
17. Affholter J, Arnold FH. Engineering a revolution. *Chemtech* 1999;29:34-39.
18. Arnold FH, Georgiou G. Directed Evolution: library creation. Humana Press, Totowa, New Jersey; 2003. 224p.
19. Wong TS, Roccatano D, Zacharias M, Schwaneberg U. A statistical analysis of random mutagenesis methods used for directed protein evolution. *J Mol Biol* 2006;355:858-871.
20. Wong TS, Tee KL, Hauer B, Schwaneberg U. Sequence saturation mutagenesis with tunable mutation frequencies. *Anal Biochem* 2005;341:187-189.

21. Eggert T, Jaeger KE, Reetz MT. Directed Evolution by random mutagenesis: a critical evaluation. In: Svendsen A, editor. Enzyme functionality. Design, engineering and screening. New York: Marcel Dekker Inc.; 2004. p. 375-387.
22. Abecassis V, Pompon D, Truan G. High efficiency family shuffling based on multi-step PCR and in vivo DNA recombination in yeast: statistical and functional analysis of a combinatorial library between human cytochrome P450 1A1 and 1A2 Nucleic Acids Res 2000;28:1-10.
23. Bulter T, Alcalde M. Preparing libraries in *Saccharomyces cerevisiae*. In: Arnold FH, Gergiou G, editors. Directed Enzyme Evolution. Totowa, New Jersey: Humana Press; 2003. p. 17-22.
24. Alcalde M, Zumárraga M, Polaina J, Ballesteros A, Plou FJ. Combinatorial saturation mutagenesis by in vivo overlap extension for the engineering of fungal laccases. Comb. Chem. High. T. Scr 2006;9:719-727
25. Okkels JS. In vivo gene shuffling in yeast: a fast and easy method for directed evolution of enzymes. In: Svendsen A, editor. Enzyme functionality: design, engineering, and screening. New York: Marcel Dekker, Inc; 2004. p 413-424.
26. Bulter T, Alcalde M, Sieber V, Meinhold P, Schlachtbauer C, Arnold FH. Functional expression of a fungal laccase in *Saccharomyces cerevisiae* by directed evolution. Appl Environ Microb 2003;69:987-995.
27. Riva S. Laccases: blue enzymes for green chemistry. Trends Biotechnol 2006;24:219-226
28. Alcalde M. Laccase: biological functions, molecular structure and industrial applications. In: Polaina J, MacCabe AP, editors. Industrial Enzymes: structure, function and applications. Heidelberg: Springer; 2007. p 459-474.

29. Alcalde M, Bulter T, Zumárraga M, García-Arellano H, Mencía M, Plou FJ, Ballesteros A. Screening mutant libraries of fungal laccases in the presence of organic solvents. *J Biomol Screen* 2005;10:624-631.
30. Fernandez Larrea J, Stahl U. Isolation and characterization of a laccase gene from *Podospora anserina*. *Mol Gen Genet* 1996;252:539-551.
31. Germann UA, Muller G, Hunziker PE, Lerch K. Characterization of 2 allelic forms of *Neurospora crassa* laccase amino-terminal and carboxyl-terminal processing of a precursor. *J Biol Chem* 1988;263:885-896.
32. Kiiskinen LL, Saloheimo M. Molecular cloning and expression in *Saccharomyces cerevisiae* of a laccase gene from the ascomycete *Melanocarpus albomyces*. *Appl Environ Microb* 2004;70:137-144.
33. Larsson T, Lindgren A, Ruzgas T. Spectroelectrochemical study of cellobiose dehydrogenase and diaphorase in a thiol-modified gold capillary in the absence of mediators. *Bioelectrochemistry* 2001;53:243-249.
34. Christenson A, Shleev S, Mano N, Heller A, Gorton L. Redox potentials of the blue copper sites of bilirubin oxidases. *BBA-Bioenergetics* 2006;1757:1634-1641.
35. Shleev S, Morozova O, Nikitina O, Gorshina E, Rusinova T, Serezhenkov V, Burbaev D, Gazaryan I, Yaropolov A. Comparison of physico-chemical characteristics of four laccases from different basidiomycetes. *Biochimie* 2004; 86:693–703.
36. Shleev S, Shumakovich G, Nikitina O, Morozova O, Pavlishko H, Gayda G, Gonchar M. Purification and characterization of alcohol oxidase from a genetically constructed over-producing strain of the methylotrophic yeast *Hansenula polymorpha*. *Biochemistry (Moscow)* 2006; 71:245-250.

37. Cammack R, Fernandez VM, Hatchikian EC. Nickel-iron hydrogenase. Method Enzymol 1994;243:43-68
38. Drummond DA, Iverson BL, Georgiou G, Arnold FH. Why high-error-rate random mutagenesis libraries are enriched in functional and improved proteins. J Mol Biol 2005;350:806-816.
39. Daugherty PS, Chen G, Iverson BL, Georgiou G. Quantitative analysis of the effect of the mutation frequency on the affinity maturation of single chain Fv antibodies. P Natl Acad Sci USA 2000;97:2029-2034.
40. Cline J, Hogrefe H. Randomize gene sequences with new PCR mutagenesis kit. Stratagene Strategies 2000;13:157-162.
41. LinGoerke JL, Robbins DJ, Burczak JD. PCR-based random mutagenesis using manganese and reduced dNTP concentration. Biotechniques 1997; 23:409-412.
42. Qian Z, Lutz S. Improving the catalytic activity of *Candida antarctica* lipase B by circular permutation. J Am Chem Soc 2005;127:13466-13467.
43. Shuster JR. Gene expression in yeast: protein secretion. Curr Opin Biotech 1991;2:685-690.
44. Xu F, Palmer AE, Yaver DS, Berka RM, Gambetta GA, Brown SH, Solomon EI. Targeted mutations in a *Trametes villosa* laccase. J Biol Chem 1999; 274:12372-12375
45. Palmer AE, Szilagyi RK, Cherry JR, Jones A, Xu F, Solomon EI. Spectroscopy characterization of the Leu513His variant of fungal laccase: effect of increased axial ligand interaction on the geometric and electronic structure of the type 1 Cu site. Inorg Chem 2003;42:4006-4017

46. Xu F, Berka RM, Wahleithner JA, Nelson BA, Shuster JR, Brown SH, Palmer AE, Solomon EI. Site-directed mutations in fungal laccase: effect on redox potential, activity and pH profile. *Biochem J* 1998;334:63-70
47. Li H, Webb SP, Ivanic J, Jensen JH. Determinants of the relative reduction potentials of type-1 copper sites in proteins. *J Am Chem Soc* 2004;126:8019
48. Hakulinen N, Kiiskinen LL, Kruus K, Saloheimo M, Paananen A, Koivula A, Rouvinen J. Crystal structure of a laccase from *Melanocarpus albomyces* with an intact trinuclear copper site. *Nat Struct Biol* 2002;9:601-605.
49. Narra HP, Ochman H. Of what use is sex to bacteria?. *Curr Biol* 2006;16:705-710.
50. Cherry JR, Lamsa MH, Schneider P, Vind J, Svendsen A, Jones A, Pedersen AH. Directed Evolution of a fungal peroxidase. *Nat Biotechnol* 1999;17:379-384.
51. Xu F. Applications of oxidoreductases: recent progress. *Ind. Biotechnol.* 2005;1:38-50.

Figure legends

Fig.1. Landscapes for libraries made under different error prone-PCR conditions. Activity of clones are plotted in descending order. Solid horizontal line shows the activity of the parent type in the assay. Dashed lines indicate the coefficient of variation of the assay (below 10 %).

Fig.2. Landscapes of generation created by *in vivo* assembly of Mutazyme/Taq libraries. Activity of clones are plotted in descending order. Dashed lines indicate the activity of parent type. a) Screening in the presence of 50 % acetonitrile; b) screening in the presence of 50 % of ethanol; c) activities of clones in acetonitrile and ethanol. Framed clones were submitted to re-screening procedures (see Method section).

Fig.3. Relative activities of parent type and 2E9 variant at different concentrations of ethanol and acetonitrile. Each point represents the average of three independent experiments.

Fig. 4. Spectral investigation of parent type and 2E9 variant. a) EPR spectra b) CD spectra. Dotted line, parent type; solid line, 2E9 variant.

Fig. 5. pH activities profiles of parent type laccase and 2E9 variant. Activities were measured in B&R buffer at different pHs with ABTS as substrate. Laccase activity is normalized to the optimum activity value. Each point represents the average of three independent experiments.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5